



Oct 11, 2022

Tissue Fixation Preparation

Stephen Fisher¹, Marielena Grijalva¹, Rong Guo¹, sarahjoh¹, Hieu Nguyen¹, John Renz², Jean G Rosario¹, Steven Rudich², Brian Gregory¹, Junhyong Kim¹, Kate O'Neill¹

¹University of Pennsylvania; ²Gift of Life Donor Program

1 Works for me

Share

dx.doi.org/10.17504/protocols.io.6qpvr61ozvmk/v1

Stephen Fisher
University of Pennsylvania

ABSTRACT

This protocol describes fixation of tissue in preparation for pathology review to assess tissue quality.

Prior to performing molecular analyses, it is critical to have experienced pathologist confirm the tissue of interest is captured, the region of interest is represented and there are no abnormalities. Fixed tissue is preferred for pathologist review as it prevents autolysis and degradation of the tissue, cells and cellular components so they can be observed both anatomically and microscopically in their most natural state following sectioning. Various reagents can be used for fixation. Formaldehyde is a water-based solution and is the most commonly used chemical fixative. Paraformaldehyde has been found to be a preferred fixative for the ovary as it maintains oocyte and follicular morphology which is often disrupted with other fixatives.

DOI

dx.doi.org/10.17504/protocols.io.6qpvr61ozvmk/v1

PROTOCOL CITATION

Stephen Fisher, Marielena Grijalva, Rong Guo, sarahjoh, Hieu Nguyen, John Renz, Jean G Rosario, Steven Rudich, Brian Gregory, Junhyong Kim, Kate O'Neill 2022. Tissue Fixation Preparation. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.6qpvr61ozvmk/v1>



FUNDERS ACKNOWLEDGEMENT

NIH
Grant ID: U54HD104392

LICENSE

_____ This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 13, 2022

LAST MODIFIED

Oct 11, 2022

PROTOCOL INTEGER ID

68610

MATERIALS TEXT

- Freshly dissected tissue block
- Ice
- 70% EtOH (made fresh; kept at 4°C)
- Ice bucket

[1.5 mL LoBind tubes](#)

- **Eppendorf Catalog #022431021**

- 4% sucrose in 1X DPBS; kept at 4°C

[DPBS \(10X\), no calcium, no magnesium](#) **Thermo Fisher**

Scientific Catalog #14200075

- 15mL conical tubes with 10% buffered formalin
- 15mL conical tubes with 4% paraformaldehyde (PFA) (fresh for 1 week, keep on ice)

[Eppendorf Conical Tube 15mL](#) **Merck Millipore**

Sigma Catalog #EP0030122151

4% PFA Tissue Fixation of Ovaries

4d

- 1 Place each pre-weighed piece of tissue in either a 1.5mL Eppendorf tube or 1.5mL cryo tube.
- 2 Keep PFA tubes on ice. Transfer tissue specimen into a tube and refrigerate at **4 °C** for ^{1d}

🕒 24:00:00 .

- 3 After 🕒 24:00:00 , transfer tissue into 4% sucrose/PBS solution and incubate for 🕒 24:00:00^{2d} at 🌡 4 °C .
- 4 After 🕒 24:00:00 , transfer tissue specimen to 70% EtOH (freshly made) and store at 🌡 4 °C^{1d} until ready for paraffin embedding.
- 5 Proceed with paraffin embedding of samples within 3 weeks.
- 6 Paraffin embedded tissue can now be used for H&E Pathology Review.

Formalin Tissue Fixation of Fallopian Tube, Uterine Body, and Cervix

2d

- 7 Place each pre-weighed piece of tissue in either a 1.5mL Eppendorf tube or 1.5mL cryo tube.
- 8 Place tissue specimen in a prepped and labeled tube. Keep at room temperature for 🕒 24:00:00 .^{1d}
- 9 After 🕒 24:00:00 , transfer tissue specimen to 70% EtOH (freshly made) and store at 🌡 4 °C^{1d} until ready for paraffin embedding.
- 10 Proceed with paraffin embedding of samples within 3 weeks.
- 11 Paraffin embedded tissue can now be used for H&E Pathology Review.