




JUL 28, 2023

Western blotting

Forked from [Western blotting to detect ATP13A2 and ATP13A3](#)

Marine Houdou¹, rosanne.wouters¹, Peter Vangheluwe¹

¹KU Leuven



rosanne.wouters

ABSTRACT

Protocol to detect proteins via Western Blotting.

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.ewov1oemklr2/v1

External link:

<https://doi.org/10.3390/biom13020337>

Protocol Citation: Marine Houdou, rosanne.wouters, Peter Vangheluwe 2023. Western blotting.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.ewov1oemklr2/v1>

MANUSCRIPT CITATION:

Houdou M, Jacobs N, Coene J, Azfar M, Vanhoutte R, Haute CVd, Eggermont J, Daniëls V, Verhelst SHL, Vangheluwe P, Novel Green Fluorescent Polyamines to Analyze ATP13A2 and ATP13A3 Activity in the Mammalian Polyamine Transport System. *Biomolecules* 13(2). doi: [10.3390/biom13020337](https://doi.org/10.3390/biom13020337)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 01, 2023

Last Modified: Jul 28, 2023

PROTOCOL integer ID:
82745

Keywords: ASAPCRN

MATERIALS

■ Antibodies:

- Goat anti-mouse IgG (H+L) secondary antibody HRP conjugated: Thermo Scientific, 31430
- Goat anti-rabbit IgG (H+L) secondary antibody HRP conjugated: Thermo Scientific, 31460
- Mouse monoclonal anti-GAPDH (lot #067M4785V, dilution 1:5,000): Sigma, G8795.
- Rabbit anti-ATP13A2 antibodies (lot #0000102992, dilution 1:1,000): Sigma, A3361.
- Rabbit anti-ATP13A3 antibody (lot # 000035781, dilution 1:2,000): Atlas Antibodies, HPA029471.

■ 0.25% Trypsin-EDTA: Gibco, 25200056

■ Dulbecco's Phosphate Buffered saline modified without calcium chloride and magnesium chloride (DPBS): Gibco, D8537

■ Micro-BCA Protein Assay Kit: Pierce BCA Protein Assay Kit, Thermo Scientific, 23225

■ NuPAGE LDS sample buffer: Invitrogen, NP0007

■ Ponceau staining: Sigma, P7170

■ Pre-cast 4-12% Bis-Tris gels: Invitrogen, NP0321BOX

■ PVDF membranes: Thermo Scientific, 88518


■ RIPA Lysis and Extraction Buffer: Invitrogen: 89900

■ SIGMAFAST Protease Inhibitor Cocktail Tablets, EDTA-Free: Sigma: S8830




■ SuperSignal West Pico PLUS chemiluminescent Substrate: Thermo Scientific, 34095

Harvesting cells

- 1 collect the cells by scrapping them with a scrapper in Dulbecco's Phosphate Buffered saline modified without calcium chloride and magnesium chloride (DPBS) or, using 0.25% Trypsin-EDTA for which stop enzymatic reaction by adding culture medium.

- 2 Centrifuge cell suspensions at 450 g, 4°C for  00:05:00 . 5m
- 3 Resuspend cell pellets with DPBS and centrifuge following the same indications as in 2. Repeat once.
- 4 Discard supernatants and keep cell pellets on ice.

Cell lysis and protein concentration determination

- 5 Resuspend cell pellets in RIPA buffer (RIPA Lysis and Extraction Buffer supplemented with protease cocktail inhibitors).
- 6 Vortex  00:00:30 and keep on ice for  00:30:00 . 30m 30s
- 7 Centrifuge at 20,000g, 4°C for  00:30:00 . 30m
- 8 Keep supernatants on ice to proceed with protein concentration determination using the micro-BCA Protein Assay Kit.

SDS-PAGE



7h 10m

9 Loading

9.1 Mix 20 µg of protein with NuPAGE LDS sample buffer and 5% β-mercaptoethanol final.


9.2 Load protein on pre-cast 4-12% Bis-Tris gels. Include at least one lane with a protein ladder.

10 Running

10.1 Run for  00:10:00 at 100V and  01:30:00 at 110-130V.

1h 40m



11 Transfer

11.1 Transfer onto PVDF membranes using a liquid transfer and following settings: 100V,  01:15:00, 4°C.

1h 15m


11.2 Scan membrane if necessary.

12 Blocking


12.1  01:00:00 Block membranes with blocking buffer (5% milk powder in 1X TBS and 0.1% Tween20 (REF)) for  01:00:00 at room temperature, 19 rpm.

2h

13 Primary antibodies


13.1 Incubate membrane with primary antibodies in solution (1% bovine serum albumin in 1X TBS-Tween20 (TBS-T) buffer),  Overnight at 4°C, 19 rpm.

1h

13.2 Wash membrane three times for  00:05:00 in TBS-T, 19 rpm.

5m

14 Secondary antibodies


14.1 Incubate membrane with peroxidase-conjugated secondary antibodies in solution (1% milk powder in 1X TBS-T) for  01:00:00 at room temperature and, 19 rpm.

1h

14.2 Wash membrane five times for  00:05:00 in TBS-T, 19 rpm.

5m

15 Detection

- 
- 16** Use a chemiluminescence reagent to detect signal and acquire with a Biorad Camera (Vilber Lourmat) and its software (ImageLab).