Apr 01, 2026 Version 3

DNA extraction (Salting out) V.3

DOI
https://dx.doi.org/10.17504/protocols.io.vv9e696

Karen Guimaraes[1]

[1]Laboratório de Genética & Biodiversidade, Instituto de Ciências da Educação, Universidade Federal do Oeste do Pará, Santarém, Pará, Brazil

Karen Guimaraes
Laboratório de Genética & Biodiversidade, Instituto de Ciênc...

OPEN ACCESS

DOI: https://dx.doi.org/10.17504/protocols.io.vv9e696

Protocol Citation: Karen Guimaraes 2026. DNA extraction (Salting out). **protocols.io** https://dx.doi.org/10.17504/protocols.io.vv9e696

Manuscript citation:

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

Created: November 26, 2018

Last Modified: April 01, 2026

Protocol Integer ID: 18081

Keywords: dna extraction, extraction, dna

Materials

NaCl; Tris-HCl; EDTA; Proteinase K; Ethanol; Sterile H_2O; RNAse; Isopropanol; Microcentrifuge capable of at least 13.000rpm; Incubator

Troubleshooting

Before start

1. In microtubes containing tissue fragments, add 440µL of lysis buffer (10mM Tris-HCl, 2mM EDTA, 400mM NaCl, 2% SDS) and 10 µL of proteinase K (10mg/mL);
2. Incubate in a water bath at 55°C for approximately 1:30h or overnight;
3. Add 300µL of NaCl (5M) and centrifuge for 10min at 10.000rpm.
4. Transfer supernatant containing the DNA to microcentrifuge tube (1,5mL);
5. Add 500µl of 100% isopropanol.
6. Centrifuge for 10min at 10.000rpm;
7. Discard the supernatant and reuse microcentrifuge tube;
8. Add 700µL of 70% ethanol;
9. Centrifuge for 3min at 13.000rpm;
10. Discard the supernatant and dry microcentrifuge tube;
11. Add sterile H_2O and 5µL of RNAse (10mg/mL);
12. Incubate at 37°C for 30 min and stored at -20ºC.