Drawing blood from a vein with a syringe

№	List of practical actions	Performance format (perform/say)[1]
	Preparation stage of the procedure	
1.	Review the prescription sheet	say/perform
2.	Greet the patient	say
3.	Introduce myself. Perform patient identification	say
4.	Inform the patient about the upcoming procedure, obtain verbal consent	say
5.	Check with the patient about the last meal	say
6.	Ask the patient to assume the required position	say
7.	Fully prepare the necessary equipment for the procedure	say/perform
8.	Check the integrity, appearance, and expiration dates of ampoules/vials with medication and medical devices before donning non-sterile gloves	perform
9.	Register the prescription in the laboratory test registration log. Mark the date, time, and registration number on the referral form. Label the tube	say/perform
10.	Remove the cap from the tube, place the tube in a rack	say/perform
11.	Perform hand hygiene according to regulatory acts. Don non-sterile medical gloves	say/perform
	Main stage of the procedure	
12.	Open the package of cotton balls	perform
13.	Prepare the syringe, the blood collection needle, check needle patency	perform
14.	Place a cushion under the patient's arm	perform
15.	Apply a tourniquet 10–12 cm above the intended injection site. Ask the patient to clench and unclench their fist several times to fill the vein	perform
16.	Select, inspect, and palpate the injection site	perform
17.	Treat the injection site with two cotton balls soaked in antiseptic in one direction	perform
18.	Place the antiseptic-soaked cotton ball into my left hand	perform
19.	Hold the syringe in my dominant hand, position the second finger on the cannula, and wrap the remaining fingers around the barrel	perform
20.	Fix the vein by stretching the skin at the venipuncture site with the first finger of my left hand	perform
21.	Puncture the skin at an angle of no more than 15 degrees, position the needle parallel to the vein, puncture the vein by advancing the needle along the vein until blood appears in the needle cannula	perform
22.	Untie the tourniquet, ask the patient to unclench their fist	perform
23.	Draw the required amount of blood into the syringe	perform
24.	Apply an antiseptic-soaked cotton ball to the venipuncture site, withdraw the needle	perform
25.	Hold the cotton ball in place for 3–5 minutes	say
26.	Correctly transfer the blood from the syringe into the tube	perform
27.	Close the tube with the cap	perform
	Final Stage of the Procedure	
28.	Correctly disinfect the cotton balls	say/perform
29.	Correctly disinfect the syringe	say/perform
30.	Correctly disinfect the needle	say/perform
31.	Place the rack with tubes into the "For transport of biological material samples" container, attaching the referral form and the inventory list in a separate bag	say/perform
32.	Remove gloves with the working surface facing inward, place them into the "Disinfection of dressing materials and PPE" container. Perform hand hygiene according to regulatory acts	say/perform
33.	Place the packages into the container	say/perform

34.	Record the medication administration in the medical documentation	say

1- Depends on physical capability to perform the action and on the station equipment.

ГОМЕЛЬСКИЙ ГОСУДАРСТВЕННЫЙ МЕДИЦИНСКИЙ УНИВЕРСИТЕТ
ГомГМУ