***Intravenous injection***

| № | List of practical actions | Performance format (perform/say)[1] |
|---|---|---|
| | **Preparation stage of the procedure** | |
| 1. | Review the prescription sheet | say/perform |
| 2. | Greet the patient | say |
| 3. | Introduce myself. Perform patient identification | say |
| 4. | Inform the patient about the upcoming procedure, obtain verbal consent | say |
| 5. | Ask the patient to assume the required position | say |
| 6. | Fully prepare the necessary equipment for the procedure | say/perform |
| 7. | Check the integrity, appearance, and expiration dates of ampoules/vials with medication and medical devices before donning non-sterile gloves | perform |
| 8. | Perform hand hygiene according to regulatory acts. Don non-sterile medical gloves | say/perform |
| | **Main stage of the procedure** | |
| 9. | Open the package of cotton balls, open the needle (0.8x40 mm), and the syringe of the required volume | perform |
| 10. | Correctly treat the ampoule/vial | perform |
| 11. | Open the ampoule/vial with the medication | perform |
| 12. | Draw the medication into the syringe, holding the ampoule/vial and syringe correctly | perform |
| 13. | Draw the prescribed amount of solvent into the syringe with the medication | perform |
| 14. | Timely place the ampoule necks and ampoules/vials into appropriate containers | perform |
| 15. | Remove the needle by hand and place it into the needle container | perform |
| 16. | Correctly prepare the syringe (dose the medication) | perform |
| 17. | Place the syringe back into its package | perform |
| 18. | Place a cushion under the patient's arm | perform |
| 19. | Apply a tourniquet 10–12 cm above the intended injection site. Ask the patient to clench and unclench their fist several times to fill the vein | perform |
| 20. | Select, inspect, and palpate the injection site | perform |
| 21. | Treat the injection site with two cotton balls soaked in antiseptic in one direction | perform |
| 22. | Place the antiseptic-soaked cotton ball into my left hand | perform |
| 23. | Hold the syringe with medication in my dominant hand, position the second finger on the cannula and wrap the remaining fingers around the barrel | perform |
| 24. | Fix the vein by stretching the skin at the venipuncture site with the first finger of my left hand | perform |
| 25. | Puncture the skin at an angle of no more than 15 degrees, position the needle parallel to the vein, puncture the vein by advancing the needle along the vein until blood appears in the needle cannula | perform |
| 26. | Pull the plunger toward myself. Announce "Blood appears in the barrel" | say/perform |
| 27. | Untie the tourniquet, ask the patient to unclench their fist | perform |
| 28. | Pull the plunger toward myself again. Announce "Blood appears in the barrel" | say/perform |
| 29. | Slowly inject the medication without changing the position of the syringe in my hand | perform |
| 30. | Leave 0.5 mL of medication in the barrel to prevent air embolism | say/perform |

| | | |
|---|---|---|
| 31. | Apply an antiseptic-soaked cotton ball to the injection site, withdraw the needle | perform |
| 32. | Hold the cotton ball in place for 3–5 minutes | say |
| | **Final Stage of the Procedure** | |
| 33. | Correctly disinfect the cotton balls | say/perform |
| 34. | Correctly disinfect the syringe | say/perform |
| 35. | Correctly disinfect the needle | say/perform |
| 36. | Remove gloves with the working surface facing inward, place them into the "Disinfection of dressing materials and PPE" container. Perform hand hygiene according to regulatory acts | say/perform |
| 37. | Place the packages into the container | say/perform |
| 38. | Record the medication administration in the medical documentation | say |
| 39. | Observe the patient for 30 minutes, checking on their well-being | say |

*1- Depends on physical capability to perform the action and on the station equipment.*

ГомГМУ
ГОМЕЛЬСКИЙ ГОСУДАРСТВЕННЫЙ МЕДИЦИНСКИЙ УНИВЕРСИТЕТ