Subcutaneous injection

№	List of practical actions	Performance format (perform/say)[1]
	Preparation stage of the procedure	
1.	Review the prescription sheet	say/perform
2.	Greet the patient	say
3.	Introduce myself. Perform patient identification	say
4.	Inform the patient about the upcoming procedure, obtain verbal consent	say
5.	Ask the patient to assume the required position	say
6.	Fully prepare the necessary equipment for the procedure	say/perform
7.	Check the integrity, appearance, and expiration dates of ampoules/vials with medication and medical devices before donning non-sterile gloves	perform
8.	Perform hand hygiene according to regulatory acts. Don non-sterile medical gloves	say/perform
	Main stage of the procedure	
9.	Open the package of cotton balls, open the needle (0.8x40 mm), and the syringe of the required volume	perform
10.	Correctly treat the ampoule/vial	perform
11.	Open the ampoule/vial with the medication	perform
12.	Draw the medication into the syringe, holding the ampoule/vial and syringe correctly	perform
13.	Timely place the ampoule neck and ampoule/vial into appropriate containers	perform
14.	Remove the needle by hand and place it into the needle container	perform
15.	Correctly prepare the syringe (dose the medication)	perform
16.	Place the syringe back into its package	perform
17.	Select, inspect, and palpate the injection site	perform
18.	Treat the injection site with two cotton balls soaked in antiseptic in one direction	perform
19.	Place the antiseptic-soaked cotton ball into my left hand	perform
20.	Hold the syringe with medication in my dominant hand, position the second finger on the cannula and wrap the remaining fingers around the barrel	perform
21.	Gather a skin fold with the first, second, and third fingers of my left hand	perform
22.	Insert the needle with a quick motion into the base of the fold, at a 45° angle to a depth of ⅔ of the needle	perform
23.	Slowly inject the medication	perform
24.	Apply an antiseptic-soaked cotton ball to the injection site, withdraw the needle	perform
25.	Hold the cotton ball in place for 2–3 minutes	say
	Final Stage of the Procedure	
26.	Correctly disinfect the cotton balls	say/perform
27.	Correctly disinfect the syringe	say/perform
28.	Correctly disinfect the needle	say/perform
29.	Remove gloves with the working surface facing inward, place them into the "Disinfection of dressing materials and PPE" container. Perform hand hygiene according to regulatory acts	say/perform
30.	Place the packages into the container	say/perform
31.	Record the medication administration in the medical documentation	say
32.	Observe the patient for 30 minutes, checking on their well-being	say

1- Depends on physical capability to perform the action and on the station equipment.