2. Blood pressure measurement

№	List of practical actions	Format of presentation (To do / To say)
	The preparatory stage of the procedure	
1.	Seat the patient in a comfortable position with a straight back and legs uncrossed.	To say
2.	Position the patient's arm on the table so that the midpoint of the upper arm is at heart level.	To say
3.	Check the blood pressure monitor (mechanical/electronic) for proper function. Select a cuff of appropriate size (width - 40% of the arm circumference). Verifiy the integrity of all monitor components.	To say
4.	Perform hand hygiene (refer to algorithm No. 1).	To do
	The main stage of the procedure	
5.	Expose the patient's shoulder by removing clothing. Position the cuff 2-3 cm above the elbow bend. Secure it so that one finger fits underneath. Center the cuff over the brachial artery.	To do
6.	Palpate the arterial pulse in the cubital fossa and place the diaphragm of the stethoscope over the pulsation site	To do[1] / To say
7.	Close the air valve. Inflate the cuff to a pressure 20-30 mmHg above the estimated systolic BP.	To do
8.	Place the stethoscope head over the arterial pulse point (in the antecubital fossa).	To do
9.	Deflate the cuff slowly at a rate of 2-3 mmHg per second.	To do
10.	Note the appearance of the first Korotkov sound (systolic BP).	To do
11.	Note the complete disappearance of Korotkov sounds (diastolic BP).	To do
12.	If necessary, repeat the measurement: fully deflate the cuff, then proceed to steps 7-11	To do
	The final stage of the procedure	
13.	State the blood pressure measurement result	To say
14.	Removed the cuff and stethoscope	To do
1- when measuring BP on a manikin, palpate the arterial pulse in the antecubital fossa and state: "Palpating the brachial artery pulse in the antecubital fossa."		