


## IN MEMORIUM: DR. VINOD BHAKUNI – PROTEIN UNFOLDING EXPERT

**Sohail Akhtar and Ashish Arora**

*Molecular and Structural Biology Division, CSIR-Central Drug Research Institute, Chattar Manzil Palace, M. G. Marg Lucknow 226 001, UP, India*

Vinod Bhakuni, the well known biophysicist passed away on July 15, 2011 of a sudden heart attack in Lucknow. With his death, the Indian scientific community lost an accomplished colleague who enriched equilibrium protein unfolding, stability and function by his contribution for about two decades. He was the Head of the department and In-charge of Molecular and Structural Biology division of the CSIR-Central Drug Research Institute, Lucknow.




Upon joining CDRI as a scientist in 1987

Vinod Bhakuni was born on May 24, 1962 at Pithoragarh, Uttar Pradesh (now in Uttaranchal). He obtained M.Sc (Spl) in Organic Chemistry from Lucknow University and Ph.D in Chemistry from CSIR-CDRI with Dr. C. M. Gupta, as advisor. Dr. Bhakuni had spent about two years working on thermodynamics of protein folding

with Ernesto Freire at Johns Hopkins University, Baltimore, USA as a visiting postdoctoral fellow. His interest remained in the field and after returning back to India in 1992, he established his group at CSIR-CDRI working on the different aspects of protein folding and stability. His deep interest in the area originated from his acute sense of understanding of the significance of folding intermediates and their stability in protein unfolding pathway and its applications. The various projects he undertook are shown schematically in Fig. 1, and are briefly discussed below.

### Folding Pathway of Proteins

The extent of unfolding of denatured states of protein under different conditions has long been of interest because of the possible relevance of their conformation to the protein folding pathway. The characterization of intermediates and the factors involved in their stability may provide important insight into the interactions responsible for their formation as well as their role



**Figure 1: Schematics of Dr. Vinod Bhakuni's Research Initiatives**

in protein folding. Dr. Bhakuni studied several model proteins to understand the different aspects of protein unfolding, stability and function. The earliest investigation of his group was on tetrameric bovine liver catalase. His group reported the presence of unique oligomeric intermediate of catalase. The enzyme at low GdmCl concentration ( $>0.3\text{M}$ ) dissociate from native tetrameric form to a functionally active dimeric form (Prakash *et al.*, 2002).

The studies on Glucose oxidase, an acidic dimeric enzyme from *Aspergillus niger*, revealed another interesting observation. In the presence of urea or GdmCl, GOD unfolded through a monomeric intermediate. However, a violation of 2-fold rule was observed (Akhtar *et al.*, 2002). Ideally, the amount of different denaturants viz. GdmCl and urea required to unfold a multimeric protein is different. In all the reports till the study was published, amount of urea required to denature oligomeric proteins was found to be twice or more than that of GdmCl. However in case of GOD, the amounts of the two denaturants required were more or less same (Table 1).

To understand the role of domains in the stability and folding of protein, a detailed study on Serine Hydroxymethyltransferases were carried out (Bhatt *et al.*, 2004). The SHMT from mesophilic *Bacillus subtilis* and thermophilic *Bacillus stearothermophilus* are highly homologous (homology-89%, identity-77%) and composed of

two domains, a large N-terminus PLP binding domain and a C-terminal domain. As expected, the bstSHMT is more stable as compared to the bsSHMT. However, their folding patterns were different as bsSHMT showed a non-cooperative unfolding. Furthermore, the domain swapping studies (the carboxy terminal domain of bs- and bst-SHMT were swapped) shows that the folding and stability of SHMTs are governed by its C-terminal domain (Fig. 2). The chimera containing N-terminus of bsSHMT and C-terminus of bstSHMT showed the change in its non-cooperative unfolding to cooperative unfolding and vice versa.

He chose various other model proteins, of which some are listed in Table 2, to investigate the folding intermediates and pathways. One of the model proteins, Serine Hydroxymethyltransferase from *Mycobacterium tuberculosis* also showed a unique stoichiometry of PLP binding. All the known SHMT have one PLP per enzyme monomer where as in case of SHM1 it is one PLP per enzyme dimer. The difference might be due to two double mutations (VS\*TTV\*HKT) in the PLP binding pocket of SHMT as compared to all the other known SHMTs where this sequence is VTTTTHKT (Chaturvedi and Bhakuni, 2003).

### Factors Governing Protein Stability


At the time when Bhakuni began his scientific career, understanding the role of different factors

**Table 1**  
**Comparison of Urea and GdmCl  $C_m$  Values for Proteins**

Protein	oligomeric state	$C_m$ (urea, M)	$C_m$ (GdmCl, M)	$C_m$ (urea)/ $C_m$ (GdMCl)
vitronectin	tetramer	6	2.8	2.1
phosphofructokinase	tetramer	2.8	0.6	4.6
triosephosphate isomerase	dimer	$>6.0$	1.4	$>4.3$
ceratinase	dimer	5.2	2.0	2.6
glucose oxidase	dimer	3.6	3	1.2
bovine growth hormone	dimer	8.3	3.8	2.1
invertase	dimer	5.25	1.6	3.28
glutathoine transferase	dimer	5.5	1.5	3.66
RNase T1	monomer	4.6	2.9	1.59
RNase A	monomer	6.6	3.0	2.2
maltose binding protein	monomer	3.5	1.05	3.33
ubiquitin	monomer	3	3.9	0.77

**Table 2**  
**List of Few Proteins for which the Protein Unfolding was Studied (Kumar and Bhakuni, 2010; Mishra et al., 2010; Tripathi et al., 2009; Mishra and Bhakuni, 2003; Chaturvedi and Bhakuni, 2003)**

Protein	Pathway/Intermediates
<i>Mycobacterium tuberculosis</i> malate synthase G	Non cooperative / monomeric intermediate
<i>Entamoeba histolytica</i> Phosphoserine aminotransferase	Non cooperative / monomeric intermediate
<i>Plasmodium vivax</i> Glutathione S transferase	Non Cooperative / functionally active dimeric intermediate
<i>E. Coli</i> Methylenetetrahydrofolate reductase	Cooperative / no intermediate



**Figure 2:** (A) Schematic diagram showing the swapping of the C-terminal domains between bst- and bs-SHMT. The numbers denote the place of amino acid in the primary sequence of the protein. (B) Equilibrium GdmCl-induced unfolding of bsSHMT and its chimera. Plot of fractional change in the wavelength of maximum fluorescence emission of PyP for bsSHMT and its chimera. (C) Plot of fractional change in the wavelength of maximum fluorescence emission of PyP for bstSHMT and its chimera

on protein stability was rather a hot area of protein folding. pH is known to influence the stability of a protein by altering the net charge on the protein. Many proteins denature at extreme pH because of the presence of destabilizing repulsive interaction between like charges in the native protein. The exact behaviour of a given protein at low or high pH is a complex interplay between varieties of stabilizing and destabilizing forces, some of which are sensitive to the environment. Salts have been known to affect the physio-chemical properties of the proteins like their solubility, stability and pKa. Dr. Bhakuni chose to understand the interplay of these forces in the subunit assembly of multimeric enzymes.

In mid nineties, his group worked on hen egg white lysozyme and bovine liver catalase. Lysozyme shows denaturation at low acidic pH where as the catalase at basic pH (Babu and Bhakuni, 1997; Prajapati *et al.*, 1998). The partially denatured proteins showed the loss in secondary and tertiary structure; however the increase in ionic strength confers a gain in its secondary and tertiary structures. During the late nineties a much cited work from his laboratory was published showing the first proof of the hydrophobic dye ANS in the refolding of proteins (Ali *et al.*, 1999).

The dual roles of ionic interactions (mono- and di-valent cations) on the stability of the different enzymes were also studied in details. The increase in ionic strength (di-valent cations) leads to the dissociation of the dimeric GOD (12). The binding of different activator cations ( $K^+$ ,  $Mg^{2+}$ ) and the inhibitor cations ( $Cd^{2+}$ ,  $Zn^{2+}$ ,  $Mg^{2+}$ ) on the same TIM barrel domain of *Mycobacterium tuberculosis* isopropylmalate synthase and its

significance in the function of the enzyme was also reported by the group recently (Singh and Bhakuni, 2008). Table 3 summarizes some of the work related to ionic perturbation of the proteins carried out using different model proteins.


### Biomolecular Studies and Molecular Structures

More recently, his interests revolved around translating his expertise in protein unfolding and stability in understanding the functional aspect of proteins especially of Hyaluronate lyases from *Streptococcus pneumonia* and *Streptococcus agalactiae*. HL is composed of three domain,  $\beta$ I domain,  $\alpha$ -domain and C-terminal domain connected through linkers. The group studied these enzymes and showed the importance of carboxy terminal domain in increasing the surface potential of the catalytic interface in recruiting the polymeric hyaluronan. The role of linker was established in regulating the C-terminal domain in a conformation suitable for maximum activity (Fig. 3). The group also observed that there is no interaction of  $Ca^{2+}$  with its C-terminal domain; instead these ions interact with the substrate hyaluronan to change its conformation. The possible inhibitor (GTC, L-NAME) of the HLs were also reported (Akhtar and Bhakuni, 2003; Akhtar *et al.*, 2006; Akhtar and Bhakuni, 2007). For the last two or three years, Bhakuni was also working on different aspects of amyloidogenesis to understand its mechanism of formation and remedy. The trimeric HL from bacteriophage at acidic pH starts aggregating as observed by thioflavin and other binding assays. HylP2 was observed to form an unusual functionally active fibril through unfolding of its N-terminal domain (Mishra *et al.*, 2006; Mishra and Bhakuni, 2009).

**Table 3**  
**List of Few Proteins for which the Protein Stability was Studied (Ref. Tripathi *et al.*, 2009; Bhatt *et al.*, 2005; Kumar and Bhakuni, 2008; Tripathi *et al.*, 2008)**

Protein	Salt or pH	Enzymatic activity	Effect on the structure
FprA	Upto 1M NaCl or KCl >10mM $MgCl_2$ / $CaCl_2$	No change Loss	Stabilize the protein 2° structure break, cofactor release
Isocitrate Lyase	$Mg^{++}$ , $Mn^{++}$ $Zn^{++}$ , $Cd^{++}$	Increase Loss	No change Loss in 2° structure
PvGST	Mono and Divalent salts	Loss	Loss in 2° structure
Monothiol glutaredoxin	Acidic pH Alkaline pH		No change Loss in 2° and 3° structure





**Figure 3: The C-terminal domain modulates the enzymatic activity of  $\alpha$ -domain.** (A) Enzymatic activity profile of different proteins at 37°C with polymeric HA and its smaller fragment (HA6/HA8/HA10). SagHL (*S. agalactiae* HL), Sag Domain-I (SagHL without C-terminal domain), SagC (C-terminal domain of SagHL; SpnHL (*S. pneumoniae* HL), SpnN (N-terminal domain of SpnHL, SpnC (C-terminal domain of SpnHL). (B) Kinetic properties of recombinant enzyme and its different domain

### Dr. Vinod Bhakuni will be Remembered

In the course of his about 20 years of stay at CSIR-CDRI, Bhakuni had more than 50 publications in SCI journals and established a sound reputation of himself and the department (2003-2011). He was the recipient of many awards including the prestigious Shanti Swaroop Bhatnagar Prize in



Receiving the Bhatnagar Prize in 2006

2006 and was conferred the fellowship of all the three national academies.

Vinod Bhakuni's demise is a deep personal loss for many others and us. He is survived by his parents, wife, son and a daughter. He will be remembered by a large body of friends and well-wishers and in some sense he would always remain with us.

### Acknowledgement

We thank our laboratory members for many contributions and comments. We apologize to colleagues whose work was not cited. This is communication number 8140 from CSIR-CDRI, Lucknow.

### Abbreviations

GdmCl, guanidinium hydrochloride; GOD, glucose oxidase; SHMTs, serine hydroxymethyltransferases; bsSHMT, SHMT from *Bacillus subtilis*; bstSHMT, SHMT from *Bacillus stearothermophilus*; SHM1, serine hydroxymethyltransferase from *Mycobacterium tuberculosis*; HL, hyaluronate lyases; SpnHL, HL from *Streptococcus pneumoniae*; SagHL, HL from *Streptococcus agalactiae*; HylP2, HL from bacteriophage; PLP, pyridoxal- 5'-phosphate; PyP, pyridoxamine-P.

### References

- [1] Akhtar, M. S., Ahmad, A., and Bhakuni, V. (2002), Guanidinium chloride- and Urea-induced unfolding of the dimeric enzyme glucose oxidase. *Biochemistry* **41**, 3819-3827.
- [2] Akhtar, M. S., Ahmad, A., and Bhakuni, V. (2002), Divalent cation induced changes in structural properties of the dimeric enzyme glucose oxidase: dual effect of dimer stabilization and dissociation with loss of cooperative interactions in enzyme monomer. *Biochemistry* **41**, 7142-7149.
- [3] Akhtar, M. S., and Bhakuni, V. (2003), *Streptococcus pneumoniae* hyaluronate lyase contains two non-cooperative independent folding/unfolding structural domains: characterization of functional domain and inhibitors of enzyme. *J. Biol. Chem.* **278**, 25509-25516.
- [4] Akhtar, M. S., and Bhakuni, V. (2007), Role of ionic interactions and linker in the domain interaction and modulation of functional activity of hyaluronate lyases. *Biochem Biophys Res Commun.* **353**, 286-292.
- [5] Akhtar, M. S., Krishnan, M. Y., and Bhakuni, V. (2006), Insights into the mechanism of action of hyaluronate lyase: role of C-terminal domain and Ca<sup>2+</sup> in the functional regulation of enzyme. *J. Biol. Chem.* **281**, 28336-28344.
- [6] Ali, V., Prakash, K., Kulkarni, S., Ahmad, A., Madhusudan, K. P., and Bhakuni, V. (1999), 8-anilino-1-naphthalene sulfonic acid (ANS) induces folding of acid unfolded cytochrome c to molten globule state as a result of electrostatic interactions. *Biochemistry* **38**, 13635-13642.

- [7] Babu, K. R., and Bhakuni, V. (1997), Ionic-strength-dependent transition of hen egg-white lysozyme at low pH to a compact state and its aggregation on thermal denaturation. *Eur J. Biochem.* **245**, 781-789.
- [8] Bhatt, A. N., Khan, M. Y. and Bhakuni, V. (2004), The C-terminal domain of dimeric serine hydroxymethyltransferase plays a key role in stabilization of the quaternary structure and cooperative unfolding of protein: domain swapping studies with enzymes having high sequence identity. *Protein Sci.* **13**, 2184-2195.
- [9] Bhatt, A. N., Shukla, N., Aliverti, A., Zanetti, G., and Bhakuni, V. (2005), Modulation of cooperativity in Mycobacterium tuberculosis NADPH-ferredoxin reductase: cation-and pH-induced alterations in native conformation and destabilization of the NADP+-binding domain. *Protein Sci.* **14**, 980-992.
- [10] Chaturvedi, S., and Bhakuni, V. (2003), Unusual structural, functional, and stability properties of serine hydroxymethyltransferase from Mycobacterium tuberculosis. *J. Biol. Chem.*, **278**, 40793-40805.
- [11] Kumar, R., and Bhakuni, V. (2010), A functionally active dimer of mycobacterium tuberculosis malate synthase G. *Eur Biophys J.* **39**, 1557-1562.
- [12] Kumar, R., and Bhakuni, V. (2008), Mycobacterium tuberculosis isocitrate lyase (MtbIcl): role of divalent cations in modulation of functional and structural properties. *Proteins* **72**, 892-900.
- [13] Mishra, P., Akhtar, M.S., and Bhakuni, V. (2006), Unusual structural features of the bacteriophage-associated hyaluronate lyase (hylp2). *J. Biol Chem.* **281**, 7143-7150.
- [14] Mishra, P., and Bhakuni, V. (2009), Self-assembly of bacteriophage-associated hyaluronate lyase (HYLP2) into an enzymatically active fibrillar film. *J. Biol. Chem.* **284**, 5240-5249.
- [15] Mishra, V., Ali, V., Nozaki, T., and Bhakuni, V. (2010). Biophysical characterization of Entamoeba histolytica phosphoserine aminotransferase (EhPSAT): role of cofactor and domains in stability and subunit assembly. *Eur Biophys J.* **40**, 599-610.
- [16] Misra, S. K., and Bhakuni, V. (2003), Unique holoenzyme dimers of the tetrameric enzyme Escherichia coli methylenetetrahydrofolate reductase: characterization of structural features associated with modulation of the enzyme's function. *Biochemistry.* **42**, 3921-3928.
- [17] Prajapati, S., Bhakuni, V., Babu, K. R., and Jain, S. K. (1998), Alkaline unfolding and salt-induced folding of bovine liver catalase at high pH. *Eur. J. Biochem.* **255**, 178-184.
- [18] Prakash, K., Prajapati, S., Ahmad, A., Jain, S. K., and Bhakuni, V. (2002), Unique oligomeric intermediates of bovine liver catalase. *Protein Sci.*, **11**, 46-57.
- [19] Singh, K., and Bhakuni, V. (2008), Toxoplasma gondii ferredoxin-NADP+ reductase: Role of ionic interactions in stabilization of native conformation and structural cooperativity. *Proteins* **1**, 1879-1888.
- [20] Tripathi, T., Na, B. K., Sohn, W. M., Becker, K., and Bhakuni, V. (2009), Structural, functional and unfolding characteristics of glutathione S-transferase of Plasmodium vivax. *Arch Biochem Biophys.* **487**, 115-122.
- [21] Tripathi, T., Rahlfs, S., Becker, K., and Bhakuni, V. (2008), Structural and stability characteristics of a monothiol glutaredoxin: glutaredoxin-like protein 1 from Plasmodium falciparum. *Biochim Biophys Acta.* **1784**, 946-952.