

SPECTRALITY OF MORAN-TYPE MEASURES WITH STAGGERED CONTRACTION RATIOS

JUN JASON LUO, LIN MAO, AND JING-CHENG LIU

ABSTRACT. Consider a Moran-type iterated function system (IFS) $\{\phi_{k,d}\}_{d \in D_{2p_k}, k \geq 1}$, where each contraction map is defined as

$$\phi_{k,d}(x) = (-1)^d b_k^{-1}(x + d),$$

with integer sequences $\{b_k\}_{k=1}^\infty$ and $\{p_k\}_{k=1}^\infty$ satisfying $b_k \geq 2p_k \geq 2$, and digit sets $D_{2p_k} = \{0, 1, \dots, 2p_k - 1\}$ for all $k \geq 1$. We first prove that this IFS uniquely generates a Borel probability measure μ . Furthermore, under the divisibility constraints

$$p_2 \mid b_2, \quad 2 \mid b_2, \quad \text{and} \quad 2p_k \mid b_k \text{ for } k \geq 3,$$

with $\{b_k\}_{k=1}^\infty$ bounded, we prove that μ is a spectral measure, that is, $L^2(\mu)$ admits an orthogonal basis of exponentials. To fully characterize the spectral properties, we introduce a multi-stage decomposition strategy for spectrums. By imposing the additional hypothesis that all parameters p_k are even, we establish a complete characterization of μ 's spectrality. This result unifies and extends the frameworks proposed in [2, 9, 32], providing a generalized criterion for such measures.

1. INTRODUCTION

A Borel set $\Omega \subset \mathbb{R}^n$ with positive, finite Lebesgue measure is termed a spectral set if the space $L^2(\Omega)$ admits an orthogonal basis of exponential functions. This concept, central to functional analysis, gained prominence with Fuglede's 1974 conjecture [12], which posited that Ω is a spectral set if and only if it tiles $\mathbb{R}^n$ translationally, i.e., there exists $\mathcal{J} \subset \mathbb{R}^n$ such that $\{\Omega + t : t \in \mathcal{J}\}$ partitions $\mathbb{R}^n$. While disproven in dimensions $n \geq 3$ by Tao, Kolountzakis and Matolcsi [31, 27, 17, 18], the conjecture remains unresolved for $n = 1, 2$. Recent breakthroughs by Lev and Matolcsi [20] established its validity under convexity constraints across all dimensions, revitalizing interest in restricted cases.

The development of harmonic analysis and fractal geometry has propelled spectral measures, generalizations of spectral sets, to the forefront of mathematical inquiry. Formally, a compactly supported Borel probability measure μ on $\mathbb{R}^n$ is called a

2010 *Mathematics Subject Classification.* Primary 28A80; Secondary 42B10, 42C05.

Key words and phrases. Moran-type measure; Spectral measure; Fourier transform.

The research was supported by the NNSF of China (No. 12071125), the Natural Science Foundation of Chongqing (No. CSTB2023NSCQ-MSX0553), the Hunan Provincial NSF (No. 2024JJ3023) and the education Department Important Foundation of Hunan province in China (No. 23A0059).

spectral measure if there exists a countable subset $\Lambda \subset \mathbb{R}^n$ (termed its *spectrum*) such that the exponential system $\{e^{2\pi i \langle \lambda, \cdot \rangle} : \lambda \in \Lambda\}$ forms an orthonormal basis for $L^2(\mu)$. Identifying such measures, particularly among singular fractal measures, constitutes a pivotal challenge in modern analysis. Seminal work by Jorgensen and Pedersen [16] revealed that the $\frac{1}{n}$ -Cantor measure $\mu_{1/n}$ is spectral if and only if n is even. Subsequent advancements extended this to self-similar measures [19], where Dai, He, and Lau [8] characterized that a self-similar measure μ_ρ with a consecutive digit set $D_N = \{0, 1, \dots, N-1\}$ is a spectral measure if and only if the reciprocal of its contraction ratio $p = \rho^{-1}$ is an integer and $N \mid p$. Parallel developments resolved long-standing questions on Bernoulli convolutions [14, 7], while self-affine measures and infinite convolutions were systematized through Hadamard triples [3, 11, 21, 22, 23].

Moran-type measures, as natural extensions of self-similar and self-affine measures, exhibit intricate infinite product structures. Strichartz [29] pioneered their spectral analysis in 2000. Later, An and He [2] concerned the Moran-type measure $\mu_{\{b_k\}, \{D_{p_k}\}}$ generated by a sequence of integers $\{b_k\}_{k=1}^\infty$ with $b_k \geq 2$ and a sequence of consecutive digit sets $\{D_{p_k}\}_{k=1}^\infty$ with $p_k \geq 2$ by proving sufficiency of divisibility conditions ($p_k \mid b_k$) for spectrality. Recent work by An, Li and Zhang [5] and Deng and Li [9] established necessity, culminating in a complete characterization. Current research explores generalized Moran-type measures under varied constraints [1, 4, 10, 13, 24, 25, 26, 28, 30, 33], demonstrating their rich spectral properties.

Notably, all preceding achievements on spectrality of fractal measures fundamentally rely on the uniformity of linear components in their defining contraction maps. A paradigm shift emerged through Wu's recent work [32], which constructed the first self-similar spectral measure $\mu_{\pm\rho}$ with staggered contraction ratios $\pm\rho$ and consecutive digit set D_{2N} . The study established a sharp characterization: $\mu_{\pm\rho}$ is spectral if and only if the reciprocal of contraction parameter $p = \rho^{-1}$ is an integer and satisfies $2N \mid p$.

Building on this conceptual breakthrough, we systematically investigate the spectrality of a novel class of Moran-type measures on $\mathbb{R}$ featuring hierarchically staggered contraction ratios. Our approach diverges from classical methodologies. We first rigorously establish the existence and uniqueness of Moran-type measures through Moran-type iterated function systems, then study their spectral properties by revealing intrinsic arithmetic constraints in non-uniform scaling systems.

Definition 1.1. Let $\{b_k\}_{k=1}^\infty, \{n_k\}_{k=1}^\infty$ be two sequences of integer numbers with all $b_k \geq n_k \geq 2$ and let $\Phi_k = \{\phi_{k,i}\}_{i=0}^{n_k-1}$ be a family of contracting similitudes on $\mathbb{R}$ for each $k \geq 1$, where

$$\phi_{k,i}(x) = (-1)^i b_k^{-1}(x + i), \quad x \in \mathbb{R} \quad (i = 0, 1, \dots, n_k - 1). \quad (1.1)$$

We call the sequence $\{\Phi_k\}_{k=1}^\infty$ a *Moran-type iterated function system (IFS)* with staggered contraction ratios.

Theorem 1.2. *Let $\{\Phi_k\}_{k=1}^\infty$ be the Moran-type IFS as in Definition 1.1, then the following two statements hold:*

- (i) *There exists a unique sequence of nonempty compact sets $\{E_k\}_{k=1}^\infty$ in $\mathbb{R}$ such that*

$$E_k = \Phi_k(E_{k+1}) := \bigcup_{i=0}^{n_k-1} \phi_{k,i}(E_{k+1}), \quad k \geq 1 \quad (1.2)$$

and

$$\lim_{l \rightarrow \infty} |\phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \cdots \circ \phi_{l,j_l}(0) - \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \cdots \circ \phi_{l,j_l}(a_{l+1})| = 0 \quad (1.3)$$

uniformly for all sequences $\{j_l : 0 \leq j_l \leq n_l - 1\}_{l=k}^\infty$ and any $a_{l+1} \in E_{l+1}$.

- (ii) *There exists a unique sequence of probability measures $\{\mu_k\}_{k=1}^\infty$ satisfying*

$$\mu_k = \frac{1}{n_k} \mu_{k+1} \circ \Phi_k^{-1} := \sum_{i=0}^{n_k-1} \frac{1}{n_k} \mu_{k+1} \circ \phi_{k,i}^{-1}, \quad k \geq 1, \quad (1.4)$$

where each μ_k is supported on E_k .

The first measure $\mu := \mu_1$ is called a *Moran-type measure with staggered contraction ratios*, which will be our research object throughout the paper. The supporting set E_1 is usually called a *Moran-type set*. The geometric fashion and Hausdorff dimension of Moran-type sets were investigated in detail by Hua *et al.* [15].

Theorem 1.3. *Let $\{\Phi_k\}_{k=1}^\infty$ be the Moran-type IFS as in Definition 1.1 with assumption that the sequence $\{b_k\}_{k=1}^\infty$ is bounded and $2 \mid n_k$ for all $k \geq 1$. Let $\mu := \mu_1$ be as in (1.4). Then μ is a spectral measure if $2 \mid b_2$, $n_2 \mid 2b_2$ and $n_k \mid b_k$ for $k \geq 3$.*

The necessity of this condition generally fails to hold, as demonstrated by the counterexample in Theorem 4.3 (Section 4). However, under the additional hypothesis that 4 divides all n_k , we give a complete characterization of the spectrality of μ in the next theorem. This result unifies and extends the frameworks proposed in [2, 9, 32], providing a generalized criterion for such measures.

Theorem 1.4. *Let $\{\Phi_k\}_{k=1}^\infty$ be the Moran-type IFS as in Definition 1.1 with assumption that the sequence $\{b_k\}_{k=1}^\infty$ is bounded and $4 \mid n_k$ for all $k \geq 1$. Let $\mu := \mu_1$ be as in (1.4). Then μ is a spectral measure if and only if $n_2 \mid 2b_2$ and $n_k \mid b_k$ for $k \geq 3$.*

The primary challenges in establishing these theorems stem from two fundamental considerations. First, unlike classical self-similar measures, the Moran-type measure μ lacks a convolutional structure, which precludes direct analysis via the framework of infinite convolution measures. Notably, by leveraging the iterative relation (1.4),

we demonstrate that its spectrality can be equivalently characterized through an infinite convolution (Proposition 4.2). Second, in addressing the necessity condition of Theorem 1.4, we introduce a multi-stage decomposition strategy for spectrums (Remark 5.2). This approach enables the extraction of hierarchical constraints by recursively partitioning the spectrum of the spectral measure.

A natural extension lies in investigating the spectral consistency across the sequence of measures $\{\mu_k\}_{k=1}^\infty$. For some typical Moran-type measures, Liu *et al.* [25] recently showed that spectrality of μ_1 propagates inductively to all μ_k ($k \geq 1$). Nevertheless, the general case remains an open question, as the interplay between scaling parameters and spectral inheritance mechanisms resists straightforward generalization.

The rest of this paper is organized as follows. Section 2 synthesizes foundational results, including the Fourier transform of measures and some criteria for identifying spectral measures. Section 3 establishes Theorem 1.2, while Sections 4 and 5 develop the proofs of Theorems 1.3 and 1.4, respectively.

2. PRELIMINARIES

Let μ be a Borel probability measure on $\mathbb{R}$, the Fourier transform of μ is defined by

$$\widehat{\mu}(t) = \int e^{2\pi itx} d\mu(x), \quad t \in \mathbb{R}.$$

We denote by $\mathcal{Z}(\widehat{\mu}) := \{t \in \mathbb{R} : \widehat{\mu}(t) = 0\}$ the zero set of $\widehat{\mu}$. Let $\Lambda \subset \mathbb{R}$ be a countable set, then the set $E(\Lambda) = \{e^{2\pi i\lambda x} : \lambda \in \Lambda\}$ is an orthogonal set of $L^2(\mu)$ if and only if

$$(\Lambda - \Lambda) \setminus \{0\} \subset \mathcal{Z}(\widehat{\mu}).$$

If this is the case, then Λ is called a *bi-zero set* of μ . Moreover, if Λ is a bi-zero set of μ but $\Lambda \cup \{\alpha\}$ is not a bi-zero set for any $\alpha \in \mathbb{R} \setminus \Lambda$, then we call Λ a *maximal bi-zero set* of μ . The following is a basic criterion for determining Λ to be a bi-zero set or spectrum of μ .

Theorem 2.1 ([16]). *Let Λ be a countable subset of $\mathbb{R}$ and*

$$Q_{\Lambda, \mu}(t) := \sum_{\lambda \in \Lambda} |\widehat{\mu}(t + \lambda)|^2, \quad t \in \mathbb{R}.$$

Then

- (i) Λ is a bi-zero set of μ if and only if $Q_{\Lambda, \mu}(t) \leq 1$ for all $t \in \mathbb{R}$;
- (ii) Λ is a spectrum of μ if and only if $Q_{\Lambda, \mu}(t) = 1$ for all $t \in \mathbb{R}$.

Lemma 2.2 ([8]). *Let $\mu = \nu * \omega$ be a convolution of two measures, and let Λ be a bi-zero set of ν . Then Λ is also a bi-zero set of μ , but it can not be a spectrum of μ when ω is not a Dirac measure.*

Lemma 2.3. *For any $c \in \mathbb{R} \setminus \{0\}$, the measure μ is a spectral measure if and only if $\mu' := \mu(c \cdot)$ is a spectral measure.*

Proof. If μ is a spectral measure with a spectrum Λ , for any $t \in \mathbb{R}$,

$$\begin{aligned} Q_{c\Lambda, \mu'}(t) &= \sum_{\lambda \in \Lambda} |\widehat{\mu'}(t + c\lambda)|^2 \\ &= \sum_{\lambda \in \Lambda} \left| \int e^{2\pi i(t+c\lambda)x} d\mu(cx) \right|^2 \\ &= \sum_{\lambda \in \Lambda} \left| \int e^{2\pi i(\frac{t}{c} + \lambda)y} d\mu(y) \right|^2 \\ &= \sum_{\lambda \in \Lambda} |\widehat{\mu}(\frac{t}{c} + \lambda)|^2 \equiv 1. \end{aligned}$$

Then μ' is also a spectral measure. Likewise for the other hand. $\square$

For any $n \in \mathbb{Z} \setminus \{0\}$, we let $v_2(n)$ be the greatest integer $k \geq 0$ such that 2^k divides n in $\mathbb{Z}$, that is, $2^{-k}n \in 2\mathbb{Z} + 1$ and let $v_2(0) = \infty$. For any rational number $\frac{m}{n} \in \mathbb{Q}$ with $m, n \in \mathbb{Z}$ where $n \neq 0$, we define

$$v_2\left(\frac{m}{n}\right) = v_2(m) - v_2(n).$$

Recently, Deng and Li [10] gave a necessary and sufficient condition for the spectrality of the Moran-type Bernoulli covolution of the form

$$\mu = \delta_{b_1^{-1}\{0, d_1\}} * \delta_{(b_1 b_2)^{-1}\{0, d_2\}} * \cdots * \delta_{(b_1 \cdots b_k)^{-1}\{0, d_k\}} * \cdots,$$

where b_k, d_k are integers.

Theorem 2.4 ([10]). *For the above μ with $|b_k| > |d_k|$ for all $k \geq 2$. Assume further that the sequence $\{|d_k|\}_{k=1}^\infty$ is bounded. Then μ is a spectral measure if and only if $s_i \neq s_j$ for all $j > i \geq 1$, where*

$$s_k = v_2\left(\frac{b_1 b_2 \cdots b_k}{2d_k}\right), \quad k = 1, 2, \dots$$

Definition 2.5. *Let R be an integer satisfying $|R| > 1$, and $B, L \subset \mathbb{Z}$ be finite integer sets with the same cardinality $\#B = \#L \geq 2$. If the matrix*

$$\frac{1}{\sqrt{\#B}} \left(e^{-2\pi i R^{-1} b \ell} \right)_{b \in B, \ell \in L}$$

is unitary, we call (R, B, L) a Hadamard triple in $\mathbb{R}$.

The Hadamard triple is an important tool in studying the spectrality of measures as it is closely related to the spectrum of a discrete measure. Let A be a finite subset

of $\mathbb{R}$, we define a discrete measure on A by

$$\delta_A = \frac{1}{\#A} \sum_{a \in A} \delta_a$$

where δ_a is the Dirac measure at a . Moreover, we also define the mask polynomial of the set A by

$$m_A(x) = \frac{1}{\#A} \sum_{a \in A} e^{2\pi i a x}, \quad x \in \mathbb{R}.$$

The mask polynomial is obviously equal to the Fourier transform of the discrete measure δ_A . Trivially, (R, B, L) is a Hadamard triple if and only if L is a spectrum of the measure $\delta_{R^{-1}B}$ [11].

Lemma 2.6 ([22]). *If $\{(N_j, B_j, L_j)\}_{j=1}^n$ are finitely many Hadamard triples in $\mathbb{R}$, let*

$$N = N_1 N_2 \cdots N_n, \quad B = (N_2 \cdots N_n) B_1 + \cdots + N_n B_{n-1} + B_n$$

and

$$L = L_1 + N_1 L_2 + \cdots + (N_1 N_2 \cdots N_{n-1}) L_n,$$

then (N, B, L) is a Hadamard triple.

Let $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ be a sequence of Hadamard triples in $\mathbb{R}$, the infinite convolution generated by the Hadamard triples is defined by

$$\mu = \delta_{N_1^{-1}B_1} * \cdots * \delta_{(N_1 \cdots N_k)^{-1}B_k} * \cdots \quad (2.1)$$

which is usually represented as a convolution of two measures

$$\mu = \mu_k * \mu_{>k}$$

where

$$\mu_k = \delta_{N_1^{-1}B_1} * \cdots * \delta_{(N_1 \cdots N_k)^{-1}B_k}$$

and

$$\mu_{>k} = \delta_{(N_1 \cdots N_{k+1})^{-1}B_{k+1}} * \delta_{(N_1 \cdots N_{k+1}N_{k+2})^{-1}B_{k+2}} * \cdots.$$

Under the assumption that $B_k \subset \{0, 1, \dots, N_k - 1\}$ for all $k \geq 1$, An *et al.* [1] found a sufficient condition for spectrality of the infinite convolution μ .

Lemma 2.7 ([1]). *Suppose that $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ is a sequence of Hadamard triples with $B_k \subset \{0, 1, \dots, N_k - 1\}$ for all $k \geq 1$. Suppose that*

$$\liminf_{k \rightarrow \infty} \#B_k < \infty.$$

Then the infinite convolution μ as in (2.1) is a spectral measure and it always admits a spectrum in $\mathbb{Z}$.

The *integral periodic zero set* of a Borel probability measure μ on $\mathbb{R}$ is defined to be the set

$$Z(\mu) = \{\xi \in \mathbb{R} : \widehat{\mu}(\xi + k) = 0 \text{ for all } k \in \mathbb{Z}\}.$$

Let $\mathcal{P}(E)$ be the set of all Borel probability measures on $E \subset \mathbb{R}$. A family of measures $\Psi \subset \mathcal{P}(E)$ is called an *admissible family* if $Z(\mu) = \emptyset$ for every $\mu \in \text{cl}(\Psi)$, the closure of Ψ with respect to the weak topology on $\mathcal{P}(E)$. We say that Ψ is *tight* if for each $\varepsilon > 0$, there exists a compact subset $K \subset \mathbb{R}$ such that $\inf_{\mu \in \Psi} \mu(K) > 1 - \varepsilon$.

Lemma 2.8 ([1]). *Let $\mu \in \mathcal{P}([0, 1])$. Then $Z(\mu) \neq \emptyset$ if and only if $\mu = \frac{1}{2}(\delta_0 + \delta_1)$.*

Let $\mu = \mu_k * \mu_{>k}$ be as in (2.1). We define

$$\omega_{>k}(\cdot) = \mu_{>k} \left(\frac{\cdot}{N_1 \cdots N_k} \right).$$

That is,

$$\omega_{>k} = \delta_{N_{k+1}^{-1}B_{k+1}} * \delta_{(N_{k+1}N_{k+2})^{-1}B_{k+2}} * \cdots.$$

Theorem 2.9 ([22]). *Let $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ be a sequence of Hadamard triples in $\mathbb{R}$, suppose that infinite convolution measure μ defined by (2.1) exists. If there exists a subsequence $\{\omega_{>n_j}\}$ which is tight and admissible, then μ is a spectral measure with a spectrum in $\mathbb{Z}$.*

Theorem 2.10 ([23]). *Let $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ be a sequence of Hadamard triples in $\mathbb{R}$, suppose that infinite convolution measure μ defined by (2.1) exists. If there exists a subsequence $\{\omega_{>n_j}\}$ which converges weakly to ω , and $Z(\omega) = \emptyset$, then μ is a spectral measure with a spectrum in $\mathbb{Z}$.*

If the sequence of Hadamard triples $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ only has a finite number of different elements, then the following consequences are clear.

Lemma 2.11 ([22]). *Let $\{(N_j, B_j, L_j)\}_{j=1}^m$ be finitely many Hadamard triples, and let $\Sigma = \{1, 2, \dots, m\}^\mathbb{N}$. For $\sigma = \sigma_1\sigma_2\cdots \in \Sigma$, define*

$$\mu_\sigma = \delta_{N_{\sigma_1}^{-1}B_{\sigma_1}} * \cdots * \delta_{(N_{\sigma_1}\cdots N_{\sigma_k})^{-1}B_{\sigma_k}} * \cdots.$$

Let $\Psi = \{\mu_\sigma : \sigma \in \Sigma\}$. Then we have $\text{cl}(\Psi) = \Psi$.

Lemma 2.12 ([23]). *Suppose that $\{(N_k, B_k, L_k)\}_{k=1}^\infty$ is chosen from a finite set of Hadamard triples. If for each $k \geq 1$,*

$$\gcd \left(\bigcup_{j=k}^\infty (B_j - B_j) \right) = 1,$$

then $Z(\mu) = \emptyset$, where μ is as in (2.1).

The final useful lemma is due to Deng and Li [9].

Lemma 2.13 ([9]). *Let $p_{i,j}$ be positive numbers such that $\sum_{j=1}^n p_{i,j} = 1$ ($i = 1, 2, \dots, m$) and $x_{i,j} \geq 0$ such that $\sum_{i=1}^m \max_{1 \leq j \leq n} \{x_{i,j}\} \leq 1$. Then $\sum_{j=1}^n \sum_{i=1}^m p_{i,j} x_{i,j} = 1$ if and only if $\sum_{i=1}^m x_{i,1} = 1$ and $x_{i,1} = x_{i,2} = \dots = x_{i,n}$ for $1 \leq i \leq m$.*

3. EXISTENCE AND UNIQUENESS OF MORAN-TYPE MEASURES

In this section, we establish the theory of Moran-type IFS and show the existence and uniqueness of the Moran-type measure by proving Theorem 1.2.

Proof of Theorem 1.2.

(i) Since $\Phi_k = \{\phi_{k,i}\}_{i=1}^{n_k}$ is a family of contracting similitudes on $\mathbb{R}$ and $b_k \geq n_k \geq 2$ ($k \geq 1$). For any $l \geq k, m \geq 1$ and for all sequences $\{j_l : 0 \leq j_l \leq n_l - 1\}_{l=k}^\infty$, we have

$$\begin{aligned} & |\phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(0) - \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l+m,j_{l+m}}(0)| \\ &= |\phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(0 - \phi_{l+1,j_{l+1}} \circ \dots \circ \phi_{l+m,j_{l+m}}(0))| \\ &= b_k^{-1} b_{k+1}^{-1} \dots b_l^{-1} \cdot \left| \sum_{i=1}^m (-1)^{j_{l+1} + \dots + j_{l+i}} b_{l+1}^{-1} \dots b_{l+i}^{-1} \cdot j_{l+i} \right| \\ &= b_k^{-1} b_{k+1}^{-1} \dots b_l^{-1} \cdot \sum_{i=1}^m b_{l+1}^{-1} \dots b_{l+i}^{-1} \cdot j_{l+i} \\ &\leq 2^{k-l+2}. \end{aligned}$$

This implies

$$\lim_{l \rightarrow \infty} \sup_{\substack{0 \leq j_i \leq n_i - 1 \\ k \leq i \leq l+m, m \geq 1}} |\phi_{k,j_k} \circ \dots \circ \phi_{l,j_l}(0) - \phi_{k,j_k} \circ \dots \circ \phi_{l+m,j_{l+m}}(0)| = 0. \quad (3.1)$$

As

$$\begin{aligned} |\phi_{k,j_k} \circ \dots \circ \phi_{l,j_l}(0)| &\leq b_k^{-1} \cdot j_k + \dots + b_k^{-1} \dots b_l^{-1} \cdot j_l \\ &< 1 + \frac{1}{2} + \dots + \frac{1}{2^{l-k}} < 2. \end{aligned} \quad (3.2)$$

Thus the sequence $\{\phi_{k,j_k} \circ \dots \circ \phi_{l,j_l}(0)\}_{l=k}^\infty$ is uniformly bounded in $[-2, 2]$. Together with (3.1), we can get that the sequence $\{\phi_{k,j_k} \circ \dots \circ \phi_{l,j_l}(0)\}_{l=k}^\infty$ converges uniformly for all sequences $\{j_l : 0 \leq j_l \leq n_l - 1\}_{l=k}^\infty$.

For each $k \geq 1$, define

$$E_k = \left\{ \lim_{l \rightarrow \infty} \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(0) : 0 \leq j_i \leq n_i - 1, k \leq i \leq l \right\}. \quad (3.3)$$

Obviously, $\{E_k\}_{k=1}^\infty$ satisfies (1.2). We just need to show the compactness of E_k .

From (3.2), it follows that $E_k \subset [-2, 2]$ for each $k \geq 1$, hence $\{E_k\}_{k=1}^\infty$ is uniformly bounded. Let $\{x_n\}_{n=1}^\infty$ be a Cauchy sequence of E_k , then for every x_n , there is a sequence $\{j_{n,i}\}_{i=k}^\infty$ such that

$$x_n = \lim_{l \rightarrow \infty} \phi_{k,j_{n,k}} \circ \dots \circ \phi_{l,j_{n,l}}(0).$$

Since $0 \leq j_{n,k} \leq n_k - 1$, there exists η_k such that the index set $\{n : j_{n,k} = \eta_k\}$ is infinite. Repeating this process, there are $\eta_{k+1}, \eta_{k+2}, \dots, \eta_l, \dots$ such that the index set $\{n : j_{n,i} = \eta_i, i = k, \dots, l\}$ is infinite for all $l \geq k$. Let

$$x = \lim_{l \rightarrow \infty} \phi_{k,\eta_k} \circ \phi_{k+1,\eta_{k+1}} \circ \dots \circ \phi_{l,\eta_l}(0).$$

By (3.1), for any $\varepsilon > 0$, there is an integer $N > k$ such that for any $m, n \geq N$,

$$|\phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{m,j_m}(0) - \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{n,j_n}(0)| < \varepsilon. \quad (3.4)$$

That is, E_k is contained in the ε -neighborhood of the set $\{\phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{N,j_N}(0) : 0 \leq j_i \leq n_i - 1, i = k, \dots, N\}$. Since $\{x_n\}_{n=1}^\infty$ is a Cauchy sequence in E_k , there exists $N' \geq N$ such that $|x_n - x_m| < \varepsilon$ for all $m, n > N'$. Choose $n > N'$ such that $j_{n,i} = \eta_i$ for $i = k, k+1, \dots, N$. Then for any $l > N'$, by (3.4), we have

$$\begin{aligned} |x - x_l| &\leq |x - x_n| + |x_n - x_l| \\ &\leq |x - x_n| + \varepsilon \\ &\leq |x - \phi_{k,\eta_k} \circ \phi_{k+1,\eta_{k+1}} \circ \dots \circ \phi_{N,\eta_N}(0)| \\ &\quad + |\phi_{k,\eta_k} \circ \phi_{k+1,\eta_{k+1}} \circ \dots \circ \phi_{N,\eta_N}(0) - x_n| + \varepsilon \\ &\leq 3\varepsilon. \end{aligned}$$

Hence $\lim_{n \rightarrow \infty} x_n = x \in E_k$, and E_k is compact. Moreover, the above argument also shows that

$$\lim_{l \rightarrow \infty} \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(E_{l+1}) = \left\{ \lim_{l \rightarrow \infty} \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(0) \right\}$$

uniformly for all sequences $\{j_i : 0 \leq j_l \leq n_l - 1\}_{i=k}^\infty$. Thus (1.3) follows and the existence is proved.

Next we prove the uniqueness. Suppose $\{F_k\}_{k=1}^\infty$ is another sequence of compact sets on $\mathbb{R}$ satisfying (1.2). Let $x \in F_k$. (1.2) shows that there exist $x_i \in F_i$ and $0 \leq j_i \leq n_i - 1$ for $i = k+1, k+2, \dots$ such that

$$x = \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(x_{l+1}), \quad l = k, k+1, \dots$$

By (1.3), $x = \lim_{l \rightarrow \infty} \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \dots \circ \phi_{l,j_l}(0)$. Then $x \in E_k$. Hence $F_k \subset E_k$. Similarly we can show $E_k \subset F_k$. Therefore $F_k = E_k, k \geq 1$, and the uniqueness follows.

(ii) For each $k \geq 1$, we denote by the symbolic space starting from the k -th level

$$\Sigma_k := \{j_k \cdots j_i \cdots : 0 \leq j_i \leq n_i - 1, i \geq k\}$$

and denote by the corresponding cylinder sets of Σ_k

$$[j_k \cdots j_i] := \{j_k \cdots j_i \sigma : \sigma \in \Sigma_{i+1}\}, \quad \text{where } i \geq k.$$

Define a metric ρ_k on Σ_k by

$$\rho_k(j_k \cdots j_i \cdots, j'_k \cdots j'_i \cdots) = 2^{-\min\{i: j_i \neq j'_i\}},$$

then (Σ_k, ρ_k) is a compact metric space, and the cylinder sets $[j_k \cdots j_i]$ are the open sets of the space. If we put a mass on every cylinder set of Σ_k as follows:

$$\omega_k([j_k \cdots j_i]) = n_k^{-1} \cdots n_i^{-1}, \quad i \geq k. \quad (3.5)$$

The ω_k constitutes the unique Borel probability measure on Σ_k . Note that (3.3) defines a continuous map $\pi_k : \Sigma_k \rightarrow E_k$. Hence $\mu_k := \omega_k \circ \pi_k^{-1}$ is a Borel probability measure on E_k .

For any Borel subset $A \subset E_k$, by (3.5),

$$\begin{aligned} \mu_k(A) &= \omega_k \circ \pi_k^{-1}(A) \\ &= \omega_k \left(\{j_k j_{k+1} \cdots j_n \cdots : \lim_{n \rightarrow \infty} \phi_{k,j_k} \circ \phi_{k+1,j_{k+1}} \circ \cdots \circ \phi_{n,j_n}(0) \in A\} \right) \\ &= \sum_{j_k=0}^{n_k-1} \frac{1}{n_k} \omega_{k+1} \left(\{j_{k+1} \cdots j_n \cdots : \lim_{n \rightarrow \infty} \phi_{k+1,j_{k+1}} \circ \cdots \circ \phi_{n,j_n}(0) \in \phi_{k,j_k}^{-1}(A)\} \right) \\ &= \sum_{i=0}^{n_k-1} \frac{1}{n_k} \mu_{k+1} \circ \phi_{k,i}^{-1}(A). \end{aligned}$$

Hence μ_k satisfies (1.4) and the support of μ_k is E_k .

Finally, we show the uniqueness of the sequence of measures $\{\mu_k\}_{k=1}^\infty$. Let $\mathcal{P}([-2, 2])$ be the set of all Borel probability measures on $[-2, 2]$, equip it with a dual Lipschitz metric

$$L(\nu, \nu') = \sup_{\text{Lip}(g) \leq 1} \left| \int g d\nu - \int g d\nu' \right|$$

where $\text{Lip}(g) = \sup_{x \neq y} \frac{|g(x) - g(y)|}{|x - y|}$ is the Lipschitz constant of g . Then $(\mathcal{P}([-2, 2]), L)$ is a compact metric space [6]. Let $g : [-2, 2] \rightarrow \mathbb{R}$ be a function with $\text{Lip}(g) \leq 1$, and let

$$h = \sum_{i=0}^{n_k-1} \frac{1}{n_k} g \circ \phi_{k,i}. \quad (3.6)$$

Then

$$\begin{aligned} |h(x) - h(y)| &= \sum_{i=0}^{n_k-1} \frac{1}{n_k} |g \circ \phi_{k,i}(x) - g \circ \phi_{k,i}(y)| \\ &\leq \sum_{i=0}^{n_k-1} \frac{1}{n_k} |g \circ \phi_{k,i}(x) - g \circ \phi_{k,i}(y)| \\ &\leq \sum_{i=0}^{n_k-1} \frac{1}{n_k} \cdot b_k^{-1} \cdot |x - y| \\ &\leq b_k^{-1} \cdot |x - y|. \end{aligned}$$

That is, $\text{Lip}(h) \leq b_k^{-1} < 1$.

Suppose that $\{\mu'_k\}_{k=1}^\infty$ is another sequence of measures satisfying (1.4), by (3.6), we have

$$\begin{aligned} \left| \int g d\mu_k - \int g d\mu'_k \right| &= \left| \int h d\mu_{k+1} - \int h d\mu'_{k+1} \right| \\ &\leq \text{Lip}(h) L(\mu_{k+1}, \mu'_{k+1}) \\ &\leq b_k^{-1} L(\mu_{k+1}, \mu'_{k+1}). \end{aligned}$$

Hence $L(\mu_k, \mu'_k) \leq b_k^{-1} L(\mu_{k+1}, \mu'_{k+1}) \leq \dots \leq b_k^{-n} L(\mu_{k+n}, \mu'_{k+n})$ for any $n \geq 1$. By letting $n \rightarrow \infty$, it concludes that $L(\mu_k, \mu'_k) = 0$. Therefore, $\mu_k \equiv \mu'_k$, proving the uniqueness. $\square$

4. PROOF OF THEOREM 1.3

In this section, we study the spectrality of the Moran-type measure $\mu := \mu_1$ defined in (1.4). Throughout the section, we assume that n_k 's are all even numbers, i.e., $n_k = 2p_k$ where $p_k \in \mathbb{N}$ for $k \geq 1$. The sequence of measures $\{\mu_k\}_{k=1}^\infty$ can be reformulated as

$$\mu_k = \sum_{j=0}^{2p_k-1} \frac{1}{2p_k} \mu_{k+1} \circ \phi_{k,j}^{-1}, \quad k \geq 1. \quad (4.1)$$

Lemma 4.1. *Let $\mu := \mu_1$ be the Moran-type measure defined by the above, then its Fourier transform is*

$$\widehat{\mu}(t) = e^{-b_1^{-1}\pi t i} \prod_{k=1}^{\infty} f_k((b_k \cdots b_1)^{-1}t), \quad t \in \mathbb{R},$$

where $f_k(t) = \frac{1}{p_k} \sum_{j=0}^{p_k-1} \cos(4j + 1 - b_{k+1}^{-1})\pi t$.

Proof. From (4.1), the Fourier transform of μ has the expression

$$\begin{aligned} \widehat{\mu}_k(t) &= \sum_{j=0}^{2p_k-1} \frac{1}{2p_k} \widehat{\mu_{k+1} \circ \phi_{k,j}^{-1}}(t) \\ &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} \int e^{2\pi i t x} d\mu_{k+1} \circ \phi_{k,j}^{-1}(x) \\ &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} \int e^{2\pi i t (-1)^j b_k^{-1}(x+j)} d\mu_{k+1}(x) \\ &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} e^{2\pi i t (-1)^j b_k^{-1} j} \int e^{2\pi i t (-1)^j b_k^{-1} x} d\mu_{k+1}(x) \\ &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} e^{2\pi i t (-1)^j b_k^{-1} j} \widehat{\mu_{k+1}}((-1)^j b_k^{-1} t), \quad t \in \mathbb{R}. \end{aligned}$$

By considering the real and imaginary parts, denoted by ‘Re’ and ‘Im’, respectively, we have

$$\widehat{\mu}_{k+1}((-1)^j b_k^{-1} t) = \operatorname{Re} \widehat{\mu}_{k+1}(b_k^{-1} t) + i(-1)^j \operatorname{Im} \widehat{\mu}_{k+1}(b_k^{-1} t).$$

Hence

$$\begin{pmatrix} \operatorname{Re} \widehat{\mu}_k(t) \\ \operatorname{Im} \widehat{\mu}_k(t) \end{pmatrix} = M_k(b_k^{-1} t) \begin{pmatrix} \operatorname{Re} \widehat{\mu}_{k+1}(b_k^{-1} t) \\ \operatorname{Im} \widehat{\mu}_{k+1}(b_k^{-1} t) \end{pmatrix} \quad (4.2)$$

where the function matrix $M_k(t) = \begin{pmatrix} a_{k,1}(t) & a_{k,2}(t) \\ a_{k,3}(t) & a_{k,4}(t) \end{pmatrix}$ is given by

$$\begin{aligned} a_{k,1}(t) &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} \cos 2\pi t j, & a_{k,2}(t) &= -\frac{1}{2p_k} \sum_{j=0}^{2p_k-1} \sin 2\pi t j, \\ a_{k,3}(t) &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} (-1)^j \sin 2\pi t j, & a_{k,4}(t) &= \frac{1}{2p_k} \sum_{j=0}^{2p_k-1} (-1)^j \cos 2\pi t j. \end{aligned}$$

By iterating (4.2), one can get

$$\begin{aligned} & \begin{pmatrix} \operatorname{Re} \widehat{\mu}_1(t) \\ \operatorname{Im} \widehat{\mu}_1(t) \end{pmatrix} \\ &= M_1(b_1^{-1} t) M_2((b_2 b_1)^{-1} t) \cdots M_k((b_k \cdots b_2 b_1)^{-1} t) \begin{pmatrix} \operatorname{Re} \widehat{\mu}_{k+1}((b_k \cdots b_2 b_1)^{-1} t) \\ \operatorname{Im} \widehat{\mu}_{k+1}((b_k \cdots b_2 b_1)^{-1} t) \end{pmatrix} \\ &= \lim_{k \rightarrow \infty} M_1(b_1^{-1} t) M_2((b_2 b_1)^{-1} t) \cdots M_k((b_k \cdots b_2 b_1)^{-1} t) \begin{pmatrix} 1 \\ 0 \end{pmatrix}. \end{aligned}$$

A simple calculation yields that $\det M_k(t) \equiv 0$ for $t \in \mathbb{R}, k \geq 1$. Thus the function matrix $M_k(t)$ can be expressed as

$$M_k(t) = \alpha(t)^T \beta_k(t),$$

where

$$\alpha(t) = (\cos \pi t, -\sin \pi t)$$

and

$$\beta_k(t) = \left(\frac{1}{p_k} \sum_{j=0}^{p_k-1} \cos(4j+1)\pi t, -\frac{1}{p_k} \sum_{j=0}^{p_k-1} \sin(4j+1)\pi t \right).$$

On the other hand, one can check that

$$\begin{aligned} \beta_k(t) \alpha(b_{k+1}^{-1} t)^T &= \frac{1}{p_k} \sum_{j=0}^{p_k-1} (\cos(4j+1)\pi t \cos b_{k+1}^{-1} \pi t + \sin(4j+1)\pi t \sin b_{k+1}^{-1} \pi t) \\ &= \frac{1}{p_k} \sum_{j=0}^{p_k-1} \cos(4j+1 - b_{k+1}^{-1}) \pi t \\ &:= f_k(t). \end{aligned}$$

Since

$$\begin{aligned}
\sum_{k=1}^{\infty} (1 - f_k((b_k \cdots b_1)^{-1}t)) &= \sum_{k=1}^{\infty} \frac{2}{p_k} \sum_{j=0}^{p_k-1} \sin^2 \frac{(4j+1 - b_{k+1}^{-1})(b_k \cdots b_1)^{-1}t}{2} \\
&\leq \sum_{k=1}^{\infty} \frac{2}{p_k} \sum_{j=0}^{p_k-1} \left(\frac{(4j+1 - b_{k+1}^{-1})(b_k \cdots b_1)^{-1}t}{2} \right)^2 \\
&\leq \sum_{k=1}^{\infty} \frac{2}{p_k} \sum_{j=0}^{p_k-1} (2p_k(b_k \cdots b_1)^{-1}t)^2 \\
&\leq \sum_{k=1}^{\infty} 2^{3-2k} t^2 < \infty,
\end{aligned}$$

the infinite product $\prod_{k=1}^{\infty} f_k((b_k \cdots b_1)^{-1}t)$ converges uniformly on each compact subset of $\mathbb{R}$.

From the above discussion, it follows that

$$\begin{aligned}
&\begin{pmatrix} \operatorname{Re} \widehat{\mu}_1(t) \\ \operatorname{Im} \widehat{\mu}_1(t) \end{pmatrix} \\
&= \lim_{k \rightarrow \infty} M_1(b_1^{-1}t) M_2((b_2 b_1)^{-1}t) \cdots M_k((b_k \cdots b_2 b_1)^{-1}t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= \lim_{k \rightarrow \infty} \alpha(b_1^{-1}t)^T \beta_1(b_1^{-1}t) \alpha((b_2 b_1)^{-1}t)^T \beta_2((b_2 b_1)^{-1}t) \cdots \alpha((b_k \cdots b_1)^{-1}t)^T \beta_k((b_k \cdots b_1)^{-1}t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= \alpha(b_1^{-1}t)^T \lim_{k \rightarrow \infty} \left(\prod_{n=1}^{k-1} \beta_n((b_n \cdots b_1)^{-1}t) \alpha((b_{n+1} \cdots b_1)^{-1}t)^T \right) \beta_k((b_k \cdots b_1)^{-1}t) \begin{pmatrix} 1 \\ 0 \end{pmatrix} \\
&= \alpha(b_1^{-1}t)^T \prod_{n=1}^{\infty} f_n((b_n \cdots b_1)^{-1}t).
\end{aligned}$$

Hence $\widehat{\mu}(t) = \widehat{\mu}_1(t) = e^{-b_1^{-1}\pi t i} \prod_{n=1}^{\infty} f_n((b_n \cdots b_1)^{-1}t)$ as desired. $\square$

Let D_n denote the consecutive digit set $\{0, 1, \dots, n-1\}$ for $n \geq 1$, and let

$$\mathcal{D}_k = D_{p_k} \oplus (p_k - \frac{1 + b_{k+1}^{-1}}{2}) D_2. \quad (4.3)$$

Define an infinite convolution measure $\nu := \nu_{\{b_k\}, \{\mathcal{D}_k\}}$ by

$$\nu = \delta_{b_1^{-1}\mathcal{D}_1} * \delta_{(b_1 b_2)^{-1}\mathcal{D}_2} * \cdots * \delta_{(b_1 \cdots b_k)^{-1}\mathcal{D}_k} * \cdots. \quad (4.4)$$

We shall show that the measure ν has the same spectrality as the Moran-type measure $\mu := \mu_1$ defined in (4.1).

Proposition 4.2. *The Moran-type measure $\mu := \mu_1$ as in (4.1) is a spectral measure if and only if ν as in (4.4) is a spectral measure.*

Proof. By using the notation in Lemma 4.1, we notice that

$$f_k(t) = \frac{1}{p_k} \operatorname{Re} \left(e^{(1-b_{k+1}^{-1})\pi it} \sum_{j=0}^{p_k-1} e^{4j\pi it} \right), \quad t \in \mathbb{R}.$$

If $t \notin \frac{1}{2}\mathbb{Z}$, then

$$f_k(t) = \frac{\sin 2p_k \pi t}{p_k \sin \pi t} \cos(2p_k - 1 - b_{k+1}^{-1})\pi t.$$

Since $\delta_{\mathcal{D}_k} = \delta_{D_{p_k}} * \delta_{(p_k - \frac{1+b_{k+1}^{-1}}{2})D_2}$, the mask polynomial of $\mathcal{D}_k$ can be expressed as

$$\begin{aligned} m_{\mathcal{D}_k}(t) &= m_{D_{p_k}}(t) \cdot m_{(p_k - \frac{1+b_{k+1}^{-1}}{2})D_2}(t) \\ &= \left(\frac{1}{p_k} \sum_{j=0}^{p_k-1} e^{2\pi i t j} \right) \left(\frac{1}{2} + \frac{1}{2} e^{2\pi i (p_k - \frac{1+b_{k+1}^{-1}}{2})t} \right) \\ &= \frac{\sin p_k \pi t}{p_k \sin \pi t} \cos(p_k - \frac{1+b_{k+1}^{-1}}{2})\pi t \cdot e^{(2p_k - \frac{3+b_{k+1}^{-1}}{2})\pi it} \\ &= f_k(t/2) e^{(2p_k - \frac{3+b_{k+1}^{-1}}{2})\pi it}, \quad t \notin \mathbb{Z}. \end{aligned} \tag{4.5}$$

By the continuity of the functions f_k and $m_{\mathcal{D}_k}$, the above result holds for all $t \in \mathbb{R}$. Then for $t \in \mathbb{R}$,

$$\begin{aligned} \widehat{\nu}(t) &= \prod_{k=1}^{\infty} m_{\mathcal{D}_k}((b_k \cdots b_1)^{-1}t) \\ &= \prod_{k=1}^{\infty} f_k((b_k \cdots b_1)^{-1}t/2) e^{(2p_k - \frac{3+b_{k+1}^{-1}}{2})(b_k \cdots b_1)^{-1}\pi t i} \\ &= \prod_{k=1}^{\infty} f_k((b_k \cdots b_1)^{-1}t/2) \cdot e^{-\pi b_1^{-1} \frac{t}{2} i} \cdot e^{\pi b t i} \\ &= \widehat{\mu}(t/2) \cdot e^{\pi b t i} \end{aligned}$$

where $b := \frac{1}{2}b_1^{-1} + \sum_{k=1}^{\infty} (2p_k - \frac{3+b_{k+1}^{-1}}{2})(b_k \cdots b_1)^{-1} < \infty$ as $b_k \geq 2p_k \geq 2$. Consequently, for any $\Lambda \subset \mathbb{R}$ and $t \in \mathbb{R}$,

$$Q_{\Lambda, \nu}(t) = \sum_{\lambda \in \Lambda} |\widehat{\nu}(\lambda + t)|^2 = \sum_{\lambda \in \frac{1}{2}\Lambda} |\widehat{\mu}(\lambda + t/2)|^2 = Q_{\frac{1}{2}\Lambda, \mu}(t/2).$$

Therefore, Λ is a spectrum of ν if and only if $\frac{1}{2}\Lambda$ is a spectrum of μ by Theorem 2.1. $\square$

According to Proposition 4.2, it suffices to consider the spectrality of ν . The following special case that $n_k \equiv 2$ (i.e., $p_k \equiv 1$) for $k \geq 1$ provides a counterexample to show that the condition in Theorem 1.3 for ν to be a spectral measure is not a necessary condition.

Theorem 4.3. Assume that $p_k \equiv 1$ for $k \geq 1$. If $2 \mid b_k$ for $k \geq 2$, then ν defined by (4.4) is a spectral measure. Conversely, if ν is a spectral measure, b_k is not always even for $k \geq 2$.

Proof. Since $p_k \equiv 1$ for $k \geq 1$, we have

$$\mathcal{D}_k = \left\{ 0, \frac{1 - b_{k+1}^{-1}}{2} \right\}.$$

Define $\tilde{D}_k := 2b_{k+1}\mathcal{D}_k = \{0, b_{k+1} - 1\}$, the measure ν defined by (4.4) can be expressed as

$$\nu = \delta_{(2b_1b_2)^{-1}\tilde{D}_1} * \delta_{(2b_1b_2b_3)^{-1}\tilde{D}_2} * \delta_{(2b_1b_2b_3b_4)^{-1}\tilde{D}_3} * \cdots$$

which is a Moran-type Bernoulli convolution.

If $2 \mid b_k$ for all $k \geq 2$. We can check the condition in Theorem 2.4 that

$$\begin{aligned} s_{k+1} &= v_2 \left(\frac{b_1 b_2 \cdots b_{k+1} b_{k+2}}{b_{k+2} - 1} \right) \\ &= v_2 (b_1 b_2 \cdots b_{k+1} b_{k+2}) \\ &> v_2 (b_1 b_2 \cdots b_{k+1}) \\ &= v_2 \left(\frac{b_1 b_2 \cdots b_{k+1}}{b_{k+1} - 1} \right) \\ &= s_k \end{aligned}$$

for all $k \geq 1$. Hence ν is a spectral measure.

On the other hand, let $b_3 = 7, b_k = 8$ for $k \neq 3$, we can get

$$\begin{aligned} s_1 &= v_2 \left(\frac{b_1 b_2}{b_2 - 1} \right) = 6, \quad s_2 = v_2 \left(\frac{b_1 b_2 b_3}{b_3 - 1} \right) = 5, \\ s_k &= v_2 \left(\frac{b_1 b_2 \cdots b_{k+1}}{b_{k+1} - 1} \right) = 3k, \quad k = 3, 4, \dots \end{aligned}$$

Then s_k 's are different from each other for all $k \geq 1$. Therefore, ν is a spectral measure but $2 \nmid b_3$. $\square$

In the rest of this section, our main concern is the general case that $p_k \not\equiv 1$. First of all, we need to modify the expression of ν in a more appropriate way. By using (4.3), we have

$$\begin{aligned} \delta_{(b_1 \cdots b_k)^{-1}\mathcal{D}_k} &= \delta_{(b_1 \cdots b_k)^{-1}D_{p_k}} * \delta_{(b_1 \cdots b_k)^{-1}(p_k - \frac{1+b_{k+1}^{-1}}{2})D_2} \\ &= \delta_{(b_1 \cdots b_k)^{-1}D_{p_k}} * \delta_{(b_1 \cdots b_k 2b_{k+1})^{-1}(b_{k+1}(2p_k-1)-1)D_2}. \end{aligned}$$

Then the measure ν in (4.4) is of the following form

$$\begin{aligned} \nu &= \delta_{b_1^{-1}D_1} * \delta_{(b_1 b_2)^{-1}D_2} * \cdots * \delta_{(b_1 \cdots b_k)^{-1}\mathcal{D}_k} * \cdots \\ &= \delta_{b_1^{-1}D_{p_1}} * \delta_{(b_1 2b_2)^{-1}(b_2(2p_1-1)-1)D_2} * \delta_{(b_1 b_2)^{-1}D_{p_2}} * \delta_{(b_1 b_2 2b_3)^{-1}(b_3(2p_2-1)-1)D_2} * \cdots * \\ &\quad \delta_{(b_1 \cdots b_k)^{-1}D_{p_k}} * \delta_{(b_1 \cdots b_k 2b_{k+1})^{-1}(b_{k+1}(2p_k-1)-1)D_2} * \cdots \end{aligned}$$

Starting from the second term, by swapping the positions of two consecutive terms in the above expression, one can get

$$\begin{aligned} \nu &= \delta_{b_1^{-1}D_{p_1}} * \delta_{b_1^{-1}b_2^{-1}D_{p_2}} * \delta_{b_1^{-1}b_2^{-1}2^{-1}(b_2(2p_1-1)-1)D_2} * \delta_{b_1^{-1}b_2^{-1}2^{-1}(\frac{b_3}{2})^{-1}D_{p_3}} * \\ &\quad \delta_{b_1^{-1}b_2^{-1}2^{-1}(\frac{b_3}{2})^{-1}2^{-1}(b_3(2p_2-1)-1)D_2} * \delta_{b_1^{-1}b_2^{-1}b_3^{-1}2^{-1}(\frac{b_4}{2})^{-1}D_{p_4}} * \cdots * \\ &\quad \delta_{b_1^{-1}b_2^{-1}\cdots b_{k-1}^{-1}2^{-1}(\frac{b_k}{2})^{-1}D_{p_k}} * \delta_{b_1^{-1}b_2^{-1}\cdots b_{k-1}^{-1}2^{-1}(\frac{b_k}{2})^{-1}2^{-1}(b_k(2p_{k-1}-1)-1)D_2} * \cdots \end{aligned} \quad (4.6)$$

By removing the terms in which $p_k = 1$, we can relabel the above infinite convolution to ensure that every digit set contains at least two elements in the following way:

- (i) if $p_1 = 1$ and $p_2 = 1$, let $b'_1 = 2b_1b_2$, $D'_1 = (b_2(2p_1 - 1) - 1)D_2$;
- (ii) if $p_1 = 1$ and $p_2 \neq 1$, let $b'_1 = b_1b_2$, $D'_1 = D_{p_2}$ and $b'_2 = 2$, $D'_2 = (b_2(2p_1 - 1) - 1)D_2$;
- (iii) if $p_1 \neq 1$ and $p_2 = 1$, let $b'_1 = b_1$, $D'_1 = D_{p_1}$ and $b'_2 = 2b_2$, $D'_2 = (b_2(2p_1 - 1) - 1)D_2$;
- (iv) if $p_1 \neq 1$ and $p_2 \neq 1$, let $b'_1 = b_1$, $D'_1 = D_{p_1}$, $b'_2 = b_2$, $D'_2 = D_{p_2}$ and $b'_3 = 2$, $D'_3 = (b_2(2p_1 - 1) - 1)D_2$.

In general, for $k \geq 3$, we set

$$m_k = 2k - 2 - \#\{n : p_n = 1, n < k\}.$$

- (v) If $p_k \neq 1$, let $b'_{m_k} = \frac{b_k}{2}$, $D'_{m_k} = D_{p_k}$ and $b'_{m_k+1} = 2$, $D'_{m_k+1} = (b_k(2p_{k-1} - 1) - 1)D_2$;
- (vi) if $p_k = 1$, let $b'_{m_k} = b_k$, $D'_{m_k} = (b_k(2p_{k-1} - 1) - 1)D_2$.

Therefore, we get a sequence of integers $\{b'_k\}_{k=1}^\infty$ and a sequence of digit sets $\{D'_k\}_{k=1}^\infty$. Accordingly, the measure ν in (4.4) can be simplified to

$$\nu = \delta_{b'_1{}^{-1}D'_1} * \delta_{(b'_1b'_2)^{-1}D'_2} * \cdots \quad (4.7)$$

where $\#D'_k \geq 2$ for all $k \geq 1$.

Decompose such ν into two parts as before:

$$\nu = \nu_k * \nu_{>k}$$

where

$$\nu_k = \delta_{b'_1{}^{-1}D'_1} * \delta_{(b'_1b'_2)^{-1}D'_2} * \cdots * \delta_{(b'_1\cdots b'_k)^{-1}D'_k}$$

and

$$\nu_{>k} = \delta_{(b'_1\cdots b'_kb'_{k+1})^{-1}D'_{k+1}} * \delta_{(b'_1\cdots b'_kb'_{k+1}b'_{k+2})^{-1}D'_{k+2}} * \cdots.$$

Moreover, we scale the measure $\nu_{>k}$ by

$$\omega_{>k}(\cdot) := \nu_{>k} \left(\frac{\cdot}{b'_1 \cdots b'_k} \right) = \delta_{b'_{k+1}{}^{-1}D'_{k+1}} * \delta_{(b'_{k+1}b'_{k+2})^{-1}D'_{k+2}} * \cdots \quad (4.8)$$

Proposition 4.4. *If $p_1 \mid b_1$, $p_2 \mid b_2$, $2 \mid b_2$ and $2p_k \mid b_k$ for $k \geq 3$, then there exists a sequence of sets $\{L_k\}_{k=1}^\infty$ in $\mathbb{Z}$ such that (b'_k, D'_k, L_k) are all Hadamard triples.*

Proof. Cases (i)-(iv) are easily verified, we only check the general cases (v) and (vi). For $k \geq m_3$, there exists $i \geq 3$ such that $m_i \leq k < m_{i+1}$.

If $p_i \neq 1$ and $k = m_i$, then $b'_k = b'_{m_i} = \frac{b_i}{2}$, $D'_k = D'_{m_i} = D_{p_i}$, we can get

$$\delta_{b'_k-1} D'_k = \delta_{(\frac{b_i}{2})^{-1} D_{p_i}}.$$

It follows that the zero set

$$\mathcal{Z}(\widehat{\delta_{b'_k-1} D'_k}) = \frac{b_i}{2p_i}(\mathbb{Z} \setminus p_i \mathbb{Z}).$$

Let $L_k = \frac{b_i}{2p_i} D_{p_i}$. By the assumption that $2p_i \mid b_i$, we have that L_k is a spectrum of $\delta_{b'_k-1} D'_k$ with $L_k \subset \mathbb{Z}$. Hence (b'_k, D'_k, L_k) is a Hadamard triple.

If $p_i \neq 1$ and $k = m_i + 1$, then $b'_k = b'_{m_i+1} = 2$, $D'_k = D'_{m_i+1} = b_i(2p_{i-1} - 1)D_2$, we can get

$$\delta_{b'_k-1} D'_k = \delta_{2^{-1}(b_i(2p_{i-1}-1)-1)D_2}.$$

The zero set is

$$\mathcal{Z}(\widehat{\delta_{b'_k-1} D'_k}) = \frac{2\mathbb{Z} + 1}{b_i(2p_{i-1} - 1) - 1}.$$

Let $L_k = D_2$. By the assumption that $2 \mid b_i$, we have that L_k is a spectrum of $\delta_{b'_k-1} D'_k$ with $L_k \subset \mathbb{Z}$. Hence (b'_k, D'_k, L_k) is a Hadamard triple.

If $p_i = 1$, $k = m_i$, then $b'_k = b_i$, $D'_k = (b_i(2p_{i-1} - 1) - 1)D_2$. we can get

$$\delta_{b'_k-1} D'_k = \delta_{b_i^{-1}(b_i(2p_{i-1}-1)-1)D_2}.$$

The zero set is

$$\mathcal{Z}(\widehat{\delta_{b'_k-1} D'_k}) = \frac{b_i(2\mathbb{Z} + 1)}{2(b_i(2p_{i-1} - 1) - 1)}.$$

Let $L_k = \frac{b_i}{2} D_2$. By the assumption that $2 \mid b_i$, it concludes that L_k is the spectrum of $\delta_{b'_k-1} D'_k$ with $L_k \subset \mathbb{Z}$. Hence (b'_k, D'_k, L_k) is also a Hadamard triple. $\square$

Theorem 4.5. *If $p_2 \mid b_2$, $2 \mid b_2$ and $2p_k \mid b_k$ for $k \geq 3$, and the sequence $\{b_k\}_{k=1}^\infty$ is bounded, then ν in (4.7) is a spectral measure.*

Proof. By Lemma 2.3, the value of b_1 does not affect the spectrality of ν , hence we can assume that $p_1 \mid b_1$.

Depending on whether p_k is equal to 1 for $k \geq 1$, we divide $\{p_k\}_{k=1}^\infty$ into the following three cases:

Case 1: $p_k \equiv 1$ for all $k \geq 1$, the desired result follows from Theorem 4.3.

Case 2: There are only finitely many k 's that make $p_k \geq 2$. Then there exists a positive number $\kappa \in \mathbb{N}$ such that $p_k \equiv 1$ for all $k \geq \kappa$. Let $\{L_k\}_{k=1}^\infty \subset \mathbb{Z}$ be as in Proposition 4.4. Write

$$B = b'_1 \cdots b'_{m_\kappa}, \quad L = L_1 + b'_1 L_2 + \cdots + (b'_1 \cdots b'_{m_\kappa-1}) L_{m_\kappa}$$

and

$$D = (b'_2 \cdots b'_{m_\kappa}) D'_1 + (b'_3 \cdots b'_{m_\kappa}) D'_2 + \cdots + b'_{m_\kappa} D'_{m_\kappa-1} + D'_{m_\kappa}.$$

Then

$$\nu_{m_\kappa} = \delta_{(b'_1)^{-1}D'_1} * \delta_{(b'_1 b'_2)^{-1}D'_2} * \cdots * \delta_{(b'_1 \cdots b'_{m_\kappa})^{-1}D'_{m_\kappa}} = \delta_{B^{-1}D}.$$

Since $\{(b'_k, D'_k, L_k)\}_{k=1}^{m_\kappa}$ are finite many Hadamard triples, from Lemma 2.6, it follows that (B, D, L) is also a Hadamard triple. That is, ν_{m_κ} is a spectral measure with a spectrum L .

On the other hand, by (4.8),

$$\begin{aligned} \omega_{>m_\kappa} &= \delta_{b'^{-1}_{m_\kappa+1}D'_{m_\kappa+1}} * \delta_{(b'_{m_\kappa+1}b'_{m_\kappa+2})^{-1}D'_{m_\kappa+2}} * \cdots \\ &= \delta_{b_{\kappa+1}^{-1}(b_{\kappa+1}-1)D_2} * \delta_{(b_{\kappa+1}b_{\kappa+2})^{-1}(b_{\kappa+2}-1)D_2} * \cdots. \end{aligned}$$

Thus $\{(b'_k, D'_k, L_k)\}_{k=m_\kappa+1}^\infty$ are Hadamard triples with $\#D'_k = 2$ and $D'_k = \{0, b'_k - 1\} \subset \{0, 1, \dots, b'_k - 1\}$ for all $k \geq m_\kappa + 1$. Since $\{b'_k\}$ is bounded, by Lemma 2.7, we have that $\omega_{>m_\kappa}$ is a spectral measure with a spectrum $\Gamma \subset \mathbb{Z}$. Hence $\nu_{>m_\kappa}$ is also a spectral measure with a spectrum $B\Gamma \subset B\mathbb{Z}$. Set

$$\Lambda = L \oplus B\Gamma.$$

For any $t \in \mathbb{R}$,

$$\begin{aligned} Q_{\Lambda, \nu}(t) &= \sum_{\eta \in \Lambda} |\widehat{\nu}(t + \eta)|^2 \\ &= \sum_{\eta \in \Lambda} |\widehat{\nu}_{m_\kappa}(t + \eta)|^2 \cdot |\widehat{\nu}_{>m_\kappa}(t + \eta)|^2 \\ &= \sum_{\lambda \in L} \sum_{\gamma \in \Gamma} |\widehat{\nu}_{m_\kappa}(t + \lambda + B\gamma)|^2 \cdot |\widehat{\nu}_{>m_\kappa}(t + \lambda + B\gamma)|^2 \\ &= \sum_{\lambda \in L} \left(|\widehat{\nu}_{m_\kappa}(t + \lambda)|^2 \cdot \sum_{\gamma \in \Gamma} |\widehat{\nu}_{>m_\kappa}(t + \lambda + B\gamma)|^2 \right) \\ &= \sum_{\lambda \in L} |\widehat{\nu}_{m_\kappa}(t + \lambda)|^2 \equiv 1. \end{aligned}$$

Therefore, ν is a spectral measure with a spectrum Λ by Theorem 2.1.

Case 3: There are infinitely many k 's that make $p_k \geq 2$. That is, for all $l, k \geq 1$, there exists an integer $N(k) > k + l$, such that $p_{N(k)} \geq 2$. We can get

$$\gcd(D'_{m_{N(k)}} - D'_{m_{N(k)}}) = \gcd(D_{p_{N(k)}} - D_{p_{N(k)}}) = 1,$$

then

$$\gcd\left(\bigcup_{j=l+k}^\infty (D'_j - D'_j)\right) = 1. \quad (4.9)$$

Since $\{b_k\}_{k=1}^\infty$ is bounded, by Proposition 4.4, the sequence $\{(b'_k, D'_k, L_k)\}_{k=1}^\infty$ is chosen from a finite set of Hadamard triples. Thus the measure $\omega_{>l}$ is generated by finitely many Hadamard triples. By Lemma 2.12 and (4.9), the integral periodic zero set $Z(\omega_{>l}) = \emptyset$ for all $l \geq 1$.

(i) If $\{\omega_{>l}\}_{l=1}^\infty$ has no weakly convergent subsequences. In this case, we have $\text{cl}(\{\omega_{>l}\}_{l=1}^\infty) = \{\omega_{>l}\}_{l=1}^\infty$. Moreover, we may find a compact subset $K \subset \mathbb{R}$ such that $\omega_{>l} \in \mathcal{P}(K)$ for all $l \geq 1$. Hence the family of measures $\{\omega_{>l}\}_{l=1}^\infty$ is admissible and tight. Theorem 2.9 implies that ν is a spectral measure with a spectrum in $\mathbb{Z}$.

(ii) If there exists a subsequence, say $\{\omega_{>l_j}\}_{j=1}^\infty$, which converges weakly to ω . Let Ψ be the set of all measures generated by finitely many Hadamard triples, and let $\Psi_0 \subset \Psi$ be the set of the measures where there are infinitely many k 's such that $p_k \geq 2$. By Lemma 2.11, we have $\omega' \in \Psi$. If $\omega' \in \Psi_0$, then $Z(\omega') = \emptyset$. If $\omega' \in \Psi \setminus \Psi_0$, then ω' belongs to Case 1 or Case 2. That is, there exists a sufficiently large $M > 0$ such that $p_k \equiv 1$ for all $k \geq M$ in the expression of $\omega'_{>M}$. The sequence $\{\omega_{>l_j+M}\}_{j=1}^\infty$ converges weakly to $\omega'_{>M} \in \Psi \setminus \Psi_0$. In this situation, we may regard $\omega'_{>M}$ as $\omega_{>k}$ for some $k \geq m_\kappa$ in Case 2. The support of measure $\omega_{>k}$ is

$$\text{spt}(\omega_{>k}) \subset \left[0, \sum_{l=k+1}^\infty (b'_{k+1} \cdots b'_l)^{-1} (b'_l - 1)\right] \subset [0, 1].$$

That means $\omega_{>k} \in \mathcal{P}([0, 1])$. Since $\omega_{>k}$ is generated by Hadamard triples, it is purely singular without atoms, $\omega_{>k}(\{1\}) = 0$. Lemma 2.8 yields that $Z(\omega_{>k}) = \emptyset$. Therefore, $Z(\omega'_{>M}) = \emptyset$. Consequently, ν is a spectral measure by Theorem 2.10. $\square$

Proof of Theorem 1.3. The theorem directly follows from Proposition 4.2 and Theorem 4.5. $\square$

5. PROOF OF THEOREM 1.4

In this section, we follow the same notation as in Section 4, that is, $n_k = 2p_k$, $k \geq 1$ and ν is defined by (4.4).

To prove Theorem 1.4, we define a new Moran-type measure generated by a sequence of positive rational numbers $\{d_k := \frac{t_k}{t_k}\}_{k=1}^\infty$ and a sequence of consecutive digit sets $\mathcal{D}_k = \{0, 1, \dots, \gamma_k - 1\}$ as follows:

$$\nu = \delta_{d_1^{-1}\mathcal{D}_1} * \delta_{(d_1 d_2)^{-1}\mathcal{D}_2} * \cdots * \delta_{(d_1 \cdots d_k)^{-1}\mathcal{D}_k} * \cdots. \quad (5.1)$$

Write

$$\omega_{>k} = \delta_{(d_{k+1})^{-1}\mathcal{D}_{k+1}} * \delta_{(d_{k+1} d_{k+2})^{-1}\mathcal{D}_{k+2}} * \delta_{(d_{k+1} d_{k+2} d_{k+3})^{-1}\mathcal{D}_{k+3}} * \cdots. \quad (5.2)$$

If $\{t_k\}_{k=1}^\infty$ and $\{\gamma_k\}_{k=1}^\infty$ both are bounded, we let $c \in \mathbb{N}$ be the common multiple of $\{t_k\}_{k=1}^\infty$ and $\{\gamma_k\}_{k=1}^\infty$. Obviously, the zero set of $\hat{\nu}$ is

$$\mathcal{Z}(\hat{\nu}) = \bigcup_{k=1}^\infty d_1 \cdots d_k \left(\frac{\mathbb{Z} \setminus \gamma_k \mathbb{Z}}{\gamma_k} \right) \subset \frac{d_1}{c} \mathbb{Z}.$$

Assuming that Λ containing 0 is a spectrum of ν , we have $\frac{c}{d_1}\Lambda \subset \mathbb{Z}$. It follows that

$$\frac{1}{d_1}\Lambda = \bigcup_{n=0}^{c-1} \left(\frac{n}{c} + \Lambda_n \right),$$

where $\Lambda_n = \mathbb{Z} \cap \left(\frac{\Lambda}{d_1} - \frac{n}{c} \right)$ and $\frac{n}{c} + \Lambda_n = \emptyset$ if $\Lambda_n = \emptyset$.

Denote by $q_k := \frac{c}{\gamma_k} \in \mathbb{N}$, $k \geq 1$. For any $n \in \{0, 1, \dots, c-1\}$, there exists a unique pair of integers $(i, j) \in \{0, 1, \dots, q_1-1\} \times \{0, 1, \dots, \gamma_1-1\}$ such that

$$n = i + q_1 j.$$

Hence we further have

$$\frac{1}{d_1}\Lambda = \bigcup_{n=0}^{c-1} \left(\frac{n}{c} + \Lambda_n \right) = \bigcup_{i=0}^{q_1-1} \bigcup_{j=0}^{\gamma_1-1} \left(\frac{i + q_1 j}{c} + \Lambda_{i+q_1 j} \right). \quad (5.3)$$

The proof of the following technical lemma is similar to that of Proposition 3.1 in [33] (also Proposition 3.3 in [9]), while d_k 's here are rational numbers.

Lemma 5.1. *Let ν be defined by (5.1) with bounded $\{t_k\}_{k=1}^\infty$ and $\{\gamma_k\}_{k=1}^\infty$. If ν is a spectral measure with a spectrum Λ containing 0, then $\omega_{>1}$ as in (5.2) is a spectral measure and for any $\{j_i : 0 \leq i \leq q_1-1\} \subset \{0, 1, \dots, \gamma_1-1\}$, the set*

$$\Gamma = \bigcup_{i=0}^{q_1-1} \left(\frac{i + q_1 j_i}{c} + \Lambda_{i+q_1 j_i} \right)$$

is a spectrum of $\omega_{>1}$ if $\Gamma \neq \emptyset$.

Proof. We first show that Γ is a bi-zero set of $\omega_{>1}$ if $\Gamma \neq \emptyset$. It is obviously true if Γ has only one element. We just have to consider that Γ has at least two elements. For any $\lambda_1 \neq \lambda_2 \in \Gamma$, there exist $i_1, i_2 \in \{0, 1, \dots, q_1-1\}$, $z_1 \in \Lambda_{i_1+q_1 j_{i_1}}$, $z_2 \in \Lambda_{i_2+q_1 j_{i_2}}$ such that

$$\lambda_1 = \frac{i_1 + q_1 j_{i_1}}{c} + z_1, \quad \lambda_2 = \frac{i_2 + q_1 j_{i_2}}{c} + z_2.$$

By the definitions of Λ and Γ , we have $d_1 \Gamma \subset \Lambda$. Since $\nu(\cdot) = \delta_{d_1^{-1} \mathcal{D}_1}(\cdot) * \omega_{>1}(d_1 \cdot)$, it follows that

$$0 = \widehat{\nu}(d_1(\lambda_1 - \lambda_2)) = m_{\mathcal{D}_1}(\lambda_1 - \lambda_2) \widehat{\omega}_{>1}(\lambda_1 - \lambda_2).$$

Since $m_{\mathcal{D}_1}$ is a $\mathbb{Z}$ -periodic function, it yields that

$$\begin{aligned} m_{\mathcal{D}_1}(\lambda_1 - \lambda_2) &= m_{\mathcal{D}_1} \left(\frac{i_1 - i_2 + q_1(j_{i_1} - j_{i_2})}{c} + z_1 - z_2 \right) \\ &= m_{\mathcal{D}_1} \left(\frac{i_1 - i_2 + q_1(j_{i_1} - j_{i_2})}{c} \right). \end{aligned}$$

If $i_1 = i_2$, then $j_{i_1} = j_{i_2}$, $m_{\mathcal{D}_1}(\lambda_1 - \lambda_2) = m_{\mathcal{D}_1}(0) = 1$. Hence $\widehat{\omega}_{>1}(\lambda_1 - \lambda_2) = 0$.

If $i_1 \neq i_2$, since $i_1 - i_2 \in [1 - q_1, q_1 - 1] \setminus \{0\}$, we have $i_1 - i_2 \notin q_1\mathbb{Z}$. Suppose that $\lambda_1 - \lambda_2 \in \mathcal{Z}(m_{\mathcal{D}_1}) = \frac{\mathbb{Z} \setminus \gamma_1\mathbb{Z}}{\gamma_1}$, then there exists $z \in \mathbb{Z} \setminus \gamma_1\mathbb{Z}$ such that

$$\frac{i_1 - i_2 + q_1(j_{i_1} - j_{i_2})}{c} = \frac{z}{\gamma_1}.$$

Thus $c = q_1\gamma_1$ shows that $i_1 - i_2 = q_1(z - j_{i_1} + j_{i_2}) \in q_1\mathbb{Z}$, which is impossible. Therefore, $m_{\mathcal{D}_1}(\lambda_1 - \lambda_2) \neq 0$ and $\widehat{\nu}_{>1}(\lambda_1 - \lambda_2) = 0$, proving that Γ is a bi-zero set of $\omega_{>1}$.

Next we show that Γ is a spectrum of $\omega_{>1}$. Denote $\Lambda_{i,j} := \Lambda_{i+q_1j}$. For any $t \in \mathbb{R}$,

$$1 = Q_{\Lambda,\nu}(d_1t) = \sum_{i=0}^{q_1-1} \sum_{j=0}^{\gamma_1-1} \sum_{\lambda \in i+q_1j+c\Lambda_{i,j}} \left| \widehat{\nu} \left(\frac{d_1\lambda}{c} + d_1t \right) \right|^2,$$

where $\sum_{\lambda \in i+q_1j+c\Lambda_{i,j}} \left| \widehat{\nu} \left(\frac{d_1\lambda}{c} + d_1t \right) \right|^2 = 0$ if $\Lambda_{i,j} = \emptyset$. Since $\Lambda_{i,j} \subset \mathbb{Z}$, we have

$$\begin{aligned} 1 &= Q_{\Lambda,\nu}(d_1t) \\ &= \sum_{i=0}^{q_1-1} \sum_{j=0}^{\gamma_1-1} \sum_{\tilde{\lambda} \in \Lambda_{i,j}} \left| m_{\mathcal{D}_1} \left(\frac{i+q_1j}{c} + \tilde{\lambda} + t \right) \right|^2 \left| \widehat{\omega}_{>1} \left(\frac{i+q_1j}{c} + \tilde{\lambda} + t \right) \right|^2 \\ &= \sum_{i=0}^{q_1-1} \sum_{j=0}^{\gamma_1-1} \left| m_{\mathcal{D}_1} \left(\frac{i+q_1j}{c} + t \right) \right|^2 \sum_{\tilde{\lambda} \in \Lambda_{i,j}} \left| \widehat{\omega}_{>1} \left(\frac{i+q_1j}{c} + \tilde{\lambda} + t \right) \right|^2. \end{aligned}$$

Let $p_{i,j} := \left| m_{\mathcal{D}_1} \left(\frac{i+q_1j}{c} + t \right) \right|^2$ and $x_{i,j} := \sum_{\tilde{\lambda} \in \Lambda_{i,j}} \left| \widehat{\omega}_{>1} \left(\frac{i+q_1j}{c} + \tilde{\lambda} + t \right) \right|^2$, we have $\sum_{i=0}^{q_1-1} \sum_{j=0}^{\gamma_1-1} p_{i,j}x_{i,j} = 1$. Let $t \in \mathbb{R} \setminus \mathbb{Q}$, since $\mathcal{Z}(\mathcal{D}_1) \subset \mathbb{Q}$, we have $p_{i,j} > 0$.

For any $i \in \{0, 1, \dots, q_1 - 1\}$, let $L_i = \{i + q_1j : 0 \leq j \leq \gamma_1 - 1\}$, then $(c, \mathcal{D}_1, L_i)$ is Hadamard triple, that is,

$$\sum_{j=0}^{\gamma_1-1} p_{i,j} = \sum_{j=0}^{\gamma_1-1} \left| m_{\mathcal{D}_1} \left(\frac{i+q_1j}{c} + t \right) \right|^2 = 1.$$

Since $\{j_i : 0 \leq i \leq q_1 - 1\} \subset \{0, 1, \dots, \gamma_1 - 1\}$, Γ is a bi-zero set of $\omega_{>1}$, then

$$\sum_{i=0}^{q_1-1} \max\{x_{i,0}, x_{i,1}, \dots, x_{i,\gamma_1-1}\} \leq 1.$$

By Lemma 2.13, we have

$$\sum_{i=0}^{q_1-1} \sum_{\tilde{\lambda} \in \Lambda_{i,j}} \left| \widehat{\omega}_{>1} \left(\frac{i+q_1j}{c} + \tilde{\lambda} + t \right) \right|^2 = 1$$

for all $j \in \{0, 1, \dots, \gamma_1 - 1\}$, and

$$\begin{aligned}
& \sum_{\tilde{\lambda} \in \Lambda_{i,0}} \left| \widehat{\omega}_{>1} \left(\frac{i}{c} + \tilde{\lambda} + t \right) \right|^2 \\
&= \sum_{\tilde{\lambda} \in \Lambda_{i,1}} \left| \widehat{\omega}_{>1} \left(\frac{i+q_1}{c} + \tilde{\lambda} + t \right) \right|^2 \\
&\vdots \\
&= \sum_{\tilde{\lambda} \in \Lambda_{i,\gamma_1-1}} \left| \widehat{\omega}_{>1} \left(\frac{i+q_1(\gamma_1-1)}{c} + \tilde{\lambda} + t \right) \right|^2
\end{aligned} \tag{5.4}$$

for all $i \in \{0, 1, \dots, q_1 - 1\}$. Hence

$$Q_{\Gamma, \omega_{>1}}(t) = \sum_{\gamma \in \Gamma} |\widehat{\omega}_{>1}(t + \gamma)|^2 = 1$$

for $t \in \mathbb{R} \setminus \mathbb{Q}$. By the continuity of $Q_{\Gamma, \omega_{>1}}$, we have $Q_{\Gamma, \omega_{>1}}(t) \equiv 1$ for $t \in \mathbb{R}$. Theorem 2.1 implies that $\omega_{>1}$ is a spectral measure with the spectrum Γ . $\square$

Remark 5.2. From the proof of Lemma 5.1, it is easy to see that $\omega_{>k}$ is a spectral measure for all $k \geq 1$ provided that ν is a spectral measure. Furthermore, if Γ_k containing 0 is a spectrum of $\omega_{>k}$, then for any $\{j_i : 0 \leq i \leq q_k - 1\} \subset \{0, 1, \dots, \gamma_k - 1\}$, the set

$$\Gamma_{k+1} = \bigcup_{i=0}^{q_k-1} \left(\frac{i + q_k j_i}{c} + \Lambda_{i+q_k j_i} \right)$$

is a spectrum of $\omega_{>k+1}$ if $\Gamma_{k+1} \neq \emptyset$, where $c = q_k \gamma_k$ and $\Lambda_{i+q_k j_i} = \mathbb{Z} \cap \left(\frac{\Gamma_k}{d_k} - \frac{i+q_k j_i}{c} \right)$.

Proposition 5.3. Let ν be defined by (5.1) is a spectral measure with bounded $\{t_k\}$ and $\{\gamma_k\}$, then

- (i) If $\gamma_i \nmid t_{i+1}$ for some i , then $\gamma_{i+1} \mid l_{i+1}$.
- (ii) If $\gamma_i \nmid t_{i+1}$, $\gcd(l_{i+1}, t_{i+1}) = 1$, and $d_{i+2} = l_{i+2}$ is an integer satisfying $t_{i+1} \mid l_{i+2}$ for some i , then $t_{i+1} \gamma_{i+2} \mid l_{i+2}$.

Proof. By Remark 5.2, we only need to prove the case that $i = 1$. Let Λ be a spectrum of ν with $0 \in \Lambda$, then Lemma 5.1 shows that

$$\Gamma = \bigcup_{i=0}^{q_1-1} \left(\frac{i + q_1 j_i}{c} + \Lambda_{i+q_1 j_i} \right) = \bigcup_{i=0}^{q_1-1} \left(\frac{i}{c} + \frac{j_i}{\gamma_1} + \Lambda_{i,j_i} \right)$$

is a spectrum of $\omega_{>1}$ if $\Gamma \neq \emptyset$.

- (i) If $\gamma_1 \nmid t_2$. We set $j_0 = 0$, and for $i \neq 0$ we set $j_i = 0$ if $\frac{l_2}{\gamma_2} \not\equiv t_2 \frac{i}{c} \pmod{1}$, $j_i = 1$ if $\frac{l_2}{\gamma_2} \equiv t_2 \frac{i}{c} \pmod{1}$. Then $\gamma_1 \nmid t_2$ implies that $\frac{l_2}{\gamma_2} \not\equiv t_2 \frac{i+q_1 j_i}{c} \pmod{1}$ for $i \neq 0$.

Using (5.3) for $\omega_{>1}$, the spectrum Γ has the following multi-stage decomposition

$$\begin{aligned}\Gamma &= \bigcup_{i=0}^{q_1-1} \left(\frac{i + q_1 j_i}{c} + \Lambda_{i, j_i} \right) \\ &= d_2 \bigcup_{i'=0}^{q_2-1} \bigcup_{j'=0}^{\gamma_2-1} \left(\frac{i' + q_2 j'}{c} + \Gamma_{i', j'} \right)\end{aligned}\quad (5.5)$$

where $\Gamma_{i', j'} = \mathbb{Z} \cap \left(\frac{\Gamma}{d_2} - \frac{i' + q_2 j'}{c} \right)$. Since $0 \in \Gamma_{0,0} \neq \emptyset$, by (5.4), we have $\Gamma_{0, j'} \neq \emptyset$ for all $0 \leq j' \leq \gamma_2 - 1$. Choose an integer $z'_0 \in \Gamma_{0,1}$, then there exists $(i, j_i) \in \{0, 1, \dots, q_1 - 1\} \times \{0, 1, \dots, \gamma_1 - 1\}$ and an integer $z_0 \in \Lambda_{i, j_i}$ such that

$$\frac{l_2}{t_2} \left(\frac{1}{\gamma_2} + z'_0 \right) = \frac{i + q_1 j_i}{c} + z_0.$$

This yields that $i = j_i = 0$ since $\frac{l_2}{\gamma_2} \not\equiv t_2 \frac{i + q_1 j_i}{c} \pmod{1}$ for any $1 \leq i \leq q_1 - 1$. Hence $\frac{l_2}{\gamma_2} + l_2 z'_0 = t_2 z_0$, showing that $\gamma_2 \mid l_2$.

(ii) If $\gamma_1 \nmid t_2$, $\gcd(l_2, t_2) = 1$, and $d_3 = l_3$ is an integer satisfying $t_2 \mid l_3$, We set $j_0 = 0$, and for $i \neq 0$ we set $j_i = 0$ if $\frac{t_2 i}{c} \notin \mathbb{Z}$, $j_i = 1$ if $\frac{t_2 i}{c} \in \mathbb{Z}$. Hence $0 \in \Lambda_{0,0} \subset \Gamma$, Γ is a spectrum of $\omega_{>1}$. Since $\gamma_1 \nmid t_2$, one can check that

$$\left(\frac{i + q_1 j_i}{c} \right) t_2 = \frac{t_2 i}{c} + \frac{t_2}{\gamma_1} j_i \notin \mathbb{Z}$$

for any $i \neq 0$. By using (5.5) and the fact that $\Lambda_{i, j_i}, \Gamma_{i', j'} \subset \mathbb{Z}$ for any i, i', j' , we have $l_2 \Gamma_{0,0} \subset t_2 \Lambda_{0,0}$. It follows from $\gcd(l_2, t_2) = 1$ that $\Gamma_{0,0} \subset t_2 \mathbb{Z}$.

For any $0 \leq \iota \leq q_2 - 1$, choose $0 \leq \tau_\iota \leq \gamma_2 - 1$ and let

$$\Lambda^* = \bigcup_{\iota=0}^{q_2-1} \left(\frac{\iota + q_2 \tau_\iota}{c} + \Gamma_{\iota, \tau_\iota} \right)$$

where $\tau_\iota = 0$ for $\iota = 0$. Then Λ^* is a spectrum of the measure $\omega_{>2}$ by Remark 5.2.

Using multi-stage decomposition again, we have

$$\begin{aligned}\Lambda^* &= \bigcup_{\iota=0}^{q_2-1} \left(\frac{\iota + q_2 \tau_\iota}{c} + \Gamma_{\iota, \tau_\iota} \right) \\ &= d_3 \bigcup_{s=0}^{q_3-1} \bigcup_{t=0}^{\gamma_3-1} \left(\frac{s + q_3 t}{c} + \Lambda_{s,t}^* \right) \\ &= l_3 \bigcup_{s=0}^{q_3-1} \bigcup_{t=0}^{\gamma_3-1} \left(\frac{s}{c} + \frac{t}{\gamma_3} + \Lambda_{s,t}^* \right)\end{aligned}$$

where $\Lambda_{s,t}^* = \mathbb{Z} \cap \left(\frac{\Lambda^*}{d_3} - \frac{s+q_3t}{c} \right)$. That $d_3 = l_3$ implies that $t_3 = 1$ and $\gamma_2 \nmid t_3$, hence $\gamma_3 \mid l_3$ by (i), it follows that

$$l_3 \left(\frac{t}{\gamma_3} + \Lambda_{0,t}^* \right) \subset \mathbb{Z}$$

for any $0 \leq t \leq \gamma_3 - 1$. It can be seen that

$$l_3 \left(\frac{t}{\gamma_3} + \Lambda_{0,t}^* \right) \subset \Gamma_{0,0} \subset t_2 \mathbb{Z}.$$

Since $t_2 \mid l_3$, let $l_3 = l'_3 t_2$, then $l'_3 \left(\frac{t}{\gamma_3} + \Lambda_{0,t}^* \right) \subset \mathbb{Z}$ for any $0 \leq t \leq \gamma_3 - 1$. Consequently, we have $\gamma_3 \mid l'_3$ and $t_2 \gamma_3 \mid l_3$, completing the proof. $\square$

Proposition 5.4. *Suppose that $2 \mid p_k$ for all $k \geq 1$. If ν in (4.4) is a spectral measure, then $2 \mid b_{k+1}$ for all $k \geq 1$.*

Proof. Since $2 \mid p_k, k \geq 1$, we get

$$D_{p_k} = D_2 \oplus D_{\frac{p_k}{2}}.$$

The measure ν can be expressed as

$$\begin{aligned} \nu &= \delta_{b_1^{-1}D_2} * \delta_{b_1^{-1}D_{\frac{p_1}{2}}} * \delta_{b_1^{-1}(p_1 - \frac{1+b_2^{-1}}{2})D_2} * \cdots * \delta_{(b_1 \cdots b_k)^{-1}D_2} * \delta_{(b_1 \cdots b_k)^{-1}D_{\frac{p_k}{2}}} * \\ &\quad \delta_{(b_1 \cdots b_k)^{-1}(p_k - \frac{1+b_{k+1}^{-1}}{2})D_2} * \delta_{(b_1 \cdots b_k b_{k+1})^{-1}D_2} * \delta_{(b_1 \cdots b_k b_{k+1})^{-1}D_{\frac{p_{k+1}}{2}}} * \cdots . \end{aligned}$$

If $2 \mid b_{k+1}$ does not hold for all $k \geq 1$, that is, there exists $n \geq 1$ such that $b_{n+1} \in 2\mathbb{Z} + 1$. Then we have

$$\mathcal{Z} \left(\widehat{\delta}_{(b_1 \cdots b_{n+1})^{-1}D_2} \right) = \frac{b_1 \cdots b_{n+1}}{2} (2\mathbb{Z} + 1) \subset \frac{b_1 \cdots b_n}{2} (2\mathbb{Z} + 1) = \mathcal{Z} \left(\widehat{\delta}_{(b_1 \cdots b_n)^{-1}D_2} \right).$$

Writing $\nu = \delta_{(b_1 \cdots b_{n+1})^{-1}D_2} * \nu'$. Since $\mathcal{Z} \left(\widehat{\delta}_{(b_1 \cdots b_{n+1})^{-1}D_2} \right) \subset \mathcal{Z}(\widehat{\nu'})$, if Λ is a spectrum of ν , then Λ is a bi-zero set of ν' . But Λ can not be a spectrum of ν by Lemma 2.2. That is a contradiction. $\square$

For the measure ν as in (4.6), we may change the digit sets in each level into the consecutive digit sets by adjusting the contraction ratios as follows:

$$\begin{aligned} \nu &= \delta_{b_1^{-1}D_{p_1}} * \delta_{b_1^{-1}b_2^{-1}D_{p_2}} * \delta_{b_1^{-1}b_2^{-1}2^{-1}(b_2(2p_1-1)-1)D_2} * \delta_{b_1^{-1}b_2^{-1}2^{-1}(\frac{b_3}{2})^{-1}D_{p_3}} * \\ &\quad \delta_{b_1^{-1}b_2^{-1}2^{-1}(\frac{b_3}{2})^{-1}2^{-1}(b_3(2p_2-1)-1)D_2} * \cdots \\ &= \delta_{b_1^{-1}D_{p_1}} * \delta_{b_1^{-1}b_2^{-1}D_{p_2}} * \delta_{b_1^{-1}b_2^{-1}(\frac{2}{b_2(2p_1-1)-1})^{-1}D_2} * \delta_{b_1^{-1}b_2^{-1}(\frac{2}{b_2(2p_1-1)-1})^{-1}(\frac{b_3(b_2(2p_1-1)-1)}{2})^{-1}D_{p_3}} * \\ &\quad \delta_{b_1^{-1}b_2^{-1}(\frac{2}{b_2(2p_1-1)-1})^{-1}(\frac{b_3(b_2(2p_1-1)-1)}{2})^{-1}(\frac{2}{b_3(2p_2-1)-1})^{-1}D_2} * \cdots . \end{aligned}$$

Let

$$\begin{aligned} b_1'' &= b_1, & b_{2k+1}'' &= \frac{2}{b_{k+1}(2p_k - 1) - 1}, \\ b_2'' &= b_2, & b_{2k+2}'' &= \frac{b_{k+2}(b_{k+1}(2p_k - 1) - 1)}{2}, \end{aligned} \quad (5.6)$$

and

$$p_1'' = p_1, \quad p_2'' = p_2, \quad p_{2k+1}'' = 2, \quad p_{2k+2}'' = p_{k+2}. \quad (5.7)$$

Then ν can be rewritten as

$$\nu = \delta_{b_1''^{-1}D_{p_1''}} * \delta_{b_1''^{-1}b_2''^{-1}D_{p_2''}} * \delta_{b_1''^{-1}b_2''^{-1}b_3''^{-1}D_{p_3''}} * \cdots. \quad (5.8)$$

Accordingly, let

$$\omega_{>k} = \delta_{b_{k+1}''^{-1}D_{p_{k+1}''}} * \delta_{b_{k+1}''^{-1}b_{k+2}''^{-1}D_{p_{k+2}''}} * \delta_{b_{k+1}''^{-1}b_{k+2}''^{-1}b_{k+3}''^{-1}D_{p_{k+3}''}} * \cdots. \quad (5.9)$$

Theorem 5.5. *Let $2 \mid p_k$ for all $k \geq 1$ and $\{b_k\}_{k=1}^\infty$ be bounded. If ν in (5.8) is a spectral measure, then $p_2 \mid b_2$ and $2p_k \mid b_k$ for $k \geq 3$.*

Proof. Proposition 5.4 implies that $2 \mid b_k$ for all $k \geq 2$. By (5.6), (5.7) and Proposition 5.3(i), we have $p_2 \mid b_2$.

From Lemma 5.1, it follows that $\omega_{>1}$ as in (5.9) is a spectral measure where

$$\begin{aligned} \omega_{>1} &= \delta_{b_2''^{-1}D_{p_2''}} * \delta_{b_2''^{-1}b_3''^{-1}D_{p_3''}} * \delta_{b_2''^{-1}b_3''^{-1}b_4''^{-1}D_{p_4''}} * \cdots \\ &= \delta_{b_2^{-1}D_{p_2}} * \delta_{b_2^{-1}(\frac{2}{b_2(2p_1-1)-1})^{-1}D_2} * \delta_{b_2^{-1}(\frac{2}{b_2(2p_1-1)-1})^{-1}(\frac{b_3(b_2(2p_1-1)-1)}{2})^{-1}D_{p_3}} * \cdots. \end{aligned}$$

Since p_2 is even and $(b_2(2p_1 - 1) - 1)$ is odd, we get $p_2 \nmid (b_2(2p_1 - 1) - 1)$ and $\gcd(2, b_2(2p_1 - 1) - 1) = 1$. Moreover, $b_2(2p_1 - 1) - 1 \mid \frac{b_3(b_2(2p_1-1)-1)}{2}$ (as $2 \mid b_3$). Therefore, Proposition 5.3(ii) yields that $p_3(b_2(2p_1 - 1) - 1) \mid \frac{b_3(b_2(2p_1-1)-1)}{2}$, and hence $p_3 \mid \frac{b_3}{2}$.

Repeating the above argument, we conclude that $p_k \mid \frac{b_k}{2}$ for all $k \geq 3$, completing the proof. $\square$

Proof of Theorem 1.4. The theorem directly follows from Proposition 4.2, Theorem 4.5 and Theorem 5.5. $\square$

REFERENCES

- [1] L.X. An, X. Fu and C.K. Lai, *On spectral Cantor-Moran measures and a variant of Bourgain's sum of sine problem*, Adv. Math. 349 (2019) 84-124.
- [2] L.X. An and X.G. He, *A class of spectral Moran measures*, J. Funct. Anal. 266 (1) (2014) 343-354.
- [3] L.X. An, X.G. He and K.S. Lau, *Spectrality of a class of infinite convolutions*, Adv. Math. 283 (2015) 362-376.
- [4] L.X. An, X.G. He and H.X. Li, *Spectrality of infinite Bernoulli convolutions*, J. Funct. Anal. 269 (2015) 1571-1590.

- [5] L.X. An, Q. Li and M.M. Zhang, *The generalized Fuglede's conjecture holds for a class of Cantor-Moran measures*, Pacific J. Math. 334 (2) (2025) 189-209. L.X. An, Q. Li and M.M. Zhang, *Characterization of spectral Cantor-Moran measures with consecutive digits*, preprint (2022).
- [6] C. Bishop and Y. Peres, *Fractals in probability and analysis*, Cambridge studies in advanced mathematics, (2017).
- [7] X.R. Dai, *When does a Bernoulli convolution admit a spectrum?*, Adv. Math. 231 (3-4) (2012) 1681-1693.
- [8] X.R. Dai, X.G. He and K.S. Lau, *On spectral N -Bernoulli measures*, Adv. Math. 259 (2014) 511-531.
- [9] Q.R. Deng and M.T. Li, *Spectrality of Moran-type self-similar measures on $\mathbb{R}$* , J. Math. Anal. Appl. 506 (1) (2022) 125547.
- [10] Q.R. Deng and M.T. Li, *Spectrality of Moran-type Bernoulli convolutions*, Bull. Malays. Math. Sci. Soc. 46 (4) (2023) 136.
- [11] D. Dutkay, J. Haussermann and C.K. Lai, *Hadamard triples generate self-affine spectral measures*, Trans. Amer. Math. Soc. 371 (2) (2019) 1439-1481.
- [12] B. Fuglede, *Commuting self-adjoint partial differential operators and a group theoretic problem*, J. Funct. Anal. 16 (1974) 101-121.
- [13] L. He and X.G. He, *On the Fourier orthonormal bases of Cantor-Moran measures*, J. Funct. Anal. 272 (5) (2017) 1980-2004.
- [14] T.Y. Hu and K.S. Lau, *Spectral property of the Bernoulli convolutions*, Adv. Math. 219 (2) (2008) 554-567.
- [15] S. Hua, H. Rao, Z. Wen and J. Wu, *On the structures and dimensions of Moran sets*, Sci. China Ser. A-Math. 43 (2000) 836-852.
- [16] P. Jorgensen and S. Pedersen, *Dense analytic subspaces in fractal L^2 -spaces*, J. Anal. Math. 75 (1998) 185-228.
- [17] M.N. Kolountzakis and M. Matolcsi, *Tiles with no spectra*, Forum Math. 18 (3) (2006) 519-528.
- [18] M.N. Kolountzakis and M. Matolcsi, *Complex Hadamard matrices and the spectral set conjecture*, Collect. Math. Extra (2006) 281-291.
- [19] I. Laba and Y. Wang, *On spectral Cantor measures*, J. Funct. Anal. 193 (2) (2002) 409-420.
- [20] N. Lev and M. Matolcsi, *The Fuglede conjecture for convex domains is true in all dimensions*, Acta Math. 228 (2) (2022) 385-420.
- [21] W.X. Li, J.J. Miao and Z.Q. Wang, *Weak convergence and spectrality of infinite convolutions*, Adv. Math. 404 (2022) 108425.
- [22] W.X. Li, J.J. Miao and Z.Q. Wang, *Spectrality of random convolutions generated by finitely many Hadamard triples*, Nonlinearity 37 (1) (2024) 015003.
- [23] W.X. Li, J.J. Miao and Z.Q. Wang, *Spectrality of infinite convolutions and random convolutions.*, J. Funct. Anal. 287 (7) (2024) 110539.
- [24] J. Liu, Z.Y. Lu and T. Zhou, *Spectrality of Moran-Sierpinski type measures*, J. Funct. Anal. 284 (6) (2023) 109820.
- [25] J.C. Liu, Q.Q. Liu and J.J. Luo and J.J. Wang, *Spectrality of a class of Moran measures on $\mathbb{R}^2$* , preprint, (2024).
- [26] Z.S. Liu, *Spectrality of homogeneous Moran measures on the plane*, Chaos Solitons Fract. 183 (2024) 114926.
- [27] M. Matolcsi, *Fuglede's conjecture fails in dimension 4*, Proc. Amer. Math. Soc. 133 (10) (2005) 3021-3026.

- [28] R. Shi, *Spectrality of a class of Cantor–Moran measures*, J. Funct. Anal. 276 (12) (2019) 3767-3794.
- [29] R.S. Strichartz, *Mock Fourier series and transforms associated with certain Cantor measures*, J. Anal. Math. 81 (2000) 209-238.
- [30] M.W. Tang and F.L. Yin, *Spectrality of Moran measures with four-element digit sets*, J. Math. Anal. Appl. 461 (1) (2018) 354-363.
- [31] T. Tao, *Fuglede’s conjecture is false in 5 and higher dimensions*, Math. Res. Let. 11 (2-3) (2004) 251-258.
- [32] H.H. Wu, *Spectral self-similar measures with alternate contraction ratios and consecutive digits*, Adv. Math. 443 (2024) 109585.
- [33] S. Wu and Y.Q. Xiao, *Spectrality of a class of infinite convolutions on $\mathbb{R}$* , Nonlinearity, 37 (5) (2024) 055015.

COLLEGE OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF NONLINEAR ANALYSIS AND ITS APPLICATIONS (CHONGQING UNIVERSITY), MINISTRY OF EDUCATION, CHONGQING UNIVERSITY, CHONGQING, 401331, P.R. CHINA

Email address: jun.luo@cqu.edu.cn

COLLEGE OF MATHEMATICS AND STATISTICS, KEY LABORATORY OF NONLINEAR ANALYSIS AND ITS APPLICATIONS (CHONGQING UNIVERSITY), MINISTRY OF EDUCATION, CHONGQING UNIVERSITY, CHONGQING, 401331, P.R. CHINA

Email address: maol@stu.cqu.edu.cn

KEY LABORATORY OF COMPUTING AND STOCHASTIC MATHEMATICS (MINISTRY OF EDUCATION), SCHOOL OF MATHEMATICS AND STATISTICS, HUNAN NORMAL UNIVERSITY, CHANGSHA, HUNAN 410081, P.R. CHINA

Email address: jcliu@hunnu.edu.cn