

# Experiments for Smooth RS-MCA Proximity Thresholds

## Fresh-Eyes Threshold Memo, Missing Theorems, and Retained Locator Ledger

RS-MCA experimental notes

June 26, 2026

### Abstract

This replacement starts with the current “fresh eyes” synthesis for the smooth Reed–Solomon Proximity Prize problem. The negative side is now quite rigid: Paper D v5 gives a prize-facing upper cap on the MCA threshold through a self-contained deep-point route, while Paper A and the list lower bounds rule out any no-slack capacity theorem. The positive side is not yet a theorem; it should be organized around two local-limit problems, L1 arbitrary-word locator fibers and M1 all-line residue packing, plus extension-line MCA, Paper D conversion/radius audits, and exact protocol-ledger rewrites.

The main new additions over the old `experiments.tex` are two compact bridge lemmas. First, for ordinary column-distance interleaving, interleaved lists inject into common feasible supports, so a uniform base locator-fiber theorem gives the same polynomial exponent for all protocol arities; there is no automatic  $n^{\mu B}$  product loss. Second, a list bound plus an MCA bound implies a concrete line-decoding bound with loss  $M + nL + 1$ . The older BCHKS provenance notes, quotient-locator dictionaries, normal forms, arithmetic gates, and finite-threshold infrastructure are retained below whenever they still look useful.

## Contents

|          |                                                                 |          |
|----------|-----------------------------------------------------------------|----------|
| <b>1</b> | <b>Fresh-Eyes Settlement Memo</b>                               | <b>3</b> |
| 1.1      | Bottom line . . . . .                                           | 4        |
| 1.2      | What is already proved or very close to proved . . . . .        | 4        |
| 1.3      | The corrected positive theorem shape . . . . .                  | 5        |
| 1.4      | What remains missing . . . . .                                  | 5        |
| <b>2</b> | <b>New Bridge I: Common-Support Interleaved Fiber Injection</b> | <b>6</b> |
| <b>3</b> | <b>New Bridge II: List plus MCA Gives Line Decoding</b>         | <b>7</b> |
| <b>4</b> | <b>Updated Positive-Theorem Package</b>                         | <b>8</b> |
| <b>5</b> | <b>Immediate Experimental Additions</b>                         | <b>8</b> |
| <b>6</b> | <b>Fresh Non-Claims</b>                                         | <b>9</b> |
| <b>7</b> | <b>Retained Previous Experiments Material</b>                   | <b>9</b> |
| <b>8</b> | <b>Status, Provenance, and What Counts as New</b>               | <b>9</b> |

|           |                                                                                     |           |
|-----------|-------------------------------------------------------------------------------------|-----------|
| <b>9</b>  | <b>BCHKS Locator Construction in Smooth-Quotient Notation</b>                       | <b>10</b> |
| 9.1       | Imported theorem/proof pattern . . . . .                                            | 10        |
| 9.2       | Notation dictionary . . . . .                                                       | 11        |
| 9.3       | The shared locator identity . . . . .                                               | 11        |
| 9.4       | List-fiber pigeonhole wrapper . . . . .                                             | 12        |
| 9.5       | Slack-two and subfield target for the older CS25 route . . . . .                    | 12        |
| 9.6       | Pasteable main-paper wording . . . . .                                              | 13        |
| <b>10</b> | <b>Triage of the AI-Generated Four-Item Packet</b>                                  | <b>13</b> |
| <b>11</b> | <b>Restricted Resonance Setup for Paper B</b>                                       | <b>15</b> |
| <b>12</b> | <b>Cycle 14 Normal Form</b>                                                         | <b>16</b> |
| <b>13</b> | <b>Cycle 18 Component Split</b>                                                     | <b>16</b> |
| <b>14</b> | <b>New Divisibility-Gate Theorem</b>                                                | <b>17</b> |
| <b>15</b> | <b>Nearby Proven Gates</b>                                                          | <b>18</b> |
| <b>16</b> | <b>Slope Map and Remaining Heuristics</b>                                           | <b>18</b> |
| <b>17</b> | <b>Next Experimental Checks</b>                                                     | <b>19</b> |
| <b>18</b> | <b>June 26 Normal-Form and Dependency Results</b>                                   | <b>19</b> |
| 18.1      | Syndrome-pencil normal form for support-wise line incidence . . . . .               | 19        |
| 18.2      | Interleaved codegree reduction . . . . .                                            | 20        |
| 18.3      | Deep-point cap dependency split . . . . .                                           | 21        |
| 18.4      | Common code-line residual budget . . . . .                                          | 22        |
| <b>19</b> | <b>Recent PR Triage: L1, F1, M1, and Challenge-Map Ledgers</b>                      | <b>23</b> |
| 19.1      | L1: proper-subgroup and monomial-prefix lanes . . . . .                             | 23        |
| 19.2      | F1: arbitrary-anchor residue clouds . . . . .                                       | 23        |
| 19.3      | M1: all-line Hankel aperiodic packet . . . . .                                      | 24        |
| 19.4      | M1: same-slope one-exchange root slices . . . . .                                   | 24        |
| 19.5      | M1: variable-line packet reduction . . . . .                                        | 34        |
| 19.6      | M1: width-one fixed-root closure . . . . .                                          | 35        |
| 19.7      | Challenge-map pullbacks and public status corrections . . . . .                     | 36        |
| <b>20</b> | <b>High-Agreement Tangent Staircase</b>                                             | <b>36</b> |
| <b>21</b> | <b>High-Agreement Adjacent Ledgers</b>                                              | <b>40</b> |
| 21.1      | No-loss CA and projective slopes . . . . .                                          | 40        |
| 21.2      | Finite-parameter curve MCA . . . . .                                                | 41        |
| 21.3      | Interleaved-list uniqueness . . . . .                                               | 42        |
| 21.4      | The $\mathbb{F}_{17^{32}}$ , $n = 512$ , $k = 256$ protocol-facing ledger . . . . . | 43        |

|                                                                                           |           |
|-------------------------------------------------------------------------------------------|-----------|
| <b>22 General High-Agreement Ledgers Beyond the <math>\mathbb{F}_{17^{32}}</math> Row</b> | <b>44</b> |
| 22.1 A coordinate tangent star . . . . .                                                  | 45        |
| 22.2 Uniform line, CA, and projective-slope threshold . . . . .                           | 45        |
| 22.3 Degree- $d$ curve MCA/CA . . . . .                                                   | 46        |
| 22.4 Interleaved-list uniqueness . . . . .                                                | 47        |
| 22.5 General finite-grid threshold calculator . . . . .                                   | 47        |
| 22.6 Prize-rate applicability . . . . .                                                   | 49        |
| <b>23 Towards-Prize Finite-Threshold Theorems</b>                                         | <b>49</b> |
| <b>24 June 29 Finite-Slope and Interleaved-List PR Packets</b>                            | <b>52</b> |
| 24.1 Boundary-off external-anchor normal form . . . . .                                   | 52        |
| 24.2 $a = 507$ adjacent bridge route cut . . . . .                                        | 53        |
| 24.3 Interleaved-list lower row at agreement 326 . . . . .                                | 53        |
| 24.4 Random simple-pole finite-slope floors . . . . .                                     | 53        |
| 24.5 Coset-packet finite-slope floors for $a = 261, \dots, 288$ . . . . .                 | 54        |
| <b>25 Non-Claims</b>                                                                      | <b>54</b> |
| <b>A BibTeX Entry to Copy into Main Papers</b>                                            | <b>55</b> |

## 1 Fresh-Eyes Settlement Memo

I would frame the current state as follows. The attached suite substantially settles the negative side and gives a plausible corrected theorem shape, but it does not yet settle the Proximity Prize threshold in the sharp “largest  $\delta^*$ ” sense. The clean prize-relevant statement is no longer “prove smooth RS reaches capacity”. It is

$$\delta = 1 - \rho - \eta, \quad \eta \text{ must clear generated-field entropy, quotient-core, MCA-floor,} \\ \text{interleaving, line-field, and challenge-field budgets.}$$

For

$$C = \text{RS}[\mathbb{F}, D, k], \quad n = |D|, \quad \rho = k/n \in \{1/2, 1/4, 1/8, 1/16\},$$

the official challenge statements use this MCA/list-threshold form; in this experimental ledger the concrete prize envelope is [1]

$$\varepsilon_* = 2^{-128}, \quad k \leq 2^{40}, \quad |\mathbb{F}| < 2^{256}.$$

The protocol-relevant soundness contribution has the schematic form

$$\varepsilon_{\text{mca}}(C, \delta) + \frac{|\Lambda(C^{\equiv \mu}, \delta)|}{q_{\text{line}}} + \text{query error.}$$

Thus a list theorem alone is not enough unless it is connected to the exact MCA, correlated-agreement, line-decoding, or curve-MCA quantity used by the protocol, and unless the denominator is the actual line or challenge field.

## 1.1 Bottom line

For the **MCA/correlated-agreement threshold**, the strongest attached result is Paper D v5’s universal cap. Through the self-contained deep-point route and the stated dyadic-divisor hypotheses,

$$\delta_{\text{MCA}}^*(2^{-128}) \leq 1 - \rho - 2^{-9} \quad (\rho = 1/2, 1/4, 1/8),$$

and

$$\delta_{\text{MCA}}^*(2^{-128}) \leq 1 - \rho - 2^{-10} \quad (\rho = 1/16).$$

Equivalently, throughout the relevant unsafe intervals Paper D gives

$$\varepsilon_{\text{mca}}(C, \delta) > \frac{1}{2k} \left(1 - \frac{n}{|\mathbb{F}|}\right) \geq 2^{-86} \gg 2^{-128}.$$

This is a real prize-facing cap, not merely a heuristic obstruction. It is not an error-one theorem; it proves failure probability on the order of  $1/k$ , which is already far above  $2^{-128}$ , but it does not characterize the full failure curve.

For the **interleaved-list threshold**, Paper A’s base list lower bound plus the base-to-interleaved monotonicity gives a clean negative cap. At quotient scale  $N \mid n$ , with  $\rho N \in \mathbb{Z}$ , there is a received word with at least

$$\binom{N}{\rho N + 1} / q, \quad q = |\mathbb{F}|,$$

base-code codewords at radius  $1 - \rho - 1/N$ . Therefore the same radius is unsafe for the interleaved-list budget whenever this exceeds  $2^{-128}q$ . In the worst case  $q < 2^{256}$ , the following choices suffice:

| $\rho$ | $N$  | list-failure radius | $\log_2 \binom{N}{\rho N + 1} - 256$ | margin over $2^{128}$ |
|--------|------|---------------------|--------------------------------------|-----------------------|
| 1/2    | 512  | $1/2 - 2^{-9}$      | 251.17                               | 123.17                |
| 1/4    | 512  | $3/4 - 2^{-9}$      | 156.33                               | 28.33                 |
| 1/8    | 1024 | $7/8 - 2^{-10}$     | 298.68                               | 170.68                |
| 1/16   | 2048 | $15/16 - 2^{-11}$   | 433.89                               | 305.89                |

The fourth column is the exponent of the lower-bound list size after dividing by the worst-case field size  $2^{256}$ ; the fifth subtracts the target exponent 128.

## 1.2 What is already proved or very close to proved

The old no-slack smooth-domain capacity claim is dead. The quotient-locator construction gives explicit bad slopes for canonical lines and, over deployed smooth prime fields such as BabyBear, KoalaBear, and  $3 \cdot 2^{30} + 1$ , yields support-wise line-MCA error one on intervals starting at  $1 - \rho - 2^{-18}$  for all four prize rates, with the sharper  $2^{-17}$  gap at rates 1/2 and 1/4. These statements rule out any theorem saying that every radius below  $1 - \rho$  has negligible MCA error.

Paper D is the most prize-relevant negative theorem because it converts a deep point/list phenomenon into MCA at the exact  $2^{-128}$ -style threshold. In v5, the headline MCA cap is routed locally through a self-contained deep-point argument rather than the Crites–Stewart list-to-agreement import. The remaining CA/list comparison and fallback statements still need their own radius, normalization, augmented-code, and import audits.

The list lower-bound side is also stronger than it is currently advertised. It should be promoted to an explicit interleaved-list negative corollary: base lists embed into ordinary interleaved lists, and the binomial numerators above already exceed the  $2^{-128}|\mathbb{F}|$  budget in the full field envelope for the displayed quotient divisors.

### 1.3 The corrected positive theorem shape

The positive theorem should use

$$a = \lceil (1 - \delta)n \rceil = k + \sigma, \quad \delta = 1 - \rho - \eta,$$

and require at least

$$\sigma \log_2 q_{\text{gen}} \geq (1 + \varepsilon) \log_2 \binom{n}{k + \sigma}, \quad \sigma \geq C \frac{n}{\log n},$$

plus explicit quotient-profile and failure-floor ledgers. The first inequality uses the generated field  $q_{\text{gen}}$ , not a larger extension or challenge field that merely appears later in the protocol. The MCA numerator must be divided by the actual line field  $q_{\text{line}}$ .

The intended list target is an arbitrary-word locator-fiber theorem

$$\sup_U |\text{Fib}_U(k + \sigma)| \leq n^B.$$

The intended MCA target is an all-line bound of the form

$$\varepsilon_{\text{mca}}(C, 1 - \rho - \eta) \leq \frac{n^{1+o(1)} + 2^{(\beta(\rho)/H_2(\rho))\mathcal{Q}_H(\eta)(1+o(1))}}{q_{\text{line}}},$$

where quotient-periodic classes are removed or charged by  $\mathcal{Q}_H(\eta)$ . The current packet contains important special cases and reductions, but not these full arbitrary-word and all-line theorems.

### 1.4 What remains missing

1. **L1: arbitrary-word locator local limit.** This is the real list upper-bound theorem. Characteristic-zero rigidity, monomial-prefix lanes, and split-prime transfer results are useful but do not yet imply the uniform polynomial-field theorem needed for arbitrary received words.
2. **M1: all-line aperiodic residue packing.** Canonical-line exactness and quotient floors are lower-bound tools, not all-line upper bounds. The positive MCA theorem needs a fixed-field, all-line, all-base residue-packing bound.
3. **F1: extension-line MCA.** The list side has a clean extension-code identity. MCA does not. Base-valued bad lines can deflate when sampled over an extension field, while Paper D's cap indicates that genuinely extension-valued bad lines exist in relevant regimes. Any extension-line lift must therefore be a corrected-reserve theorem, not a blanket subfield statement.
4. **Paper D conversion audit.** Paper D v5's headline MCA cap is no longer load-bearing on the direct Crites–Stewart conversion. The self-contained deep-point step, support-wise monotonicity, integer-radius conventions, and the remaining CA/list import statements should still be independently checked.
5. **Protocol-ledger rewrite.** Deployed FRI/WHIR/IOP analyses must be rewritten with exact terms

$$|\Lambda(C^{\equiv \mu}, \delta)|/q_{\text{line}} \quad \text{and} \quad \varepsilon_{\text{mca}} \text{ or line-decoding over the actual line field.}$$

The survey toy protocol already has this shape; deployed analyses still need the same ledger.

## 2 New Bridge I: Common-Support Interleaved Fiber Injection

**Status: New-local wrapper.** The older ledger treated interleaving conservatively through a safe product bound  $L_\mu(\delta) \leq L_1(\delta)^\mu$ . For ordinary column-distance RS interleaving, the following support-level injection gives the sharper upper bound: once the one-row locator-fiber theorem is proved, interleaving costs no extra exponent.

For  $S \subseteq D$ , let

$$L_S(X) = \prod_{x \in S} (X - x).$$

For a received word  $U \in \mathbb{F}^D$ , the condition  $\deg(U \bmod L_S) < k$  means that the unique polynomial of degree less than  $|S|$  interpolating  $U|_S$  has degree less than  $k$ .

**Definition 1** (one-row and common interleaved fibers). *Let  $C = \text{RS}[\mathbb{F}, D, k]$ , let  $a \geq k$ , and let  $\mathbf{U} = (U_1, \dots, U_\mu) \in (\mathbb{F}^\mu)^D$ . Define*

$$\text{Fib}_{\mathbf{U}}^{(\mu)}(a) = \{S \subseteq D : |S| = a, \deg(U_i \bmod L_S) < k \text{ for every } i = 1, \dots, \mu\}.$$

*For one row, write  $\text{Fib}_{U_i}(a)$  for the same condition with only  $U_i$ .*

**Lemma 1** (common-support interleaved fiber injection). *Let  $C = \text{RS}[\mathbb{F}, D, k]$ ,  $n = |D|$ , and let  $a = \lceil (1 - \delta)n \rceil \geq k$ . For every received  $\mu$ -row word  $\mathbf{U} \in (\mathbb{F}^\mu)^D$ , ordinary column-distance interleaving satisfies*

$$|\Lambda(C^{\equiv \mu}, \delta, \mathbf{U})| \leq |\text{Fib}_{\mathbf{U}}^{(\mu)}(a)| \leq \min_{1 \leq i \leq \mu} |\text{Fib}_{U_i}(a)|.$$

*Consequently,*

$$\sup_{\mathbf{U}} |\Lambda(C^{\equiv \mu}, \delta, \mathbf{U})| \leq \sup_U |\text{Fib}_U(a)|.$$

*In particular, if L1 proves  $\sup_U |\text{Fib}_U(k + \sigma)| \leq n^B$ , then*

$$|\Lambda(C^{\equiv \mu}, 1 - \rho - \sigma/n)| \leq n^B$$

*for every ordinary column-distance protocol arity  $\mu$ .*

*Proof.* Fix a total order on  $D$ . A listed interleaved codeword  $(P_1, \dots, P_\mu) \in C^{\equiv \mu}$  at column distance at most  $\delta$  from  $\mathbf{U}$  agrees with  $\mathbf{U}$  on at least  $a$  common columns. Choose the first  $a$ -subset  $S$  of those agreeing columns. For this  $S$ , every row  $U_i|_S$  is interpolated by the degree- $< k$  polynomial  $P_i$ , hence  $S \in \text{Fib}_{\mathbf{U}}^{(\mu)}(a)$ .

For fixed  $S$ , the tuple  $(P_1, \dots, P_\mu)$  is uniquely determined, because each  $P_i$  has degree less than  $k$  and is fixed by its values on  $|S| \geq k$  points. Thus the canonical-support map from listed interleaved codewords to common feasible supports is injective. The second inequality is immediate because every common feasible support is feasible for each individual row. Taking suprema gives the final claim.  $\square$

**Remark 1** (impact). *For ordinary column-distance RS interleaving, L2 should be demoted from a separate product-exponent problem to a corollary of L1. The hard list theorem remains the base arbitrary-word locator local limit.*

### 3 New Bridge II: List plus MCA Gives Line Decoding

**Status: New-local bridge.** The next proposition turns a line-decoding proof obligation into a finite-constant budget once L1 and M1 are available. It uses finite slopes and the support-wise MCA normalization used in this packet.

**Proposition 1** (list/MCA bridge to line decoding). *Let  $C = \text{RS}[\mathbb{F}, D, k]$ , let  $|D| = n$ , and let  $a = \lceil (1 - \delta)n \rceil > k$ . Suppose that every allowed received line  $f + \gamma g$  has at most  $M$  support-wise MCA-bad finite slopes at radius  $\delta$ , equivalently  $\varepsilon_{\text{mca}}(C, \delta) \leq M/|\mathbb{F}|$  for the chosen finite slope sampler. Suppose also that the two-row interleaved list size satisfies*

$$|\Lambda(C^{\equiv 2}, \delta, W)| \leq L \quad \text{for every } W \in (\mathbb{F}^2)^D.$$

*Then  $C$  is line-decodable with parameters*

$$(\delta, M + nL + 1, n + 1)$$

*in the following sense: if an allowed received line  $f + \gamma g$  has more than  $M + nL$  finite slopes  $\gamma$  that are  $\delta$ -close to selected codewords  $U(\gamma) \in C$ , then there are codewords  $c_1, c_2 \in C$  such that*

$$U(\gamma) = c_1 + \gamma c_2$$

*for at least  $n + 1$  of those slopes.*

*Proof.* For each close slope  $\gamma$ , choose a support  $S_\gamma \subseteq D$  of size  $a$  on which  $f + \gamma g$  agrees with the selected codeword  $U(\gamma)$ . Discard the at most  $M$  support-wise MCA-bad slopes. For every remaining slope, support-wise containment gives codewords  $c_{1,\gamma}, c_{2,\gamma} \in C$  such that

$$c_{1,\gamma}|_{S_\gamma} = f|_{S_\gamma}, \quad c_{2,\gamma}|_{S_\gamma} = g|_{S_\gamma}.$$

Thus  $(c_{1,\gamma}, c_{2,\gamma})$  is a two-row interleaved codeword  $\delta$ -close to the received two-row word  $(f, g)$ , using the common support  $S_\gamma$ . There are at most  $L$  possible pairs.

If the original line has more than  $M + nL$  close slopes, then after discarding bad slopes more than  $nL$  slopes remain. By pigeonhole, one pair  $(c_1, c_2)$  occurs for at least  $n + 1$  slopes. For each such slope, the codewords  $U(\gamma)$  and  $c_1 + \gamma c_2$  both agree with  $f + \gamma g$  on  $S_\gamma$ , hence agree with each other on  $a > k$  positions. Since both have degree less than  $k$ , they are identical codewords. This proves the stated line-decoding conclusion.  $\square$

**Remark 2** (finite-constant consequence). *If M1 gives  $M = n^{1+o(1)}$  and L1 plus Lemma 1 gives  $L = n^B$ , then Proposition 1 gives*

$$M + nL + 1 \leq n^{B+1+o(1)}.$$

*For  $n \leq 2^{44}$  and  $|\mathbb{F}| < 2^{256}$ , a raw list numerator  $n^B$  is compatible with a  $2^{-128}$  field budget only for about  $B \lesssim 2.9$ , and the line-decoding bridge is compatible only for about  $B \lesssim 1.9$ . Thus the bridge is useful, but it leaves a real finite-constant target.*

## 4 Updated Positive-Theorem Package

The clean theorem to aim for is the following finite-reserve package. Let

$$C = \text{RS}[\mathbb{F}_{\text{line}}, D, k], \quad n = |D|, \quad a = k + \sigma, \quad \delta = 1 - a/n = 1 - \rho - \eta.$$

Assume

$$\sigma \log_2 q_{\text{gen}} \geq (1 + \varepsilon) \log_2 \binom{n}{k + \sigma}, \quad \sigma \geq C_{\rho, B, \varepsilon} \frac{n}{\log n},$$

the quotient profile  $\mathcal{Q}_H(\eta)$  is empty or explicitly budgeted,  $\eta$  lies above all proved failure floors including Paper D's universal-cap floor, and the MCA or line experiment is proved over  $q_{\text{line}}$ , not merely over  $q_{\text{chal}}$ .

**Conjecture 1** (corrected finite reserve package). *Under the preceding hypotheses, the desired conclusions are*

$$|\Lambda(C^{\equiv \mu}, \delta)| \leq n^B$$

for all ordinary column-distance protocol arities  $\mu$ , by Lemma 1; and

$$\varepsilon_{\text{mca}}(C, \delta) \leq \frac{n^{1+o(1)} + 2^{(\beta(\rho)/H_2(\rho))\mathcal{Q}_H(\eta)(1+o(1))}}{q_{\text{line}}},$$

or the same bound over  $q_{\text{gen}}$  plus a proved extension-line lift. If the consumed primitive is line decoding, then Proposition 1 should give

$$C \text{ is } (\delta, M + nL + 1, n + 1) \text{ line-decodable,}$$

with  $M$  and  $L$  supplied by M1 and L1.

This package would turn the current negative cap into a sharp threshold program: below the cap one cannot hope for  $2^{-128}$  MCA soundness, while above the corrected reserve one would have finite numerators that can be checked against  $q_{\text{line}}$ .

## 5 Immediate Experimental Additions

1. Add `experimental/interleaved_fiber_injection.md`, containing Lemma 1. Update certificate logic so an L1 locator-fiber theorem gives  $L_\mu = n^B$ , not  $n^{\mu B}$ , for ordinary column-distance RS interleaving.
2. Add `experimental/list_mca_to_linedecoding.md`, containing Proposition 1. This gives a concrete M2 bridge and turns line decoding into a finite-constant ledger problem.
3. Build a certificate scanner reporting generated-field entropy margin, quotient profile after  $k = \rho n - r$ , Paper D cap floor, base and interleaved list numerators, MCA numerator, line-decoding numerator, field separation among  $q_{\text{gen}}$ ,  $q_{\text{line}}$ , and  $q_{\text{chal}}$ , extension status, and exact failure audit.
4. Keep CA/list statements that use Crites–Stewart marked CONDITIONAL until audited. Paper D v5's MCA cap should instead be tracked as PROVED when the divisor, binomial, and subfield hypotheses are verified.



## 6 Fresh Non-Claims

This revision does not claim an exact settlement of the smooth-case Proximity Prize threshold. It claims that the shape is now rigid:

No unslacked capacity theorem; prove a reserve theorem at  $1 - \rho - \eta$ .

The theorem-backed negative side gives a draft universal MCA cap at constant gap  $2^{-9}$  for  $\rho = 1/2, 1/4, 1/8$  and  $2^{-10}$  for  $\rho = 1/16$ . The positive side should be pursued as

L1 locator local limit + M1 residue-line packing + extension-line theorem + protocol-ledger rewrite.

The interleaved-list problem is less fundamental than it looked in the old file: for column-distance interleaving, a uniform locator-fiber theorem immediately bounds the interleaved list at the same exponent.

## 7 Retained Previous Experiments Material

The remaining sections are retained from the previous `experiments.tex` whenever they still provide useful provenance, notation, normal forms, finite certificate gates, or non-claims. The new front matter above should be read as the current threshold-level guide; the older material below remains a toolbox.

## 8 Status, Provenance, and What Counts as New

This file is the place where we collect new things, but it must separate new repository contributions from imported constructions. The status labels used below are:

|           |                                                                  |
|-----------|------------------------------------------------------------------|
| IMPORTED  | an external theorem, proof pattern, or conversion;               |
| WRAPPER   | a repository repackaging, dictionary, or pigeonhole consequence; |
| NEW-LOCAL | a restricted algebraic theorem proved in this note;              |
| TARGET    | a precise proof obligation not yet promoted;                     |
| HEURISTIC | evidence or a search direction, not a theorem.                   |

**Hard attribution rule.** Any locator-polynomial argument derived from the subgroup construction of Ben-Sasson–Carmon–Haböck–Kopparty–Saraf must cite [2, Thm. 7.1] where the construction is stated or proved. Any admissible subgroup instantiation must also cite [2, Thm. 1.13]. The citation belongs at the theorem statement and again at the proof entry point; a bibliography entry or acknowledgement alone is not enough.

**What is new in this note.** The following items are the repository-side contributions collected here.

- A smooth-quotient notation dictionary translating BCHKS subgroup notation into the RS-MCA notation  $Q = D^{c_{\text{nb}}}$  and the dyadic/FFT-domain ledgers used in Papers A–D.
- A list-fiber pigeonhole wrapper: once a locator construction assigns many subset certificates to a smaller slope set, one heavy slope has many locator certificates, and, after a distinctness check, many nearby codewords.

- A slack-two and subfield-pigeonhole proof target for the older Crites–Stewart/CA route: the imported locator construction should be run at the augmented-code rung needed by that conversion, and slopes should be pigeonholed over the generated/base field when the construction really takes values there.
- A NEW-LOCAL divisibility gate in the restricted Paper B window  $B = \mathbb{F}_p$ ,  $F = \mathbb{F}_{p^2}$ ,  $D = \mathbb{F}_p$ ,  $t = \sigma = 2$ ,  $j = 3$ . This is independent of the BCHKS priority issue and is the main theorem proved in Section 14.

**What is not new.** The polynomial identity behind the item-(d) locator proof is not new here. In BCHKS notation it is the expansion of a vanishing polynomial over a complement of a chosen subset; in smooth-quotient notation it is the same expansion of  $\prod_{b \in A} (X^{c_{\text{fb}}} - b)$ , up to choosing a subset or its complement. The use of  $\ell$  for the chosen subset size also follows the external presentation closely enough that it should either be cited or renamed to  $\ell_{\text{pg}}$ .

**Source map.** The main papers remain the authoritative documents.

- Paper A, `tex/RS_disproof_v3.tex`, refutes the old no-slack smooth-domain support-wise line-MCA statement. If it uses the locator construction, it should cite BCHKS and call the repository contribution a smooth-domain/list-fiber corollary, not a new locator proof.
- Paper B, `tex/slackMCA_v3.tex`, is the main target for the local resonance material in Sections 11–17.
- Paper C, `tex/snarks_v4.tex`, records protocol ledgers. Nothing in this note may be consumed by a protocol without a separate conversion theorem.
- Paper D, `tex/cs25_cap_v4.tex`, gives the universal cap via an imported list-to-agreement route. Its locator-side input should cite BCHKS explicitly if it uses the quotient-locator construction.

## 9 BCHKS Locator Construction in Smooth-Quotient Notation

**Status: Imported plus Wrapper.** This section is a provenance-safe restatement and dictionary. It is included so later main-paper edits have a clean place to copy from.

### 9.1 Imported theorem/proof pattern

BCHKS prove a general multiplicative-subgroup construction [2, Thm. 7.1]. In a compact form, the imported statement is as follows.

**Theorem 1** (BCHKS quotient-locator construction, imported). *Let  $G \subseteq H \subseteq \mathbb{F}_q^*$  be multiplicative subgroups and let  $\Phi : H \rightarrow G$  be  $x \mapsto x^c$ , where  $c = |H|/|G|$ . Let  $E \subseteq G$ , and let  $\ell_{\text{pg}}, a \geq 1$  satisfy*

$$|E^{(+\ell_{\text{pg}})}| = \#\{e_1 + \cdots + e_{\ell_{\text{pg}}} : e_i \in E \text{ distinct}\} \geq a.$$

*Let  $D = \Phi^{-1}(E)$ ,  $n = |D|$ , and*

$$C = \text{RS}[\mathbb{F}_q, D, n - (\ell_{\text{pg}} + 2)c].$$

Then, for  $\gamma = \ell_{\text{pg}}c/n$ , there are functions  $f, g : D \rightarrow \mathbb{F}_q$  such that

$$\#\{z \in \mathbb{F}_q : \Delta(f + zg, C) \leq \gamma\} \geq a,$$

while

$$\Delta([f, g], C^2) \geq (\ell_{\text{pg}} + 1)c/n.$$

*Reference.* This is exactly the imported construction of [2, Thm. 7.1]; the proof is not reproduced as a repository contribution. BCHKS define the line endpoints by the monomials  $x^{n-\ell_{\text{pg}}c}$  and  $x^{n-(\ell_{\text{pg}}+1)c}$ , then expand a vanishing polynomial over the complement of a chosen  $\ell_{\text{pg}}$ -subset. The coefficient of  $X^{n-(\ell_{\text{pg}}+1)c}$  is an affine transform of a restricted sum from  $E^{(+\ell_{\text{pg}})}$ , giving the exceptional slope values.  $\square$

The BCHKS Theorem 1.13 is the admissible-subgroup specialization of this construction: take  $E = G$ , take  $H = D$ , set  $c = |D|/|G|$ , and use  $\ell_{\text{pg}} = |G|/2$ . Any RS-MCA statement corresponding to that specialization should cite [2, Thm. 1.13] directly.

## 9.2 Notation dictionary

| BCHKS notation                                                       | RS-MCA notation                                                               | Meaning                                                                                                                    |
|----------------------------------------------------------------------|-------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------|
| $H$                                                                  | evaluation domain $D$                                                         | Large multiplicative domain on which codewords are evaluated.                                                              |
| $G$                                                                  | quotient $Q = D^{c_{\text{fib}}}$                                             | Image of the power map; one point in $Q$ has $c_{\text{fib}}$ preimages in $D$ .                                           |
| $c =  H / G $                                                        | $c_{\text{fib}} = n/ Q $                                                      | Fiber size of the quotient map. Avoid calling this $a$ , because BCHKS also uses $a$ for the number of exceptional slopes. |
| $\Phi(x) = x^c$                                                      | $x \mapsto x^{c_{\text{fib}}}$                                                | Quotient/fiber map.                                                                                                        |
| $E \subseteq G$                                                      | active quotient subset $E \subseteq Q$                                        | Quotient-level support used by the locator.                                                                                |
| $E^{(+\ell_{\text{pg}})}$                                            | restricted coefficient/slope set                                              | first- Sums of distinct chosen quotient elements; slopes are affine transforms of these sums.                              |
| $\ell_{\text{pg}}$                                                   | $\ell_{\text{pg}}$ or renamed subset size                                     | Number of chosen quotient elements. Use $\ell_{\text{pg}}$ in RS-MCA text to avoid collision with other $\ell$ 's.         |
| $f = x^{n-\ell_{\text{pg}}c}, \quad g = x^{n-(\ell_{\text{pg}}+1)c}$ | locator line endpoints                                                        | Same monomial line after quotient notation is substituted.                                                                 |
| $P_S(X) = \prod_{\alpha \in S'} (X^c - \alpha)$                      | $\prod_{b \in A} (X^{c_{\text{fib}}} - b)$ , possibly after complementing $A$ | The locator polynomial.                                                                                                    |
| exceptional $z$ 's                                                   | bad slopes / heavy list fibers                                                | Slope parameters for which a line word is close to the RS code.                                                            |

## 9.3 The shared locator identity

The following identity is written down because it is the exact place where priority can be lost. The identity is the BCHKS identity in RS-MCA notation.

**Lemma 2** (Locator expansion, imported identity). *Let  $D$  map  $c_{\text{fib}}$ -to-one onto a quotient set  $E$  by  $x \mapsto x^{c_{\text{fib}}}$ , and let  $A \subseteq E$  have size  $\ell_{\text{pg}}$ . Put  $A^c = E \setminus A$  and define*

$$P_A(X) = \prod_{\alpha \in A^c} (X^{c_{\text{fib}}} - \alpha).$$

*Then*

$$P_A(X) = X^{n - \ell_{\text{pg}} c_{\text{fib}}} - e_1(A^c) X^{n - (\ell_{\text{pg}} + 1) c_{\text{fib}}} + \text{terms of degree at most } n - (\ell_{\text{pg}} + 2) c_{\text{fib}}.$$

*Equivalently, since  $e_1(A^c) = e_1(E) - e_1(A)$ , the coefficient of the next monomial is an affine transform of the restricted sum  $e_1(A)$ .*

*Reference.* This is the elementary expansion used inside [2, Thm. 7.1]. We record it only to fix notation. It should not be cited as a new lemma of the repository unless the BCHKS provenance is present at the statement.  $\square$

## 9.4 List-fiber pigeonhole wrapper

The next proposition is the repository-side wrapper around the imported identity. It is intentionally stated in certificate form, because distinctness of the nearby codewords is a separate check in concrete instantiations.

**Proposition 2** (List-fiber pigeonhole wrapper). *Let  $\mathcal{A}$  be a family of  $\ell_{\text{pg}}$ -subsets of a quotient set  $E$ . Suppose the BCHKS locator construction assigns to each  $A \in \mathcal{A}$  a slope  $z(A)$  lying in a finite set  $S$ , and a locator certificate showing that the corresponding word  $f + z(A)g$  is within radius  $\gamma$  of a locator codeword  $P_A \in C$ . Then some slope  $z_0 \in S$  has at least*

$$\frac{|\mathcal{A}|}{|S|}$$

*locator certificates above it. If the codewords  $P_A$  attached to the certificates over  $z_0$  are distinct, then the Hamming ball of radius  $\gamma$  around  $f + z_0 g$  contains at least  $|\mathcal{A}|/|S|$  codewords of  $C$ .*

*Proof.* The first assertion is the pigeonhole principle applied to the map  $A \mapsto z(A)$ . The second assertion is only the definition of list size, after the distinctness check removes possible duplicate locator codewords. The construction that supplies the certificates is imported from Theorem 1.  $\square$

**Remark 3** (What the wrapper adds). *The new part here is not the locator polynomial. The wrapper isolates the extra bookkeeping needed by RS-MCA: which slope set pays the pigeonhole bill, whether the slope set is the ambient field  $\mathbb{F}$  or a generated/base subfield  $B$ , and whether locator certificates remain distinct after specialization. Those are the checks needed before using the construction in Papers A, B, or D.*

## 9.5 Slack-two and subfield target for the older CS25 route

**Status: Target.** The older CS25/CA route needs a locator-heavy list fiber in the augmented code rung  $C^+ = \text{RS}[\mathbb{F}, D, k + 1]$ , because that is the object fed into the Crites–Stewart list-to-agreement conversion. Paper D v5 no longer uses this as the load-bearing MCA-cap route, but the target remains useful for CA/list comparison audits. The proof obligation should be stated as follows.

**Proposition 3** (Slack-two/subfield locator target). *Let  $D \subseteq B \subseteq \mathbb{F}$ , and suppose a BCHKS-type quotient locator construction over  $D$  produces slopes in  $B$ , not merely in  $\mathbb{F}$ . If the one-rung/augmented-code version produces distinct locator codewords in  $C^+ = \text{RS}[\mathbb{F}, D, k+1]$  for a family  $\mathcal{A}$  of certificates, then there exists a slope  $z_0 \in B$  with list size at least*

$$\frac{|\mathcal{A}|}{|B|}$$

*inside the appropriate Hamming ball for  $C^+$ .*

*Bookkeeping only.* This is Proposition 2 with  $S = B$ . The nontrivial work is external or elsewhere: one must verify the BCHKS locator hypotheses at the augmented-code rung, verify that the slope coefficients lie in  $B$ , and verify distinctness of the produced codewords. Until those checks are present, this proposition is a target, not a promoted Paper D theorem.  $\square$

## 9.6 Pasteable main-paper wording

**Paper A wording.** The polynomial locator identity used here is the quotient-fiber form of the construction of Ben-Sasson–Carmon–Haböck–Kopparty–Saraf [2, Thm. 7.1]; their Theorem 1.13 is the admissible subgroup instantiation. Thus the locator proof itself is not a new contribution of this manuscript. The present use is the smooth-domain specialization, the field/divisor bookkeeping, and the list-fiber pigeonhole packaging.

**Paper D wording.** The locator-side input is the BCHKS quotient-locator construction [2, Thm. 7.1; cf. Thm. 1.13], written in the smooth-domain quotient notation  $Q = D^{\text{cfib}}$ . The additional bookkeeping needed here is the slack-two augmented-code rung and, when available, subfield pigeonholing over  $B$  rather than over the ambient extension field  $\mathbb{F}$ .

## 10 Triage of the AI-Generated Four-Item Packet

**Status: Audit.** The four advertised items should be positioned as follows before any promotion into a main paper. The original AI packet used the labels (a)–(d) without making them self-contained in this note, so we decode them here.

| Item | Short name                     | Meaning in this ledger                                                                                                                                                                                                                                                                                                                                                                                                                                               |
|------|--------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (a)  | Weak-slack positive regime     | A positive-looking proximity/list or MCA-adjacent statement that only reaches slack on the order of $1/\sqrt{n}$ . It is useful as a comparison theorem, but it is far from the constant/reserve book-keeping needed for the corrected RS-MCA target.                                                                                                                                                                                                                |
| (b)  | Finite Fermat-prime packet     | A special finite-domain instantiation over only three Fermat-prime cases, in the current experimental ledger represented by the $p \in \{17, 257, 65537\}$ style checks. It is evidence or a worked computation, not a uniform theorem.                                                                                                                                                                                                                              |
| (c)  | Exponential-field construction | A large-field variant whose proof requires the ambient field to be exponential in the domain/parameter size. This can clarify the mechanism but does not match the polynomial-field or challenge-field regimes needed by the Proximity Prize ledger.                                                                                                                                                                                                                 |
| (d)  | BCHKS quotient-locator packet  | The interesting locator/proximity-gap construction: choose a quotient of a multiplicative subgroup, use $\ell_{\text{pg}}$ -subsets and restricted sums to create many exceptional slopes or nearby locator certificates. This is the BCHKS construction of [2, Thm. 7.1], with the admissible subgroup instance in [2, Thm. 1.13]; the repository contribution is only the smooth-domain dictionary, list-fiber wrapper, and later slack-two/subfield book-keeping. |

Item (d) should not be confused with the NEW-LOCAL divisibility-gate theorem in Section 14. The former is an imported locator/proximity construction; the latter is an independent restricted Paper B resonance calculation.

| Item | Limitation                                                 | Repository handling                                                                                                                                                                |
|------|------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| (a)  | Works only at slack on the order of $1/\sqrt{n}$ .         | Use as a weak-slack comparison point only. It does not clear the corrected-reserve MCA objective.                                                                                  |
| (b)  | Applies only to three Fermat-prime cases.                  | Treat as finite-prime evidence or a worked computation, not as a uniform smooth-domain theorem.                                                                                    |
| (c)  | Requires exponential field size.                           | Treat as a large-field sanity check. It is not directly usable for polynomial-field or prize-ledger parameters.                                                                    |
| (d)  | Most relevant, but the locator proof is essentially BCHKS. | Import, cite, and adapt from [2, Thm. 7.1 and Thm. 1.13]. The repository contribution is only the smooth-domain/list-fiber/slack-two packaging after exact hypotheses are checked. |

**Promotion checklist.** Before any statement using item (d) moves to Papers A–D, check the following.

- Does the statement cite BCHKS at the theorem and proof entry point?
- Is the object a proximity-gap statement, a list statement, CA, MCA, line-decoding, or a protocol-facing theorem?
- Which field pays the slope/list pigeonhole bill:  $B$ ,  $\mathbb{F}$ , or another generated field?
- Are the locator codewords distinct after specialization?
- Is the claimed slack in the same normalization as the consumed theorem?
- Are items (a), (b), and (c) kept out of asymptotic/protocol claims unless their limitations are explicitly part of the theorem statement?

## 11 Restricted Resonance Setup for Paper B

**Status:** New-local for the divisibility gate; otherwise local normal-form bookkeeping.

The remaining sections preserve and reorganize the Cycle 14–18 algebraic material. This branch is independent of the BCHKS priority correction.

The common restricted setting is

$$B = \mathbb{F}_p, \quad F = \mathbb{F}_{p^2}, \quad D = \mathbb{F}_p, \quad t = \sigma = 2, \quad j = 3,$$

and we work off the tangent/global endpoint

$$R_0 = \{ [W]_E \wedge [B_{\text{num}}]_E = 0 \}.$$

The field ledgers remain distinct:

$$q_{\text{gen}} = p, \quad q_{\text{line}} = p^2.$$

The regime is sub-reserve, since  $\eta = \sigma/n = 2/n$ . Thus none of the following proves a corrected-reserve list, CA, MCA, line-decoding, SNARK, or protocol theorem.

The base/generated field is  $B = \mathbb{F}_p$ . The line field is the quadratic extension  $F = \mathbb{F}_{p^2} = B(\alpha)$ , with basis  $\{1, \alpha\}$  over  $B$  and  $\alpha^2 = \nu \in B$ . The domain is  $D = \mathbb{F}_p$ , so  $n = |D| = p$ .

A degree-two residue-line datum consists of

$$E \in F[X], \quad \deg E = 2, \quad E(x) \neq 0 \text{ for all } x \in D,$$

and a numerator

$$B_{\text{num}} \in F[X], \quad \deg B_{\text{num}} < 2.$$

We write

$$A = F[X]/E, \quad [P]_E \in A, \quad b = [B_{\text{num}}]_E.$$

The anchor word on  $D$  is interpolated by a polynomial  $W$ . The operation  $u \wedge v$  means the determinant of the coordinate vectors of  $u, v \in A$  in a chosen  $F$ -basis of the two-dimensional  $F$ -space  $A$ . Off  $R_0$ , the pair  $\{[W]_E, b\}$  is such a basis.

The co-support has size  $j = 3$ . If  $T \subset D$  is the three-point co-support, define

$$\tau_1 = e_1(T), \quad \tau_2 = e_2(T), \quad \tau_3 = e_3(T).$$

All valid co-supports have  $\tau_i \in B$ . The quantity  $C_2$  denotes the number of distinct slopes produced by the relevant landing branch after restricting to valid split triples  $T \subset \mathbb{F}_p$ . Bounding landing points therefore bounds  $C_2$ , but a sharp slope theorem may require more because many landing points can share a slope.

## 12 Cycle 14 Normal Form

The Cycle 14 audit reduces the  $t = 2, j = 3$  landing equation to two affine forms in the elementary co-support parameters

$$\tau = (\tau_1, \tau_2, \tau_3).$$

**Lemma 3** (Affine wedge normal form). *Assume  $\{[W]_E, b\}$  is an  $F$ -basis of  $A$ . Then the landing equation can be written in the form*

$$\iota(\tau) = A_0(\tau_1, \tau_2) - \tau_3[W]_E, \quad \mu(\tau) = B_0(\tau_1, \tau_2) - \tau_3 b,$$

where

$$A_0 = p_1[W]_E + p_2 b, \quad B_0 = q_1[W]_E + q_2 b,$$

and  $p_i, q_i \in F$  are affine-linear functions of  $(\tau_1, \tau_2)$ . If the basis is normalized so  $[W]_E \wedge b = 1$ , then

$$\Delta = \iota \wedge \mu = (p_1 - \tau_3)(q_2 - \tau_3) - p_2 q_1.$$

*Proof.* The Cycle 14 quotient calculation, which is a concrete instance of Paper B's residue-line normal form, gives the displayed affine forms. Expanding in the ordered basis  $\{[W]_E, b\}$ , the coordinate vector of  $\iota$  is  $(p_1 - \tau_3, p_2)$ , while the coordinate vector of  $\mu$  is  $(q_1, q_2 - \tau_3)$ . Their wedge determinant is therefore

$$(p_1 - \tau_3)(q_2 - \tau_3) - p_2 q_1.$$

□

## 13 Cycle 18 Component Split

Choose a basis  $F = B + \alpha B$  with  $\alpha^2 = \nu \in B$ , and write

$$p_1 = p_{10} + \alpha p_{11}, \quad p_2 = p_{20} + \alpha p_{21}, \quad q_1 = q_{10} + \alpha q_{11}, \quad q_2 = q_{20} + \alpha q_{21}.$$

Split

$$\Delta = \Delta_0 + \alpha \Delta_1, \quad \Delta_0, \Delta_1 \in B[\tau_1, \tau_2, \tau_3].$$

**Theorem 2** (Monic quadratic and linear alpha component). *In the setting above,*

$$\begin{aligned} \Delta_0 &= \tau_3^2 - (p_{10} + q_{20})\tau_3 + p_{10}q_{20} + \nu p_{11}q_{21} - p_{20}q_{10} - \nu p_{21}q_{11}, \\ \Delta_1 &= -(p_{11} + q_{21})\tau_3 + p_{10}q_{21} + p_{11}q_{20} - p_{20}q_{11} - p_{21}q_{10}. \end{aligned}$$

Consequently  $\Delta_0$  is monic quadratic in  $\tau_3$ , while  $\Delta_1$  has degree at most one in  $\tau_3$ .



*Proof.* Substitute the coordinate expressions for  $p_i, q_i$  into the identity of Lemma 3 and use  $\alpha^2 = \nu$ . Collecting the coefficient of 1 gives the displayed formula for  $\Delta_0$ , and collecting the coefficient of  $\alpha$  gives the displayed formula for  $\Delta_1$ . The coefficient of  $\tau_3^2$  in  $\Delta_0$  is 1, and the  $\alpha$ -component has no  $\tau_3^2$ -term.  $\square$

**Corollary 1** (Graph reduction outside the zero-alpha branch). *If  $\Delta_1$  is not the zero polynomial, write*

$$\Delta_1 = s(\tau_1, \tau_2)\tau_3 + h(\tau_1, \tau_2).$$

*On the open locus  $s \neq 0$ , the common-zero condition  $\Delta_0 = \Delta_1 = 0$  is supported on the graph*

$$\tau_3 = -h/s.$$

*Proof.* This is immediate from solving the linear equation  $\Delta_1 = 0$  for  $\tau_3$  on  $s \neq 0$ .  $\square$

## 14 New Divisibility-Gate Theorem

The next theorem is the main NEW-LOCAL theorem extracted from the Cycle 18 reduction. It is stronger than merely saying that the branch is a graph: it shows that any two-dimensional family must satisfy an explicit divisibility identity.

**Theorem 3** (Divisibility gate for the graph branch). *Assume the restricted setting above and assume  $\Delta_1 \neq 0$ . Write*

$$\Delta_1 = s(\tau_1, \tau_2)\tau_3 + h(\tau_1, \tau_2),$$

*and define the denominator-cleared graph polynomial*

$$G(\tau_1, \tau_2) = s(\tau_1, \tau_2)^2 \Delta_0(\tau_1, \tau_2, -h(\tau_1, \tau_2)/s(\tau_1, \tau_2)).$$

*Then  $\deg G \leq 4$ . If  $G$  is not the zero polynomial, then*

$$\#\{\tau \in B^3 : \Delta_0(\tau) = \Delta_1(\tau) = 0\} = O(p).$$

*Consequently the corresponding number of distinct slopes satisfies*

$$C_2 = O(p) = O(n).$$

*Therefore a growing-prime family with*

$$C_2 = \Theta(p^2) = \Theta(q_{\text{line}})$$

*in the nonzero- $\Delta_1$  branch can occur only if  $G \equiv 0$ .*

*Proof.* Because the  $p_i, q_i$  are affine-linear in  $(\tau_1, \tau_2)$ , the coefficient  $s$  has degree at most 1, and  $h$  has degree at most 2. Since  $\Delta_0$  is monic quadratic in  $\tau_3$ , clearing the denominators after substituting  $\tau_3 = -h/s$  gives a polynomial  $G$  of degree at most 4.

On  $s \neq 0$ , the equation  $\Delta_1 = 0$  forces  $\tau_3 = -h/s$ . Hence every common zero on this open locus gives a base pair  $(\tau_1, \tau_2)$  with  $G(\tau_1, \tau_2) = 0$ . If  $G$  is nonzero, Schwartz–Zippel gives at most  $4p$  such base pairs, and each base pair gives at most one value of  $\tau_3$ .

It remains to count the exceptional locus  $s = 0$ . There  $\Delta_1 = 0$  also forces  $h = 0$ . If  $s$  is a nonzero affine form, this is contained in an affine line; if  $s \equiv 0$ , then  $h$  is a nonzero polynomial of degree at most 2. In either case there are  $O(p)$  base pairs. For each such pair,  $\Delta_0$  is a monic quadratic in  $\tau_3$ , so it gives at most two values of  $\tau_3$ . The total common-zero count is  $O(p)$ .

Finally, the number of distinct slopes is bounded by the number of landing points  $\tau$ . Thus  $C_2 = O(p)$  whenever  $G \not\equiv 0$ , and any  $\Theta(p^2)$  family in this branch must satisfy  $G \equiv 0$ .  $\square$

**Remark 4.** *The identity  $G \equiv 0$  says that, after clearing the  $s$ -denominator,  $\Delta_0$  is divisible by the linear equation  $\Delta_1$  on the graph branch. This is the first exact algebraic gate that a large slope family in this restricted window must pass.*

## 15 Nearby Proven Gates

The same restricted lane contains two other useful gates. They are included here because they explain where Theorem 3 fits.

**Proposition 4** (Base-component complete-intersection gate). *Suppose  $\Delta_0, \Delta_1 \in B[\tau_1, \tau_2, \tau_3]$  have no common surface component. Then*

$$\#V(\Delta_0, \Delta_1)(B) = O(p),$$

*and hence this branch contributes  $C_2 = O(p)$  slopes.*

*Proof.* The two components have bounded degree. If they share no surface component, their common zero set in  $\mathbb{A}_B^3$  is a curve of bounded degree. A bounded-degree affine curve over  $\mathbb{F}_p$  has  $O(p)$  rational points, and slopes are bounded by landing points.  $\square$

**Proposition 5** (Cycle 16 determinant gate). *Let  $Q(z_0, z_1)$  be the Cycle 15–16 determinant consistency polynomial for the affine system*

$$L_z(\tau) = \iota(\tau) - z\mu(\tau) = 0, \quad z = z_0 + \alpha z_1.$$

*If  $Q$  is not identically zero, then in the restricted  $D = \mathbb{F}_p$ ,  $t = \sigma = 2$ ,  $j = 3$ , off- $R_0$  window,*

$$C_2 \leq 4p = O(p) = O(n).$$

*Proof.* Cycle 16 gives  $\deg Q \leq 4$  and shows that every landing slope must satisfy  $Q(z_0, z_1) = 0$ . If  $Q \neq 0$ , Schwartz–Zippel over  $B = \mathbb{F}_p$  gives at most  $4p$  possible pairs  $(z_0, z_1)$ , hence at most  $4p$  slopes.  $\square$

## 16 Slope Map and Remaining Heuristics

On the graph branch, the landing equation  $\iota = z\mu$  gives

$$p_1 - \tau_3 = zq_1, \quad p_2 = z(q_2 - \tau_3).$$

Where  $q_1 \neq 0$ , substituting  $\tau_3 = -h/s$  yields

$$z(\tau_1, \tau_2) = \frac{p_1 + h/s}{q_1}.$$

Equivalently, eliminating  $\tau_3$  recovers the Cycle 14 quadratic

$$q_1 z^2 - (p_1 - q_2)z - p_2 = 0,$$

so the residual map is controlled by

$$(\tau_1, \tau_2) \mapsto [q_1 : (p_1 - q_2) : p_2] \in \mathbb{P}^2(F).$$

**Heuristic 1** (Exceptional identity should be rare). *For generic source data, the polynomial  $G$  should not vanish identically. Thus the graph branch should usually be curve-sized, and the first possible large-slope families should come from algebraically special data satisfying the divisibility identity  $G \equiv 0$  or from the zero-alpha branch  $\Delta_1 \equiv 0$ .*

**Heuristic 2** (What remains after the gate). *If  $G \equiv 0$ , the problem is no longer to show that the common-zero set is small. It may be surface-sized. The remaining question is whether the projective coefficient map has one-dimensional image on every source-valid surface, giving  $C_2 = O(p)$ , or whether some growing-prime source-valid family has two-dimensional image and  $C_2 = \Theta(p^2)$ .*

## 17 Next Experimental Checks

A useful scanner should compute, for each source-valid candidate,

$$\Delta_1 = s\tau_3 + h, \quad G = s^2\Delta_0(\tau_1, \tau_2, -h/s),$$

then record whether  $G \equiv 0$ , whether  $\Delta_1 \equiv 0$ , the exceptional locus  $s = h = 0$ , the image size of

$$[q_1 : (p_1 - q_2) : p_2],$$

and the direct slope count over distinct split triples  $T \subset \mathbb{F}_p$ .

The certificate should keep at least

$$p, q_{\text{gen}}, q_{\text{line}}, E, B_{\text{num}}, \text{seed}, \text{stratum}, R_0\text{-status}, \\ \Delta_1 \equiv 0, \quad G \equiv 0, \quad \#\{s = h = 0\}, \quad \#\text{image}, \quad C_2, \quad \#\{T \text{ split distinct}\}.$$

## 18 June 26 Normal-Form and Dependency Results

**Status: mixed.** This section records the parts of the June 26 PR batch that survived a statement-level validity check. The statements below are useful proof infrastructure, not prize claims. In particular, they do not improve the current Cycle116/119 numerator 52,747,567,092 over  $\mathbb{F}_{1732}$ , and they do not prove the corrected Paper B MCA conjecture.

### 18.1 Syndrome-pencil normal form for support-wise line incidence

**Status: New-local.** This is the cleanest reusable F1 contribution from the batch. It turns support-wise line incidence into a Hankel-pencil locator equation over the actual line field.

Let  $D = \{x_1, \dots, x_n\} \subset F$  have distinct points and let  $C = \text{RS}[F, D, k]$  with redundancy  $r = n - k$ . For each  $x_i$ , put

$$\lambda_i = \left( \prod_{h \neq i} (x_i - x_h) \right)^{-1}.$$

For a word  $y : D \rightarrow F$ , define

$$\text{Syn}(y)_m = \sum_i \lambda_i x_i^m y(x_i), \quad 0 \leq m < r.$$

Then  $\text{Syn}(y) = 0$  if and only if  $y \in C$ . Fix  $T \subset D$  with  $|T| = j$ , assume  $0 \leq j \leq r$ , and set  $t = r - j$ . Write

$$L_T(X) = \prod_{x \in T} (X - x) = \ell_0 + \ell_1 X + \dots + \ell_j X^j, \quad \ell_T = (\ell_0, \dots, \ell_j)^T.$$

For  $w \in F^r$ , define the  $t \times (j+1)$  Hankel window

$$H_{t,j}(w)_{a,b} = w_{a+b}, \quad 0 \leq a < t, \quad 0 \leq b \leq j.$$

**Theorem 4** (Hankel-pencil test). *Let  $f, g : D \rightarrow F$ , put  $u = \text{Syn}(f)$  and  $v = \text{Syn}(g)$ , and let  $S = D \setminus T$ . For a finite slope  $z \in F$ , the word  $f + zg$  is explained by a codeword of  $C$  on  $S$  if and only if*

$$(H_{t,j}(u) + zH_{t,j}(v))\ell_T = 0.$$

*Moreover, this explanation is support-wise noncontained for the line  $f + zg$  on  $S$  if and only if, in addition,*

$$H_{t,j}(v)\ell_T \neq 0.$$

*Consequently the support-wise MCA bad slopes at agreement  $k + t$  are exactly the slopes for which there is a squarefree  $D$ -split locator  $L_T$  of degree  $j = r - t$  satisfying the displayed incidence and noncontainment tests.*

*Proof.* Let  $W_T \subset F^r$  be the span of the parity-check columns indexed by  $T$ :

$$W_T = \text{span}_F\{\lambda_x(1, x, \dots, x^{r-1}) : x \in T\}.$$

Equivalently,  $w \in W_T$  means that  $w$  is the syndrome of a word supported on  $T$ . For every  $w \in W_T$ , write

$$w_m = \sum_{x \in T} c_x x^m.$$

Then, for  $0 \leq a < t$ ,

$$(H_{t,j}(w)\ell_T)_a = \sum_{b=0}^j \ell_b w_{a+b} = \sum_{x \in T} c_x x^a L_T(x) = 0.$$

Conversely, since  $L_T$  is monic, the recurrence  $\sum_{b=0}^j \ell_b w_{a+b} = 0$  for  $0 \leq a < r - j$  determines all remaining coordinates from the first  $j$  coordinates, so its solution space has dimension at most  $j$ . The  $j$  columns indexed by the distinct points of  $T$  form a Vandermonde system of rank  $j$ , hence the recurrence space is exactly  $W_T$ .

The word  $f + zg$  is explained on  $S$  if and only if there is  $c \in C$  such that  $f + zg - c$  is supported on  $T$ . Taking syndromes, this is equivalent to  $u + zv \in W_T$ , which is the first displayed equation by the recurrence criterion. Under that equation, simultaneous explanation of  $f$  and  $g$  on the same  $S$  is equivalent to  $u, v \in W_T$ . Since  $u = (u + zv) - zv$ , this is equivalent to  $v \in W_T$ , hence to  $H_{t,j}(v)\ell_T = 0$ . Noncontainment is therefore exactly the negation of that condition.  $\square$

## 18.2 Interleaved codegree reduction

**Status: New-local reduction; saving remains conditional.** This result is useful for the L2 interleaved-list ledger because it replaces the naive Cartesian product by a recursion through higher-agreement tails.

Let  $C = \text{RS}[F, D, k]$ . For a word  $V : D \rightarrow F$  and codeword  $c \in C$ , write

$$A_V(c) = \{x \in D : c(x) = V(x)\}, \quad \text{Fib}_V(s) = \{c \in C : |A_V(c)| \geq s\},$$

and  $M_V(s) = |\text{Fib}_V(s)|$ . For a  $\mu$ -row word  $U = (U_1, \dots, U_\mu)$ , define

$$\Lambda_\mu^{(s)}(U) = \#\{(c_1, \dots, c_\mu) \in C^\mu : |\cap_i A_{U_i}(c_i)| \geq s\}.$$

**Theorem 5** (two-regime codegree bound). *For every agreement threshold  $a$ ,*

$$\Lambda_2^{(a)}(U_1, U_2) \leq M_{U_2}(a) + M_{U_2}(2a - k) M_{U_1}(a).$$

*More generally,*

$$\Lambda_\mu^{(a)}(U_1, \dots, U_\mu) \leq \Lambda_{\mu-1}^{(a)}(U_2, \dots, U_\mu) + \Lambda_{\mu-1}^{(2a-k)}(U_2, \dots, U_\mu) M_{U_1}(a).$$

*Proof.* For the two-row case, sum over  $c_2 \in \text{Fib}_{U_2}(a)$ , and put  $N_2 = |A_{U_2}(c_2)|$ . The possible  $c_1$ 's are codewords agreeing with  $U_1$  on at least  $a$  points of  $A_{U_2}(c_2)$ . If  $N_2 < 2a - k$ , then two such  $c_1$ 's would agree with each other on more than  $k$  points, forcing equality. Thus this punctured list has size at most one. If  $N_2 \geq 2a - k$ , use the trivial bound  $M_{U_1}(a)$ . Summing these two regimes gives the displayed two-row inequality.

For the  $\mu$ -row case, fix  $(c_2, \dots, c_\mu)$  and put

$$S = \bigcap_{i=2}^{\mu} A_{U_i}(c_i).$$

The listed  $c_1$ 's must agree with  $U_1$  on at least  $a$  points of  $S$ . If  $|S| < 2a - k$ , the same unique-decoding argument gives at most one such  $c_1$ . If  $|S| \geq 2a - k$ , use the bound  $M_{U_1}(a)$ . Counting the two classes of  $(c_2, \dots, c_\mu)$  gives the recursion.  $\square$

**Remark 5.** *The reduction is unconditional. The desired L2 saving still requires an L1-family input saying that the tail lists at agreement  $2a - k$  are small after quotient-periodic contributions are budgeted.*

### 18.3 Deep-point cap dependency split

**Status: Wrapper / Audit.** This result does not edit Paper D, but it explains why the headline MCA cap can be routed locally without using the imported Crites–Stewart conversion.

**Proposition 6** (CS25-free route to the Paper D MCA cap). *Assume the finite-field, coset, divisibility, and binomial hypotheses used in Paper D's universal cap theorem. Let  $C = \text{RS}[F, D, k]$ ,  $C^+ = \text{RS}[F, D, k+1]$ ,  $q = |F|$ , and suppose the elementary quotient-locator fiber construction gives a  $C^+$ -list of size*

$$L \geq \binom{N}{\rho N + 2} / |B| \geq q/k + 1$$

*at the endpoint  $\delta_N = 1 - \rho - 2/N$ . Then there is a support-wise MCA-bad line for  $C$  with bad-slope density strictly larger than*

$$\frac{1}{2k} \left(1 - \frac{n}{q}\right).$$

*By support-wise MCA monotonicity, the same lower bound holds for all  $\delta_N \leq \delta < 1 - \rho$ .*

*Proof.* Let  $U : D \rightarrow F$  be the received word with a  $C^+$ -list  $\{P_1, \dots, P_L\}$  at agreement  $m = k + 2n/N$ . For  $\alpha \in F \setminus D$ , form

$$f_\alpha(x) = \frac{U(x)}{x - \alpha}, \quad g_\alpha(x) = -\frac{1}{x - \alpha}.$$

For every listed  $P_i$ , the slope  $z = P_i(\alpha)$  gives the codeword

$$Q_i(X) = \frac{P_i(X) - P_i(\alpha)}{X - \alpha} \in F[X]_{<k}$$

on the same agreement support. Conversely every such slope comes from a listed  $P_i$ . Since  $m > k$ , the direction  $g_\alpha$  cannot be explained by a degree- $< k$  polynomial on the same support, so these slopes are support-wise MCA-bad.

Averaging over  $\Omega = F \setminus D$ , two distinct degree- $< k + 1$  polynomials collide at at most  $k$  points of  $\Omega$ . Therefore some  $\alpha$  has at least

$$M \geq \frac{L}{1 + k(L-1)/(q-n)}$$

distinct values among  $P_1(\alpha), \dots, P_L(\alpha)$ . Its bad-slope density is at least

$$\frac{L(q-n)}{q(q-n+k(L-1))}.$$

This is strictly greater than  $(q-n)/(2kq)$  exactly when

$$2kL > q - n + k(L-1).$$

The lower bound  $L \geq q/k + 1$  gives  $kL \geq q + k$ , so the left-minus-right quantity is at least  $n + 2k > 0$ . This proves the endpoint bound, and monotonicity in  $\delta$  extends it to the stated interval.  $\square$

## 18.4 Common code-line residual budget

**Status: New-local finite MDS theorem.** This M2 lemma is a guardrail for line-decoding imports: a close common code-line exception still pays for residual coordinates outside the common support.

**Proposition 7** (common code-line residual budget). *Let  $C \subseteq F^D$  be an MDS linear code of dimension  $k$ , so every nonzero codeword has fewer than  $k$  zeros on  $D$ . Let  $\ell_z = f + zg$  be a received line and suppose that for some codewords  $c_f, c_g \in C$  there is a common support  $S_0 \subseteq D$  of size  $b$  such that  $f = c_f$  and  $g = c_g$  on  $S_0$ . Fix an agreement threshold  $a$  with  $a + b - n \geq k$ . Put  $\Omega = D \setminus S_0$ ,*

$$f' = f - c_f, \quad g' = g - c_g, \quad h = \max(1, a - b),$$

and

$$c_0 = \#\{x \in \Omega : f'(x) = g'(x) = 0\}.$$

Every support-wise noncontained slope at agreement  $a$  satisfies

$$\#\{x \in \Omega : f'(x) + zg'(x) = 0\} \geq h.$$

Consequently, if  $h > c_0$ , then the number of such slopes is at most

$$\left\lfloor \frac{|\Omega| - c_0}{h - c_0} \right\rfloor.$$

*Proof.* If  $\ell_z$  is explained on a support  $S$  of size at least  $a$ , then

$$|S \cap S_0| \geq a - |\Omega| = a + b - n \geq k.$$

On  $S \cap S_0$ , the explaining codeword agrees with  $c_f + zc_g$ . By the MDS zero-rigidity condition, the two codewords are equal. Thus, outside  $S_0$ , the line can only be explained where  $f' + zg' = 0$ , and at least  $h = \max(1, a - b)$  such outside zeros are needed for a noncontained support.

Each non-common residual coordinate  $x \in \Omega$  with  $(f'(x), g'(x)) \neq (0, 0)$  can vanish for at most one finite slope  $z$ . A bad slope needs at least  $h - c_0$  private non-common residual zeros after the  $c_0$  common zeros are counted. Charging these private coordinates to slopes gives the stated packing bound.  $\square$

## 19 Recent PR Triage: L1, F1, M1, and Challenge-Map Ledgers

**Status: Audit / Experimental / bounded Proved notes.** The open PR batch of June 27, 2026, adds useful proof infrastructure but no new unconditional prize-scale threshold beyond the tangent-floor consequences already in the high-agreement ledger. The promotion rule used here is conservative: bounded toy theorems and exact normal forms may enter the experimental notes, while computation-dependent numerators and generated proof programs remain labelled as such.

### 19.1 L1: proper-subgroup and monomial-prefix lanes

Scott Hughes'  $d = 3$  proper-subgroup note records the actual  $H^{2k}$  collision object for the odd moments 1, 3, 5. For  $p > 5$ ,  $H \leq \mathbb{F}_p^*$  of order  $n$ , and

$$S_H(a_1, a_3, a_5) = \sum_{h \in H} \psi(a_1 h + a_3 h^3 + a_5 h^5),$$

the exact Fourier identity is

$$V_{H,k}^{(3)} = p^{-3} \sum_{a_1, a_3, a_5 \in \mathbb{F}_p} |S_H(a_1, a_3, a_5)|^{2k}.$$

After splitting frequencies into the linear, cubic, and quintic strata, and using the stated one-variable Gauss/Katz inputs, the note gives

$$0 \leq V_{H,k}^{(3)} - \frac{n^{2k}}{p^3} \leq \frac{p-1}{p^3} p^k + \frac{p(p-1)}{p^3} (3\sqrt{p})^{2k} + \frac{p^2(p-1)}{p^3} (5\sqrt{p})^{2k}.$$

This is useful L1 infrastructure, not a reserve-scale arbitrary-word locator local limit.

Vadim Avdeev's monomial-prefix note settles one bounded toy lane over

$$F = \mathbb{F}_{17}[z]/(z^{32} - 3), \quad H = \langle z \rangle, \quad |H| = 512,$$

with degree convention  $\deg P \leq 256$ . In the range  $258 \leq A \leq 512$ , the admissible monomial agreement sizes are

$$258, 259, 260, 262, 264, 268, 272, 280, \\ 288, 304, 320, 352, 384, 512.$$

The proof combines a local length-16 imbalance check over  $\mathbb{F}_{17}$ , a dyadic power-sum descent gate, and quotient-complement survivor families. This should be cited only as a monomial-prefix benchmark, not as an arbitrary-word list theorem.

### 19.2 F1: arbitrary-anchor residue clouds

The F1 arbitrary-anchor update splits the balanced residue-cloud problem from the residual-list problem. If  $t < \sigma$ , residue-line bad slopes inject into an ordinary list for  $\text{RS}[F, D, k+t]$  at residual slack  $\sigma - t$ . In the balanced case  $t = \sigma$ , arbitrary anchors have a real free-packing floor:

$$\Lambda_{\text{NC}} \geq \left\lfloor \frac{|D| - k}{\sigma} \right\rfloor.$$

Moreover this is the exact ceiling for ordered  $k$ -degenerate support families, so any larger arbitrary-anchor packet must use dense overlaps. The dense-overlap gate is explicit: for two supports  $S, T$  with overlap  $J$ , compatibility of distinct slopes is controlled by whether the residue direction  $N/E$  is degree- $< k$  on  $J$ . This is a useful F1 warning and search normal form, not an extension-line lift theorem.

### 19.3 M1: all-line Hankel aperiodic packet

The large M1 PR is not promoted wholesale. The useful distillation is the exact Hankel-pencil object. For a line  $(f, g)$ , syndromes  $u = \text{Syn}(f)$  and  $v = \text{Syn}(g)$ , and a split complement  $T$  with locator vector  $\ell_T$ , finite bad slopes are governed by

$$(H_{t,j}(u) + zH_{t,j}(v))\ell_T = 0, \quad H_{t,j}(v)\ell_T \neq 0.$$

In the  $t = 2$  window this becomes

$$b_T \neq 0, \quad \det[a_T \ b_T] = 0, \quad a_T = H_{2,j}(u)\ell_T, \quad b_T = H_{2,j}(v)\ell_T.$$

Same-slope one-exchange collisions lie in fixed-slope root slices, and the residual one-exchange graph can be further organized into quadratic companion slices and top packets. These reductions define a promising M1 proof program, but they do not yet prove the desired uniform polynomial bound on residual aperiodic slope images.

### 19.4 M1: same-slope one-exchange root slices

The first local reduction in the all-line Hankel packet is elementary and proved in `experimental/notes/m1/m1_same_slope_root_slice_lemma.md`. Let

$$L_z = H_{t,j}(u) + zH_{t,j}(v)$$

be the finite-slope landing map. Fix a  $(j-1)$ -core  $R$ , and write

$$T_y = R \cup \{y\}, \quad \ell_{T_y} = (X - y)\ell_R.$$

If two distinct one-exchange extensions  $T_{y_1}$  and  $T_{y_2}$  satisfy

$$L_z \ell_{T_{y_1}} = L_z \ell_{T_{y_2}} = 0,$$

then subtracting gives

$$(y_2 - y_1)L_z \ell_R = 0.$$

Thus  $L_z \ell_R = 0$ , and substituting into either endpoint equation gives

$$L_z(X\ell_R) = 0.$$

Consequently

$$L_z \ell_{T_y} = 0 \quad \text{for every } y.$$

So every same-slope one-exchange edge lies in the full fixed-slope root slice with core  $R$ . After fixed-slope root slices have been charged, the residual one-exchange graph has only different-slope edges. This is a local reduction, not a bound for the remaining different-slope codegree. The verifier `experimental/scripts/verify_m1_same_slope_root_slice_lemma.py` checks the subtraction identity and its row-wise linear-map consequence in small fields.

The same two padded equations also lift the root slice to a higher-slack Hankel core. With  $w_z = u + zv$ , the equations

$$H_{t,j}(w_z)(\ell_R, 0)^T = 0, \quad H_{t,j}(w_z)(0, \ell_R)^T = 0$$

are exactly the first  $t$  and last  $t$  rows of

$$H_{t+1,j-1}(w_z)\ell_R = 0.$$



Thus a same-slope root slice is not a fresh  $t$ -level residual multiplicity: it belongs to the lifted  $(t+1, j-1)$  Hankel-core ledger at the same finite slope.

The same mechanism has a two-exchange full-plane form. For a  $(j-2)$ -core  $R$ , write a two-root locator in elementary coordinates

$$\ell_{T_{s,p}} = (X^2 - sX + p)\ell_R, \quad s = x + y, \quad p = xy.$$

For fixed finite slope  $z$ ,

$$L_z \ell_{T_{s,p}} = L_z(X^2 \ell_R) - sL_z(X \ell_R) + pL_z(\ell_R)$$

is affine-linear in  $(s, p)$ . Hence three affinely non-collinear same-slope points in the two-root plane force

$$L_z \ell_R = L_z(X \ell_R) = L_z(X^2 \ell_R) = 0,$$

equivalently

$$H_{t+2, j-2}(u + zv)\ell_R = 0.$$

Thus full two-exchange planes are charged to the lifted  $(t+2, j-2)$  Hankel ledger. After fixed-root lines and full planes are removed, any residual same-slope two-exchange component through  $R$  is a line packet in the  $(s, p)$ -plane.

The full-plane statement is the  $h=2$  case of a general elementary-packet lift. For a  $(j-h)$ -core  $R$ , write

$$\ell_{R,c} = (X^h + c_{h-1}X^{h-1} + \cdots + c_0)\ell_R, \quad c \in F^h.$$

For fixed finite slope  $z$ , the map  $c \mapsto L_z \ell_{R,c}$  is affine-linear. Hence if  $h+1$  affinely independent coefficient points are killed at the same slope, then

$$L_z(X^m \ell_R) = 0, \quad 0 \leq m \leq h.$$

These  $h+1$  padded equations are exactly the row blocks of

$$H_{t+h, j-h}(u + zv)\ell_R = 0.$$

Thus every full affine-rank same-slope  $h$ -exchange packet is charged losslessly to the lifted  $(t+h, j-h)$  Hankel-core ledger. Applying the same argument separately to  $u$  and  $v$  gives the simultaneous-kernel version: full affine-rank packets in  $K_{r,d}(u, v)$  lift to  $K_{r+h, d-h}(u, v)$ . The lower-rank case is also exact. If  $C \subset F^h$  is any killed coefficient packet and  $\text{aff}(C) = c_* + W$ , then

$$L_z(X^h \ell_R) + \sum_m c_{*,m} L_z(X^m \ell_R) = 0, \quad \sum_m w_m L_z(X^m \ell_R) = 0 \quad (w \in W).$$

Conversely these equations kill the whole affine subpacket  $c_* + W$ . Thus after full-rank packets are charged, the residual same-slope elementary packets are precisely lower-rank affine subpackets; for  $h=2$  these are the lines classified next. Among coefficient hyperplanes, the fixed-root ones are exactly the evaluation hyperplanes. Namely

$$b + \sum_{m=0}^{h-1} a_m c_m = 0$$

is the locus  $P_c(\alpha) = 0$  for a fixed finite root  $\alpha$  iff, up to a nonzero scalar,  $(a_0, \dots, a_{h-1}, b) = (1, \alpha, \dots, \alpha^h)$ . Such hyperplanes are charged to the  $(h-1)$ -exchange root-slice ledger. For  $h=2$ ,

this is  $p - \alpha s + \alpha^2 = 0$ , equivalently  $(x - \alpha)(y - \alpha) = 0$ . More generally, every coefficient hyperplane has a one-root fiber dichotomy. Fix a monic  $(h - 1)$ -root factor

$$Q_d(X) = X^{h-1} + d_{h-2}X^{h-2} + \cdots + d_0$$

and write  $d_{h-1} = 1, d_{-1} = 0$ . The extension

$$P_y = (X - y)Q_d$$

has coefficients  $c_m(y) = d_{m-1} - yd_m$ , so the hyperplane equation restricts to

$$b + \sum_m a_m c_m(y) = \left(b + \sum_m a_m d_{m-1}\right) - y \sum_m a_m d_m.$$

Thus a fixed  $(h - 1)$ -core has at most one extension in the hyperplane unless the whole affine one-root line lies there. In the killed same-slope packet setting, such a full line is exactly a one-exchange root slice and lifts to the  $(t + 1, j - 1)$  Hankel core. After these full one-root fibers are charged, a codimension-one coefficient-hyperplane packet has no residual one-exchange edges. The same 0/1/all dichotomy holds for any affine rank-defect packet  $A = c_* + W \subset F^h$ . For the same one-root fiber, put

$$d_{\text{vec}} = (d_0, \dots, d_{h-1}).$$

If two distinct roots  $y_1, y_2$  have  $c(y_i) \in A$ , then

$$c(y_2) - c(y_1) = -(y_2 - y_1)d_{\text{vec}} \in W,$$

so  $d_{\text{vec}} \in W$  and the whole affine line  $c(y)$  lies in  $A$ . Thus, after full one-root fibers are root-slice charged, no residual affine rank-defect packet has one-exchange edges. The two-root fiber has the same affine form. For a fixed monic  $(h - 2)$ -root factor  $Q_e$ , write

$$P_{s,p} = (X^2 - sX + p)Q_e, \quad c_m(s, p) = e_{m-2} - se_{m-1} + pe_m.$$

If an affine rank-defect packet contains three non-collinear points of this  $(s, p)$ -plane, then both plane directions lie in  $W$ , so the whole two-root plane is contained in the packet and is charged to the  $(t + 2, j - 2)$  full-plane lift. After full planes are charged, every residual two-root fiber intersection is at most a line packet. In general, for a moving monic  $r$ -root factor  $B_a$  multiplying a fixed  $(h - r)$ -root factor  $Q$ , the map  $a \mapsto B_a Q$  is an affine embedding of  $F^r$  into coefficient space. The preimage of  $A = c_* + W$  is therefore an affine subspace of  $F^r$ : if it has full affine rank  $r$ , the whole moving  $r$ -root fiber is contained in  $A$  and is charged to the  $(t + r, j - r)$  full-packet lift; otherwise the residual intersection has dimension at most  $r - 1$ . Over a finite field of size  $Q_F$ , this gives the formal residual count

$$\#\{a \in F^r : \text{coeff}(B_a Q) \in A\} \leq Q_F^{r-1}$$

on every fixed  $r$ -root fiber after the full-fiber charge; split, domain, quotient, tangent, and non-containment filters only shrink this count. Consequently the residual  $r$ -exchange graph inside one affine packet has maximum degree at most

$$\binom{h}{r} (Q_F^{r-1} - 1),$$

because each vertex has only  $\binom{h}{r}$  fixed  $(h - r)$ -subfactors to complete.

The residual two-root lines are exactly the expected models: if

$$As + Bp + C = 0$$

and  $B = 0$ , then  $x + y = s_0$  is a fixed-sum packet; if  $B \neq 0$ , then after writing  $c = -A/B$ ,  $\beta = -C/B$ , the equation becomes

$$(x - c)(y - c) = c^2 + \beta.$$

The zero right-hand side is a fixed-root line, and the nonzero right-hand side is a product-Mobius packet. Hence after fixed-root lines are charged, the only residual line packets are fixed-sum and nondegenerate product-Mobius packets.

In the actual  $t = 2$  Hankel window, these non-fixed line packets cannot be constant-slope residuals. If

$$H_{2,j}(u + zv)\ell_{R,s,p} = (d_2 - sd_1 + pd_0, d_3 - sd_2 + pd_1)$$

vanishes identically on a fixed-sum line  $s = s_0$ , then the coefficient of  $p$  gives  $(d_0, d_1) = 0$ , and the Hankel overlap forces  $d_2 = d_3 = 0$ . If it vanishes identically on a nondegenerate product-Mobius line  $p = cs - c^2 + \mu$ ,  $\mu \neq 0$ , then  $(d_1, d_2) = c(d_0, d_1)$  and  $(d_2, d_3) = (c^2 - \mu)(d_0, d_1)$ , so  $\mu d_0 = 0$  and again all four  $d_i$  vanish. Thus every constant-slope non-fixed line packet is already charged to the full-plane lift  $H_{4,j-2}(u + zv)\ell_R = 0$ . After fixed-root and full-plane charges, the active finite-slope map on each surviving fixed-sum or product-Mobius packet is injective.

In the  $t = 2$  Hankel window the same one-exchange core has a quadratic determinant gate. Put

$$a_y = H(u)\ell_{T_y} = a_X - ya_0, \quad b_y = H(v)\ell_{T_y} = b_X - yb_0 \quad \text{in } F^2.$$

Then the finite-slope determinant condition is

$$\Delta_R(y) = \det(a_y, b_y) = 0, \quad b_y \neq 0,$$

where

$$\Delta_R(y) = \det(a_X, b_X) - y(\det(a_0, b_X) + \det(a_X, b_0)) + y^2 \det(a_0, b_0).$$

Thus a nonzero  $\Delta_R$  admits at most two anchors  $y$ . If three distinct anchors pass the determinant gate, then  $\Delta_R \equiv 0$ , so the core is a ruled determinant branch. The ruled branch first has the abstract affine dichotomy: if

$$\det(a_X - ya_0, b_X - yb_0) \equiv 0,$$

then either there is a fixed finite slope  $z_0$  with

$$a_y + z_0 b_y = 0 \quad \text{for every } y,$$

or  $b_y = 0$  for every  $y$ , or all four vectors

$$a_X, a_0, b_X, b_0$$

lie in a single output line. In the last case write

$$a_y = \alpha(y)c, \quad b_y = \beta(y)c.$$

For arbitrary affine pencils this last case could be a moving-slope branch, but the Hankel shift structure rules it out here. If

$$c_i(w) = \text{row}_i(H_{3,j-1}(w)\ell_R), \quad i = 0, 1, 2,$$

then

$$H_{2,j}(w)\ell_{T_y} = (c_1(w) - yc_0(w), c_2(w) - yc_1(w)).$$

A common output line imposes two recurrence equations on  $(c_0(w), c_1(w), c_2(w))$ , whose solution space is one-dimensional. Applying this to  $w = u$  and  $w = v$  makes the two Hankel pencils proportional, unless  $b_y = 0$  identically. Therefore every active ruled Hankel core is already a fixed finite-slope core; after fixed-slope root slices are charged, ruled cores contribute no residual one-exchange edges.

The same note also records the exact one-exchange triangle dichotomy in the Johnson graph. Three  $j$ -set locators that are pairwise one-exchange are either a star

$$T_i = R \cup \{y_i\}, \quad |R| = j - 1,$$

or a top-packet triangle

$$T_i = U \setminus \{x_i\}, \quad |U| = j + 1.$$

Thus every active star triangle is precisely a three-anchor event through one  $(j - 1)$ -core, hence a ruled determinant core by the quadratic gate above. Since ruled Hankel cores collapse to fixed-slope or inactive cores, residual one-exchange triangles after fixed-slope root-slice charging are therefore top-packet triangles.

Top packets then lift losslessly to a  $t = 1$  Hankel kernel. If

$$T_x = U \setminus \{x\}, \quad \ell_U = (X - x)\ell_{T_x},$$

then for every syndrome vector  $w$ ,

$$H_{1,j+1}(w)\ell_U = \text{row}_1(H_{2,j}(w)\ell_{T_x}) - x \text{row}_0(H_{2,j}(w)\ell_{T_x}).$$

Thus a  $t = 2$  incidence for  $T_x$  at slope  $z$  implies

$$(H_{1,j+1}(u) + zH_{1,j+1}(v))\ell_U = 0.$$

If two members of the same top packet have distinct slopes, subtracting the two lifted scalar equations gives

$$H_{1,j+1}(u)\ell_U = H_{1,j+1}(v)\ell_U = 0.$$

Therefore every residual top-packet edge after fixed-slope charging lies over a common lifted  $t = 1$  kernel. This turns the top-packet triangle residual into a higher-slack kernel target rather than a fresh one-exchange multiplicity phenomenon.

Writing

$$K_{\text{top}}(u, v) = \{U \subset D : |U| = j + 1, H_{1,j+1}(u)\ell_U = H_{1,j+1}(v)\ell_U = 0\},$$

the top-packet branch has the explicit compression ledger

$$E_{\text{top}}(A_{\text{res}}) \leq \binom{j+1}{2} |K_{\text{top}}(u, v)|, \quad \text{Tri}_1(A_{\text{res}}) \leq \binom{j+1}{3} |K_{\text{top}}(u, v)|.$$

Indeed a residual top edge recovers its union  $U$ , and (because residual one-exchange edges have distinct slopes) the lift above forces  $U \in K_{\text{top}}(u, v)$ . Inside such a packet,

$$H_{2,j}(w)\ell_{U \setminus \{x\}} = \rho_x(w)(1, x), \quad w \in \{u, v\},$$

so the remaining slope label is the scalar ratio  $\rho_x(u) + z\rho_x(v) = 0$ ,  $\rho_x(v) \neq 0$ . Thus top-packet edges and triangles are charged to the lifted  $t = 1$  top-kernel family.

The top-kernel family also has an exact root-slice recursion. More generally, for  $r \geq 1$  and locator size  $d$ , put

$$K_{r,d}(u, v) = \{ U \subset D : |U| = d, H_{r,d}(u)\ell_U = H_{r,d}(v)\ell_U = 0 \}.$$

If  $U_{y_i} = R \cup \{y_i\} \in K_{r,d}(u, v)$  for two distinct anchors through the same  $(d-1)$ -core  $R$ , then the same subtraction argument applied separately to  $u$  and  $v$  gives

$$H_{r,d}(w)\ell_R = H_{r,d}(w)(X\ell_R) = 0, \quad w \in \{u, v\}.$$

These are exactly the first  $r$  and last  $r$  rows of

$$H_{r+1,d-1}(w)\ell_R = 0.$$

Hence  $R \in K_{r+1,d-1}(u, v)$ . So one-exchange collisions inside the lifted top-kernel ledger are charged losslessly to the next simultaneous Hankel kernel; after those root slices are paid, the residual  $K_{r,d}$  family has no one-exchange edges. This records the residual-depth frontier shift as an identity rather than a multiplicative loss source.

The same Hankel shift also classifies the one-outside external-anchor ruled branch from the boundary-off audit. For a shadow  $S \subset D$ ,  $|S| = j-1$ , and an external anchor  $\beta \notin D$ ,

$$\ell_{S,\beta} = (X - \beta)\ell_S.$$

The  $t = 2$  determinant gate is again quadratic in  $\beta$ . If it is nonzero, at most two external anchors pass. If it vanishes identically, then with

$$c_i(w) = \text{row}_i(H_{3,j-1}(w)\ell_S)$$

one has

$$H_{2,j}(w)\ell_{S,\beta} = (c_1(w) - \beta c_0(w), c_2(w) - \beta c_1(w)).$$

The same recurrence argument as above forces the ruled external-anchor branch to be inactive or fixed finite slope. In the fixed-slope case it is a boundary root slice and lifts to

$$H_{3,j-1}(u + z_0 v)\ell_S = 0.$$

Thus, after fixed-slope boundary slices are charged, each shadow  $S$  supports at most two active non-ruled external anchors. This classifies the ruled  $\text{Boundary}_{\text{off}}$  branch locally, but it is not yet a global bound for the one-outside target image. Equivalently, projecting a residual one-outside target  $B_\beta = S \cup \{\beta\}$  to its domain shadow  $S$  has fibers of size at most two. If two external anchors over the same shadow had the same finite slope  $z$ , then subtraction gives the lifted boundary root slice

$$H_{3,j-1}(u + zv)\ell_S = 0,$$

so after that charge the remaining two-anchor shadow fibers have distinct finite slopes. Hence

$$|\text{Boundary}_{\text{off}}^{\text{res}}| \leq 2|\text{Shadow}_{\text{off}}^{\text{res}}|.$$

This reduces the live one-outside object to a shadow-image problem rather than an external-anchor multiplicity problem. The shadow condition itself is intrinsic. Writing

$$c_i = \text{row}_i(H_{3,j-1}(u + zv)\ell_S), \quad 0 \leq i \leq 2,$$

there is an external anchor  $\beta$  with  $H_{2,j}(u + zv)\ell_{S,\beta} = 0$  iff either  $(c_0, c_1, c_2) = (0, 0, 0)$ , the already charged lifted boundary slice, or

$$c_0 \neq 0, \quad c_1^2 = c_0 c_2, \quad \beta = c_1/c_0.$$

Thus the residual boundary shadow image lies in a rank-one scalar Hankel locus with recovered anchor  $c_1/c_0 \notin D$ , plus the usual active filter. As  $z$  varies, write  $c_i(z) = a_i + zb_i$ . The recovered-anchor gate is the quadratic

$$Q_S(z) = c_1(z)^2 - c_0(z)c_2(z).$$

If  $Q_S$  is nonzero, the fixed shadow contributes at most two candidate finite slopes and hence at most two recovered anchors. If  $Q_S \equiv 0$ , the line  $(c_0, c_1, c_2)(z) = a + zb$  lies on one generator of the rank-one cone  $x_1^2 = x_0 x_2$ . Thus any finite recovered anchor is independent of  $z$ , and both  $H_{2,j}(u)\ell_{S,\beta}$  and  $H_{2,j}(v)\ell_{S,\beta}$  vanish there. The active filter removes this degenerate branch, while zero triples are already charged to the lifted boundary slice. The remaining boundary-shadow problem is therefore a nonzero quadratic-root problem over the shadows. Equivalently, with  $r_\beta = (1, \beta, \beta^2)$  and  $h_\beta(c) = (c_1 - \beta c_0, c_2 - \beta c_1)$ , define the anchor gate

$$A_S(\beta) = \det(a, b, r_\beta).$$

Then  $A_S(\beta) = \det(h_\beta(a), h_\beta(b))$ . If  $h_\beta(b) \neq 0$ , a root  $A_S(\beta) = 0$  gives a unique finite slope  $z$  with  $h_\beta(a + zb) = 0$ ; nonzero triples are exactly the recovered rank-one anchors above, while zero triples are charged. If  $A_S \equiv 0$ , then  $a, b$  are proportional and the branch is inactive or charged. Thus the same residual target is a nondegenerate conic-secant incidence: a nonzero quadratic in the outside anchor  $\beta$ , with the active filter imposed. Fixing  $\beta$  gives a further one-root reduction. For  $|R| = j - 2$ , put  $P_{R,\beta} = (X - \beta)\ell_R$  and  $\ell_{R,\beta,y} = (X - y)P_{R,\beta}$ . The determinant

$$\det(H_{2,j}(u)\ell_{R,\beta,y}, H_{2,j}(v)\ell_{R,\beta,y})$$

is quadratic in  $y$ . A nonzero determinant has at most two domain roots; an identically zero determinant is inactive or fixed finite slope, and the fixed-slope case lifts to

$$H_{3,j-1}(u + zv)P_{R,\beta} = 0.$$

After charging these one-outside lifted boundary-core slices, projection  $(R \cup \{y\}, \beta) \mapsto (R, \beta)$  has residual fibers of size at most two. Combining this with the shadow-fiber bound, for each fixed  $(j - 2)$ -core  $R$  the residual boundary incidences between external anchors  $\beta$  and domain roots  $y \in D \setminus R$  form a bipartite graph of degree at most two on both sides. Hence each fixed  $R$  supports at most  $2(n - j + 2)$  residual external anchors. Thus the next boundary object is the lower-dimensional boundary-core image, or still further the active  $(j - 2)$ -core image. For fixed  $R$ , write

$$U_m = H_{2,j}(u)(X^m \ell_R), \quad V_m = H_{2,j}(v)(X^m \ell_R), \quad 0 \leq m \leq 2.$$

Then the remaining fixed-core incidence is cut out by the symmetric bidegree  $(2, 2)$  determinant

$$\Delta_R(\beta, y) = \det(U_2 - (\beta + y)U_1 + \beta y U_0, V_2 - (\beta + y)V_1 + \beta y V_0).$$

The fixed-shadow and fixed-anchor quadratics are the two coordinate specializations of this same determinant. Thus the active core image is a concrete bidegree-two incidence target, not an uncontrolled external-anchor search. Equivalently,  $\Delta_R(\beta, y) = F_R(\beta + y, \beta y)$ , where

$$F_R(s, p) = \det(U_2 - sU_1 + pU_0, V_2 - sV_1 + pV_0)$$

is the ordinary two-root elementary determinant. Fixing either coordinate is the fixed-root line  $p = \alpha s - \alpha^2$  in the  $(s, p)$ -plane, so the boundary-core target is a mixed-domain slice of the same determinant geometry used by the fixed-root, fixed-sum, and product-Mobius line-packet ledger. Moreover the mixed slice of each surviving non-fixed line packet is explicit: on a fixed-sum line  $x + y = s_0$  one has  $\beta = s_0 - y$ , while on a nondegenerate product-Mobius line  $(x - c)(y - c) = \mu \neq 0$  one has  $\beta = c + \mu/(y - c)$ . Thus, over a fixed core  $R$ , such a packet contributes at most  $n - j + 2$  one-outside boundary-core pairs, namely the domain roots whose packet partner leaves  $D$ ; the  $\mu = 0$  case is the fixed-root line already charged. The same elementary-plane form also controls slope multiplicity on the remaining conic branch. For fixed finite slope  $z$ , put  $W_m = U_m + zV_m$ . The same-slope equations are

$$W_2 - sW_1 + pW_0 = 0,$$

a pair of affine-linear equations in  $(s, p)$ . Hence the same-slope fiber is empty, a point, an affine line, or the full plane. A repeated slope on two distinct elementary points therefore forces a constant-slope affine line, or the full-plane lift. Fixed-root lines are already boundary/root-slice charges, and non-fixed constant-slope lines collapse to the full-plane charge above. Thus after these charges the finite-slope label is injective on the live fixed-core boundary conic, so the quartic point count below also controls the number of live slopes. For the complementary conic case, write

$$F_R(s, p) = f_{20}s^2 + f_{11}sp + f_{02}p^2 + f_{10}s + f_{01}p + f_{00}.$$

Then for fixed  $y$ ,

$$F_R(\beta + y, \beta y) = A_R(y)\beta^2 + B_R(y)\beta + C_R(y),$$

where

$$\begin{aligned} A_R(y) &= f_{20} + f_{11}y + f_{02}y^2, \\ B_R(y) &= f_{10} + (2f_{20} + f_{01})y + f_{11}y^2, \\ C_R(y) &= f_{00} + f_{10}y + f_{20}y^2. \end{aligned}$$

In odd characteristic the nondegenerate anchor count is controlled by the quartic discriminant  $B_R(y)^2 - 4A_R(y)C_R(y)$ ; a fully zero quadratic is the fixed-root component  $p = ys - y^2$ , already charged. Thus the remaining non-line boundary-core target is a quartic discriminant/Kummer trace over domain roots, before outside-domain and active filters. If this quartic discriminant is identically zero, then coefficient comparison shows that either  $F_R = \lambda L^2$  for an affine line  $L$ , already a fixed-sum, fixed-root, or product-Mobius line-packet branch, or  $F_R = \lambda(s^2 - 4p)$ . The latter restricts to  $\lambda(\beta - y)^2$  on the boundary slice, hence contributes only  $\beta = y$ , which is removed by the outside-domain/distinct-root filters. So the live conic target after line-packet and fixed-root charges has nonzero quartic discriminant. If  $A_R \not\equiv 0$ , then away from at most two roots of  $A_R$  the quadratic formula identifies anchor roots with the double cover

$$W_R^2 = B_R(y)^2 - 4A_R(y)C_R(y), \quad W_R = 2A_R(y)\beta + B_R(y).$$

After squarefree reduction this cover has genus at most one, since the discriminant has degree at most four. If  $A_R \equiv 0$ , the determinant is affine-linear in  $(s, p)$  and has already returned to the line-packet branch. Thus the live non-line boundary target has uniformly bounded genus/conductor, up to at most two linear exceptional fibers. There is also an exact full-subgroup Kummer gate. If  $D \leq \mathbb{F}_p^*$  has index  $e$ ,  $\chi$  has kernel  $D$ , and  $\ell = \text{lcm}(e, 2)$ , then away from the charged fixed-root fibers and the at most two  $A_R = 0$  exceptions the full-subgroup count expands as

$$N_R(D) = |D| + \frac{1}{e} \sum_{a=0}^{e-1} \sum_{y \in \mathbb{F}_p^*} \chi^a(y) \chi_2(\text{Disc}_R(y)) + O(1).$$

Combining  $\chi$  and  $\chi_2$  through a character of order  $\ell$ , the  $a$ -term is geometrically trivial only if

$$\operatorname{div}\left(y^{a\ell/e} \operatorname{Disc}_R(y)^{\ell/2}\right) \equiv 0 \pmod{\ell}.$$

Equivalently, all nonzero roots of  $\operatorname{Disc}_R$  have even multiplicity and the zero-root parity matches  $a$ . Hence the squarefree part of a no-cancellation quartic is forced to be either 1 or  $y$ :

$$\operatorname{Disc}_R(y) = cG(y)^2 \quad \text{or} \quad \operatorname{Disc}_R(y) = cyG(y)^2.$$

The first case forces  $a = 0$ , while the second forces  $e$  even and  $a = e/2$ . If this power gate fails, the standard one-dimensional Kummer-Weil bound on  $\mathbb{P}^1$  has support contained in the four discriminant roots plus 0 and  $\infty$ , so the term has depth-independent conductor and is bounded by  $4\sqrt{p}$ . Thus any no-cancellation boundary quartic is an explicit rational graph or fixed square-root rational graph branch, not a hidden growing-conductor family. It follows that the full-subgroup live count over a fixed core satisfies

$$N_R^{\text{live}}(D) \leq |D| + 4\sqrt{p} + O(1)$$

when the power gate is absent, and

$$N_R^{\text{live}}(D) \leq 2|D| + 4\sqrt{p} + O(1)$$

in general. There is at most one geometrically trivial Fourier term: the principal term in the  $cG^2$  case or the quadratic subgroup term in the  $cyG^2$  case. The outside-domain and active filters only delete points, and the same-slope fiber injection above turns this point count into the same live slope count for the fixed core. Combining with the degree-two graph bound gives the ledger-facing non-line conic estimate

$$N_R^{\text{conic}}(D) \leq \min\{2(n-j+2), |D| + 4\sqrt{p} + O(1)\}$$

when the power gate is absent, and

$$N_R^{\text{conic}}(D) \leq \min\{2(n-j+2), 2|D| + 4\sqrt{p} + O(1)\}$$

in general. Reducible fixed-sum and product-Mobius line packets remain in the separate graph/packet ledger; the non-line fixed-core geometry has no remaining uncontrolled multiplicity. The same fixed-core target also has a slope-side recurrence chart. Write

$$a_i = \operatorname{row}_i(H_{4,j-2}(u)\ell_R), \quad b_i = \operatorname{row}_i(H_{4,j-2}(v)\ell_R), \quad c_i(z) = a_i + zb_i.$$

For  $P(X) = X^2 - sX + p$ , the equation

$$H_{2,j}(u + zv)(P\ell_R) = 0$$

is exactly

$$c_2 - sc_1 + pc_0 = 0, \quad c_3 - sc_2 + pc_1 = 0.$$

If  $Q_R(z) = c_0c_2 - c_1^2 \neq 0$ , then

$$s = \frac{c_0c_3 - c_1c_2}{Q_R(z)}, \quad p = \frac{c_1c_3 - c_2^2}{Q_R(z)},$$



and the split-root condition is governed by the quartic numerator

$$(c_0c_3 - c_1c_2)^2 - 4(c_0c_2 - c_1^2)(c_1c_3 - c_2^2).$$

If  $Q_R(z) = 0$  and a solution exists, then the four  $c_i$ 's are either all zero, giving the full-plane lift, or form a geometric progression  $c_i = c_0\alpha^i$ , in which case every solution has  $P(\alpha) = 0$  and is a fixed-root line. Thus the residual non-line root-core target lives in the nonzero-denominator recurrence chart. Set

$$A_R = c_0c_3 - c_1c_2, \quad B_R = c_1c_3 - c_2^2, \quad \Theta_R = A_R^2 - 4Q_RB_R.$$

Because the  $c_i$ 's are linear in  $z$ , the polynomials  $Q_R, A_R, B_R$  have degree at most two and  $\Theta_R$  has degree at most four. On  $Q_R \neq 0$ , a split lift with roots  $r_1, r_2$  gives

$$y = Q_R(z)(r_1 - r_2), \quad y^2 = \Theta_R(z),$$

and conversely a point on  $Y^2 = \Theta_R(z)$  recovers the formal roots  $(s_R(z) \pm y/Q_R(z))/2$ . Therefore, after the denominator-zero fixed-root/full-plane cases are charged, the residual split root-core slopes are filtered points on a degree-four double cover of the slope line. Its squarefree reduction has genus at most one; if  $\Theta_R$  is a square the roots are rational functions of  $z$ , and otherwise the standard Kummer-Weil bound has depth-independent conductor. The domain/outside filter does not change this conductor scale. If  $D = (\mathbb{F}_p^*)^e$  and  $\chi$  is the quotient character of order  $e$ , then on the cover, away from  $Q_R = 0$ ,

$$r_{\pm}(z, Y) = \frac{A_R(z) \pm Y}{2Q_R(z)}, \quad 1_D(x) = \frac{1}{e} \sum_{a=0}^{e-1} \chi^a(x),$$

with characters extended by zero at  $x = 0$ . Hence the condition  $r_+ \in D, r_- \notin D$  is an explicit finite sum of Kummer traces

$$\sum_{(z,Y)} \chi^a(r_+(z, Y)) \chi^b(r_-(z, Y))$$

on the same genus-zero/genus-one cover. The finite poles lie over the at most two roots of  $Q_R$ ; finite zeros lie over the at most two roots of  $B_R$ ; the remaining support is at infinity and at the at-most-four branch points of  $\Theta_R$ . Thus every non-power summand has  $O_e(\sqrt{p})$  size by the standard curve Kummer-Weil bound with conductor independent of depth. The only no-cancellation cases are explicit power-divisor congruences for  $r_+^a r_-^b$  on this bounded divisor support.

Consequently, after fixed-slope root slices are removed, the residual  $t = 2$  one-exchange graph has maximum degree at most  $j$ : each locator has only  $j$  possible  $(j-1)$ -cores, and each core supplies at most one residual neighbor. In particular the unordered residual edge count is at most  $j|A|/2$ , and also at most  $\binom{|D|}{j-1}$  by counting cores. Plugging this into the average support-collinearity ledger `experimental/notes/m1/m1_average_support_collinearity.md`, and writing the line-field size as  $Q$ , gives for the residual family  $A_{\text{res}}, M = |A_{\text{res}}|$ ,

$$B_2^{\max}(A_{\text{res}}) \leq \frac{1 - p_z}{Mp_z} + \frac{4jQ}{M}, \quad p_z = Q^{-2}(1 - Q^{-2}).$$

Thus this branch contributes  $o(1)$  to the average missing-slope density whenever  $Mp_z \rightarrow \infty$  and  $M/(jQ) \rightarrow \infty$ . The remaining M1 obstruction must come from small residual family size, root-slice charges, or higher packet/two-exchange structure; this is not yet a worst-case packing theorem.

The same substitution has a packet-level higher-exchange form. Inside one same-slope affine  $h$ -exchange packet, after full moving  $r$ -root fibers have been charged, the residual codegree bound

$\Gamma_r \leq \binom{h}{r}(Q_F^{r-1} - 1)$  gives, for average-ledger support slack  $\tau$ ,

$$B_\tau^{\max}(A_{\text{res}}) \leq \frac{1-p_z}{Mp_z} + \frac{4}{M} \sum_{r=1}^{\min(h, \tau-1)} \binom{h}{r} (Q_F^{r-1} - 1) Q_F^{\tau-r}, \quad p_z = Q_F^{-\tau} (1 - Q_F^{-\tau}).$$

This is still a local packet ledger input; a global M1 estimate must also sum over packet choices and the quotient, tangent/contained, split-root, and fixed-slope ledgers.

## 19.5 M1: variable-line packet reduction

The follow-up variable-line packet note isolates a local reduction inside the same Hankel-pencil branch. Fix a  $(j-2)$ -core  $R$  and a non-fixed proper one-dimensional determinant component  $L$  in the two-root plane

$$T = R \cup \{x, y\}, \quad s = x + y, \quad p = xy.$$

The constant-line collapse above verifies the needed variable-slope hypothesis for every surviving fixed-sum or nondegenerate product-Mobius packet after fixed-root and full-plane charges. Thus the active-new mass  $r_L$  on such an  $L$  obeys

$$r_L \leq \mathbf{1}_{d_L=m_L=r_L=1} + (d_L - m_L) + 2 \binom{m_L}{2}.$$

The unordered packet pairs in the final term inject into the global different-slope two-exchange codegree ledger, so summing over non-fixed line packets gives

$$\sum_L r_L \leq S_{\text{dom}} + Q_{\text{def}} + 2E_{\text{pkt}}.$$

The remaining domain-singleton term is reduced to contained/tangent and one-outside target images:

$$S_{\text{dom}} \leq Z_0 + 2 \binom{j}{2} \binom{n-j}{2} |\text{Target}_{\text{cb}}| + (j-1) \binom{n-j+1}{2} |\text{Target}_{\text{off}}|.$$

Writing  $a = n - j$ , the exceptional zero-lower term vanishes whenever

$$a > \frac{n+1}{2},$$

and hence throughout the positive-slack rate-half range  $k \geq n/2$ ,  $a = k + t$ ,  $t \geq 1$ . Consequently the non-fixed variable-line branch is reduced to two explicit residual estimates: the active different-slope two-exchange codegree and the one-outside boundary shadow image, after quotient defects, contained/tangent ledgers, and fixed-slope boundary slices have been paid. Indeed the shadow-fiber bound gives

$$|\text{Target}_{\text{off}}^{\text{res}}| \leq 2|\text{Shadow}_{\text{off}}^{\text{res}}|.$$

After the fixed-anchor boundary-core slices are also charged, the sharper fiber bound

$$(j-1)|\text{Target}_{\text{off}}^{\text{res}}| \leq 2|\text{Core}_{\text{off}}^{\text{res}}|$$

may be substituted directly into the singleton target term. The boundary contribution

$$(j-1) \binom{n-j+1}{2} |\text{Target}_{\text{off}}^{\text{res}}|$$

is then bounded by

$$2 \binom{n-j+1}{2} |\text{Core}_{\text{off}}^{\text{res}}|,$$

so a polynomial bound for the boundary-core image closes this branch with only an  $n^2$  bookkeeping loss, rather than the older  $n^3$  boundary-image loss. The fixed-core graph form also gives

$$|\text{Core}_{\text{off}}^{\text{res}}| \leq 2(n-j+2) |\text{Root}_{\text{off}}^{\text{res}}|,$$

so bounding the active  $(j-2)$ -core image closes the same branch with an  $n^3$  boundary loss after all boundary-core root slices have been charged.

**Status.** This is a proved local packet lemma under the Hankel-pencil normal form after the same-slope, fixed-root, full-plane, and boundary-slice charges recorded above. It is not the final all-line M1 theorem and does not add a leaderboard row.

## 19.6 M1: width-one fixed-root closure

PR #136, due to AllenGrahamHart, extracts a compact closure criterion for the width-one critical-tail branch. At a base-free canonical  $b = 2$  node, let

$$V = \langle P, Q \rangle \subset F[X]_{<q}, \quad |D^{\text{can}}| = q + s.$$

A root-free width-one certificate is a direction  $A \in V$  whose root shadow on  $D^{\text{can}}$  has maximal size  $q - 1$ . Writing  $Z = Z(A)$  and  $O = D^{\text{can}} \setminus Z$ , one has

$$A = c\ell_Z, \quad |Z| = q - 1, \quad |O| = s + 1.$$

Equivalently,  $O$  supports a width-one certificate if and only if

$$L_O = \ell_{D^{\text{can}} \setminus O} \in V,$$

or equivalently all  $3 \times 3$  coefficient minors of  $P, Q, L_O$  vanish. Thus width-one certificates are bounded-complement rank tests rather than an unstructured projective-direction family.

The large width-one cube descends losslessly under fixed-root absorption. If  $P_m \subset Z$ , then

$$A^{P_m} = A / \ell_{P_m} = c\ell_{Z \setminus P_m}, \quad D^{\text{can}} \setminus P_m = (Z \setminus P_m) \sqcup O,$$

so the descended width remains one and the bounded complement  $O$  is unchanged. With  $a = \lfloor (q-2)/2 \rfloor$ , the co-small flag cube has the first-root partition

$$\sum_{e=1}^a \binom{q-1}{e} = \sum_{x \in Z} \sum_{f=0}^{a-1} \binom{|Z_{>x}|}{f}.$$

After exposing the first root  $x$ , every nonempty width-one flag injects into a one-root absorbed fixed-divisor/root-slice family. Quantitatively, for

$$M_x = \#\{F \subset Z_{>x} : |F| \leq a-1\},$$

one gets

$$\max_{x \in Z} M_x \geq \frac{1}{q-1} \sum_{e=1}^a \binom{q-1}{e} \geq \frac{2^{q-1}}{2q^2} \quad (q \geq 4).$$

Thus a large width-one cube already produces a large one-root slice; it cannot hide as a many-first-root averaging phenomenon.

Consequently, in fixed surplus  $s_0 \leq \sigma$ , small nodes are polynomial by the active-tree bound, and on  $q_S > s_S + 2$  there is at most one width-one shadow per node. The local closure criterion is

$$WO_1(A_0) \leq \text{FixedRootOneRoot}_{r1}(A_0) + O_\sigma(Q^{\sigma+1}),$$

where  $Q$  is the initial quotient width. This closes the width-one critical-tail branch only after a polynomial bound for the displayed one-root fixed-root/root-slice ledger is proved or imported. It is not the final all-line M1 theorem and does not add a leaderboard row.

## 19.7 Challenge-map pullbacks and public status corrections

The adjacent-ledger update adds the missing pullback rule from slope ledgers to protocol challenge maps. For a challenge-to-slope map  $\phi : K \rightarrow Z$ , a bad set  $B \subseteq Z$  pulls back with probability

$$\frac{|\phi^{-1}(B)|}{|K|}.$$

If the nonempty fibers have sizes  $s_1 \geq s_2 \geq \dots$ , the adversarial envelope for  $|B| = L$  is  $(s_1 + \dots + s_L)/|K|$ . For an  $\mathbb{F}_q$ -linear map of rank  $r$ , the effective denominator is  $q^r$ , not the ambient challenge space size. This prevents accidental multiplication of denominators in protocol ledgers.

The public-board status corrections are:

| row           | $a$ | tangent floor | status                                      |
|---------------|-----|---------------|---------------------------------------------|
| Cycle116 gate | 262 | 251           | gate unconditional; exact count conditional |
| Cycle119 gate | 263 | 250           | gate unconditional; exact count conditional |
| reserve272    | 272 | 241           | proved but subsumed                         |
| reserve288    | 288 | 225           | proved but subsumed                         |
| reserve313    | 313 | 200           | proved but subsumed                         |

The exact Cycle84 numerator 52,747,567,092 remains a computation-dependent record; only the weaker  $> 2^{-128}$  gate follows from the generic tangent floor without that finite census.

## 20 High-Agreement Tangent Staircase

**Status: New-local finite MDS theorem.** This section records a generic moving-root tangent floor and, in the very-high-agreement range, a matching upper bound. It is independent of quotient-periodic structure and does not use smoothness. It is relevant to the finite row ledger because the bad slopes are paid with the actual line field  $q_{\text{line}}$ .

For a polynomial  $P \in F[X]$ , write

$$P = P_{<k} + P_{\geq k}, \quad \deg P_{<k} < k,$$

where  $P_{\geq k}$  is the sum of the terms of degree at least  $k$ . Let  $\text{LD}_{\text{sw}}(C, a)$  denote the largest possible number of finite slopes on a received line that have a support-wise noncontained explanation on some support of size at least  $a$ .

**Theorem 6** (moving-root tangent floor). *Let  $D \subset F$  have  $|D| = n$ , and let  $C = \text{RS}[F, D, k]$ . For every integer  $a$  with  $k + 1 \leq a \leq n$ ,*

$$\text{LD}_{\text{sw}}(C, a) \geq n - a + 1.$$

*Proof.* Choose a subset  $A \subset D$  of size  $a - 1$ , and put

$$B_A(X) = \prod_{x \in A} (X - x).$$

Define the received line

$$f = (XB_A)_{\geq k}, \quad g = -(B_A)_{\geq k}.$$

For every  $t \in D \setminus A$ , set

$$L_t(X) = (X - t)B_A(X).$$

Then  $L_t$  is the locator of the support  $A \cup \{t\}$ , which has size  $a$ , and linearity of truncation gives

$$f + tg = (L_t)_{\geq k}.$$

On every root of  $L_t$ , the word represented by  $(L_t)_{\geq k}$  agrees with  $-(L_t)_{< k}$ , a codeword of  $C$ . Thus the slope  $t$  is explained on  $A \cup \{t\}$ .

It remains to check support-wise noncontainment on this same support. On  $A$ , since  $B_A = 0$ , we have

$$g = -(B_A)_{\geq k} = (B_A)_{< k}.$$

If  $g$  were explained on  $A \cup \{t\}$  by a degree- $< k$  polynomial  $q$ , then  $q = (B_A)_{< k}$ , because  $|A| = a - 1 \geq k$ . At the remaining point  $t$ , this would force

$$-(B_A)_{\geq k}(t) = (B_A)_{< k}(t),$$

that is,  $B_A(t) = 0$ , contrary to  $t \notin A$ . Hence  $g$  is not separately explained on  $A \cup \{t\}$ , so the slope is support-wise noncontained. The slopes  $t \in D \setminus A$  are distinct, giving  $n - a + 1$  bad finite slopes.  $\square$

**Theorem 7** (very-high-agreement tangent staircase). *Let  $C = \text{RS}[F, D, k]$  with  $|D| = n$ . If*

$$3a - 2n \geq k,$$

*then*

$$\text{LD}_{\text{sw}}(C, a) = n - a + 1.$$

*Proof.* The lower bound is Theorem 6. For the upper bound, put  $s = n - a$ . If a received line has at most one support-wise noncontained finite slope at agreement at least  $a$ , then the claimed upper bound is immediate. Otherwise pick two such slopes  $z_1 \neq z_2$ , with explaining codewords  $p_1, p_2 \in C$  on supports  $S_1, S_2$ , respectively. Since  $|S_i| \geq a$ ,

$$|S_1 \cap S_2| \geq 2a - n.$$

On this intersection,

$$g = (p_1 - p_2)/(z_1 - z_2), \quad f = p_1 - z_1 g.$$

The condition  $3a - 2n \geq k$  and the inequality  $a \leq n$  imply  $2a - n \geq k$ . Therefore these two right-hand sides extend uniquely to codewords  $c_g, c_f \in C$ . Let  $S_0$  be a maximal common support on which  $f = c_f$  and  $g = c_g$ , and put  $b = |S_0|$  and  $d = n - b$ . Then

$$b \geq 2a - n, \quad d \leq 2(n - a) = 2s.$$

By maximality, outside  $S_0$  there is no coordinate at which both residuals  $f - c_f$  and  $g - c_g$  vanish.

The residual-budget proposition, Proposition 7, is applicable because

$$a + b - n \geq a + (2a - n) - n = 3a - 2n \geq k.$$

Here its common-zero parameter is  $c_0 = 0$ . If  $b \geq a$ , then every noncontained slope needs at least one residual zero outside  $S_0$ , and there are only  $d \leq s$  such coordinates. If  $b < a$ , then every noncontained slope needs at least

$$h = a - b = d - s$$

private residual zeros outside  $S_0$ . Since  $s < d \leq 2s$ , the residual budget gives

$$\#\{\text{bad slopes}\} \leq \left\lfloor \frac{d}{d-s} \right\rfloor \leq s + 1 = n - a + 1.$$

This proves the matching upper bound.  $\square$

**Corollary 2** (tangent-star extremizers in the exact range). *Let  $C = \text{RS}[F, D, k]$ ,  $|D| = n$ , and suppose that*

$$k + 1 \leq a \leq n, \quad 3a - 2n \geq k.$$

*If a received line  $f + zg$  has exactly  $n - a + 1$  support-wise noncontained finite bad slopes at agreement at least  $a$ , and if  $n - a + 1 \geq 2$ , then the line is tangent-star extremal in the following sense. There are codewords  $c_f, c_g \in C$  and a common support  $S_0 \subseteq D$  of size  $a - 1$  such that, after replacing  $f, g$  by  $f - c_f, g - c_g$ , every bad slope is obtained by choosing exactly one residual coordinate in  $D \setminus S_0$ . More precisely, writing  $\Omega = D \setminus S_0$ , the map*

$$x \mapsto -\frac{f(x) - c_f(x)}{g(x) - c_g(x)}$$

*is a bijection from  $\Omega$  to the bad-slope set.*

*Proof.* Choose two distinct bad slopes and form the maximal common code-line support  $S_0$  as in the proof of Theorem 7; write  $b = |S_0|$ ,  $s = n - a$ , and  $d = n - b$ . Equality in the upper bound from that proof cannot occur when  $b \geq a$ , because then there are at most  $n - a$  bad slopes. Hence  $b < a$ . The residual-budget bound is

$$\#\{\text{bad slopes}\} \leq \left\lfloor \frac{d}{d-s} \right\rfloor \leq s + 1 = n - a + 1.$$

For equality, the second inequality must be sharp, so  $d - s = 1$ , equivalently  $b = a - 1$ . The maximality of  $S_0$  gives no common residual zeros outside  $S_0$ . Since  $|\Omega| = n - a + 1$  and each residual coordinate can vanish for at most one finite slope, equality forces every residual coordinate to be used exactly once. This is precisely the displayed residual-root bijection.  $\square$

**Corollary 3** (exact high-agreement gate for the  $\mathbb{F}_{17^{32}}$ , rate-1/2 row). *Let  $H \subset \mathbb{F}_{17^{32}}^*$  be any multiplicative subgroup of order 512, and let*

$$C = \text{RS}[\mathbb{F}_{17^{32}}, H, 256].$$

*For every  $a \geq 427$ ,*

$$\text{LD}_{\text{sw}}(C, a) = 513 - a.$$

*In particular,*

$$\text{LD}_{\text{sw}}(C, 506) = 7, \quad \text{LD}_{\text{sw}}(C, 507) = 6.$$

Since

$$6 \cdot 2^{128} < 17^{32} < 7 \cdot 2^{128},$$

we have

$$\frac{\text{LD}_{\text{sw}}(C, 507)}{17^{32}} < 2^{-128} \quad \text{but} \quad \frac{\text{LD}_{\text{sw}}(C, 506)}{17^{32}} > 2^{-128}.$$

Thus, under the finite-slope support-wise MCA convention, the  $2^{-128}$  integer agreement staircase for this row is pinned between 506 and 507. The largest safe integer Hamming radius is 5, while integer radius 6 is already unsafe. The corresponding normalized grid radii are

$$5/512 \quad \text{and} \quad 6/512 = 3/256.$$

As a real-radius supremum, the transition is  $3/256$ , but the closed endpoint  $\delta = 3/256$  is on the unsafe side.

*Proof.* Here  $n = 512$  and  $k = 256$ . The hypothesis  $3a - 2n \geq k$  becomes  $3a - 1024 \geq 256$ , hence  $a \geq 427$ . The exact formula is Theorem 7. The arithmetic inequalities are

$$17^{32} - 6 \cdot 2^{128} = 326217393234836465063858652730341978625 > 0$$

and

$$7 \cdot 2^{128} - 17^{32} = 14064973686101998399515954701426232831 > 0.$$

□

**Corollary 4** (no seventh finite-slope branch past agreement 506). *For  $C = \text{RS}[\mathbb{F}_{17^{32}}, H, 256]$  with  $|H| = 512$ , no finite-slope support-wise MCA mechanism can add a seventh bad slope at agreement  $a \geq 507$ . At equality, any extremal line is tangent-star in the sense of Corollary 2.*

*Proof.* By Corollary 3,

$$\text{LD}_{\text{sw}}(C, a) = 513 - a \quad (a \geq 427).$$

For every  $a \geq 507$ , the right-hand side is at most 6. Thus a seventh finite support-wise bad slope is impossible in this range. The extremizer classification is Corollary 2. □

**Remark 6** (effect on the earlier agreement-353 target). *The agreement-353 target is no longer a frontier for the rate-1/2,  $\mathbb{F}_{17^{32}}$ ,  $n = 512$  row: Theorem 6 already gives*

$$\text{LD}_{\text{sw}}(C, 353) \geq 512 - 353 + 1 = 160.$$

*The quotient-core strict352 packet remains a useful mechanism record, but the generic tangent floor gives a stronger lower bound at agreement 352 and beyond. Corollary 4 rules out a seventh finite-slope branch past agreement 506. The adjacent-ledger section below treats the high-agreement projective-slope, CA, curve-MCA, and interleaved-list coding objects; protocol-facing conversions still require separate challenge-field, extension-lift, folding, query, and cryptographic ledgers.*

## 21 High-Agreement Adjacent Ledgers

**Status: New-local finite Reed–Solomon theorem package, one MDS interleaved-list theorem, and a Conditional protocol-ledger corollary.** The tangent staircase of Section 20 settles more than finite-slope support-wise MCA. In the same high-agreement range it also pins the no-loss correlated-agreement line count, the projective-slope line count, the finite-parameter degree- $d$  curve count, and the interleaved-list size. The protocol statement at the end of this section is only a ledger corollary: it applies to reductions whose coding error is exactly one of the listed terms, plus an explicitly added query/folding error.

Except in the interleaved-list theorem,  $C = \text{RS}[F, D, k]$  with  $|D| = n$ , dimension  $k$ , and  $q = |F|$ . The interleaved-list uniqueness statement only uses the MDS property. The agreement parameter is an integer  $a$ , so the corresponding closed Hamming radius is  $1 - a/n$ .

### 21.1 No-loss CA and projective slopes

Let  $\text{LD}_{\text{ca}}(C, a)$  denote the maximum number of finite slopes  $z \in F$  on an affine received line  $f + zg$  such that  $f + zg$  has a  $C$ -explanation on at least  $a$  positions while the pair  $(f, g)$  has no  $C^2$ -explanation on any support of size at least  $a$ . This is the integer-agreement version of no-loss CA. Let  $\text{LD}_{\text{sw,proj}}(C, a)$  be the corresponding support-wise MCA count when slopes are points of  $\mathbb{P}^1(F)$ , so the point at infinity is also allowed.

**Theorem 8** (high-agreement CA and projective tangent staircase). *Let  $C = \text{RS}[F, D, k]$  with  $|D| = n$ . If*

$$3a - 2n \geq k,$$

*then*

$$\text{LD}_{\text{ca}}(C, a) = \text{LD}_{\text{sw,proj}}(C, a) = n - a + 1.$$

*Proof.* The finite-slope support-wise count is  $n - a + 1$  by Theorem 7. Every no-loss CA-bad finite slope is support-wise MCA-bad: if the line word is explained on a support  $S$  of size at least  $a$  and  $(f, g)$  were simultaneously explained on that same  $S$ , then  $(f, g)$  would be  $C^2$ -close at the no-loss CA radius. Thus

$$\text{LD}_{\text{ca}}(C, a) \leq n - a + 1.$$

The moving-root line from Theorem 6 gives the matching CA lower bound. Indeed, with  $A \subset D$ ,  $|A| = a - 1$ , put  $B_A(X) = \prod_{x \in A} (X - x)$ ,

$$f = (XB_A)_{\geq k}, \quad g = -(B_A)_{\geq k}.$$

For each  $t \in D \setminus A$ , the slope  $t$  is explained on  $A \cup \{t\}$ . Suppose  $(f, g)$  were explained by two degree- $< k$  polynomials on some support  $T$  of size at least  $a$ . Since

$$|T \cap A| \geq a - (n - a + 1) = 2a - n - 1,$$

and the hypothesis  $3a - 2n \geq k$  implies  $2a - n - 1 \geq k$  when  $a < n$  (the case  $a = n$  gives  $n - 1 \geq k$ ), the two explaining polynomials must be the unique degree- $< k$  polynomials agreeing with  $f$  and  $g$  on  $A$ , namely  $-(XB_A)_{< k}$  and  $(B_A)_{< k}$ . At any point  $x \in D \setminus A$ , simultaneous agreement with those two polynomials would force both  $XB_A(x) = 0$  and  $B_A(x) = 0$ , impossible. Hence  $(f, g)$  is globally CA-far at agreement  $a$ , and the  $n - a + 1$  slopes are CA-bad.

It remains to pass from finite to projective slopes. A projective bad-slope set cannot be all of  $\mathbb{P}^1(F)$ : otherwise, after choosing any projective point as infinity, the remaining  $q$  points would be



finite bad slopes in an affine coordinate, contradicting  $q > n - a + 1$  since  $D \subseteq F$  and  $a > k$ . Choose a nonbad projective point and send it to infinity by a change of basis of the two-dimensional received-line span. All bad projective points are then finite slopes in the new coordinate, so Theorem 7 bounds their number by  $n - a + 1$ . The affine moving-root construction gives the matching projective lower bound.  $\square$

## 21.2 Finite-parameter curve MCA

Fix an integer  $d \geq 1$ . A degree- $d$  received curve is a family

$$W_\gamma = f_0 + \gamma f_1 + \cdots + \gamma^d f_d, \quad \gamma \in F.$$

Let  $\text{CurveLD}_{\text{sw}}^{(d)}(C, a)$  be the maximum number of parameters  $\gamma \in F$  for which  $W_\gamma$  has a  $C$ -explanation on a support of size at least  $a$ , while the tuple  $(f_0, \dots, f_d)$  is not simultaneously explained by  $C^{d+1}$  on that support. Define  $\text{CurveLD}_{\text{ca}}^{(d)}(C, a)$  similarly, replacing the support-wise condition by the no-loss global CA condition that  $(f_0, \dots, f_d)$  is not  $C^{d+1}$ -close at agreement  $a$ .

**Theorem 9** (degree- $d$  curve tangent staircase). *Let  $C = \text{RS}[F, D, k]$ ,  $|D| = n$ ,  $|F| = q$ , and  $d \geq 1$ . If*

$$(d + 2)a - (d + 1)n \geq k,$$

*then*

$$\text{CurveLD}_{\text{sw}}^{(d)}(C, a) = \text{CurveLD}_{\text{ca}}^{(d)}(C, a) = \min\{q, d(n - a + 1)\}.$$

*Proof.* Put  $m = n - a + 1$ . For the lower bound, choose  $A \subset D$  with  $|A| = a - 1$  and let  $\Omega = D \setminus A$ , so  $|\Omega| = m$ . Choose  $M = \min\{q, dm\}$  distinct parameters in  $F$ , partition them among the points  $x \in \Omega$  into sets  $R_x$  of size at most  $d$ , and for each  $x \in \Omega$  choose a nonzero polynomial

$$P_x(T) = \sum_{i=0}^d p_{i,x} T^i$$

whose root set contains  $R_x$ . Define  $f_i = 0$  on  $A$  and  $f_i(x) = p_{i,x}$  for  $x \in \Omega$ . If  $\gamma \in R_x$ , then  $W_\gamma$  vanishes on  $A \cup \{x\}$ , so it is explained by the zero codeword on a support of size  $a$ . On this same support the tuple  $(f_0, \dots, f_d)$  is not simultaneously explained: every component vanishes on  $A$ , which has at least  $k$  points, so the only possible explaining codewords are all zero, but  $P_x$  is not the zero polynomial and hence some component is nonzero at  $x$ .

The same construction is CA-far. If the tuple were globally explained on a support  $T$  of size at least  $a$ , then

$$|T \cap A| \geq 2a - n - 1.$$

The curve-range hypothesis implies  $2a - n - 1 \geq k$  when  $a < n$ , and the case  $a = n$  is immediate from  $n - 1 \geq k$ . Thus all explaining codewords would be zero, which is impossible on every point of  $\Omega$  by construction. Hence all  $M$  parameters are curve-CA-bad and support-wise curve-MCA-bad.

For the upper bound, it is enough to prove the support-wise statement, since curve-CA-bad parameters are support-wise bad. If there are at most  $d$  bad parameters, the bound is immediate. Otherwise pick  $d + 1$  distinct bad parameters  $\gamma_0, \dots, \gamma_d$ , with explaining codewords on supports  $S_0, \dots, S_d$ . Their intersection has size at least

$$b_0 \geq (d + 1)a - dn.$$

On this intersection, interpolation in the parameter  $\gamma$  expresses each component  $f_i$  as an  $F$ -linear combination of the  $d + 1$  explaining codewords, hence as a codeword  $c_i \in C$ . Let  $S_*$  be a maximal support on which all  $f_i = c_i$ , and put  $b = |S_*|$ . Then  $b \geq b_0$ , and the hypothesis gives

$$a + b - n \geq a + b_0 - n = (d + 2)a - (d + 1)n \geq k.$$

Now consider any bad parameter  $\gamma$ , with explaining codeword  $p_\gamma$  on a support  $S$  of size at least  $a$ . On  $S \cap S_*$ , whose size is at least  $a + b - n \geq k$ , the codeword  $p_\gamma$  agrees with

$$c_\gamma = \sum_{i=0}^d \gamma^i c_i \in C.$$

Therefore the residual word  $W_\gamma - c_\gamma$  must vanish on at least  $h = \max\{1, a - b\}$  points of  $D \setminus S_*$ . By maximality of  $S_*$ , for each residual coordinate  $x \in D \setminus S_*$  the polynomial

$$R_x(T) = \sum_{i=0}^d T^i (f_i(x) - c_i(x))$$

is not identically zero, so it has at most  $d$  roots in  $F$ . Counting incidences  $(\gamma, x)$  with  $R_x(\gamma) = 0$  gives

$$\#\{\text{bad parameters}\} \cdot h \leq d(n - b).$$

If  $b \geq a$ , then  $h = 1$  and the right side is at most  $d(n - a) \leq d(n - a + 1)$ . If  $b < a$ , then

$$\#\{\text{bad parameters}\} \leq d \frac{n - b}{a - b} = d \left( 1 + \frac{n - a}{a - b} \right) \leq d(n - a + 1),$$

because  $a - b \geq 1$ . The trivial field-size cap gives the additional upper bound  $q$ . This proves the exact formula.  $\square$

### 21.3 Interleaved-list uniqueness

For  $\mu \geq 1$ , let  $\Lambda_\mu(C, a)$  denote the maximum, over all  $\mu$ -row received words  $U = (U_1, \dots, U_\mu)$ , of the number of tuples  $(c_1, \dots, c_\mu) \in C^\mu$  that agree with  $U$  on at least  $a$  common positions.

**Theorem 10** (high-agreement interleaved-list uniqueness). *Let  $C \subseteq F^D$  be MDS of length  $n$  and dimension  $k$ . If*

$$2a - n \geq k,$$

*then for every  $\mu \geq 1$ ,*

$$\Lambda_\mu(C, a) = 1.$$

*Proof.* The lower bound is obtained by taking  $U$  itself to be a code tuple. For the upper bound, suppose two tuples  $c = (c_1, \dots, c_\mu)$  and  $c' = (c'_1, \dots, c'_\mu)$  both agree with  $U$  on supports of size at least  $a$ . The supports intersect in at least  $2a - n \geq k$  positions. On this intersection, each row  $c_i$  agrees with  $c'_i$ . Since  $C$  is MDS, two codewords that agree on  $k$  positions are equal. Thus  $c_i = c'_i$  for every row  $i$ , so the tuples are identical.  $\square$

## 21.4 The $\mathbb{F}_{17^{32}}$ , $n = 512$ , $k = 256$ protocol-facing ledger

**Corollary 5** (high-agreement adjacent ledgers for the  $17^{32}$  row). *Let  $H \subset \mathbb{F}_{17^{32}}^*$  have order 512, and let*

$$C = \text{RS}[\mathbb{F}_{17^{32}}, H, 256].$$

*Put  $q = 17^{32}$  and*

$$B_* = \left\lfloor q/2^{128} \right\rfloor = 6.$$

*Then the following statements hold.*

1. *For every  $a \geq 427$ ,*

$$\text{LD}_{\text{ca}}(C, a) = \text{LD}_{\text{sw,proj}}(C, a) = \text{LD}_{\text{sw}}(C, a) = 513 - a.$$

*Thus the affine/projective line numerator is 6 at  $a = 507$  and 7 at  $a = 506$ .*

2. *For every  $\mu \geq 1$  and every  $a \geq 384$ ,*

$$\Lambda_\mu(C, a) = 1.$$

3. *For degree- $d$  finite-parameter curves, whenever*

$$a \geq \left\lceil \frac{(d+1)512 + 256}{d+2} \right\rceil,$$

*we have*

$$\text{CurveLD}_{\text{sw}}^{(d)}(C, a) = \text{CurveLD}_{\text{ca}}^{(d)}(C, a) = \min\{17^{32}, d(513 - a)\}.$$

*Proof.* The first item is Corollary 3 together with Theorem 8. The second item is Theorem 10, because  $2a - 512 \geq 256$  is equivalent to  $a \geq 384$ . The third item is Theorem 9 after substituting  $n = 512$  and  $k = 256$ . The integer  $B_* = 6$  follows from

$$6 \cdot 2^{128} < 17^{32} < 7 \cdot 2^{128}.$$

□

**Corollary 6** (conditional high-agreement protocol ledger). *Consider a protocol reduction for the row of Corollary 5 whose coding-theoretic error at agreement  $a$  is exactly*

$$\frac{N_{\text{geom}}(a)}{17^{32}} + \frac{\Lambda_\mu(C, a)}{17^{32}} + \varepsilon_{\text{query}},$$

*where  $N_{\text{geom}}(a)$  is one affine/projective line CA/MCA numerator or one degree- $d$  finite-parameter curve numerator, and where no additional fold, hash, Fiat–Shamir, or extension-lift loss is hidden in the notation. In the high-agreement ranges above, this ledger becomes the following explicit integer test.*

*For affine/projective line CA/MCA,*

$$N_{\text{geom}}(a) + \Lambda_\mu(C, a) = (513 - a) + 1 = 514 - a \quad (a \geq 427).$$

*With  $\varepsilon_{\text{query}} = 0$ , this is at most the  $2^{-128}$  budget iff  $a \geq 508$ . Thus the combined line-plus-list ledger is safe at agreement 508 and unsafe at agreement 507.*

For degree- $d$  finite-parameter curve MCA/CA,

$$N_{\text{geom}}(a) + \Lambda_{\mu}(C, a) = d(513 - a) + 1$$

in the curve-staircase range. With  $\varepsilon_{\text{query}} = 0$ , the safe grid points are exactly those satisfying

$$d(513 - a) + 1 \leq 6.$$

In particular:

| $d$        | curve-plus-list safe grid             | curve term alone safe grid                  |
|------------|---------------------------------------|---------------------------------------------|
| 1          | $a \geq 508$ ( $r \leq 4$ )           | $a \geq 507$ ( $r \leq 5$ )                 |
| 2          | $a \geq 511$ ( $r \leq 1$ )           | $a \geq 510$ ( $r \leq 2$ )                 |
| 3          | $a = 512$ ( $r = 0$ )                 | $a \geq 511$ ( $r \leq 1$ )                 |
| 4          | $a = 512$ ( $r = 0$ )                 | $a = 512$ ( $r = 0$ )                       |
| 5          | $a = 512$ ( $r = 0$ )                 | $a = 512$ ( $r = 0$ )                       |
| 6          | no safe grid point with the list term | $a = 512$ ( $r = 0$ )                       |
| $d \geq 7$ | no safe grid point with the list term | no safe grid point for the curve term alone |

where  $r = n - a$  is the integer Hamming radius.

*Proof.* By Corollary 5, the interleaved-list numerator is 1 throughout the displayed high-agreement range. The affine/projective line numerator is  $513 - a$ , and the degree- $d$  curve numerator is  $d(513 - a)$ . Since

$$\lfloor 17^{32}/2^{128} \rfloor = 6,$$

the code-level numerator must be at most 6 when  $\varepsilon_{\text{query}} = 0$ . The displayed inequalities are exactly the resulting integer conditions.  $\square$

**Remark 7** (what this does and does not settle). *Theorems 8, 9, and 10 are unconditional finite statements under the hypotheses printed above: the CA/projective/curve statements are Reed–Solomon evaluation-domain statements, while interleaved uniqueness is MDS. Corollary 6 is conditional on the protocol really consuming only the printed coding terms over the printed field. It does not cover reductions with additional curve families, extension-line lifts, folding losses, query errors, or cryptographic errors unless those terms are added to the ledger.*

## 22 General High-Agreement Ledgers Beyond the $\mathbb{F}_{17^{32}}$ Row

**Status: New-local finite Reed–Solomon/MDS theorem package plus Conditional protocol ledger.** This section generalizes the high-agreement tangent-star method away from the special field  $\mathbb{F}_{17^{32}}$ . The statements below are finite-grid coding statements. They do not use smoothness, except when one wants to combine them with the quotient-core and generated-field ledgers elsewhere in this repository. The line lower bound may be proved either by the RS locator construction of Theorem 6 or by the coordinate tangent-star construction below; the upper bounds use only the MDS zero-rigidity property and linearity of the code.

Throughout this section let  $C = \text{RS}[F, D, k]$  with  $D \subset F$ ,  $|D| = n$ , and dimension  $k$ . The upper-bound arguments below use only the MDS zero-rigidity property of Reed–Solomon codes, so they apply verbatim to any linear MDS code for which the relevant received words and parameter sets are defined. Put

$$R = n - k, \quad r = n - a.$$

Thus  $r$  is the integer Hamming radius and  $a = n - r$  is the agreement threshold. We only discuss the high-agreement range  $a \geq k + 1$ , equivalently  $0 \leq r \leq R - 1$ .

## 22.1 A coordinate tangent star

The next construction is useful because it removes all dependence on the particular field  $\mathbb{F}_{1732}$ , on multiplicative smoothness, and on the locator-polynomial model. It uses only coordinate values, linearity, and the MDS zero-rigidity of Reed–Solomon codes.

**Lemma 4** (coordinate tangent-star lower bound). *Let  $C = \text{RS}[F, D, k]$  with  $D \subset F$ . For every  $a$  with  $k + 1 \leq a \leq n$ ,*

$$\text{LD}_{\text{sw}}(C, a) \geq n - a + 1.$$

*If in addition  $2a - n - 1 \geq k$ , then the same construction gives*

$$\text{LD}_{\text{ca}}(C, a) \geq n - a + 1.$$

*Proof.* Choose  $A \subset D$  with  $|A| = a - 1$ , and write  $T = D \setminus A$ , so  $|T| = n - a + 1$ . Choose an injection  $\theta : T \hookrightarrow F$ , which exists because  $T \subseteq D \subset F$ . Define two received words  $f, g : D \rightarrow F$  by

$$f(x) = g(x) = 0 \quad (x \in A), \quad f(t) = -\theta(t), \quad g(t) = 1 \quad (t \in T).$$

For a point  $t \in T$ , the finite slope  $z = \theta(t)$  gives  $f + zg = 0$  on  $A \cup \{t\}$ , a support of size  $a$ . Hence this slope is explained by the zero codeword.

The same support is support-wise noncontained. Indeed, if both  $f$  and  $g$  were separately explained on  $A \cup \{t\}$ , then the explaining codeword for  $g$  would vanish on the  $a - 1 \geq k$  points of  $A$ , hence would be the zero codeword by the MDS property. But  $g(t) = 1$ , a contradiction.

The pair  $(f, g)$  is also globally far from  $C^2$  at agreement  $a$ . The same argument applied to the word  $g$  shows that no codeword of  $C$  agrees with  $g$  on any support of size  $a$ : such a support contains at least  $a - |T| = 2a - n - 1$  points of  $A$ , and in the high-agreement applications below we will have  $2a - n - 1 \geq k$ . Alternatively, for the lower bound in the line case one may use the RS locator construction of Theorem 6, which gives the no-loss CA witness directly for  $\text{RS}[F, D, k]$ . The displayed MCA lower bound holds unconditionally for  $a \geq k + 1$ , and the CA lower bound holds in the range  $2a - n - 1 \geq k$ , in particular throughout the exact staircase range  $3a - 2n \geq k$ .  $\square$

**Remark 8** (scope). *The support-wise MCA construction is coordinate-level. It uses MDS rigidity and the inclusion  $D \subset F$ . The no-loss CA part needs global farness; in the exact range below, the needed support-count condition follows from the staircase hypothesis.*

## 22.2 Uniform line, CA, and projective-slope threshold

**Theorem 11** (uniform high-agreement line staircase). *Let  $C = \text{RS}[F, D, k]$  with  $|D| = n$ , and let  $a = n - r$ . If*

$$3a - 2n \geq k \quad \text{equivalently} \quad r \leq \frac{n - k}{3},$$

*then*

$$\text{LD}_{\text{sw}}(C, a) = \text{LD}_{\text{ca}}(C, a) = \text{LD}_{\text{sw,proj}}(C, a) = r + 1 = n - a + 1.$$

*Here  $\text{LD}_{\text{sw,proj}}$  allows slopes in  $\mathbb{P}^1(F)$ .*

*Proof.* The finite-slope support-wise upper bound is Theorem 7, whose proof uses only MDS zero-rigidity and the common residual budget. The lower bound is Lemma 4, or, for Reed–Solomon codes, Theorem 6. No-loss CA-bad slopes are a subset of support-wise MCA-bad slopes, while the same tangent-star witness gives  $r + 1$  no-loss CA-bad slopes in the present range, so the CA numerator is also  $r + 1$ .

For projective slopes, suppose a projective bad set had size larger than  $r + 1$ . If there is a nonbad projective point, use it as the point at infinity; all projective bad points then become finite bad slopes, contradicting the finite-slope bound. If every point of  $\mathbb{P}^1(F)$  is bad, choose any point as infinity. The remaining  $|F|$  finite slopes are bad. Since  $|F| \geq n > r + 1$ , this again contradicts the finite-slope bound. Thus the projective numerator is also  $r + 1$ , and the finite lower-bound construction shows equality.  $\square$

### 22.3 Degree- $d$ curve MCA/CA

For an integer  $d \geq 1$ , a finite-parameter power curve is a received family

$$W_\gamma = f_0 + \gamma f_1 + \cdots + \gamma^d f_d, \quad \gamma \in F.$$

The support-wise curve-MCA predicate asks for parameters  $\gamma$  for which  $W_\gamma$  has an explanation on a support of size at least  $a$ , while the  $(d + 1)$ -tuple  $(f_0, \dots, f_d)$  is not simultaneously explained on that same support. The no-loss curve-CA predicate replaces support-wise noncontainment by global farness of the  $(d + 1)$ -tuple from  $C^{d+1}$ .

**Theorem 12** (degree- $d$  high-agreement curve staircase). *Let  $C = \text{RS}[F, D, k]$  with  $|D| = n$ , and let  $a = n - r$ . If*

$$(d + 2)a - (d + 1)n \geq k \quad \text{equivalently} \quad r \leq \frac{n - k}{d + 2},$$

then

$$\text{CurveLD}_{\text{sw}}^{(d)}(C, a) = \text{CurveLD}_{\text{ca}}^{(d)}(C, a) = \min\{|F|, d(r + 1)\}.$$

*Proof.* First prove the lower bound. Choose  $A \subset D$  with  $|A| = a - 1$  and put  $T = D \setminus A$ , so  $|T| = r + 1$ . Choose disjoint parameter sets  $\Gamma_t \subseteq F$ ,  $t \in T$ , with  $|\Gamma_t| \leq d$  and  $\sum_t |\Gamma_t| = \min\{|F|, d(r + 1)\}$ . For each  $t$ , choose a nonzero polynomial

$$h_t(Y) = h_{t,0} + h_{t,1}Y + \cdots + h_{t,d}Y^d$$

of degree exactly  $d$ , with leading coefficient nonzero, that vanishes on all parameters in  $\Gamma_t$ . Define received words by

$$f_i(x) = 0 \quad (x \in A), \quad f_i(t) = h_{t,i} \quad (t \in T), \quad 0 \leq i \leq d.$$

For  $\gamma \in \Gamma_t$ , the curve word  $W_\gamma$  vanishes on  $A \cup \{t\}$ , hence is explained by the zero codeword on a support of size  $a$ . The tuple is not simultaneously explained on that support: on the  $a - 1 \geq k$  points of  $A$ , every explaining endpoint codeword would have to be zero, but at  $t$  the coefficient vector of  $h_t$  is not zero. This gives the support-wise curve-MCA lower bound.

The same construction is globally far in the exact range. The last endpoint  $f_d$  is zero on  $A$  and nonzero on every point of  $T$ . Any support of size  $a$  contains at least  $a - |T| = 2a - n - 1$  points of  $A$ . The hypothesis  $r \leq (n - k)/(d + 2)$ , with  $d \geq 1$ , implies  $2a - n - 1 \geq k$ . Therefore any codeword agreeing with  $f_d$  on such a support would have to be the zero codeword, but the support must contain at least one point of  $T$ , where  $f_d \neq 0$ . Thus the endpoint tuple is not  $C^{d+1}$ -close at agreement  $a$ , proving the no-loss curve-CA lower bound.

For the upper bound, it is enough to rule out more than  $d(r + 1)$  bad parameters, because the trivial cap is  $|F|$ . If a curve had more than  $d(r + 1)$  bad parameters, then in particular it would have at least  $d + 1$  bad parameters. Select  $d + 1$  distinct bad parameters  $\gamma_0, \dots, \gamma_d$  and supports  $S_i$  of size at least  $a$ . Their intersection has size at least

$$(d + 1)a - dn.$$

On this intersection the  $d + 1$  explained codewords interpolate, as functions of  $\gamma$ , to codewords  $c_0, \dots, c_d \in C$  explaining the endpoints  $f_0, \dots, f_d$ . Let  $S_0$  be a maximal common support for this endpoint tuple, put  $b = |S_0|$ , and set  $m = n - b$ . Then

$$m \leq (d + 1)(n - a) = (d + 1)r.$$

For any further bad parameter, its explaining codeword must equal  $c_0 + \gamma c_1 + \dots + \gamma^d c_d$ , because

$$a + b - n \geq (d + 2)a - (d + 1)n \geq k.$$

Thus outside  $S_0$  it needs at least  $h = \max(1, a - b)$  zeros of the residual coordinate polynomials. By maximality of  $S_0$ , no residual coordinate polynomial is identically zero, and each has degree at most  $d$ , hence can be charged to at most  $d$  bad parameters. If  $b \geq a$ , then  $m \leq r$  and the count is at most  $dm \leq dr \leq d(r + 1)$ . If  $b < a$ , then  $h = a - b = m - r$  and  $r < m \leq (d + 1)r$ ; the charging bound gives

$$\#\{\text{bad parameters}\} \leq \left\lfloor \frac{dm}{m - r} \right\rfloor \leq d(r + 1).$$

The trivial parameter-space cap gives the additional upper bound  $|F|$ . Together with the lower bound this proves equality.  $\square$

## 22.4 Interleaved-list uniqueness

**Theorem 13** (uniform interleaved-list uniqueness). *Let  $C \subseteq F^D$  be a linear MDS code of dimension  $k$ , and let  $\mu \geq 1$ . If*

$$2a - n \geq k \quad \text{equivalently} \quad r \leq \frac{n - k}{2},$$

*then every  $\mu$ -row received word has at most one  $\mu$ -row codeword tuple agreeing with it on at least  $a$  common coordinates. Since equality is attained by taking the received word itself to be a code tuple,*

$$\Lambda_\mu(C, a) = 1.$$

*Proof.* If two  $\mu$ -row codeword tuples agree with the same received word on supports  $S$  and  $S'$  of size at least  $a$ , then they agree with each other on  $S \cap S'$ , whose size is at least  $2a - n \geq k$ . Row by row, the MDS property forces the two codewords to be equal. Hence the tuples are equal. The lower bound  $\Lambda_\mu(C, a) \geq 1$  is obtained by choosing the received word to be a codeword tuple.  $\square$

## 22.5 General finite-grid threshold calculator

Let  $\varepsilon_* = 2^{-128}$  and let  $Q$  be the denominator of the sampler or challenge field used by a particular ledger. Define the integer budget

$$B_Q = \lfloor \varepsilon_* Q \rfloor.$$

The high-agreement theorems above convert coding ledgers into integer arithmetic. For example, in the exact line range  $r \leq \lfloor R/3 \rfloor$ , a single affine line, no-loss CA, or projective-line numerator is

$$N_{\text{line}}(r) = r + 1.$$

Thus the line term alone is safe whenever  $r + 1 \leq B_Q$ , and the first unsafe radius is  $r = B_Q$ , provided  $B_Q \leq \lfloor R/3 \rfloor$ . If  $B_Q > \lfloor R/3 \rfloor$ , then the tangent-star method proves safety throughout the exact high-agreement range but does not locate the eventual threshold.

For one degree- $d$  finite-parameter curve term in its exact range  $r \leq \lfloor R/(d+2) \rfloor$ , the numerator is

$$N_{\text{curve},d}(r) = \min\{|F|, d(r+1)\}.$$

If all coding terms use the same denominator  $Q$ , a protocol ledger with curve terms of degrees  $d_1, \dots, d_s$  and  $\ell$  independent interleaved-list terms has high-agreement numerator

$$N_{\text{total}}(r) = \ell + \sum_{i=1}^s d_i(r+1),$$

valid for

$$r \leq \min \left\{ \left\lfloor \frac{R}{2} \right\rfloor \text{ if } \ell > 0, \min_i \left\lfloor \frac{R}{d_i + 2} \right\rfloor \right\}.$$

The ledger is certified at target  $\varepsilon_*$  whenever

$$N_{\text{total}}(r) \leq B_Q.$$

With different denominators  $Q_i$ , one should not combine numerators; the correct test is the rational inequality

$$\sum_i \frac{N_i(r)}{Q_i} + \varepsilon_{\text{query}} \leq \varepsilon_*.$$

**Corollary 7** (row-independent closed-form gates). *Let  $R = n - k$ ,  $B_Q = \lfloor Q/2^{128} \rfloor$ , and assume the relevant radius lies in the exact high-agreement range.*

1. *A single affine/projective line, no-loss CA, or support-wise MCA term is safe for every integer radius*

$$r \leq B_Q - 1.$$

*The first unsafe radius is  $r = B_Q$ , if  $B_Q \leq \lfloor R/3 \rfloor$ .*

2. *A single line/CA/MCA term plus one interleaved-list term over the same field is safe for*

$$r \leq B_Q - 2.$$

*The first unsafe radius is  $r = B_Q - 1$ , if  $B_Q - 1 \leq \lfloor R/3 \rfloor$ .*

3. *A single degree- $d$  curve term is safe for*

$$r \leq \left\lfloor \frac{B_Q}{d} \right\rfloor - 1.$$

*The first unsafe radius is  $r = \lfloor B_Q/d \rfloor$ , if this radius is at most  $\lfloor R/(d+2) \rfloor$ .*

4. *A single degree- $d$  curve term plus one interleaved-list term over the same field is safe for*

$$r \leq \left\lfloor \frac{B_Q - 1}{d} \right\rfloor - 1.$$

*The first unsafe radius is  $r = \lfloor (B_Q - 1)/d \rfloor$ , if this radius is in range.*

*Negative right-hand sides mean that even radius zero is not certified for that ledger and denominator.*



## 22.6 Prize-rate applicability

For the Proximity Prize rates  $\rho \in \{1/2, 1/4, 1/8, 1/16\}$ , with  $k \leq 2^{40}$ , the exact line range reaches integer radii

$$r \leq \frac{n-k}{3} = \frac{(1-\rho)n}{3} = \frac{1-\rho}{3\rho}k.$$

Thus, for a field with  $|F| \approx 2^\lambda$ , the high-agreement tangent gate can pin the line threshold only when roughly

$$2^{\lambda-128} \lesssim \frac{n-k}{3}.$$

At the maximal allowed dimension  $k = 2^{40}$ , this gives the following approximate largest field-bit sizes for which the line threshold can still be pinned inside the exact tangent range:

| $\rho$ | $(n-k)/3$ at $k = 2^{40}$ | $\lambda$ up to about |
|--------|---------------------------|-----------------------|
| 1/2    | $2^{40}/3$                | 166.4                 |
| 1/4    | $2^{40}$                  | 168.0                 |
| 1/8    | $7 \cdot 2^{40}/3$        | 169.2                 |
| 1/16   | $5 \cdot 2^{40}$          | 170.3                 |

For substantially larger challenge fields, such as  $2^{192}$  or  $2^{256}$ , the tangent numerator is below the  $2^{-128}$  budget throughout the exact high-agreement range for all prize-size rows. The threshold, if any, must then be controlled by the lower-agreement quotient-core, generated-field entropy, curve/folding, or aperiodic local-limit ledgers; the tangent-star method alone no longer pins it.

## 23 Towards-Prize Finite-Threshold Theorems

**Status: New-local finite theorem infrastructure plus subsumed strict264 proof record.**

The results in this section are deliberately finite and certificate-facing. They are meant to move the experimental ledger from lower-bound numerators toward the integer agreement staircase needed by the Proximity Prize. They do not prove the corrected Paper B local-limit conjecture. The later high-agreement tangent staircase supersedes the old strict264/265 search target as a threshold-pinning route for the  $\mathbb{F}_{1732}$ ,  $n = 512$ ,  $k = 256$  row; the strict264 material below remains useful as certificate language and as a record of quotient-core mechanisms.

For a received line  $\ell_z = f + zg$ , write

$$\text{Bad}_{f,g}(C, a)$$

for the set of finite slopes  $z \in F$  for which there is a support  $S \subseteq D$ ,  $|S| \geq a$ , and a codeword  $c \in C$  satisfying  $c|_S = (f + zg)|_S$ , but there is no pair of codewords  $c_f, c_g \in C$  with  $c_f|_S = f|_S$  and  $c_g|_S = g|_S$ . Put

$$\text{LD}_{\text{sw}}^{f,g}(C, a) = |\text{Bad}_{f,g}(C, a)|, \quad \text{LD}_{\text{sw}}(C, a) = \max_{f,g} \text{LD}_{\text{sw}}^{f,g}(C, a),$$

where the maximum is over the line family allowed by the sampler under consideration. The field paying the denominator is always  $q_{\text{line}}$ , not silently  $q_{\text{gen}}$  or  $q_{\text{chal}}$ .

**Proposition 8** (certificate-to- $\text{LD}_{\text{sw}}$  gate). *Let  $C = \text{RS}[F, D, k]$ , let  $a \geq k$ , and fix a received line  $f + zg$ . Suppose there is a finite set  $\mathcal{C}$  of tuples*

$$(z_i, S_i, P_i), \quad z_i \in F, \quad S_i \subseteq D, \quad |S_i| \geq a, \quad P_i \in F[X]_{<k},$$

*with the following properties:*

1. the slopes  $z_i$  are pairwise distinct;
2.  $P_i(x) = f(x) + z_i g(x)$  for every  $x \in S_i$ ;
3. there are no  $P_{i,f}, P_{i,g} \in F[X]_{<k}$  satisfying  $P_{i,f}|_{S_i} = f|_{S_i}$  and  $P_{i,g}|_{S_i} = g|_{S_i}$ .

Then

$$\text{LD}_{\text{sw}}^{f,g}(C, a) \geq |\mathcal{C}|, \quad \text{LD}_{\text{sw}}(C, a) \geq |\mathcal{C}|.$$

*Proof.* Each tuple certifies that the slope  $z_i$  is explained on a support of size at least  $a$ , and the third condition is exactly support-wise noncontainment on that support. Pairwise distinctness of the slopes prevents multiple tuples from paying for the same bad slope. Thus all  $z_i$ 's lie in  $\text{Bad}_{f,g}(C, a)$ , giving the first inequality. The second follows from the definition of  $\text{LD}_{\text{sw}}(C, a)$  as a maximum over lines.  $\square$

**Corollary 8** (strict264 certificate target). *For the row*

$$C = \text{RS}[\mathbb{F}_{17^{32}}, H, 256], \quad |H| = 512,$$

*a strict264 certificate consists of seven tuples in Proposition 8 with  $a = 264$ . Such a certificate proves*

$$\text{LD}_{\text{sw}}(C, 264) \geq 7.$$

*Proof.* Apply Proposition 8 with  $|\mathcal{C}| = 7$  and  $a = 264$ .  $\square$

**Proposition 9** (fixed-locator unique-slope principle). *Use the notation of Theorem 4. Fix  $a \geq k$ , put  $t = a - k$ ,  $j = n - a = r - t$ , and fix a squarefree  $D$ -split locator  $L_T$  of degree  $j$ . Let*

$$A_T = H_{t,j}(u)\ell_T, \quad B_T = H_{t,j}(v)\ell_T.$$

*If  $B_T \neq 0$ , then this locator contributes at most one finite support-wise noncontained slope. More precisely, a slope exists exactly when  $A_T$  is a scalar multiple of  $B_T$ , in which case the unique slope is*

$$z_T = -A_{T,m}/B_{T,m}$$

*for any coordinate  $m$  with  $B_{T,m} \neq 0$ , and the same value must be obtained from every nonzero coordinate of  $B_T$ . Consequently*

$$\text{LD}_{\text{sw}}^{f,g}(C, a) \leq \#\{T \subset D : |T| = n - a, L_T \text{ squarefree and } D\text{-split}, B_T \neq 0, A_T \in FB_T\}.$$

*Proof.* For the fixed locator, Theorem 4 gives the incidence equation

$$A_T + zB_T = 0$$

and the noncontainment condition  $B_T \neq 0$ . If  $B_T \neq 0$ , any solution  $z$  is forced by any coordinate where  $B_T$  is nonzero. Existence is exactly the consistency condition that all coordinates give the same scalar, equivalently  $A_T \in FB_T$ . Counting possible locators gives the final bound; different locators may still collide to the same slope, so the displayed quantity is an upper bound on distinct slopes.  $\square$

**Remark 9** (why this helps the 265 upper bound). *For the current  $n = 512, k = 256$  row, this proposition remains useful for duplicate-aware locator searches. The old proposed upper bound  $\text{LD}_{\text{sw}}(C, 265) \leq 6$  is now known to be false by Theorem 6, which already gives  $\text{LD}_{\text{sw}}(C, 265) \geq 248$ . Future upper-bound searches should therefore start beyond the exact tangent gate, at agreement 507 and higher, or explicitly isolate non-tangent mechanisms.*

**Theorem 14** (subfield confinement for base-valued lines). *Let  $K \subseteq F$  be a subfield, let  $D \subset K$ , and let  $C_F = \text{RS}[F, D, k]$ . Suppose  $f, g : D \rightarrow K$ , and let  $S \subseteq D$  have  $|S| \geq k$ . If  $z \in F \setminus K$  and there is a polynomial  $P \in F[X]_{<k}$  with*

$$P(x) = f(x) + zg(x) \quad (x \in S),$$

*then there are polynomials  $P_f, P_g \in K[X]_{<k}$  such that*

$$P_f|_S = f|_S, \quad P_g|_S = g|_S.$$

*Consequently, for  $K$ -valued received lines and agreement  $a \geq k$ , every support-wise noncontained finite slope lies in  $K$ .*

*Proof.* Choose a subset  $S_0 \subseteq S$  of size  $k$ . Interpolate the values of  $f$  and  $g$  on  $S_0$  to obtain unique polynomials  $P_f, P_g \in K[X]_{<k}$ . On  $S_0$ , the polynomial  $P_f + zP_g$  agrees with  $P$ . Since both have degree less than  $k$  and agree on  $k$  points, they are equal as polynomials over  $F$ . For every  $x \in S$ , therefore,

$$0 = f(x) + zg(x) - P(x) = (f(x) - P_f(x)) + z(g(x) - P_g(x)).$$

Both coefficients in the final expression lie in  $K$ . Since  $z \notin K$ , the elements  $1, z$  are linearly independent over  $K$ . Hence both coefficients vanish for every  $x \in S$ , proving the separate explanations of  $f$  and  $g$ . Thus any explained slope outside  $K$  is support-wise contained, so a noncontained slope must lie in  $K$ .  $\square$

**Remark 10** (extension-field warning). *Theorem 14 is intentionally one-sided. It only treats  $K$ -valued received lines. It does not rule out genuinely  $F$ -valued residue-line witnesses, and therefore must not be used as an extension-field MCA theorem without an additional hypothesis forcing the line to be  $K$ -valued.*

**Theorem 15** (seven-slope arithmetic gate over  $\mathbb{F}_{17^{32}}$ ). *Let  $q_{\text{line}} = 17^{32}$ . If a row over  $\mathbb{F}_{17^{32}}$  has  $\text{LD}_{\text{sw}}(C, 264) \geq 7$ , then at radius*

$$\delta = 1 - \frac{264}{512} = \frac{31}{64}$$

*its support-wise MCA error is strictly larger than  $2^{-128}$ . In particular, for the  $n = 512, k = 256$  row, any strict264 seven-slope certificate proves*

$$\varepsilon_{\text{mca}}(C, 31/64) > 2^{-128}.$$

*Proof.* The MCA sampler over finite slopes has denominator  $q_{\text{line}} = 17^{32}$ . Seven bad slopes therefore give error at least  $7/17^{32}$ . The exact integer comparison is

$$7 \cdot 2^{128} - 17^{32} = 14064973686101998399515954701426232831 > 0.$$

Thus  $7/17^{32} > 2^{-128}$ . Agreement 264 on a domain of size 512 corresponds to relative radius  $1 - 264/512 = 31/64$ , giving the claim.  $\square$

**Theorem 16** (one-step staircase pinning criterion). *Fix a finite-slope sampler of denominator  $q_{\text{line}}$  and a target  $\varepsilon_* = 2^{-128}$ . Let*

$$B(a) = \text{LD}_{\text{sw}}(C, a)$$

*for the allowed line family, and let  $\delta_a = 1 - a/n$ . If, for some integer  $a \geq k$ ,*

$$B(a) > \varepsilon_* q_{\text{line}} \quad \text{and} \quad B(a+1) \leq \varepsilon_* q_{\text{line}},$$

then the closed-ball agreement staircase is unsafe at radius  $\delta_a$  and safe on the next open interval below it, including the grid radius  $\delta_{a+1}$ . Equivalently, the largest grid radius certified safe is  $\delta_{a+1}$ , while the continuous supremum of certified safe radii is  $\delta_a$  and is not attained under the closed endpoint convention.

Before the tangent staircase was isolated, the special case

$$B(264) \geq 7, \quad B(265) \leq 6$$

would have pinned the finite row to an unsafe/safe transition between

$$\delta_{265} = 247/512 \quad \text{and} \quad \delta_{264} = 31/64.$$

For the  $\mathbb{F}_{17^{32}}$ ,  $n = 512$ ,  $k = 256$  row, the premise  $B(265) \leq 6$  is false by the moving-root tangent floor. The actual finite-slope staircase gate supplied by Corollary 3 is  $B(506) = 7$ ,  $B(507) = 6$ .

*Proof.* The function  $B(a)$  is nonincreasing in  $a$ , because raising the agreement threshold only removes valid supports. At radius  $\delta_a$ , the closed-ball predicate allows agreement at least  $a$ , so the error is  $B(a)/q_{\text{line}}$ , which is larger than  $\varepsilon_*$  by assumption. At radius  $\delta_{a+1}$ , the predicate requires agreement at least  $a + 1$ , so the error is  $B(a + 1)/q_{\text{line}}$ , which is at most  $\varepsilon_*$ .

More generally, any radius  $\delta < \delta_a$  sufficiently close to  $\delta_a$  has integer distance bound at most  $n - a - 1$ , hence agreement at least  $a + 1$ . Thus the closed endpoint at  $\delta_a$  is the first unsafe grid step, while the safe side is immediately below that endpoint. For  $q_{\text{line}} = 17^{32}$ , the inequalities

$$6 \cdot 2^{128} < 17^{32} < 7 \cdot 2^{128}$$

convert the printed integer conditions into the two displayed error inequalities, and  $1 - 265/512 = 247/512$ ,  $1 - 264/512 = 31/64$ .  $\square$

## 24 June 29 Finite-Slope and Interleaved-List PR Packets

This section records the mathematically useful parts of PRs #131–#135. The branches were not merged wholesale, because they were stale relative to the current Paper D v6 and public-site state. The integrated source files are the compact notes and verifiers listed in `experimental/notes/triage/pr-triage-2026-06-29.md`.

### 24.1 Boundary-off external-anchor normal form

PR #131, due to AllenGrahamHart, contributes a local M1 proof-program normal form. For a shadow  $S \subset D$  of size  $j - 1$  and an external anchor  $\beta \notin D$ , the one-outside locator satisfies

$$\ell_{S,\beta} = \ell_S^+ - \beta \ell_S^0.$$

Therefore the Hankel landing matrix

$$M_S(\beta) = [H(u)\ell_{S,\beta} \quad H(v)\ell_{S,\beta}]$$

has columns affine-linear in  $\beta$ , and the rank-one condition is cut out by quadratic minors in  $\beta$ . Thus, for fixed  $S$ , either a nonzero quadratic minor gives at most two external anchors, or all minors vanish identically and the branch is a ruled rank-one pencil. The later same-slope/root-slice note now classifies this ruled  $t = 2$  Hankel branch: it is inactive or fixed finite slope, and the fixed-slope case lifts to a boundary root-slice core  $H_{3,j-1}(u + z_0 v)\ell_S = 0$ . Therefore after fixed-slope boundary slices are charged, each shadow has at most two active non-ruled external anchors. This is recorded as PROVED-LOCAL / RULED-BRANCH CLASSIFIED / PROOF-PROGRAM / AUDIT, not as a proof of M1.

## 24.2 $a = 507$ adjacent bridge route cut

PR #132, due to Scott Hughes, records a useful non-promotion result for the  $\mathbb{F}_{17^{32}}, n = 512, k = 256$  row. The tangent-star theorem gives

$$\text{LD}_{\text{sw}}(C, a) = 513 - a \quad (a \geq 427),$$

so in particular

$$\text{LD}_{\text{sw}}(C, 507) = 6.$$

Any bridge that converted an adjacent line-plus-interleaved-list “6+1 = 7” ledger into one additional same-denominator finite-slope support-wise event would imply  $\text{LD}_{\text{sw}}(C, 507) \geq 7$ , contradicting the exact tangent-star bound. Hence the adjacent arithmetic remains a separate BRIDGE-NEEDED ledger and is not a new MCA row.

## 24.3 Interleaved-list lower row at agreement 326

PR #133, due to Scott Hughes, belongs to the interleaved-list track, not the MCA slope track. It proves high-agreement uniqueness

$$\Lambda_{\mu}(C, a) = 1 \quad (384 \leq a \leq 512)$$

and a support-occupancy route cut excluding seven witnesses for

$$373 \leq a \leq 383.$$

It also gives lower-bound constructions at agreements 291, 292, 320, and, most usefully for the board, a hybrid quotient-residual construction with

$$\Lambda_{\mu}(C, 326) \geq 7.$$

Since

$$6 \cdot 2^{128} < 17^{32} < 7 \cdot 2^{128},$$

this clears the separate interleaved-list numerical gate over denominator  $|F| = 17^{32}$ . The packet does not claim an MCA numerator, protocol soundness, ordinary list-decoding exactness, or an exact  $\delta_C^*$ .

## 24.4 Random simple-pole finite-slope floors

PR #134, due to Vadim Avdeev, proves a general random simple-pole lower bound. For  $C = \text{RS}[\mathbb{F}, D, k]$ ,  $|F| = q$ ,  $|D| = n$ ,  $\beta \in F \setminus D$ , and  $k < a \leq n$ , put

$$C_a = \binom{n}{a}, \quad s = a - k, \quad R_a = \sum_{v=0}^{a-k-1} \binom{a}{v} \binom{n-a}{v} q^{-v}.$$

Then there is a simple-pole line with at least

$$\left\lceil \frac{qC_a}{C_a + R_a q^s} \right\rceil$$

finite bad slopes at agreement  $a$ . In the row

$$F = \mathbb{F}_{17}[z]/(z^{32} - 3), \quad H = \langle z \rangle, \quad |H| = 512, \quad k = 256,$$

the verified floors are

| $a$ | finite bad-slope lower bound                    |
|-----|-------------------------------------------------|
| 257 | $17^{32}$                                       |
| 258 | $17^{32}$                                       |
| 259 | $17^{32} - 68,904$                              |
| 260 | 33,439,260,151,101,646,297,506,087,371,119,470. |

The direction is a genuine simple-pole direction: it is not a degree- $< 256$  codeword on any support of size  $> 256$ .

## 24.5 Coset-packet finite-slope floors for $a = 261, \dots, 288$

PR #135, also due to Vadim Avdeev, continues the finite-slope floor beyond the range where the random simple-pole second-moment bound is useful. The construction uses full dyadic cosets  $K_M \leq H$ , whose locators have the form

$$\prod_{x \in hK_M} (X - x) = X^M - h^M,$$

so unions of full  $K_M$ -cosets share low elementary coefficients. After fixing a small partial coset when needed, this produces common-prefix support families. A separating deep anchor then turns the corresponding numerator polynomials into distinct finite bad slopes.

For the same  $\mathbb{F}_{17^{32}}, n = 512, k = 256$  row, the certified floors are

| $a$         | finite bad-slope lower bound |
|-------------|------------------------------|
| 261 ... 263 | $\binom{63}{32}$             |
| 264         | $\binom{64}{33}$             |
| 265 ... 271 | $\binom{31}{16}$             |
| 272         | $\binom{32}{17}$             |
| 273 ... 287 | $\binom{15}{8}$              |
| 288         | $\binom{16}{9}$ .            |

Every displayed row beats the moving-root tangent floor  $513 - a$ , but these are still lower-bound rows only. They do not prove a safe-side theorem, an ordinary list-decoding theorem, an interleaved-list statement, or protocol soundness.

## 25 Non-Claims

This note does not claim:

- a proof of the corrected MCA conjecture;
- a positive theorem above corrected reserve;
- a  $q_{\text{gen}}$  local-limit theorem;
- a list, CA, MCA, line-decoding, SNARK, or protocol consequence;
- a large  $\Theta(p^2)$  counterpacket;

- a new prize-worthy numerator beyond the current Cycle116/119 52,747,567,092 finite/source-scoped record;
- priority for the BCHKS quotient-locator proof pattern.

The mathematical value is narrower. On the imported side, the file gives a provenance-safe dictionary and wrapper for using BCHKS without claiming it as new. On the local Paper B side, the possible large branch in the restricted quadratic-extension window must satisfy an explicit algebraic identity before it can even be surface-sized.

## A BibTeX Entry to Copy into Main Papers

```
@misc{BCHKS25,
  author = {Eli Ben-Sasson and Dan Carmon and Ulrich Habock and
            Swastik Kopparty and Shubhangi Saraf},
  title  = {On Proximity Gaps for Reed--Solomon Codes},
  year   = {2025},
  note   = {Manuscript dated November 11, 2025},
  url    = {https://www.math.toronto.edu/swastik/rs-proximity-gaps-2025.pdf}
}
```

## References

- [1] The Proximity Prize. *The Proximity Prize: Reed–Solomon Challenge*. Preliminary version, accessed June 26, 2026. <https://proximityprize.org/>.
- [2] Eli Ben-Sasson, Dan Carmon, Ulrich Haböck, Swastik Kopparty, and Shubhangi Saraf. *On Proximity Gaps for Reed–Solomon Codes*. Manuscript dated November 11, 2025. <https://www.math.toronto.edu/swastik/rs-proximity-gaps-2025.pdf>.