

# Homework Assignment 2

## Due Friday March 4th midnight

SDS 384-11 Theoretical Statistics

1. Remember Hoeffding's Lemma? We proved it with a weaker constant in class using a symmetrization type argument. Now we will prove the original version. Let  $X$  be a bounded r.v. in  $[a, b]$  such that  $E[X] = \mu$ . Let  $f(\lambda) = \log E[e^{\lambda(X-\mu)}]$ . Show that  $f''(\lambda) \leq (b-a)^2/4$ . Now use the fundamental theorem of calculus to write  $f(\lambda)$  in terms of  $f''(\lambda)$  and finish the argument.
2. Consider a r.v.  $X$  such that for all  $\lambda \in \mathbb{R}$

$$E[e^{\lambda X}] \leq e^{\frac{\lambda^2 \sigma^2}{2} + \lambda \mu} \quad (1)$$

Prove that:

- (a)  $E[X] = \mu$ .
  - (b)  $\text{var}(X) \leq \sigma^2$ .
  - (c) If the smallest value of  $\sigma$  satisfying the above equation is chosen, is it true that  $\text{var}(X) = \sigma^2$ ? Prove or give a counter example.
3. Given a positive semidefinite matrix  $Q \in \mathbb{R}^{n \times n}$ , consider  $Z = \sum_{i,j} Q_{ij} X_i X_j$ . When  $X_i \sim N(0, 1)$ , prove the Hanson-Wright inequality.

$$P(Z \geq \text{trace}(Q) + t) \leq \exp\left(-\min\left\{c_1 t / \|Q\|_{op}, c_2 t^2 / \|Q\|_F^2\right\}\right),$$

where  $\|Q\|_{op}$  and  $\|Q\|_F$  denote the operator and frobenius norms respectively. *Hint: The rotation-invariance of the Gaussian distribution and sub-exponential nature of  $\chi^2$ -variables could be useful.*

4. We will prove properties of subgaussian random variables here. Prove that:

- (a) Moments of a mean zero subgaussian r.v.  $X$  with variance proxy  $\sigma^2$  satisfy:

$$E[|X^k|] \leq k 2^{k/2} \sigma^k \Gamma(k/2), \quad (2)$$

where  $\Gamma$  is the gamma function.

- (b) If  $X$  is a mean 0 subgaussian r.v. with variance proxy  $\sigma^2$ , prove that,  $X^2 - E[X^2]$  is a subexponential  $(c_1 \sigma^2, c_2 \sigma^2)$  (we are using the  $(\nu, b)$  parametrization of subexponentials we did in class, so  $\nu^2$  is the variance proxy). Here  $c_1, c_2$  are positive constants.
- (c) Consider two independent mean zero subgaussian r.v.s  $X_1$  and  $X_2$  with variance proxies  $\sigma_1^2$  and  $\sigma_2^2$  respectively. Show that  $X_1 X_2$  is a subexponential r.v. with parameters  $(d_1 \sigma_1 \sigma_2, d_2 \sigma_1 \sigma_2)$ . Here  $d_1, d_2$  are positive constants.