

# SDS 385: Stat Models for Big Data

## Lecture 11: Bootstrap and subsampling

---

Purnamrita Sarkar  
Department of Statistics and Data Science  
The University of Texas at Austin  
<https://psarkar.github.io/teaching>

- So far we have talked about estimation, and ways to estimate statistical quantities quickly
- But often, you are interested in quantifying the variability of your estimate
- You can do this using the variance of your estimate or by producing a confidence interval
- What is a confidence interval?

# Confidence Interval

- Data  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} P$
- Some estimator  $\hat{\theta}$  of parameter of interest  $\theta$ .
- For some coverage  $\alpha$ , want to produce a lower and upper bound such that:

$$P\left(\hat{a} \leq \theta \leq \hat{b}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

# Confidence Interval

- Data  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} P$
- Some estimator  $\hat{\theta}$  of parameter of interest  $\theta$ .
- For some coverage  $\alpha$ , want to produce a lower and upper bound such that:

$$P\left(\hat{a} \leq \theta \leq \hat{b}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

- Say you know the distribution of  $(\hat{\theta} - \theta)/\hat{\sigma}$

# Confidence Interval

- Data  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} P$
- Some estimator  $\hat{\theta}$  of parameter of interest  $\theta$ .
- For some coverage  $\alpha$ , want to produce a lower and upper bound such that:

$$P\left(\hat{a} \leq \theta \leq \hat{b}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

- Say you know the distribution of  $(\hat{\theta} - \theta)/\hat{\sigma}$
- Then you will just return:

$$P\left(\hat{\theta} - \kappa_{\alpha}\hat{\sigma} \leq \theta \leq \hat{\theta} + \kappa_{1-\alpha}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

where  $\kappa_{\alpha}, \kappa_{1-\alpha}$  are the quantiles of  $(\hat{\theta} - \theta)/\hat{\sigma}$

- The distribution of  $(\hat{\theta} - \theta)/\hat{\sigma}$  depends on  $P$ .

# Confidence Interval

- Data  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} P$
- Some estimator  $\hat{\theta}$  of parameter of interest  $\theta$ .
- For some coverage  $\alpha$ , want to produce a lower and upper bound such that:

$$P\left(\hat{a} \leq \theta \leq \hat{b}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

- Say you know the distribution of  $(\hat{\theta} - \theta)/\hat{\sigma}$
- Then you will just return:

$$P\left(\hat{\theta} - \kappa_{\alpha}\hat{\sigma} \leq \theta \leq \hat{\theta} + \kappa_{1-\alpha}\hat{\sigma}\right) \geq 1 - 2\alpha,$$

where  $\kappa_{\alpha}, \kappa_{1-\alpha}$  are the quantiles of  $(\hat{\theta} - \theta)/\hat{\sigma}$

- The distribution of  $(\hat{\theta} - \theta)/\hat{\sigma}$  depends on  $P$ .
- Often this distribution is normal, but with unknown parameters.

# If we were omniscient

- The trouble is we don't know  $P$ .
- What will we do if we did know  $P$ ?

# If we were omniscient

- The trouble is we don't know  $P$ .
- What will we do if we did know  $P$ ?
- Draw  $B$  datasets of size  $n$  from  $P$



# If we were omniscient

- The trouble is we don't know  $P$ .
- What will we do if we did know  $P$ ?
- Draw  $B$  datasets of size  $n$  from  $P$
- For the  $i^{th}$  dataset, calculate  $\hat{\theta}^{(i)}$

## If we were omniscient

- The trouble is we don't know  $P$ .
- What will we do if we did know  $P$ ?
- Draw  $B$  datasets of size  $n$  from  $P$
- For the  $i^{th}$  dataset, calculate  $\hat{\theta}^{(i)}$
- Now get the distribution of  $\hat{\theta}^{(1)}, \dots, \hat{\theta}^{(B)}$  and get the C.I.

- The trouble is we don't know  $P$ .
- All we have at hand is the  $n$  datapoints  $x_1, \dots, x_n$

- The trouble is we don't know  $P$ .
- All we have at hand is the  $n$  datapoints  $x_1, \dots, x_n$
- So we put a  $1/n$  mass on each datapoint to get an empirical distribution  $\hat{P}$

- The trouble is we don't know  $P$ .
- All we have at hand is the  $n$  datapoints  $x_1, \dots, x_n$
- So we put a  $1/n$  mass on each datapoint to get an empirical distribution  $\hat{P}$
- Drawing  $n$  points from this distribution boils down to?

- The trouble is we don't know  $P$ .
- All we have at hand is the  $n$  datapoints  $x_1, \dots, x_n$
- So we put a  $1/n$  mass on each datapoint to get an empirical distribution  $\hat{P}$
- Drawing  $n$  points from this distribution boils down to?
- Sampling with replacement!

# Bootstrap: plug in principle

True model	Bootstrapped model
------------	--------------------

$\hat{\theta}$	$\hat{\theta}^*$
----------------	------------------

$\hat{\sigma}$	$\hat{\sigma}^*$
----------------	------------------

$\frac{\hat{\theta} - \theta}{\hat{\sigma}}$	$\frac{\hat{\theta}^* - \hat{\theta}}{\hat{\sigma}^*}$
--	--

# Empirical bootstrap

How do you estimate  $P$ ?

Empirical Bootstrap  $\hat{P} = \frac{1}{n} \sum_i \delta(x_i)$

Generate  $m$  samples  $(X_1^*, \dots, X_n^*)^{(j)}$ ,  $j = 1 : m$ .

Each giving a  $(\hat{\theta}^*, \hat{\sigma}^*)$  pair.

Compute the  $\kappa_\alpha$  quantile

of the distribution of  $\frac{\hat{\theta}^* - \hat{\theta}}{\hat{\sigma}^*}$

Parametric bootstrap  $\hat{P} = P_{\hat{\theta}}$



# Empirical bootstrap

Lets try the simplest setting with  $\theta = \mu := E[X_1]$

- What is the expectation of the bootstrapped mean  $\bar{X}^*$  given the data?

# Empirical bootstrap

Lets try the simplest setting with  $\theta = \mu := E[X_1]$

- What is the expectation of the bootstrapped mean  $\bar{X}^*$  given the data?

$$\begin{aligned} E[\bar{X}^* | X_1, \dots, X_n] &= E \left[ \frac{1}{n} \sum_i X_i^* | X_1, \dots, X_n \right] \\ &= E[X_1^* | X_1, \dots, X_n] \\ &= \sum_{i=1}^n X_i \times \frac{1}{n} = \bar{X} \end{aligned}$$

# Empirical bootstrap

Lets try the simplest setting with  $\theta = \mu := E[X_1]$

- What is the variance of the bootstrapped mean  $\bar{X}^*$  given the data?

# Empirical bootstrap

Lets try the simplest setting with  $\theta = \mu := E[X_1]$

- What is the variance of the bootstrapped mean  $\bar{X}^*$  given the data?

$$\begin{aligned}\text{var}[\bar{X}^* | X_1, \dots, X_n] &= \text{var} \left[ \frac{1}{n} \sum_{i=1}^n X_i^* | X_1, \dots, X_n \right] \\ &= \frac{1}{n} \text{var} [X_1^* | X_1, \dots, X_n] \\ &= \frac{1}{n} \left( E[(X_1^*)^2 | X_1, \dots, X_n] - \bar{X}^2 \right) \\ &= \frac{1}{n} \underbrace{\left( \frac{1}{n} \sum_i X_i^2 - \bar{X}^2 \right)}_{\text{Sample Variance}}\end{aligned}$$

# Empirical bootstrap

Lets try the simplest setting with  $\theta = \mu := E[X_1]$

- What is the variance of the bootstrapped mean  $\bar{X}^*$  given the data?

$$\begin{aligned}\text{var}[\bar{X}^* | X_1, \dots, X_n] &= \text{var} \left[ \frac{1}{n} \sum_{i=1}^n X_i^* | X_1, \dots, X_n \right] \\ &= \frac{1}{n} \text{var} [X_1^* | X_1, \dots, X_n] \\ &= \frac{1}{n} \left( E[(X_1^*)^2 | X_1, \dots, X_n] - \bar{X}^2 \right) \\ &= \frac{1}{n} \underbrace{\left( \frac{1}{n} \sum_i X_i^2 - \bar{X}^2 \right)}_{\text{Sample Variance}}\end{aligned}$$

- This makes sense, since the sample variance converges to the true variance, and we all know that the variance of  $\bar{X}$  is exactly  $\sigma^2/n$

But what's the point then? Can't you just do some Taylor approximation or something, and get the final distribution?

But what's the point then? Can't you just do some Taylor approximation or something, and get the final distribution? Not always, let's take the median.

- What is the asymptotic distribution of the median of  $n$  i.i.d r.v.s drawn from  $P$ ?

But whats the point then? Can't you just do some Taylor approximation or something, and get the final distribution? Not always, lets take the median.

- What is the asymptotic distribution of the median of  $n$  i.i.d r.v.s drawn from  $P$ ?
- Its a normal, of course, like a lot of other estimators.



But whats the point then? Can't you just do some Taylor approximation or something, and get the final distribution? Not always, lets take the median.

- What is the asymptotic distribution of the median of  $n$  i.i.d r.v.s drawn from  $P$ ?
- Its a normal, of course, like a lot of other estimators.
- With variance  $\frac{1}{4nf(\tilde{\mu})^2}$ , where  $\tilde{\mu}$  is the population median and  $f$  is the density of  $P$

But whats the point then? Can't you just do some Taylor approximation or something, and get the final distribution? Not always, lets take the median.

- What is the asymptotic distribution of the median of  $n$  i.i.d r.v.s drawn from  $P$ ?
- Its a normal, of course, like a lot of other estimators.
- With variance  $\frac{1}{4nf(\tilde{\mu})^2}$ , where  $\tilde{\mu}$  is the population median and  $f$  is the density of  $P$
- If we don't know  $P$ , we can't evaluate the above.

Does it always work?

Does it always work? Lets try the maximum of  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} U([0, \theta])$

- What is the true limiting distribution?

Does it always work? Lets try the maximum of  $X_1, \dots, X_n \stackrel{\text{iid}}{\sim} U([0, \theta])$

- What is the true limiting distribution?

$$P\left(\frac{n(\theta - X_{(n)})}{\theta} > x\right) = P\left(X_{(n)} \leq \theta(1 - x/n)\right) = (1 - x/n)^n \rightarrow e^{-x}$$

- The bootstrapped limiting distribution

$$P\left(\frac{n(X_{(n)} - X_{(n)}^*)}{X_{(n)}} = 0\right) = P(X_{(n)}^* = X_{(n)}) = (1 - (1 - 1/n)^n) \rightarrow 1 - 1/e$$

Does it always work?

Does it always work?

- Rule of thumb: when the asymptotic distribution is normal.
- Another con is it will take forever if  $n$  is large, even if you parallelize
- What do you do when its not?

# Subsampling

- Starts with the realization that instead of drawing with replacement, its better to draw without replacement smaller samples
- This is in some sense, a more honest representation or approximation of the unknow distribution  $P$



# Subsampling

- Starts with the realization that instead of drawing with replacement, its better to draw without replacement smaller samples
- This is in some sense, a more honest representation or approximation of the unknow distribution  $P$ 
  - Draw  $B$  size  $b$  subsamples **without** replacement
  - For each, compute your estimator  $\hat{\theta}$
  - Now get confidence intervals or variance of this distribution

# Subsampling

- Starts with the realization that instead of drawing with replacement, its better to draw without replacement smaller samples
- This is in some sense, a more honest representation or approximation of the unknow distribution  $P$ 
  - Draw  $B$  size  $b$  subsamples **without** replacement
  - For each, compute your estimator  $\hat{\theta}$
  - Now get confidence intervals or variance of this distribution
- But now everything is on a different scale!
- For example, the standard dev. of the mean decays at a rate of  $1/\sqrt{n}$
- If you use subsampling, the numbers you will get will be  $1/\sqrt{b}$

# Subsampling

- Starts with the realization that instead of drawing with replacement, its better to draw without replacement smaller samples
- This is in some sense, a more honest representation or approximation of the unknow distribution  $P$ 
  - Draw  $B$  size  $b$  subsamples **without** replacement
  - For each, compute your estimator  $\hat{\theta}$
  - Now get confidence intervals or variance of this distribution
- But now everything is on a different scale!
- For example, the standard dev. of the mean decays at a rate of  $1/\sqrt{n}$
- If you use subsampling, the numbers you will get will be  $1/\sqrt{b}$
- What to do? You will need to analytically correct the variability.

# Subsampling - pros and cons

## Pros

- Very fast, specially you have a super-linear estimation algorithm
- Works for statistics which bootstrap doesnt work for, i.e. requires far less conditions, as long as  $b$  grows to infinity with  $n$ , but at a slower rate.


## Cons

- Very sensitive to the choice of  $b$  (next two slides)
- You need to know the scaling factor to correct for using  $b < n$

## Subsampling - cons

- Multivariate linear regression with  $d = 100$  and  $n = 50,000$  on synthetic data.
- $x$  coordinates sampled independently from StudentT(3).
- $y = w^T x + \varepsilon$ , where  $w$  in  $\mathbb{R}^d$  is a fixed weight vector and  $\varepsilon$  is Gaussian noise.
- Estimate  $\theta_n = w_n$  in  $\mathbb{R}^d$  via least squares.
- Compute a marginal confidence interval for each component of  $w_n$  and assess accuracy via relative mean (across components) absolute deviation from true confidence interval size.
- For subsampling, use  $b(n) = n^\gamma$  for various values of  $\gamma$ .
- Similar results obtained with Normal and Gamma data generating distributions, as well as if estimate a misspecified model.

## Subsampling - cons



## Next class

Bag of little bootstraps