

SDS 321: Introduction to Probability and Statistics

Lecture 10: Expectation and Variance



Purnamrita Sarkar
Department of Statistics and Data Science
The University of Texas at Austin

www.cs.cmu.edu/~psarkar/teaching

The Poisson random variable


- ▶ A Poisson random variable takes non-negative integers as values. It has a nonnegative parameter λ .
- ▶ $P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$, for $k = 0, 1, 2, \dots$
- ▶ $\sum_{k=0}^{\infty} P(X = k) = e^{-\lambda} (1 + \lambda + \frac{\lambda^2}{2!} + \frac{\lambda^3}{3!} + \dots) = 1$. (Exponential series!)

The PMF is monotonically decreasing for $\lambda=0.5$




The PMF is increasing and then decreasing for $\lambda=3$


Poisson random variable



Binomial(5,0.6)



Binomial(100,0.03)



Poisson(3)

- ▶ When n is very large and p is very small, a binomial random variable can be well approximated by a Poisson with $\lambda = np$.
- ▶ In the above figure we increased n and decreased p so that $np = 3$.
- ▶ See how close the PMF's of the Binomial(100,0.03) and Poisson(3) are!
- ▶ More formally, we see that $\binom{n}{k} p^k (1-p)^{n-k} \approx \frac{e^{-\lambda} \lambda^k}{k!}$ when n is large, k is fixed, and p is small and $\lambda = np$.

Example

Assume that on a given day 1000 cars are out in Austin. On an average three out of 1000 cars run into a traffic accident per day.

1. What is the probability that we see at least two accidents in a day?
4. If you know there is at least one accident, what is the probability that the total number of accidents is at least two?

Example

Assume that on a given day 1000 cars are out in Austin. On an average three out of 1000 cars run into a traffic accident per day.

1. What is the probability that we see at least two accidents in a day?
2. Use poisson approximation!
- 3.
4. If you know there is at least one accident, what is the probability that the total number of accidents is at least two?

Example

Assume that on a given day 1000 cars are out in Austin. On an average three out of 1000 cars run into a traffic accident per day.

1. What is the probability that we see at least two accidents in a day?
2. Use poisson approximation!
3. $P(X \geq 2) = 1 - P(X = 0) - P(X = 1) = 1 - e^{-3}(1 + 3) = 0.8$
4. If you know there is at least one accident, what is the probability that the total number of accidents is at least two?

Example

Assume that on a given day 1000 cars are out in Austin. On an average three out of 1000 cars run into a traffic accident per day.

1. What is the probability that we see at least two accidents in a day?
2. Use poisson approximation!
3. $P(X \geq 2) = 1 - P(X = 0) - P(X = 1) = 1 - e^{-3}(1 + 3) = 0.8$
4. If you know there is at least one accident, what is the probability that the total number of accidents is at least two?
5. $P(X \geq 1) = 1 - P(X = 0) = 1 - e^{-3} = 0.950$.
 $P(X \geq 2|X \geq 1) = P(X \geq 2)/P(X \geq 1) = 0.8/0.950 = 0.84$

Conditional PMF

- ▶ We have talked about conditional probabilities.
- ▶ We can also talk about conditional PMF's. Let A be an event with positive probability.
- ▶ The rules are the same. $P(X = x|A) = \frac{P(\{X = x\} \cap A)}{P(A)}$
- ▶ The conditional PMF is a valid PMF. $\sum_x P(X = x|A) = 1$

Conditional PMF-Example

- ▶ $X \sim \text{Geometric}(p)$
- ▶ What is $P(X = k | X > 1)$ for different values of k ?

Conditional PMF-Example

- ▶ $X \sim \text{Geometric}(p)$
- ▶ What is $P(X = k | X > 1)$ for different values of k ?
- ▶


$$P(X = k | X > 1) = \begin{cases} 0 & \text{If } k = 1 \\ \frac{P(X=k)}{P(X>1)} & \text{Otherwise} \\ = \frac{(1-p)^{k-1}p}{(1-p)} \\ = (1-p)^{k-2}p = P(X = k-1) \end{cases}$$

Cumulative Distribution Functions

- ▶ For any random variable the cumulative distribution function is defined as:

$$F_X(a) = \sum_{x \leq a} p(x)$$

- ▶ Can you work out the PMF of the following random variable?



Function of a random variable

- ▶ A function of a random variable is also a random variable.
- ▶ Let X be the number of heads in 5 fair coin tosses.
- ▶ We know that X has the Binomial($5, 1/2$) distribution.
- ▶ Define $Y = X \bmod 4$. Whats its PMF?

Function of a random variable

- ▶ A function of a random variable is also a random variable.
- ▶ Let X be the number of heads in 5 fair coin tosses.
- ▶ We know that X has the Binomial(5,1/2) distribution.
- ▶ Define $Y = X \bmod 4$. Whats its PMF?

$$\text{▶ } Y = \begin{cases} 0 & X \in \{0, 4\} \\ 1 & X \in \{1, 5\} \\ 2 & X = 2 \\ 3 & X = 3 \end{cases}$$

Function of a random variable

- ▶ A function of a random variable is also a random variable.
- ▶ Let X be the number of heads in 5 fair coin tosses.
- ▶ We know that X has the Binomial(5,1/2) distribution.
- ▶ Define $Y = X \bmod 4$. Whats its PMF?

$$\text{▶ } Y = \begin{cases} 0 & X \in \{0, 4\} \\ 1 & X \in \{1, 5\} \\ 2 & X = 2 \\ 3 & X = 3 \end{cases}$$

- ▶ Lets write down the PMF of Y .

- ▶ $P(Y = 0) = P(X = 0) + P(X = 4) = (1/2)^5 + \binom{5}{4}(1/2)^5$
- ▶ $P(Y = 1) = P(X = 1) + P(X = 5)$...and so on.

Function of a random variable

- ▶ A function of a random variable is also a random variable.
- ▶ Let X be the number of heads in 5 fair coin tosses.
- ▶ We know that X has the Binomial(5,1/2) distribution.
- ▶ Define $Y = X \bmod 4$. Whats its PMF?

$$\text{▶ } Y = \begin{cases} 0 & X \in \{0, 4\} \\ 1 & X \in \{1, 5\} \\ 2 & X = 2 \\ 3 & X = 3 \end{cases}$$

- ▶ Lets write down the PMF of Y .

- ▶ $P(Y = 0) = P(X = 0) + P(X = 4) = (1/2)^5 + \binom{5}{4}(1/2)^5$
- ▶ $P(Y = 1) = P(X = 1) + P(X = 5) \dots$ and so on.
- ▶ More formally, if $Y = g(X)$ then we have:

$$p_Y(y) = \sum_{\{x|g(x)=y\}} p_X(x).$$

Function of a random variable-examples

- ▶ $X \sim \text{Bernoulli}(p)$.
 - ▶ What is the PMF of X^2 ?
 - ▶ What is the PMF of X^3 ?
- ▶ $X \sim \text{Binomial}(n, p)$.
 - ▶ What is the distribution of $n - X$?

Summing up

- ▶ Last time, we looked at the probability of a random variable taking on a given value:

$$p_X(x) = P(X = x)$$

- ▶ We also looked at plots of various PMFs of Uniform, Bernoulli, Binomial, Poisson and Geometric.
- ▶ Often, we want to make predictions for the value of a random variable
 - ▶ How many heads do I expect to get if I toss a fair coin 10 times?
 - ▶ How many lottery tickets should Alice expect buy until she wins the jackpot?
- ▶ We may also be interested in how far, on average, we expect our random variable to be from these predictions.
- ▶ Today we will talk about means and variances of these random variables.

Mean

You want to calculate average grade points from hw1. You know that 20 students got 30/30, 30 students got 25/30, and 50 students got 20/30.

Whats the average?

- ▶ The average grade point is

$$\frac{30 \times 20 + 25 \times 30 + 20 \times 50}{100} = 30 \times 0.2 + 25 \times 0.3 + 20 \times 0.5$$

- ▶ Let X be a random variable which represents grade points of hw1.
- ▶ How will you calculate $P(X = 30)$?
 - ▶ See how many out of 100 students got 30 out of 30 points.
 - ▶ $P(X = 30) \approx 0.2$
 - ▶ $P(X = 25) \approx 0.3$
 - ▶ $P(X = 20) \approx 0.5$
- ▶ So roughly speaking,
average grade $\approx 30 \times P(X = 30) + 25 \times P(X = 25) + 20 \times P(X = 20)$

Expectation

We define the expected value (or expectation or mean) of a discrete random variable X by

$$E[X] = \sum_x xP(X = x).$$

- ▶ X is a Bernoulli random variable with the following PMF:

$$P(X = x) = \begin{cases} p & X = 1 \\ 1 - p & X = 0 \end{cases}$$

So $E[X] =$.

Expectation

We define the expected value (or expectation or mean) of a discrete random variable X by

$$E[X] = \sum_x xP(X = x).$$

- ▶ X is a Bernoulli random variable with the following PMF:

$$P(X = x) = \begin{cases} p & X = 1 \\ 1 - p & X = 0 \end{cases}$$

So $E[X] = 1 \times p + 0 \times (1 - p) = p$.

- ▶ Expectation of a Bernoulli random variable is just the probability that it is one.
- ▶ You will also see notation like μ_X .

Expectation: example

You are tossing 4 fair coins independently. Let X denote the number of heads. What is $E[X]$?

- ▶ Any guesses? Well, on an average we should see about 2 coin tosses. No?
- ▶ Lets write down the PMF first.

Expectation: example

You are tossing 4 fair coins independently. Let X denote the number of heads. What is $E[X]$?

- ▶ Any guesses? Well, on an average we should see about 2 coin tosses. No?
- ▶ Lets write down the PMF first.

$$\text{▶ } P(X = x) = \begin{cases} 1/2^4 & X = 0 \\ 4/2^4 & X = 1 \\ 6/2^4 & X = 2 \\ 4/2^4 & X = 3 \\ 1/2^4 & X = 4 \end{cases}$$

Expectation: example

You are tossing 4 fair coins independently. Let X denote the number of heads. What is $E[X]$?

- ▶ Any guesses? Well, on an average we should see about 2 coin tosses. No?
- ▶ Lets write down the PMF first.

$$\text{▶ } P(X = x) = \begin{cases} 1/2^4 & X = 0 \\ 4/2^4 & X = 1 \\ 6/2^4 & X = 2 \\ 4/2^4 & X = 3 \\ 1/2^4 & X = 4 \end{cases}$$


$$\text{▶ So } E[X] = \frac{4}{2^4} + 2\frac{6}{2^4} + 3\frac{4}{2^4} + 4\frac{1}{2^4} = \frac{32}{16} = 2.$$

Expectation of a function of a random variable

Lets say you want to compute $E[g(X)]$. Example, I know average temperature in Fahrenheit, but I now want it in Celsius.

- ▶ $E[g(X)] = \sum_x g(x)P(X = x)$.
- ▶ Follows from the definition of PMF of functions of random variables.
- ▶ Look at page 15 of Bersekas-Tsitsiklis and derive it at home!
- ▶ So $E[X^2] = \sum_x x^2 P(X = x)$. **Second moment of X**
- ▶ So $E[X^3] = \sum_x x^3 P(X = x)$. **Third moment of X**
- ▶ So $E[X^k] = \sum_x x^k P(X = x)$. **k^{th} moment of X**
- ▶ We are assuming "under the rugs" that all these expectations are well defined.

Expectation



- ▶ Think of expectation as center of gravity of the PMF or a representative value of X .
- ▶ How about the spread of the distribution? Is there a number for it?

Variance

Often, you may want to know the spread or variation of the grade points for homework1.

- ▶ If everyone got the same grade point, then variation is?
- ▶ If there is high variation, then we know that many students got grade points very different from the average grade point in class.
- ▶ Formally we measure this using variance of a random variable X .
- ▶ $\text{var}(X) = E[(X - E[X])^2]$ or $E[(X - \mu)^2]$.
- ▶ The standard deviation of X is given by $\sigma_X = \sqrt{\text{var}X}$.
- ▶ Its easier to think about σ_X , since its on the same scale.
- ▶ The grade points have average 20 out of 30 with a standard deviation of 5 grade points. Roughly this means, most of the students have grade points within $[20 - 5, 20 + 5]$.

Variance

Often, you may want to know the spread or variation of the grade points for homework1.

- ▶ If everyone got the same grade point, then variation is? 0
- ▶ If there is high variation, then we know that many students got grade points very different from the average grade point in class.
- ▶ Formally we measure this using variance of a random variable X .
- ▶ $\text{var}(X) = E[(X - E[X])^2]$ or $E[(X - \mu)^2]$.
- ▶ The standard deviation of X is given by $\sigma_X = \sqrt{\text{var}X}$.
- ▶ Its easier to think about σ_X , since its on the same scale.
- ▶ The grade points have average 20 out of 30 with a standard deviation of 5 grade points. Roughly this means, most of the students have grade points within $[20 - 5, 20 + 5]$.

Computing the variance

- ▶ $\text{var}(X) = E[(X - \mu)^2] = \sum_x (x - \mu)^2 P(X = x)$
- ▶ Always remember! $E[X]$ or $E[g(X)]$ **do not depend on any particular value of x** . You can treat it as a constant. It only depends on the PMF of X .
- ▶ This can actually be made simpler.
- ▶ $\text{var}(X) = E[X^2] - \mu^2$.
- ▶ So you can calculate $E[X^2]$ (second moment) and then subtract the square of $E[X]$ to get the variance!

A tiny bit of algebra

$$\text{var}(X) =$$

A tiny bit of algebra

$$\text{var}(X) = \sum_x (x - \mu)^2 P(X = x) = \sum_x (x^2 + \mu^2 - 2x\mu) P(X = x)$$

A tiny bit of algebra

$$\text{var}(X) = \sum_x (x - \mu)^2 P(X = x) = \sum_x (x^2 + \mu^2 - 2x\mu) P(X = x)$$

$$= \sum_x x^2 P(X = x) + \sum_x \mu^2 P(X = x) - 2 \sum_x x\mu P(X = x)$$

A tiny bit of algebra

$$\text{var}(X) = \sum_x (x - \mu)^2 P(X = x) = \sum_x (x^2 + \mu^2 - 2x\mu) P(X = x)$$

$$= \sum_x x^2 P(X = x) + \sum_x \mu^2 P(X = x) - 2 \sum_x x\mu P(X = x)$$

$$= \sum_x x^2 P(X = x) + \underbrace{\mu^2 \sum_x P(X = x)}_{\mu^2 \text{ does not depend on } x} - 2\mu \underbrace{\sum_x x P(X = x)}_{\text{This is } \mu}$$

A tiny bit of algebra

$$\begin{aligned}\text{var}(X) &= \sum_x (x - \mu)^2 P(X = x) = \sum_x (x^2 + \mu^2 - 2x\mu) P(X = x) \\&= \sum_x x^2 P(X = x) + \sum_x \mu^2 P(X = x) - 2 \sum_x x\mu P(X = x) \\&= \sum_x x^2 P(X = x) + \underbrace{\mu^2 \sum_x P(X = x)}_{\mu^2 \text{ does not depend on } x} - 2\mu \underbrace{\sum_x x P(X = x)}_{\text{This is } \mu} \\&= \sum_x x^2 P(X = x) + \mu^2 - 2\mu^2 = E[X^2] - \mu^2\end{aligned}$$

Some simple rules– Expectation

Say you are looking at a linear function (or transformation) of your random variable X .

- ▶ $Y = aX + b$. Remember celsius to fahrenheit conversions? They are linear too!
- ▶ $E[Y] = E[aX + b] = aE[X] + b$, as simple as that! why?

$$\begin{aligned}\text{▶ } E[aX + b] &= \sum_x (ax + b)P(X = x) \\ &= a \sum_x xP(X = x) + b \sum_x P(X = x)\end{aligned}$$

Some simple rules– Expectation

Say you are looking at a linear function (or transformation) of your random variable X .

- ▶ $Y = aX + b$. Remember celsius to fahrenheit conversions? They are linear too!
- ▶ $E[Y] = E[aX + b] = aE[X] + b$, as simple as that! why?

$$\begin{aligned}\text{▶ } E[aX + b] &= \sum_x (ax + b)P(X = x) \\ &= a \sum_x xP(X = x) + b \sum_x P(X = x) \\ &= aE[X] + b\end{aligned}$$

Some simple rules– Expectation

How about $E[Y]$ for $Y = aX^2 + bX + c$?

- ▶ $E[Y] = E[aX^2 + bX + c] = aE[X^2] + bE[X] + c$, as simple as that!
why?

- ▶
$$\begin{aligned} E[aX^2 + bX + c] &= \sum_x (ax^2 + bx + c)P(X = x) \\ &= a \sum_x x^2 P(X = x) + b \sum_x x P(X = x) + c \sum_x P(X = x) \end{aligned}$$

Some simple rules– Expectation

How about $E[Y]$ for $Y = aX^2 + bX + c$?

- ▶ $E[Y] = E[aX^2 + bX + c] = aE[X^2] + bE[X] + c$, as simple as that!
why?

- ▶
$$\begin{aligned} E[aX^2 + bX + c] &= \sum_x (ax^2 + bx + c)P(X = x) \\ &= a \sum_x x^2 P(X = x) + b \sum_x x P(X = x) + c \sum_x P(X = x) \\ &= aE[X^2] + bE[X] + c \end{aligned}$$

Some simple rules– Expectation

How about $E[Y]$ for $Y = aX^2 + bX + c$?

- ▶ $E[Y] = E[aX^2 + bX + c] = aE[X^2] + bE[X] + c$, as simple as that!
why?

- ▶
$$\begin{aligned} E[aX^2 + bX + c] &= \sum_x (ax^2 + bx + c)P(X = x) \\ &= a \sum_x x^2 P(X = x) + b \sum_x x P(X = x) + c \sum_x P(X = x) \\ &= aE[X^2] + bE[X] + c \end{aligned}$$

- ▶ $Y = aX^3 + bX^2 + cX + d$. Can you guess what $E[Y]$ is?

Some simple rules– Expectation

How about $E[Y]$ for $Y = aX^2 + bX + c$?

- ▶ $E[Y] = E[aX^2 + bX + c] = aE[X^2] + bE[X] + c$, as simple as that!
why?


$$\begin{aligned}\text{▶ } E[aX^2 + bX + c] &= \sum_x (ax^2 + bx + c)P(X = x) \\ &= a \sum_x x^2 P(X = x) + b \sum_x x P(X = x) + c \sum_x P(X = x) \\ &= aE[X^2] + bE[X] + c\end{aligned}$$

- ▶ $Y = aX^3 + bX^2 + cX + d$. Can you guess what $E[Y]$ is?

$$E[Y] = aE[X^3] + bE[X^2] + cE[X] + d.$$



Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?





Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?



Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?



- ▶ Intuitively? Well you are just shifting everything by the same number.
- ▶ So? the spread of the numbers should stay the same!
- ▶ Prove it at home.

Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?

Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?

Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?

► Proof:

►
$$\text{var}(X + b) = E[(X + b)^2] - (E[X + b])^2$$

Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?

► Proof:

$$\begin{aligned}\text{var}(X + b) &= E[(X + b)^2] - (E[X + b])^2 \\ &= E[X^2 + 2bX + b^2] - (E[X] + b)^2\end{aligned}$$

Some simple rules– Variance

Let $Y = X + b$. What is $\text{var}(Y)$?

► Proof:

$$\begin{aligned}\text{var}(X + b) &= E[(X + b)^2] - (E[X + b])^2 \\ &= E[X^2 + 2bX + b^2] - (E[X] + b)^2 \\ &= E[X^2] + 2bE[X] + b^2 - ((E[X])^2 + 2bE[X] + b^2)\end{aligned}$$

Some simple rules– Variance


Let $Y = X + b$. What is $\text{var}(Y)$?

► Proof:

$$\begin{aligned}\text{var}(X + b) &= E[(X + b)^2] - (E[X + b])^2 \\ &= E[X^2 + 2bX + b^2] - (E[X] + b)^2 \\ &= E[X^2] + 2bE[X] + b^2 - ((E[X])^2 + 2bE[X] + b^2) \\ &= E[X^2] - (E[X])^2 = \text{var}(X)\end{aligned}$$



Some simple rules– Variance

Let $Y = aX$. What is $\text{var}(Y)$?





Some simple rules– Variance

Let $Y = aX$. What is $\text{var}(Y)$?



Some simple rules– Variance

Let $Y = aX$. What is $\text{var}(Y)$?



- ▶ Intuitively? Well you are just scaling everything by the same number.
- ▶ So? the spread should increase if $a > 1$!

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

- ▶ Proof:

- ▶

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

► Proof:

►
$$\text{var}(aX) = E[(aX)^2] - (E[aX])^2$$

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

► Proof:

$$\begin{aligned}\text{var}(aX) &= E[(aX)^2] - (E[aX])^2 \\ &= E[a^2 X^2] - (aE[X])^2\end{aligned}$$

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

► Proof:

$$\begin{aligned}\text{var}(aX) &= E[(aX)^2] - (E[aX])^2 \\ &= E[a^2 X^2] - (aE[X])^2 \\ &= a^2 E[X^2] - a^2 (E[X])^2\end{aligned}$$

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

► Proof:

$$\begin{aligned}\text{var}(aX) &= E[(aX)^2] - (E[aX])^2 \\ &= E[a^2 X^2] - (aE[X])^2 \\ &= a^2 E[X^2] - a^2 (E[X])^2 \\ &= a^2 (E[X^2] - (E[X])^2) = a^2 \text{var}(X)\end{aligned}$$

Some simple rules– Variance

Let $Y = aX$. Turns out $\text{var}(Y) = a^2 \text{var}(X)$.

► Proof:

$$\begin{aligned}\text{var}(aX) &= E[(aX)^2] - (E[aX])^2 \\ &= E[a^2 X^2] - (aE[X])^2 \\ &= a^2 E[X^2] - a^2 (E[X])^2 \\ &= a^2 (E[X^2] - (E[X])^2) = a^2 \text{var}(X)\end{aligned}$$

► In general we can show that $\text{var}(aX + b) = a^2 \text{var}(X)$.

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values:

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values: 0 and 1
 - ▶ $P(X^2 = 1)$

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values: 0 and 1
 - ▶ $P(X^2 = 1) = P(X = 1) = p$.

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values: 0 and 1
 - ▶ $P(X^2 = 1) = P(X = 1) = p$. $P(X^2 = 0) = P(X = 0) = 1 - p$.

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values: 0 and 1
 - ▶ $P(X^2 = 1) = P(X = 1) = p$. $P(X^2 = 0) = P(X = 0) = 1 - p$.
 - ▶ X and X^2 have identical PMF's! **They are identically distributed.**

Mean and Variance of Bernoulli

X is a Bernoulli random variable with $P(X = 1) = p$. We saw that $E[X] = p$. What is $\text{var}(X)$?

- ▶ First let's get $E[X^2]$. This is

$$E[X^2] = (1^2 \times P(X = 1) + 0^2 \times P(X = 0)) = p$$

We see that $E[X^2] = E[X]$. Is this surprising?

- ▶ Well, what is the PMF of X^2 ?
 - ▶ X^2 can take two values: 0 and 1
 - ▶ $P(X^2 = 1) = P(X = 1) = p$. $P(X^2 = 0) = P(X = 0) = 1 - p$.
 - ▶ X and X^2 have identical PMF's! **They are identically distributed.**
- ▶ $\text{var}(X) = E[X^2] - (E[X])^2 = p - p^2 = p(1 - p)$.

Mean and Variance of a Binomial

Let $X \sim \text{Bin}(n, p)$.

- ▶ $E[X] = np$ and $\text{var}(X) = np(1 - p)$.
- ▶ We will derive these in the next class.

Mean and Variance of a Poisson

X has a $\text{Poisson}(\lambda)$ distribution. What is its mean and variance?

- ▶ One can use algebra to show that $E[X] = \lambda$ and also $\text{var}(X) = \lambda$.
- ▶ How do you remember this?
- ▶ Hint: mean and variance of the Binomial approach that of a Poisson when n is large and p is small, such that $np \approx \lambda$? Anything yet?

Mean and variance of a geometric

- ▶ The PMF of a geometric distribution is $P(X = k) = (1 - p)^{k-1}p$.
 - ▶ $E[X] = 1/p$
 - ▶ $\text{var}(X) = (1 - p)/p^2$
 - ▶ We will also prove this later.

Conditional expectation

- ▶ We have done conditional probability and PMF.
- ▶ How about conditional expectation?
- ▶ Conditional expectation of random variable X conditioned on event A is written as $E[X|A]$
- ▶
$$E[X|A] = \sum_x xP(X = x|A)$$

Conditional expectation

- ▶ Say $X \sim \text{geometric}(p)$. What is $E[X|X > 1]$?

Conditional expectation

- ▶ Say $X \sim \text{geometric}(p)$. What is $E[X|X > 1]$?
- ▶ Remember $P(X = k|X > 1) = P(X = k - 1)$

Conditional expectation

- ▶ Say $X \sim \text{geometric}(p)$. What is $E[X|X > 1]$?
- ▶ Remember $P(X = k|X > 1) = P(X = k - 1)$

$$E[X|X > 1] = \sum_{k=2}^{\infty} kP(X = k|X > 1)$$

$$= \sum_{k=2}^{\infty} kP(X = k - 1)$$

- ▶ So

$$= \sum_{j=1}^{\infty} (j + 1)P(X = j)$$

$$= \sum_{j=1}^{\infty} jP(X = j) + 1 = E[X] + 1$$

The total expectation theorem

I am calculating the average combinatorics HW score in my class. I see that the average score of students who have taken combinatorics before is 90% whereas students who have not taken combinatorics before have an average of 70%. About 10% of the class has taken combinatorics before. How do I calculate the class average?

The total expectation theorem

I am calculating the average combinatorics HW score in my class. I see that the average score of students who have taken combinatorics before is 90% whereas students who have not taken combinatorics before have an average of 70%. About 10% of the class has taken combinatorics before. How do I calculate the class average?

- ▶ Should I do $(.9 + .7)/2$?

The total expectation theorem

- ▶ Consider disjoint events $\{A_1, \dots, A_n\}$ which form a partition of the sample space.
- ▶ The total probability theorem says
$$P(X = k) = \sum_i P(X = k|A_i)P(A_i).$$
- ▶ Similarly the total expectation theorem says:
$$E[X] = \sum_i E[X|A_i]P(A_i).$$
- ▶ How?

The total expectation theorem

I am calculating the average combinatorics HW score in my class. I see that the average score of students who have taken combinatorics before is 90% whereas students who have not taken combinatorics before have an average of 75%. About 10% of the class has taken combinatorics before. How do I calculate the class average?

- ▶ $C = \{\text{A student has taken combinatorics}\}$. $P(C) = .1$.
- ▶ $E[X|C] = ?$
- ▶ $E[X|C^c] = ?$
- ▶ $E[X] = E[X|C]P(C) + E[X|C^c]P(C^c) = ?$

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? .

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? **1.**

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? **1.**
- ▶ What is $P(X = 1)$? **.**

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? **1.**
- ▶ What is $P(X = 1)$? **p .**
- ▶ What is $P(X > 1)$? **.**

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? 1 .
- ▶ What is $P(X = 1)$? p .
- ▶ What is $P(X > 1)$? $1 - p$.
- ▶ What is $E[X|X > 1]$? $E[X] + 1$.

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? **1.**
- ▶ What is $P(X = 1)$? **p .**
- ▶ What is $P(X > 1)$? **$1 - p$.**
- ▶ What is $E[X|X > 1]$? **$E[X] + 1$.**
- ▶ So the whole thing is $E[X] = p + (1 + E[X])(1 - p)$

Mean of $X \sim \text{geometric}(p)$

Last time we derived the mean using differentiation. But once we know the total expectation theorem we do it much more quickly!

- ▶ Define two disjoint events $\{X = 1\}$ (first trial is success) and $\{X > 1\}$
- ▶ We have: $E[X] = E[X|X = 1]P(X = 1) + E[X|X > 1]P(X > 1)$.
- ▶ What is $E[X|X = 1]$? **1.**
- ▶ What is $P(X = 1)$? **p .**
- ▶ What is $P(X > 1)$? **$1 - p$.**
- ▶ What is $E[X|X > 1]$? **$E[X] + 1$.**
- ▶ So the whole thing is $E[X] = p + (1 + E[X])(1 - p)$
- ▶ Move things around: $E[X] = 1/p$.

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + \underbrace{E[X^2|X>1]P(X>1)}_{1-p}$$

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + \underbrace{E[X^2|X>1]P(X>1)}_{1-p}$$

\blacktriangleright Now,

$$E[X^2|X>1] = \sum_{k=2}^{\infty} k^2 P(X=k|X>1) = \sum_{k=2}^{\infty} k^2 \underbrace{P(X=k-1)}_{\text{memoryless property}}.$$

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + \underbrace{E[X^2|X>1]P(X>1)}_{1-p}$$

► Now,

$$E[X^2|X>1] = \sum_{k=2}^{\infty} k^2 P(X=k|X>1) = \sum_{k=2}^{\infty} k^2 \underbrace{P(X=k-1)}_{\text{memoryless property}} .$$

$$\blacktriangleright \text{Substitute } i = k - 1 \text{ to get: } E[X^2|X>1] = \sum_{i=1}^{\infty} (i+1)^2 P(X=i)$$

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + \underbrace{E[X^2|X>1]P(X>1)}_{1-p}$$

► Now,

$$E[X^2|X>1] = \sum_{k=2}^{\infty} k^2 P(X=k|X>1) = \sum_{k=2}^{\infty} k^2 \underbrace{P(X=k-1)}_{\text{memoryless property}}.$$

► Substitute $i = k - 1$ to get: $E[X^2|X>1] = \sum_{i=1}^{\infty} (i+1)^2 P(X=i) =$

$$E[(X+1)^2] = E[X^2 + 2X + 1] = E[X^2] + 2E[X] + 1 = E[X^2] + \frac{2}{p} + 1.$$

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + \underbrace{E[X^2|X>1]P(X>1)}_{1-p}$$

\blacktriangleright Now,

$$E[X^2|X>1] = \sum_{k=2}^{\infty} k^2 P(X=k|X>1) = \sum_{k=2}^{\infty} k^2 \underbrace{P(X=k-1)}_{\text{memoryless property}}.$$

\blacktriangleright Substitute $i = k - 1$ to get: $E[X^2|X>1] = \sum_{i=1}^{\infty} (i+1)^2 P(X=i) =$

$$E[(X+1)^2] = E[X^2 + 2X + 1] = E[X^2] + 2E[X] + 1 = E[X^2] + \frac{2}{p} + 1.$$

\blacktriangleright So plugging everything in I have:

$$E[X^2] = p + E[(1+X)^2](1-p) = p + (E[X^2] + \frac{2}{p} + 1)(1-p)$$

Variance of $X \sim \text{Geometric}(p)$

$$\blacktriangleright E[X^2] = \underbrace{E[X^2|X=1]P(X=1)}_{1 \times p} + E[X^2|X>1] \underbrace{P(X>1)}_{1-p}$$

\blacktriangleright Now,

$$E[X^2|X>1] = \sum_{k=2}^{\infty} k^2 P(X=k|X>1) = \sum_{k=2}^{\infty} k^2 \underbrace{P(X=k-1)}_{\text{memoryless property}}.$$

$$\blacktriangleright \text{Substitute } i = k - 1 \text{ to get: } E[X^2|X>1] = \sum_{i=1}^{\infty} (i+1)^2 P(X=i) =$$

$$E[(X+1)^2] = E[X^2 + 2X + 1] = E[X^2] + 2E[X] + 1 = E[X^2] + \frac{2}{p} + 1.$$

\blacktriangleright So plugging everything in I have:

$$E[X^2] = p + E[(1+X)^2](1-p) = p + (E[X^2] + \frac{2}{p} + 1)(1-p)$$

$$\blacktriangleright \text{Solving, we get } E[X^2] = \frac{2}{p^2} - \frac{1}{p}$$

$$\blacktriangleright \text{So } \text{var}(X) = E[X^2] - (E[X])^2 = \frac{1-p}{p^2}$$