

MAST90026 Computational Differential Equations: Week 2

Jesse Collis


Modified from Hailong Guo (2022)

Semester 1 2024

The University of Melbourne



Method of weighted residual



Weighted Residual Method

	Global basis	Piecewise polynomial
Collocation	Spectral collocation	Spline collocation
Galerkin	Spectral Galerkin	Finite element method

Idea of spectral collocation

Three step idea:

Step 1: Choose a trial space V_N ($N + 1$ dimensional). Write $u_N \in V_N$ as

$$u_N = \sum_{j=0}^N c_j b_j(x).$$

Step 2: Choose collocation points $\{x_0, x_1, \dots, x_N\}$.

Step 3: Determine c_j using the collocation condition:

$$Lu_N(x_j) = f(x_j), \quad j = 0, 1, \dots, N.$$

Choose of trial space and collocation points

Periodic boundary conditions:

- Space of trigonometric functions

$$u_N(x) = \sum_{j=0}^N a_j \cos(jx) + \sum_{j=1}^N b_j \sin(jx) = \sum_{j=-N}^N c_j e^{ijx}.$$

- Collocation points: $x_j = \frac{j\pi}{N}$, $j = 0, 1, \dots, 2N$.

Nonperiodic boundary conditions:

- Space of polynomials of degree N (order $N + 1$).

$$u_N(x) = p_N(x) \in \mathbb{P}_{N+1}.$$

- Collocation points are roots of orthogonal polynomial of degree $N + 1$.

Definition of orthogonal

Definition : (Orthogonal)

Given an open interval $I := (a, b)$ ($-\infty \leq a < b \leq +\infty$), and a generic weight function ω such that

$$\omega(x) > 0, \forall x \in I \text{ and } \omega \in L^1(I)$$

two different functions f and g are said to be orthogonal to each other in $L^2_\omega(a, b)$ or orthogonal with respect to ω if

$$(f, g)_\omega := \int_a^b f(x)g(x)\omega(x)dx = 0$$

Definition

A sequence of polynomials $\{p_n\}_{n=0}^{\infty}$ with $\deg(p_n) = n$ is said to be orthogonal in $L^2_{\omega}(a, b)$ if

$$(p_n, p_m)_{\omega} = \int_a^b p_n(x)p_m(x)\omega(x)dx = \gamma_n\delta_{mn}$$

where the constant $\gamma_n = \|p_n\|_{\infty}^2$ is nonzero, and δ_{mn} is the Kronecker delta.

Existence and uniqueness of orthogonal polynomial

Theorem

For any given positive weight function $\omega \in L^1(a, b)$, there exists a unique sequence of monic orthogonal polynomials $\{\bar{p}_n\}$ with $\deg(\bar{p}_n) = n$, which can be constructed as follows

$$\bar{p}_0 = 1, \bar{p}_1 = x - \alpha_0$$

$$\bar{p}_{n+1} = (x - \alpha_n) \bar{p}_n - \beta_n \bar{p}_{n-1}, \quad n \geq 1$$

where

$$\alpha_n = \frac{(x \bar{p}_n, \bar{p}_n)_\omega}{\|\bar{p}_n\|_\omega^2}, \quad n \geq 0,$$

$$\beta_n = \frac{\|\bar{p}_n\|_\omega^2}{\|\bar{p}_{n-1}\|_\omega^2}, \quad n \geq 1.$$

Example of orthogonal polynomials

Consider $(a, b) = (-1, 1)$ and $\omega = 1$. Legendre polynomials:

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_{n+1}(x) = \frac{2n+1}{n+1}xP_n(x) - \frac{n}{n+1}P_{n-1}(x),$$

Consider $(a, b) = (-1, 1)$ and $\omega = (1 - x^2)^{-\frac{1}{2}}$. Chebyshev polynomials of the first kind:

$$P_0(x) = 1, \quad P_1(x) = x, \quad P_{n+1}(x) = 2xP_n(x) - P_{n-1}(x),$$

More general. Consider $(a, b) = (-1, 1)$ and $\omega = (1 - x)^\alpha(1 + x)^\beta$ with $\alpha, \beta > -1$. The obtained orthogonal polynomials are Jacobi polynomials

Other definition of Legendre and Chebyshev polynomials

Legendre polynomials:

- Eigenfunction of

$$-\frac{d}{dx} \left((1-x^2) \frac{d\phi(x)}{dx} \right) = k(k+1)\phi(x)$$

- Rodrigue's formula

$$P_n(x) = \frac{1}{2^n n!} \frac{d^n}{dx^n} (x^2 - 1)^n.$$

Chebyshev polynomial of the first kind:

- Eigenfunction of

$$-\frac{d}{dx} \left(\sqrt{1-x^2} \frac{d\phi(x)}{dx} \right) = \frac{k^2}{\sqrt{1-x^2}} \phi(x)$$

- Explicit formula

$$P_n(x) = \cos(n\theta), \text{ where } \theta = \arccos(x)$$

Zeros of orthogonal polynomials

Theorem

The zeros of p_n are all real, simple, and lie in the interval (a, b) .

Numerical quadrature

The basis problem is to find quadrature nodes $\{x_j\}$ and weights $\{\omega_j\}$ such that

$$\int_a^b f(x)\omega(x)dx \approx \sum_{j=0}^N f(x_j)\omega_j.$$

Definition

The quadrature have algebraic accuracy of degree p if

$$\int_a^b p(x)\omega(x)dx = \sum_{j=0}^N p(x_j)\omega_j, \quad p(x) \in \mathbb{P}_p.$$

In general, we want choose $\{x_j\}$ and $\{\omega_j\}$ such that we have highest degree of accuracy.

Theorem (Gauss quadrature)

Let $\{x_j\}_{j=0}^N$ be the set of zeros of the orthogonal polynomial p_{N+1} . Then there exists a unique set of quadrature weights $\{\omega_j\}_{j=0}^N$, defined by (3.36), such that

$$\int_a^b p(x)\omega(x)dx = \sum_{j=0}^N p(x_j)\omega_j, \quad \forall p \in P_{2N+1}$$

Orthogonal Polynomials and Related Approximation Results where the quadrature weights are all positive and given by

$$\omega_j = \frac{k_{N+1}}{k_N} \frac{\|p_N\|_{\omega}^2}{p_N(x_j)p'_{N+1}(x_j)}, \quad 0 \leq j \leq N,$$

where k_j is the leading coefficient of the polynomial p_j .

Choices of collocation points

We have many different options for collocation points that suit different scenarios

Chebyshev points:

- Chebyshev-Gauss: $x_j = \cos \frac{(2j+1)\pi}{2N+2}$, $0 \leq j \leq N$.
- Chebyshev-Gauss-Radau: $x_j = \cos \frac{2j\pi}{2N+1}$, $0 \leq j \leq N$.
- Chebyshev-Gauss-Lobatto: $x_j = \cos \frac{j\pi}{N}$, $0 \leq j \leq N$.

Legendre points:

- Legendre-Gauss: x_j are the zeros of $P_{N+1}(x)$.
- Legendre-Gauss-Radau: x_j are the zeros of $P_N(x) + P_{N+1}(x)$.
- Legendre-Gauss-Lobatto: $x_0 = -1$, $x_N = 1$, $\{x_j\}_{j=1}^N$ are zeros of $P'_{N-1}(x)$.

Question: Why don't we just use equally distributed points?

Chebyshev-Gauss quadrature

The quadrature rules for Chebyshev points are particularly simple

1. Chebyshev-Gauss:

$$\int_{-1}^1 \frac{p(x)}{\sqrt{1-x^2}} dx = \frac{\pi}{N+1} \sum_{j=0}^N p(x_j), \quad \forall p \in P_{2N+1}$$

2. Chebyshev-Gauss-Radau:

$$\int_{-1}^1 \frac{p(x)}{\sqrt{1-x^2}} dx = \frac{\pi}{2N+1} p(1) + \frac{\pi}{N+\frac{1}{2}} \sum_{j=1}^N p(x_j), \quad \forall p \in P_{2N}$$

3. Chebyshev-Gauss-Lobatto:

$$\int_{-1}^1 \frac{p(x)}{\sqrt{1-x^2}} dx = \frac{\pi}{2N} (p(1) + p(-1)) + \frac{\pi}{N} \sum_{j=1}^{N-1} p(x_j), \quad \forall p \in P_{2N-1}$$

Illustration of spectral collocation method

Example: Consider the BVP $u''(x) + p(x)u'(x) + q(x) = r(x)$ over $[a, b]$.

- Map the interval $[a, b]$ to $[-1, 1]$.
 $\rightarrow u''(t) + \bar{p}(t)u'(t) + \bar{q}u(t) = \bar{r}(t)$. Here we still denote the mapped functions as $u(t)$.
- Let

$$u_N(x) = \sum_{j=0}^N u_j \ell_j(x),$$

where $\ell_j(x)$ be the Lagrange interpolating polynomials. So,

$$u'_N(x) = \sum_{j=0}^N u_j \ell'_j(x),$$

$$u''_N(x) = \sum_{j=0}^N u_j \ell''_j(x).$$

Differentiation matrix

Since we are collocating, we only need to find u' and u'' at the collocation points.

$$u'_N(x_k) = \sum_{j=0}^N u_j \ell'_j(x_k) = \sum_{j=0}^N D_{kj} u_j, \quad \text{and}$$
$$u''_N(x_k) = \sum_{j=0}^N u_j \ell''_j(x_k) = \sum_{j=0}^N D_{kj}^{(2)} u_j,$$

where $D_{kj} \equiv \left. \frac{d}{dx} \ell_j(x) \right|_{x_k}$, $D_{kj}^{(2)} = \left. \frac{d^2}{dx^2} \ell_j(x) \right|_{x_k}$, and ℓ_j is the j th Lagrange interpolation polynomial.

Note that $D^{(2)} = D^2$ (squared using matrix multiplication).

How to compute the Differentiation matrix

Theorem

The entries of D are determined by

$$d_{kj} = \ell'_j(x_k) = \begin{cases} \frac{Q'(x_k)}{Q'(x_j)} \frac{1}{x_k - x_j}, & \text{if } k \neq j, \\ \frac{Q''(x_k)}{2Q'(x_k)}, & \text{if } k = j, \end{cases}$$

where

$$Q(x) = p_{N+1}(x), (x - a)q_N(x), (x - a)(b - x)z_{N-1}(x)$$

are the quadrature polynomials of the Gauss, Gauss-Radau and Gauss-Lobatto quadrature, respectively. We can write $p_{N+1} = (x - x_0)(x - x_1) \cdots (x - x_N)$, $q_N = (x - x_1) \cdots (x - x_N)$, and $z_{N-1} = (x - x_1) \cdots (x - x_{N-1})$

How to compute the Differentiation matrix

Proof

The Lagrange basis polynomials can be expressed as

$$\ell_j(x) = \frac{Q(x)}{Q'(x_j)(x - x_j)}, \quad 0 \leq j \leq N$$

Differentiating it and using the fact that $Q(x_j) = 0$ leads to

$$d_{kj} = \ell'_j(x_k) = \frac{Q'(x_k)}{Q'(x_j)} \frac{1}{x_k - x_j}, \quad \forall k \neq j.$$

Applying L'Hopital's rule twice yields

$$d_{kk} = \lim_{x \rightarrow x_k} \ell'_k(x) = \frac{1}{Q'(x_k)} \lim_{x \rightarrow x_k} \frac{Q'(x)(x - x_k) - Q(x)}{(x - x_k)^2} = \frac{Q''(x_k)}{2Q'(x_k)}.$$

This completes the proof.

Chebyshev differentiation matrix

This code gives the differentiation matrix for Chebyshev-Gauss-Lobatto points using Lagrange basis polynomials

```
% CHEB  compute D = differentiation matrix, x = Chebyshev grid

function [D,x] = cheb(N)
if N==0, D=0; x=1; return, end
x = cos(pi*(0:N)/N)';
c = [2; ones(N-1,1); 2].*(-1).^(0:N)';
X = repmat(x,1,N+1);
dX = X-X';
D = (c*(1./c)') ./ (dX+(eye(N+1)));           % off-diagonal entries
D = D - diag(sum(D'))';                       % diagonal entries
```

Go back to example

Collocate at the collocation points and

$$u_N''(x_k) + \bar{p}(x_k)u_N'(x_k) + \bar{q}(x_k)u_N(x_k) = \bar{r}(x_k), \quad k = 0, 1, \dots, N,$$

becomes

$$D^2 \vec{u} + \begin{pmatrix} \bar{p}_0 & 0 & \cdots & 0 \\ 0 & \bar{p}_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \bar{p}_N \end{pmatrix} D \vec{u} + \begin{pmatrix} \bar{q}_0 & 0 & \cdots & 0 \\ 0 & \bar{q}_1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \bar{q}_N \end{pmatrix} \vec{u} = \begin{pmatrix} \bar{r}_0 \\ \bar{r}_1 \\ \vdots \\ \bar{r}_N \end{pmatrix} = \vec{r}.$$

And you can solve this for \vec{u} .

Impose boundary condition

Since we know the value of u at $x = \pm 1$, ($k = 0, N$), we don't need to collocate at these points. So we do not need the first and last rows of D and D^2 .

So for the rest of our rows we have:

$$\begin{aligned} u'_k &= \sum_{j=0}^N D_{kj} u_j \quad (k = 1, \dots, N-1) \\ &= \sum_{j=1}^{N-1} D_{kj} u_j + u_0 D_{k0} + u_N D_{kN}. \end{aligned}$$

Summary of spectral collocation method

Pros and cons

- High-order method, easy to implement
- Superior to spectral-Galerkin method in dealing with variable coefficients and/or nonlinear problems
- Differential matrix is ill-conditioned and full

End of week 2!