

Towards Structured Use of Bayesian Sequential Monitoring in Clinical Trials

Evan Kwiatkowski[†], Eugenio Andraca-Carrera[‡],
Mat Soukup[‡], Matthew A. Psioda^{†*}

[†] Department of Biostatistics, University of North Carolina,
McGavran-Greenberg Hall, CB#7420,
Chapel Hill, North Carolina, U.S.A.

[‡] Division of Biometrics VII, Office of Biostatistics
Center for Drug Evaluation and Research,
US Food and Drug Administration,
Silver Spring, Maryland, USA

June 14, 2019

Abstract

The text of your abstract. 200 or fewer words.

Keywords: 3 to 6 keywords, that do not appear in the title

*The authors gratefully acknowledge *please remember to list all relevant funding sources in the unblinded version*

1 Introduction

Things to discuss:

- 21st Century Cures Act (MATT)
- PDUFA VI reauthorization (MATT)
- Expansive work already done on sequential monitoring (EVAN – draft on 6/21)
- Our majors contribution (EVAN – as early as possible in introduction without having the flow appear weird – draft on 6/21)
- Outline for the remaining section of the paper (EVAN – draft on 6/21)

2 Methods

As you introduce ideas that come from or extend other ideas in the literature, cite the relevant literature.

2.1 Monitoring versus Estimation Priors (EVAN – draft on 6/21)

- Define generally in terms of $\boldsymbol{\theta} = (\gamma, \boldsymbol{\psi})$ where γ is a parameter of interest and $\boldsymbol{\psi}$ is a nuisance parameter (possible vector valued).
- Define *Monitoring* Priors and *Inference* Priors.
- Make connection between Inference priors and two-part mixture prior and BMA.
- Define *Skeptical* and *Enthusiastic* monitoring priors and how each would be used.

- I would have a generic graphic to illustrate the types of priors and the mixture.

2.2 Futility Monitoring Using Probability of Success (EVAN – draft on 6/21)

- Futility monitoring using POS is about stopping early when there is a high likelihood of a study being inconclusive at the end of the study.
- Since the final analysis uses the *Inference* prior, POS should be based on the inference prior.
- Develop the framework for POS and show how it is a weighted average POS based on the skeptical and enthusiastic priors.

3 Examples – (EVAN)

3.1 Single-Arm Oncology Proof-of-Activity Trial w/ Binary Endpoint

3.2 Parallel Two-Group Superiority Trial /w Continuous Binary Endpoint

3.3 Three-Arm, Placebo Controlled Non-Inferiority Trial w/ Continuous Endpoint

4 Discussion – (MATT/EVAN)

SUPPLEMENTARY MATERIAL

5 BibTeX

References