



DESCRIPTION

Miraten® is a preparation of Butamirate Citrate. Miraten® belongs to a group of medications called central action antitussive drugs. Miraten® exerts antitussive, expectorant, moderate bronchodilatory action with anti-inflammatory effects. Miraten® lowers the resistance of airways and improves blood oxygenation and spirometery indexes.

INDICATIONS

Dry cough of different origins

Cough associated with pertussis

Cough suppression during pre and post-operative surgical interventions (bronchoscopy)

DOSAGE AND ADMINISTRATION

Miraten® Syrup:

Adults: 3 teaspoonful (15 mL) 4 times daily

Children 12 years and over : 3 teaspoonful (15 mL) 3 times daily
Children 6-12 years of age: 2 teaspoonful (10 mL) 3 times daily
Children 3-6 years of age: 1 teaspoonful (5 mL) 3 times daily

Miraten® 15 mL Paediatric Drops:

Children: 1-3 years of age: 0.75 mL 4 times daily

Children: 2 months-1 year of age: 0.5 mL 4 times daily

Miraten® should be used before meal.

CONTRAINDICATIONS

lt is contraindicated to patients with hypersensitivity to Butamirate

Citrate

Breastfeeding mothers (lactation)

SIDE EFFECTS

Butamirate Citrate is well tolerated. The most common side effects include-

Dizziness

Nausea, diarrhoea

Skin rash and itching (in rare cases)

PRECAUTION AND WARNING

By inhibition of the cough reflex by Butamirate Citrate, concomitant use of expectorants should be avoided.

USE IN PREGNANCY AND LACTATION

Butamirate Citrate should not be used during pregnancy, especially during the first three months of pregnancy. If absolutely necessary, Butamirate Citrate can be used during the last six months of pregnancy with the advice of the doctor.

PHARMACEUTICAL PRECAUTION

For Syrup: Keep away from light and do not store in wet place. Keep out of reach of children

For Paediatric Drops: Keep away from light and wet place. Do not store in refrigerator. Keep out of reach of children.

PACKAGING

Miraten® 100 mL Syrup: Bottle containing 100 mL of syrup. Each 5 mL contains Butamirate Citrate INN 7.5 mg.

 $\pmb{\text{Miraten}} = \pmb{200} \text{ mL Syrup:}$ Bottle containing 200 mL of syrup. Each 5 mL contains Butamirate Citrate INN 7.5 mg.

Miraten® Paediatric Drops: Bottle containing 15 mL paediatric drops. Each mL

SK+F

Manufactured by

ESKAYEF BANGLADESH LIMITED

GAZIPUR, BANGLADESH ® REGD. TRADEMARK PM03889 V03