

# Mebidal®

Mebhydrolin tablet

## DESCRIPTION

Mebidal® is a round white tablet containing Mebhydrolin napadisylate. Mebidal diminishes or abolishes the main actions of histamine in the body by competitive reversible blockade of histamine receptor sites on tissues.

## INDICATIONS

Mebidal® is an antihistamine recommended for the treatment of hay fever, urticaria, rhinitis and all other allergic disorders.

## DOSAGE AND ADMINISTRATION

Adults : 1 to 2 tablets three times daily.

Children :

Over 10 years : 1 to 2 tablets three times daily.

Under 10 years : 1 to 4 tablets daily according to age in divided doses or as directed by the physician.

## CONTRAINDICATIONS

None known, but previously known hypersensitivity is a definite contra-indication.

## SIDE-EFFECTS

Very rare cases of granulocytopenia or agranulocytosis associated with the use of mebhydrolin have been reported. Sedation, drowsiness & dryness of mouth may occur.

## PRECAUTIONS

The substance may cause drowsiness, if affected, do not drive or operate machinery. Avoid alcoholic drinks.

No embryotoxic or teratogenic effects have been reported, but on principle caution should be exercised when prescribing any new medication during pregnancy and lactation.

## PHARMACEUTICAL PRECAUTION

Store in a dry place, keep out of reach of children.

## PACKAGING

Box containing 20 strips of 10 tablets. Each tablet contains mebhydrolin napadisylate BPC equivalent to 50 mg mebhydrolin base.

# SK+F

Manufactured by

**ESKAYEF BANGLADESH LTD.**

DHAKA, BANGLADESH

® REGD. TRADEMARK

M/PM 00244 V01

# মেবিডাল®

## মেবহাইড্রলিন ট্যাবলেট

### বর্ণনা

মেবিডাল® গোলাকৃতি সাদা ট্যাবলেট যার প্রতিটিতে রয়েছে মেবহাইড্রলিন নাপাডাইসাইলেট। মেবিডাল শরীরে হিস্টামিনের প্রধান কার্যকারিতাকে কমায়ে বা বিলুপ্ত করে। ইহা শরীরের কোষ কলায় হিস্টামিন রিসেপ্টারকে অবরুদ্ধ করে এবং এই ক্রিয়া প্রতিযোগিতামূলক ও উভয়মুখী।

### রোগ নির্দেশ

মেবিডাল® একটি এন্টিহিস্টামিন। ইহা হে-ফিভার, আরটিকেরিয়া, রাইনেটিস ও অন্যান্য এ্যালার্জিক ব্যাধির চিকিৎসায় নির্দেশিত হয়।

### সেবনমাত্রা ও প্রয়োগ

প্রাপ্ত বয়স্ক : ১ থেকে ২ ট্যাবলেট দিনে ৩ বার।

দশ বছরের উর্ধ্বে শিশু : ১ থেকে ২ ট্যাবলেট দিনে ৩ বার।

দশ বছরের নীচে : বয়স অনুযায়ী দৈনিক ১ থেকে ৪টি ট্যাবলেট বিভক্ত মাত্রায় অথবা চিকিৎসকের পরামর্শ অনুযায়ী সেব্য।

### বিরুদ্ধ ব্যবহার

মেবিডালের কোন বিরুদ্ধ ব্যবহার জানা নেই। তবে অতিসংবেদনশীলতার অতীত রেকর্ড সম্পন্ন রোগীদের দেয়া যাবে না।

### পার্শ্ব প্রতিক্রিয়া

মেবহাইড্রলিনের ব্যবহারে থ্রেনুলোসাইটোপেনিয়া ও এথ্রেনুলোসাইটোসিস এর রিপোর্ট কদাচিৎ পাওয়া যায়। কিমুনি, মুখ শুকানো এবং নিদ্রালুতা হতে পারে।

### সতর্কতা

ইহা সেবনে নিদ্রালুতা হতে পারে এবং সে ক্ষেত্রে গাড়ী অথবা যন্ত্রপাতি চালানো উচিত নয়। এ্যালকোহল জাতীয় পানীয় পরিহার করা উচিত।

ইহার এমব্রাইওটক্সিক ও টেরাটোজেনিক ক্রিয়ার কোন রিপোর্ট পাওয়া যায় নাই। কিন্তু সাধারণ নীতি অনুযায়ী গর্ভাবস্থায় ও দুগ্ধদানকালে কোন নতুন ঔষধ ব্যবহারে সতর্কতা অবলম্বন করা উচিত।

### ফার্মাসিউটিক্যাল সাবধানতা

শুকনো স্থানে রাখুন, শিশুদের নাগালের বাহিরে রাখুন।

### প্যাকেজিং

প্রতি বাক্সে ১০টি ট্যাবলেটের ২০টি স্ট্রীপ আছে। প্রতি ট্যাবলেটে ৫০ মি.গ্রা. মেবহাইড্রলিন বেইস এর সমতুল্য মেবহাইড্রলিন নাপাডাইসাইলেট বি.পি.সি. রয়েছে।

# SK+F

প্রস্তুতকারক

এসকে এফ বাংলাদেশ লিঃ

ঢাকা, বাংলাদেশ।

® রেজিস্টার্ড ট্রেডমার্ক