

DESCRIPTION

Osticare bis the combination of Diacerein and Glucosamine Sulfate. Diacerein is an anti inflammatory agent and Glucosamine is an amino-monosaccharide. The combination used to treat osteoarthritis, rheumatoid arthritis and bone and joint pain. Diacerein has analgesic and anti inflammatory activity. It inhibits interleukin-1 synthesis and reduces finger joint and knee osteoarthritis. Glucosamine stimulates formation of collagen which is the protein portion of bone joints. Collagen is the main component of the shock absorbing cushion i.e articular cartilage.

INDICATIONS

Osticare so is indicated for the treatment of osteoarthritis, rheumatoid arthritis, bone and joint pain.

DOSAGE AND ADMINISTRATION

Adult and Children aged 12 years and older: The usual dose of Osticare is 1 tablet twice daily (12 hourly) taken with food.

In case of patients with creatinine clearance less than 30 mL/min, the dose of Diacerein is to be reduced. No data regarding pediatric and neonatal dosage details is available.

CONTRAINDICATIONS

Patients with hypersensitivity to Diacerein or Glucosamine should avoid this medication.

SIDE EFFECTS

No serious adverse effect has been reported. Diarrhea, epigastric pain, nausea, vomiting, intense yellow coloring of urine may be occurred.

USE IN PREGNANCY AND LACTATION

Osticare® is contraindicated during pregnancy and breastfeeding.

PRECAUTIONS

Patients with renal impairment should monitor complete blood count, Liver function tests and urinalysis every 6 months.

STORAGE

Keep away from wet place and light. Keep out of reach of children.

PACKAGING

Osticare® Tablet: Box containing 5 strips of 6 tablets each. Each tablet contains Diacerein INN 50 mg and Glucosamine Sulfate Potassium Chloride USP equivalent to Glucosamine Sulfate 750 mg.



Manufactured by

ESKAYEF BANGLADESH LIMITED

GAZIPUR, BANGLADESH ® REGD. TRADEMARK PM04848 V01