

Species occurrence as a function of both emergent biological traits and environmental context

Peter D. Smits^{1,*}

1. University of Chicago, Chicago, Illinois 60637.

* Corresponding author; e-mail: psmits@uchicago.edu.

Manuscript elements:

Keywords:

Manuscript type: Article

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Introduction

2 How do species pools change over time as species are recruited or go extinct? When are ecotypes
enriched or depleted? How does global and regional environmental context affect the distribution of
4 species ecotypes (e.g. guilds) in a regional species pool?

A regional species pool is the set of species which form communities in a specific region; local
6 communities are subsets of the regional pool. The composition of a regional species pool changes
over time due to speciation, migration, extinction. Local scale processes like resource competition
8 only affect the regional species pool if all communities are affected.

Valentine and Bambach how they presented guilds in paleobiology. Bush and Bambach presented an
10 ecocube to describe what how marine invertebrates partition space and resources (Bambach et al.,
2007; Bush and Bambach, 2011; Bush et al., 2007). Unique combinations represent what possible
12 ecotypes are observable. The distribution of ecocube occupancy is then normally analyzed as raw
counts of unique combinations or using ordination methods and the change in disparity over time is
14 estimated (Bambach et al., 2007; Bush and Bambach, 2011; Bush et al., 2007).

One of the greatest challenges with analyzing species occurrence data is the inherent incompleteness
16 of any sample (Foote, 2001; Foote and Sepkoski, 1999; Lloyd et al., 2011; Royle and Dorazio, 2008;
Royle et al., 2014; Wang and Marshall, 2016). In the modern, only presences are certain as an
18 absence can be caused by both the species being truly absent or the species never having been
sampled (Royle and Dorazio, 2008; Royle et al., 2014). For paleontological data in the context of
20 this study, the incomplete preservation of fossil communities combined with the incomplete
sampling of what fossils there are means that the true times of origination or extinction may not be
22 observed (Foote, 2001; Foote and Sepkoski, 1999; Wang et al., 2016; Wang and Marshall, 2016)

Smits (2015) found several systematic differences in mammal species durations associated with
24 various species traits. Omnivorous taxa were found to have, on average, a greater duration than
other dietary categories. Additionally, arboreal taxa were found to have a shorter duration than
26 other locomotor categories.

An unresolved question from Smits (2015) is whether the greater extinction risk faced by arboreal is
constant over time or if there was a change in extinction risk at the Paleogene/Neogene boundary.
Specifically, the question is whether the extinction risk arboreal taxa increased in the Neogene,
driving the loss of arboreal taxa and average extinction risk of arboreal taxa down.

There are no observed massive cross-taxonomic turnover events in the North American record,
unlike the Neogene record Europe (Alroy, 1996, 2009; Alroy et al., 2000; Eronen et al., 2015; Janis,
1993).

The effect of climate on diversity and the diversification process has been the focus of considerable
research with many analyses favoring diversification being more biologically-mediated than
climate-mediated (Alroy, 1996; Alroy et al., 2000; Clyde and Gingerich, 1998; Figueirido et al.,
2012). Scale of analysis makes a big difference in interpretation of results, both temporal and
geographic. For example when the mammal fossil record analyzed at small temporal and geographic
scales a correlation between diversity and climate are observable (Clyde and Gingerich, 1998).
However, when the record is analyzed at the scale of the continent and the Cenozoic there is no
correlation with diversity and climate (Alroy et al., 2000). This results, however, does not go
against the idea that there may be short periods of correlation and that this correlation change or
reverse direction over time; instead this result means that there is no single direction of correlation
between diversity and climate (Figueirido et al., 2012).

In the case of a fluctuating correlation between diversity and climate it is hard to make the
argument of an actual causal link between the two without understanding the ecological differences
in mammalian fauna over time; when this analysis is based on diversity or taxonomy alone no
mechanisms are possible to infer. After all, taxonomy conflates many potential factors that could
affect diversification into a single variable; by separating the effects of shared common ancestry (i.e.
phylogeny) from species ecology the subtle differences in the diversification process can be observed
(Smits, 2015).

There are many candidate climatic events that may have influenced the distribution of mammal
ecotypes regionally, if not globally (Blois and Hadly, 2009; Janis, 1993; Zachos et al., 2008, 2001).

54 The Paleocene-Eocene Thermal Maximum is associated with species dwarfing and rearrangement
of local communities, though regional effects are less known CITATION. The Mid-Miocene
56 climactic optimum is associated with WHAT CITATION. The

The general cooling throughout the Cenozoic and the development of ice-caps in the Neogene. The
58 Oligo-Miocene boundary.

One of the most stunning environmental transitions of the Cenozoic in North America was gradual
60 “opening-up” of the landscape with the shift from closed or partially forested environments of the
Paleogene to the savannah and grasslands environments that characterize the Neogene (Blois and
62 Hadly, 2009; Janis, 1993; Janis et al., 2000; Strömberg, 2005).

Fourth-corner modeling an approach to explaining the patterns of either species abundance or
64 presence/absence as a product of species traits, environmental factors, and the interaction between
traits and environment CITATION. In modern ecological studies, what is being modeled is species
66 occurrences at localities distributed across a region CITATION. In this study, what is being
modeled is the pattern of species occurrence over time for most of the Cenozoic in North America
68 (Fig. 1). These two approaches, modern and paleontological, are different views of the same
three-dimensional pattern: species at localities over time. The temporal limitations of modern
70 ecological studies and difficulties with uneven spatial occurrences of fossils in paleontological studies
means that these approaches are complimentary but reveal different patterns of how species are
72 distributed in time and space.

Materials and Methods

74 Taxon occurrences and species-level information

All fossil occurrence information was downloaded from the Paleobiology Database.
76 Occurrences (PBDB) were restricted to all Mammalia sampled in North America between the
Maastrichtian and Gelasian stages. Taxonomic, stratigraphic, and ecological metadata for each

occurrence was included. The raw data is available for download at <http://goo.gl/2slgeU>.

This raw data was then sorted, cleaned, and manipulated programmatically prior to analysis.

Species taxonomic assignments given by the PBDB were updated for accuracy and consistency. For example, species classified in the order Artiodactyla were reclassified as Cetartiodactyla. These re-assignments follow Smits (2015) and were Janis et al. (2008, 1998) and the Encyclopedia of Life WEBSITE. Additionally, Taxa whose life habit was classified as either volant (i.e. Chiroptera) or aquatic (e.g. Cetacea) were excluded from this analysis because of both differences in fossilization potential and applicability to the study of terrestrial species pools.

The life habit and dietary categories provided through the PBDB were coarsened to increase per ecotype sample size; this coarsening follows the same procedure as Smits (2015). Additionally, life habit category was further modified to break-up the vague “ground-dwelling” category; re-classifying these species by ankle posture gives more precise information about that species’ environmental context. Ground-dwelling taxa were reassigned following ? by species taxonomic context. Species ecotype is defined as the interaction between life habit and diet categories. Ecotype categories with less than 10 species having ever been that combination were excluded, yielding a total of 18 of 21 possible ecotypes.

Table 1: Species trait assignments in this study are a coarser version of the information available in the PBDB. Information was coarsened to improve per category sample size and uniformity and followed this table.

This study		PBDB categories
Diet	Carnivore	Carnivore
	Herbivore	Browser, folivore, granivore, grazer, herbivore.
	Insectivore	Insectivore.
	Omnivore	Frugivore, omnivore.
Locomotor	Arboreal	Arboreal.
	Ground dwelling	Fossorial, ground dwelling, semifossorial, saltatorial.
	Scansorial	Scansorial.

Table 2: Posture assignment based on taxonomy

Order	Family	Stance
	Ailuridae	plantigrade
	Allomyidae	plantigrade
	Amphicyonidae	plantigrade
	Amphilemuridae	plantigrade
	Anthracotheriidae	digitigrade
	Antilocapridae	unguligrade
	Apheliscidae	plantigrade
	Aplodontidae	plantigrade
	Apternodontidae	scansorial
	Arctocyonidae	unguligrade
	Barbourofelidae	digitigrade
	Barylambdidae	plantigrade
	Bovidae	unguligrade
	Camelidae	unguligrade
	Canidae	digitigrade
	Cervidae	unguligrade
	Cimolodontidae	scansorial
	Coryphodontidae	plantigrade
	Cricetidae	plantigrade
	Cylindrodontidae	plantigrade
	Cyriacotheriidae	plantigrade
	Dichobunidae	unguligrade
	Dinocerata	unguligrade
	Dipodidae	digitigrade
Continued on next page		

Table 2 – continued from previous page

Order	Family	Stance
	Elephantidae	digitigrade
	Entelodontidae	unguligrade
	Eomyidae	plantigrade
	Erethizontidae	plantigrade
	Erinaceidae	plantigrade
	Esthonychidae	plantigrade
	Eutypomyidae	plantigrade
	Felidae	digitigrade
	Florentiamyidae	plantigrade
	Gelocidae	unguligrade
	Geolabididae	plantigrade
	Glyptodontidae	plantigrade
	Gomphotheriidae	unguligrade
	Hapalodectidae	plantigrade
	Heteromyidae	digitigrade
	Hyaenidae	digitigrade
	Hyaenodontidae	digitigrade
	Hypertragulidae	unguligrade
	Ischyromyidae	plantigrade
	Jimomyidae	plantigrade
	Lagomorpha	digitigrade
	Leptictidae	plantigrade
	Leptochoeridae	unguligrade
	Leptomerycidae	unguligrade
	Mammutidae	unguligrade
Continued on next page		

Table 2 – continued from previous page

Order	Family	Stance
Mesonychia	Megalonychidae	plantigrade
	Megatheriidae	plantigrade
	Mephitidae	plantigrade
	Merycoidodontidae	digitigrade
		unguligrade
	Mesonychidae	digitigrade
	Micropternodontidae	plantigrade
	Mixodectidae	plantigrade
	Moschidae	unguligrade
	Muridae	plantigrade
	Mustelidae	plantigrade
	Mylagaulidae	fossorial
	Mylodontidae	plantigrade
	Nimravidae	digitigrade
Notoungulata	Nothrotheriidae	plantigrade
		unguligrade
	Oromerycidae	unguligrade
	Oxyaenidae	digitigrade
	Palaeomerycidae	unguligrade
	Palaeoryctidae	plantigrade
	Pampatheriidae	plantigrade
	Pantolambdidae	plantigrade
	Periptychidae	digitigrade
		unguligrade
Perissodactyla		unguligrade
	Phenacodontidae	unguligrade
Continued on next page		

Table 2 – continued from previous page

Order	Family	Stance
Primates		plantigrade
	Procyonidae	plantigrade
	Proscalopidae	plantigrade
	Protoceratidae	unguligrade
	Reithroparamyidae	plantigrade
	Sciuravidae	plantigrade
	Sciuridae	plantigrade
	Simimyidae	plantigrade
	Soricidae	plantigrade
	Suidae	digitigrade
	Talpidae	fossorial
	Tayassuidae	unguligrade
	Tenrecidae	plantigrade
	Titanoideidae	plantigrade
	Ursidae	plantigrade
	Viverravidae	plantigrade
	Zapodidae	plantigrade

94

Species mass information was gathered from multiple different sources where a plurality of the body
size estimates are from the PBDB. Body part measurements for many species are also available
through the PBDB. Just as with Smits (2015), these measurements and corresponding regression
equations were used to get mass estimates for more species. Additional mass estimates and body
part measurements were sourced from numerous publications and the Neogene Old World Database;
see the supplementary material to Smits (2015) for details. Mass was log-transformed and then
mean-centered and rescaled by dividing by two-times its standard deviation; this insures that the

magnitude of effects for both continuous and discrete covariates are comparable (Gelman, 2008; Gelman and Hill, 2007).

Table 3: Regression equations used in this study for estimating body size. Equations are presented with reference to taxonomic grouping, part name, and reference.

Group	Equation	log(Measurement)	Source
General	$\log(m) = 1.827x + 1.81$	lower m1 area	Legendre (1986)
General	$\log(m) = 2.9677x - 5.6712$	mandible length	?
General	$\log(m) = 3.68x - 3.83$	skull length	?
Carnivores	$\log(m) = 2.97x + 1.681$	lower m1 length	?
Insectivores	$\log(m) = 1.628x + 1.726$	lower m1 area	?
Insectivores	$\log(m) = 1.714x + 0.886$	upper M1 area	?
Lagomorph	$\log(m) = 2.671x - 2.671$	lower toothrow area	Tomiya (2013)
Lagomorph	$\log(m) = 4.468x - 3.002$	lower m1 length	Tomiya (2013)
Marsupials	$\log(m) = 3.284x + 1.83$	upper M1 length	?
Marsupials	$\log(m) = 1.733x + 1.571$	upper M1 area	?
Rodentia	$\log(m) = 1.767x + 2.172$	lower m1 area	Legendre (1986)
Ungulates	$\log(m) = 1.516x + 3.757$	lower m1 area	?
Ungulates	$\log(m) = 3.076x + 2.366$	lower m2 length	?
Ungulates	$\log(m) = 1.518x + 2.792$	lower m2 area	?
Ungulates	$\log(m) = 3.113x - 1.374$	lower toothrow length	?

All fossil occurrences from 64 to 2 million years ago (Mya) were binned into 31 2 million year (My) bins. This temporal length was chosen because it is approximately the resolution of the North American mammal fossil record.

Environmental and temporal covariates

The group-level covariates in this study are descriptors of species' environmental context, specifically global temperature estimates and Graham's floral intervals CITATION. Global temperature across most of the Cenozoic was calculated from Mg/Ca isotope record from deep sea carbonates (Cramer et al., 2011). Mg/Ca based temperature estimates are preferable to the frequently used $\delta^{18}\text{O}$ temperature proxy (Alroy et al., 2000; Figueirido et al., 2012; Zachos et al., 2008, 2001) because Mg/Ca estimates do not conflate temperature with ice sheet volume and depth/stratification changes; this makes it preferable as an estimate of global temperature for macroevolutionary and macroecological studies (Ezard et al., 2016).

Two aspects of the Mg/Ca-based temperature curve were included in this analysis: mean and range. Both were calculated as the mean of all respective estimates for each 2 My temporal bins. Both mean and range were then rescaled as above: subtract mean, divide by twice the standard deviation.

The other major set of environmental factors included in this study are Graham’s Cenozoic plant phases CITATION. Graham’s plant phases are holistic descriptors of the taxonomic composition of which plants were present at a given time and their relative modernity, with younger phases representing increasingly modern taxa CITATION. Graham CITATION defines four intervals from the Cretaceous to the Pliocene, though only three of these intervals are included in this analysis. Graham’s plant phases CITATION was included as a series of “dummy variables” encoding the three phases included in this analysis. This means that the first phase is synonymous with the intercept and phases

Modelling species occurrence

Two different models were used in this study: a pure-presence model and a birth-death model. Both models at their core are hidden Markov model where the latent aspect of the process has an absorbing state (Allen, 2011). The difference between these two models is if the probability of a species origination and survival are considered equal or different (Table 4). Something that is important to realize is that while there are only two state “codes” in a presence-absence matrix (i.e. 0/1), there are in fact three states in a birth-death model: never having originated, extant, and extinct. The last of these is the absorbing state, as once a species has gone extinct it cannot re-originate (Allen, 2011); this is made obvious in the transition matrices as the probability of an extinct species changing states is 0 (Table 4). See below for parameter explanations (Tables 6, and 7).

Data augmentation

All presence/absence observations are incomplete. The hidden Markov model at the core of this analysis allows for observed absences to be used meaningfully to estimate the number of unobserved

		State at $t + 1$		
		0_{never}	1	$0_{extinct}$
State at t	0_{never}	$1 - \theta$	θ	0
	1	0	θ	$1 - \theta$
	$0_{extinct}$	0	0	1

(a) Pure-presence

		State at $t + 1$		
		0_{never}	1	$0_{extinct}$
State at t	0_{never}	$1 - \phi$	ϕ	0
	1	0	π	$1 - \pi$
	$0_{extinct}$	0	0	1

(b) Birth-death

Table 4: Transition matrices for the pure-presence (4a) and birth-death (4b) models. Both of these models share the core machinery of discrete-time birth-death processes but make distinct assumptions about the equality of originating and surviving (Eq. 2, and 3). Note also that while there are only two state “codes” (0, 1), there are in fact three states: never having originated 0_{never} , present 1, extinct $0_{extinct}$ (Allen, 2011).

species. Of specific concern in this analysis is the unknown “true” size of the dataset; how many species could have actually been observed? While many species have been observed, the natural incompleteness of all observations, especially in the case of paleontological data, there are obviously many species which were never sampled (Royle and Dorazio, 2008; Royle et al., 2007).

Let N be the total number of observed species, M be the upper limit of possible species that could have existed given a model of species presence, and N^* is the all-zero histories where $N^* = M - N$. This approach assumes that $\hat{N} \sim \text{Binomial}(M, \psi)$ where \hat{N} is the estimated “true” number of species and ψ is the probability that any augmented species should actually be “present.” Because M is user defined, this approach effectively gives ψ a uniform prior over N to M (Royle and Dorazio, 2008). For this study, $M = N + \lfloor N/4 \rfloor$.

Data imputation is the process of estimating missing data for partially observed covariates (Gelman and Hill, 2007; Rubin, 1996), this is simple in a Bayesian context because data are also parameters (Gelman et al., 2013). Augmented species also have no known mass so a mass estimate must be imputed for each possible species (Royle and Dorazio, 2012). This procedure assumes that mass values for augmented species are from the same distribution as observed species. The distribution of observed mass values is estimated as part of the model, and new mass values are then generated from this distribution. This approach is an example of imputing data missing completely at random (Gelman and Hill, 2007; Royle and Dorazio, 2012). Because log mass values are rescaled as a part of this study, the body mass distribution is already known ($\mathcal{N}(0, 0.5)$); augmented species body mass

Table 5: Observation parameters

Parameter	dimensions	explanation
y	$N \times T$	observed species presence/absence
z	$N \times T$	“true” species presence/absence
p	T	probability of observing a species that is present at time t
m	N	species log mass, rescaled
α_0	1	average log-odds of p
α_1	1	change in average log-odds of p per change mass
r	T	difference from α_0 associated with time t
σ	1	standard deviation of r

160 just simply drawn from this distribution.

In addition to body mass information, the augmented species need an ecotype classification. Because
 162 these species are completely unknown, they were all classified as “augmented,” an additional
 grouping indicating their unknown biology. This classification has no biological interpretation.

164 **Observation process**

The type of hidden Markov model used in this study has three characteristic probabilities:
 166 probability p of observing a species given that it is present, probability ϕ of a species surviving from
 one time to another, and probability π of a species first appearing (Royle and Dorazio, 2008). In
 168 this formulation, the probability of a species going extinct is $1 - \pi$. For the pure-presence model
 $\phi = \pi$, while for the birth-death model $\phi \neq \pi$.

170 The probability of observing a species that is present p is modeled as a logistic regression was a
 time-varying intercept and species mass as a covariate. The effect of species mass on p was assumed
 172 linear and constant over time and given a prior reflecting a possible positive relationship; these
 assumptions are reflected in the structure of the model Equation 1. The parameters associated with
 174 this part of the model are described in Table 5.

Table 6: Parameters for the model of presence in the pure-presence model

Parameter	dimensions	explanation
z	$N \times T$	“true” species presence/absence
θ	$N \times T - 1$	probability of $z = 1$
a	$T - 1 \times D$	ecotype-varying intercept; mean value of log-odds of θ
m	N	species log mass, rescaled
b_1	1	effect of species mass on log-odds of θ
b_2	1	effect of species mass, squared, on log-odds of θ
U	$T \times D$	matrix of group-level covariates
γ	$U \times D$	matrix of group-level regression coefficients
Σ	$D \times D$	covariance matrix of a
Ω	$D \times D$	correlation matrix of a
τ	D	vector of standard deviations for each ecotype a_d

$$\begin{aligned}
y_{i,t} &\sim \text{Bernoulli}(p_{i,t}z_{i,t}) \\
p_{i,t} &= \text{logit}^{-1}(\alpha_0 + \alpha_1 m_i + r_t) \\
r_t &\sim \mathcal{N}(0, \sigma)
\end{aligned} \tag{1}$$

Pure-presence process

For the pure-presence model there is only a single probability dealing with the presence of a species θ (Table 4a). This probability was modeled as multi-level logistic regression with both species-level and group-level covariates (Gelman et al., 2013; Gelman and Hill, 2007). The parameters associated with pure-presence model are presented in Table 6 and the full sampling statement in Equation 2.

The species-level of the model (Eq. 2) is a logistic regression with varying-intercept that varies by ecotype. Additionally, species mass was included as a covariate associated with two regression coefficients allowing a quadratic relationship with log-odds of occurrence. This assumption is based on the known distribution of mammal body masses where species with intermediate mass values are more common than either small or large bodied species. These assumptions are also reflected in the choice of priors for these regression coefficients.

The values of each ecotype’s intercept are themselves modeled as regressions using the group-level covariates associated with environmental context. Each of these regressions has an associated

188 variance of possible values of each ecotype’s intercept (Gelman and Hill, 2007). In addition, the
 covariances between ecotype intercepts, given this group-level regression, are modeled (Gelman and
 190 Hill, 2007).

All parameters not modeled elsewhere were given weakly informative priors (Gelman et al., 2013)
 192 CITATION STAN MANUAL STATISTICAL RETHINKING. Weakly informative means that
 priors do not necessarily encode actual prior information but instead help regularize or weakly
 194 constrain posterior estimates. These priors have a concentrated probability density around and near
 zero; this has the effect of tempering our estimates and help prevent overfitting the model to the
 196 data (Gelman et al., 2013) CITATION STAN MANUAL STATISTICAL RETHINKING.

$$\begin{aligned}
 y_{i,t} &\sim \text{Bernoulli}(p_{i,t}z_{i,t}) & \alpha_0 &\sim \mathcal{N}(0, 1) \\
 p_{i,t} &= \text{logit}^{-1}(\alpha_0 + \alpha_1 m_i + r_t) & \alpha_1 &\sim \mathcal{N}(1, 1) \\
 r_t &\sim \mathcal{N}(0, \sigma) & \sigma &\sim \mathcal{N}^+(1) \\
 z_{i,1} &\sim \text{Bernoulli}(\rho) & b_1 &\sim \mathcal{N}(0, 1) \\
 z_{i,t} &\sim \text{Bernoulli}(\theta_{i,t}) & b_2 &\sim \mathcal{N}(-1, 1) \\
 \theta_{i,t} &= \text{logit}^{-1}(a_{t,j[i]} + b_1 m_i + b_2 m_i^2) & \gamma &\sim \mathcal{N}(0, 1) \\
 a &\sim \text{MVN}(u\gamma, \Sigma) & \tau &\sim \mathcal{N}^+(1) \\
 \Sigma &= \text{diag}(\tau)\Omega\text{diag}(\tau) & \Omega &\sim \text{LKJ}(2)
 \end{aligned} \tag{2}$$

Birth-death process

198 In the birth-death model, $\phi \neq \pi$ and so each of these probabilities are modeled separately but in a
 similar manner to how θ is modeled in the pure-presence model (Eq. 2, Table 4b). The parameters
 200 associated with the birth-death presence model are presented in Table 7 and the full sampling
 statement, including observation (Eq. 1), is described in Equation 3.

202 Similar to the pure-presence model, both ϕ and π are modeled as logistic regressions with

Table 7: Parameters for the model of presence in the pure-presence model

Parameter	dimensions	explanation
z	$N \times T$	“true” species presence/absence
ϕ	$N \times T$	probability of $z_{-,t} = 1 z_{-,t-1} = 0$
π	$N \times T - 1$	probability of $z_{-,t} = 1 z_{-,t-1} = 1$
a^ϕ	$T - 1 \times D$	ecotype-varying intercept; mean value of log-odds of θ
a^π	$T - 1 \times D$	ecotype-varying intercept; mean value of log-odds of θ
m	N	species log mass, rescaled
b_1^ϕ	1	effect of species mass on log-odds of ϕ
b_1^π	1	effect of species mass on log-odds of π
b_2^ϕ	1	effect of species mass, squared, on log-odds of ϕ
b_2^π	1	effect of species mass, squared, on log-odds of π
U	$T \times D$	matrix of group-level covariates
γ^ϕ	$U \times D$	matrix of group-level regression coefficients
γ^π	$U \times D$	matrix of group-level regression coefficients
Σ^ϕ	$D \times D$	covariance matrix of a^ϕ
Σ^π	$D \times D$	covariance matrix of a^π
Ω^ϕ	$D \times D$	correlation matrix of a^ϕ
Ω^π	$D \times D$	correlation matrix of a^π
τ^ϕ	D	vector of standard deviations for each ecotype a_d^ϕ
τ^π	D	vector of standard deviations for each ecotype a_d^π

varying-intercept and one covariate associated with two parameters. The possible relationships

204 between mass and both ϕ and π are reflected in the parameterization of the model and choice of priors (Eq. 3).

206 The intercepts of ϕ and π both vary by species ecotype and those values are themselves the product of group-level regression using environmental factors as covariates (Eq. 3); this is identical to the

208 pure presence model (Eq. 2).

$$\begin{aligned}
y_{i,t} &\sim \text{Bernoulli}(p_{i,t} z_{i,t}) & \Sigma^\phi &= \text{diag}(\tau^\phi) \Omega^\phi \text{diag}(\tau^\phi) \\
p_{i,t} &= \text{logit}^{-1}(\alpha_0 + \alpha_1 m_i + r_t) & \Sigma^\pi &= \text{diag}(\tau^\pi) \Omega^\pi \text{diag}(\tau^\pi) \\
r_t &\sim \mathcal{N}(0, \sigma) & \rho &\sim \text{U}(0, 1) \\
\alpha_0 &\sim \mathcal{N}(0, 1) & b_1^\phi &\sim \mathcal{N}(0, 1) \\
\alpha_1 &\sim \mathcal{N}(1, 1) & b_1^\pi &\sim \mathcal{N}(0, 1) \\
\sigma &\sim \mathcal{N}^+(1) & b_2^\phi &\sim \mathcal{N}(-1, 1) \\
z_{i,1} &\sim \text{Bernoulli}(\phi_{i,1}) & b_2^\pi &\sim \mathcal{N}(-1, 1) \\
z_{i,t} &\sim \text{Bernoulli}\left(z_{i,t-1} \pi_{i,t} + \sum_{x=1}^t (1 - z_{i,x}) \phi_{i,t}\right) & \gamma^\phi &\sim \mathcal{N}(0, 1) \\
& & \gamma^\pi &\sim \mathcal{N}(0, 1) \\
\phi_{i,t} &= \text{logit}^{-1}(a_{t,j[i]}^\phi + b_1^\phi m_i + b_2^\phi m_i^2) & \tau^\phi &\sim \mathcal{N}^+(1) \\
\pi_{i,t} &= \text{logit}^{-1}(a_{t,j[i]}^\pi + b_1^\pi m_i + b_2^\pi m_i^2) & \tau^\pi &\sim \mathcal{N}^+(1) \\
a^\phi &\sim \text{MVN}(U \gamma^\phi, \Sigma^\phi) & \Omega^\phi &\sim \text{LKJ}(2) \\
a^\pi &\sim \text{MVN}(U \gamma^\pi, \Sigma^\pi) & \Omega^\pi &\sim \text{LKJ}(2)
\end{aligned} \tag{3}$$

Posterior inference and model adequacy

210 Programs that implement joint posterior inference for the above models (Eqs. 2, 3) were
implemented in the probabilistic programming language Stan CITATION. The models used here
212 both feature latent discrete parameters in the large matrix z (Tables 5, 6, 7; Eqs. 1, 2, 3). All
methods for posterior inference implemented in Stan are derivative based which causes
214 complications for actually implementing the above models because integers do not have derivatives.
Instead of implementing a latent discrete parameterization, the posterior probabilities of all possible
216 states of the latent parameters z were estimated (i.e. marginalized).

Species durations at minimum range-through from the FAD to the LAD, but the incompleteness of

all observations means that the actual time of origination or extinction is unknown. The marginalization approach used here means that the probabilities all possible histories for a species are calculated, from the end members of the species having existed for the entire study interval and the species having only existed between the directly observed FAD and LAD to all possible intermediaries CITATION STAN MANUAL.

The combined size of the dataset and large number of parameters in both models (Eqs. 2, 3), specifically the total number of latent parameters that are the matrix z , means that stochastic approximate posterior inference is computationally very slow even using HMC. Instead, an approximate Bayesian approach was used: variational inference. A recently developed automatic variational inference algorithm called “automatic differentiation variational inference” (ADVI) is implemented in Stan and was used here CITATION. ADVI assumes that the posterior is Gaussian but still yields a true Bayesian posterior; this assumption is similar to quadratic approximation of the likelihood function used in maximum likelihood inference CITATION. The principal limitation of assuming the joint posterior is Gaussian is that the true topology of the log-posterior isn’t represented; this is a particular burden for scale parameters which are bound to be positive (e.g. standard deviation).

After fitting both models (Eqs. 2, 3) using ADVI, model adequacy and quality of fit was assessed using a series of posterior predictive checks CITATION CITATION. Because all Bayesian models are inherently generative, simulations of new data sets is “free” CITATION. By simulating many theoretical data sets using the observed covariate information the congruence between predictions made by the model and the observed empirical data can be assessed. By combining multiple posterior predictive tests of congruence between empirical and simulated values of interest, the holistic adequacy of the model can be analyzed CITATION.

An example posterior predictive check used in this study was comparing the observed average number of observations per species to a distribution of simulated averages; if the empirically observed value sits in the middle of the distribution than the model is adequate in reproducing the observed number of occurrences per species.

Posterior simulations for time series are start with the values at $t = 1$ and then just simulating
 246 forward.

Given parameter estimates, diversity and diversification rates are estimated through posterior
 248 predictive simulations. Given the observed presence-absence matrix y , estimates of the true
 presence-absence matrix z can be simulated and the distribution of possible occurrence histories
 250 can be analyzed. This is conceptually similar to marginalization where the probability of each
 possible occurrence history is estimated (Fig. 2).

252 The posterior distribution of z gives the estimate of standing diversity N_t^{stand} for all time points as

$$N_t^{stand} = \sum_{i=1}^M z_{i,t}. \quad (4)$$

Given estimates of N^{stand} for all time points, the estimated number of originations O_t are be
 254 estimated as

$$O_t = \sum_{i=1}^M z_{i,t} = 1 | z_{i,t-1} = 0 \quad (5)$$

and number of extinctions E_t estimated as

$$E_t = \sum_{i=1}^M z_{i,t} = 0 | z_{i,t-1} = 1. \quad (6)$$

256 Per-captia growth D^{rate} , origination O^{rate} and extinction E^{rate} rates are then calculated as

$$\begin{aligned} O_t^{rate} &= \frac{O_t}{N_{t-1}^{stand}} \\ E_t^{rate} &= \frac{E_t}{N_{t-1}^{stand}} \\ D_t^{rate} &= O_t^{rate} - E_t^{rate}. \end{aligned} \quad (7)$$

Results

Comparing the fits of the pure-presence and birth-death models

Comparison of the posterior predictive performance of the pure-presence and birth-death models reveals a striking difference in quality of the models' fits to the data (Fig. 3 and 4). The birth-death model is clearly able to reproduce the observed average number of occurrence, in contrast to the pure-birth model which greatly underestimates the observed average number of occurrences. The interpretation of these results is that the results of the birth-death model are more representative of the data than the pure-presence model, though further inspection of the posterior parameter estimates can provide further insight into why these models give different posterior predictive results (Gelman et al., 2013).

Posterior results take one of two forms: direct inspection of parameter estimates, and downstream estimates of diversity and diversification rates.

Analysis of diversity

Acknowledgements

I would like to thank K. Angielczyk, M. Foote, P. D. Polly, and R. Ree for helpful discussion and advice. This entire study would not have been possible without the Herculean effort of the many contributors to the Paleobiology Database. In particular, I would like to thank J. Alroy and M. Uhen for curating most of the mammal occurrences recorded in the PBDB. This is Paleobiology Database publication XXX.

References

Allen, L. J. S. 2011. An introduction to stochastic processes with applications to biology. 2nd ed. Chapman and Hall/CRC, Boca Raton, FL.

Alroy, J. 1996. Constant extinction, constrained diversification, and uncoordinated stasis in North
280 American mammals. *Palaeogeography, Palaeoclimatology, Palaeoecology* 127:285–311.

———. 2009. Speciation and extinction in the fossil record of North American mammals. Pages
282 302–323 *in* R. K. Butlin, J. R. Bridle, and D. Schluter, eds. *Speciation and patterns of diversity*.
Cambridge University Press, Cambridge.

284 Alroy, J., P. L. Koch, and J. C. Zachos. 2000. Global climate change and North American
mammalian evolution. *Paleobiology* 26:259–288.

286 Bambach, R. K., A. M. Bush, and D. H. Erwin. 2007. Autecology and the filling of ecospace: Key
metazoan radiations. *Palaeontology* 50:1–22.

288 Blois, J. L., and E. A. Hadly. 2009. Mammalian Response to Cenozoic Climatic Change. *Annual
Review of Earth and Planetary Sciences* 37:181–208.

290 Brown, A. M., D. I. Warton, N. R. Andrew, M. Binns, G. Cassis, and H. Gibb. 2014. The
fourth-corner solution - using predictive models to understand how species traits interact with
292 the environment. *Methods in Ecology and Evolution* 5:344–352.

Bush, A. M., and R. K. Bambach. 2011. *Paleoecologic Megatrends in Marine Metazoa*, vol. 39.

294 Bush, A. M., R. K. Bambach, and G. M. Daley. 2007. Changes in theoretical ecospace utilization in
marine fossil assemblages between the mid-Paleozoic and late Cenozoic. *Paleobiology* 33:76–97.

296 Clyde, W. C., and P. D. Gingerich. 1998. Mammalian community response to the latest Paleocene
thermal maximum: an isotaphonomic study in the northern Bighorn Basin, Wyoming. *Geology*
26:1011–1014.
298

Cramer, B. S., K. Miller, P. Barrett, and J. Wright. 2011. Late Cretaceous-Neogene trends in deep
300 ocean temperature and continental ice volume: Reconciling records of benthic foraminiferal
geochemistry ($\delta^{18}\text{O}$ and Mg/Ca) with sea level history. *Journal of Geophysical Research: Oceans*
302 116:1–23.

Eronen, J. T., C. M. Janis, C. P. Chamberlain, and A. Mulch. 2015. Mountain uplift explains

304 differences in Palaeogene patterns of mammalian evolution and extinction between North
America and Europe. *Proceedings of the Royal Society B: Biological Sciences* 282:20150136.

306 Ezard, T. H. G., A. Purvis, and H. Morlon. 2016. Environmental changes define ecological limits to
species richness and reveal the mode of macroevolutionary competition. *Ecology Letters*
308 19:899–906.

Figueirido, B., C. M. Janis, J. A. Pérez-Claros, M. De Renzi, and P. Palmqvist. 2012. Cenozoic
310 climate change influences mammalian evolutionary dynamics. *Proceedings of the National
Academy of Sciences* 109:722–727.

312 Foote, M. 2001. Inferring temporal patterns of preservation, origination, and extinction from
taxonomic survivorship analysis. *Paleobiology* 27:602–630.

314 Foote, M., and J. J. Sepkoski. 1999. Absolute measures of the completeness of the fossil record.
Nature 398:415–7.

316 Gelman, A. 2008. Scaling regression inputs by dividing by two standard deviations. *Statistics in
Medicine* pages 2865–2873.

318 Gelman, A., J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari, and D. B. Rubin. 2013. *Bayesian
data analysis*. 3rd ed. Chapman and Hall, Boca Raton, FL.

320 Gelman, A., and J. Hill. 2007. *Data Analysis using Regression and Multilevel/Hierarchical Models*.
Cambridge University Press, New York, NY.

322 Janis, C. M. 1993. Tertiary mammal evolution in the context of changing climates, vegetation, and
tectonic events. *Annual Review of Ecology and Systematics* 24:467–500.

324 Janis, C. M., J. Damuth, and J. M. Theodor. 2000. Miocene ungulates and terrestrial primary
productivity: where have all the browsers gone? *Proceedings of the National Academy of Sciences*
326 97:7899–904.

Janis, C. M., G. F. Gunnell, and M. D. Uhen. 2008. Evolution of Tertiary mammals of North

- America. Vol. 2. Small mammals, xenarthrans, and marine mammals. Cambridge University Press, Cambridge.
- Janis, C. M., K. M. Scott, and L. L. Jacobs. 1998. Evolution of Tertiary mammals of North America. Vol. 1. Terrestrial carnivores, ungulates, and ungulatelike mammals. Cambridge University Press, Cambridge.
- Legendre, S. 1986. Analysis of mammalian communities from the Late Eocene and Oligocene of Southern France. *Paleovertebrata* 16:191–212.
- Lloyd, G. T., J. R. Young, and A. B. Smith. 2011. Taxonomic Structure of the Fossil Record is Shaped by Sampling Bias. *Systematic Biology* 61:80–89.
- Royle, J. A., and R. M. Dorazio. 2008. Hierarchical modeling and inference in ecology: the analysis of data from populations, metapopulations and communities. Elsevier, London.
- . 2012. Parameter-expanded data augmentation for Bayesian analysis of capture-recapture models. *Journal of Ornithology* 152:521–537.
- Royle, J. A., R. M. Dorazio, and W. a. Link. 2007. Analysis of Multinomial Models With Unknown Index Using Data Augmentation. *Journal of Computational and Graphical Statistics* 16:67–85.
- Royle, J. A., J. D. Nichols, M. Kéry, E. Ranta, and M. Kery. 2014. detection is of species when Modelling occurrence and abundance imperfect 110:353–359.
- Rubin, D. B. 1996. Multiple imputation after 18+ years. *Journal of the American Statistical Association* 91:473–489.
- Smits, P. D. 2015. Expected time-invariant effects of biological traits on mammal species duration. *Proceedings of the National Academy of Sciences* 112:13015–13020.
- Strömberg, C. A. E. 2005. Decoupled taxonomic radiation and ecological expansion of open-habitat grasses in the Cenozoic of North America. *Proceedings of the National Academy of Sciences of the United States of America* 102:11980–4.

- 352 Tomiya, S. 2013. Body Size and Extinction Risk in Terrestrial Mammals Above the Species Level.
The American Naturalist 182:196–214.
- 354 Wang, S. C., P. J. Everson, H. J. Zhou, D. Park, and D. J. Chudzicki. 2016. Adaptive credible
intervals on stratigraphic ranges when recovery potential is unknown. Paleobiology 42:240–256.
- 356 Wang, S. C., and C. R. Marshall. 2016. Estimating times of extinction in the fossil record. Biology
Letters 12:20150989.
- 358 Warton, D. I., B. Shipley, and T. Hastie. 2015. CATS regression - a model-based approach to
studying trait-based community assembly. Methods in Ecology and Evolution 6:389–398.
- 360 Zachos, J. C., G. R. Dickens, and R. E. Zeebe. 2008. An early Cenozoic perspective on greenhouse
warming and carbon-cycle dynamics. Nature 451:279–283.
- 362 Zachos, J. C., M. Pagani, L. Sloan, E. Thomas, and K. Billups. 2001. Trends, rhythms, and
aberrations in global climate 65 Ma to present. Science 292:686–693.




Figure 1: Conceptual diagram of the paleontological fourth corner problem. The observed presence matrix (orange) is the empirical presence/absence pattern for all species for all time points; this matrix is an incomplete observation of the “true” presence/absence pattern (purple). The estimated true presence matrix is modeled as a function of both environmental factors over time (blue) and multiple species traits (green). Additionally, the affect of environmental factors on species traits are also modeled as traits are expected to mediate the effects of a species environmental context. This diagram is based partially on material presented in Brown et al. (2014) and Warton et al. (2015).

	Time Bin							
	1	2	3	4	5	6	7	8
Observed	0	0	0	1	0	1	1	0
Certain	?	?	?	1	1	1	1	?
Potential	0	0	0	1	1	1	1	0
Potential	0	0	1	1	1	1	1	0
Potential	0	1	1	1	1	1	1	0
Potential	1	1	1	1	1	1	1	0
Potential	0	0	0	1	1	1	1	1
Potential	0	0	1	1	1	1	1	1
Potential	0	1	1	1	1	1	1	1
Potential	1	1	1	1	1	1	1	1

Figure 2: Conceptual figure of all possible occurrence histories for an observed species. The first row represents the observed presence/absence pattern for a single species at eight time points. The second row corresponds to the known aspects of the “true” occurrence history of that species. The remaining rows correspond to all possible occurrence histories that are consistent with the observed data. The process of parameter marginalization described in the text




Figure 3: Comparison of the average observed number of occurrences per species (blue line) to the average number of occurrences from 100 posterior predictive datasets using the posterior estimates from the pure-presence model.




Figure 4: Comparison of the average observed number of occurrences per species (blue line) to the average number of occurrences from 100 posterior predictive datasets using the posterior estimate from the birth-death model.




Figure 5: Probability of a mammal ecotype occurring over time as estimated from the pure-presence model. Each panel depicts 100 random samples from the model's posterior. The columns are by locomotor category and rows by dietary category; their intersections are the observed and analyzed ecotypes. Panels with no lines are ecotypes not observed in the dataset.




Figure 6: Probability of a mammal ecotype origination probabilities at each time point as estimated from the birth-death model. Each panel depicts 100 random samples from the model's posterior. The columns are by locomotor category and rows by dietary category; their intersections are the observed and analyzed ecotypes. Panels with no lines are ecotypes not observed in the dataset.




Figure 7: Probability of a mammal ecotype survival probabilities at each time point as estimated from the birth-death model. Each panel depicts 100 random samples from the model's posterior. The columns are by locomotor category and rows by dietary category; their intersections are the observed and analyzed ecotypes. Panels with no lines are ecotypes not observed in the dataset.




Figure 8: Estimates of the effect of species mass on probability of observing a present species (p). Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units. Estimates are from the pure-presence model.




Figure 9: Estimates of the effect of species mass on probability of observing a present species (p). Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units. Estimates are from the birth-death model.




Figure 10: Mean estimate of the effect of species mass on the probability of a species occurrence for each of the three plant phases. The effect of mass is considered constant over time and that the only aspect of the model that changes with plant phase is the intercept of the relationship between mass and occurrence. The three plant phases are indicated by the color of the line. Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units.




Figure 11: Mean estimate of the effect of species mass on the probability of a species originating for each of the three plant phases. The effect of mass is considered constant over time and that the only aspect of the model that changes with plant phase is the intercept of the relationship between mass and origination. The three plant phases are indicated by the color of the line. Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units.




Figure 12: Mean estimate of the effect of species mass on the probability of a species survival for each of the three plant phases. The effect of mass is considered constant over time and that the only aspect of the model that changes with plant phase is the intercept of the relationship between mass and survival. The three plant phases are indicated by the color of the line. Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units.




Figure 13: Estimated effects of the group-level covariates describing environmental context on log-odds of species occurrence. These estimates are from the pure-presence model.




Figure 14: Estimated effects of the group-level covariates describing environmental context on log-odds of species origination. These estimates are from the birth-death model.




Figure 15: Estimated effects of the group-level covariates describing environmental context on log-odds of species survival. These estimates are from the birth-death model.




Figure 16: Posterior of standing log-diversity of North American mammals for the Cenozoic as estimated from the birth-death model; 100 posterior draws are plotted to indicate the uncertainty in these estimates. The dramatic differences between diversity estimates at the first and second time points in this series are caused by a well known edge effect in discrete-time birth-death models caused by $\phi_{-,t=1}$ being partially undefined (Royle and Dorazio, 2008).




Figure 17: Posterior estimates of North American mammal diversification rates for the Cenozoic; 100 estimates are plotted to indicate the uncertainty in these estimates. As a reminder, diversification rate is the difference between origination and extinction rates and is in units of species gained per species present per time unit (2 My).