

The macroecology behind macroevolution: North American
mammal functional diversity and its relation to environmental
change

Peter D. Smits^{1,*}

1. University of California – Berkeley, California, 94720.

* Corresponding author; e-mail: psmits@berkeley.edu.

Manuscript elements:

Keywords: macroecology, macroevolution, paleobiology, species pool, community assembly

Manuscript type: Article

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

The set of species in a region changes over time as new species enter through speciation or immigration and as species leave the system through extinction and extirpation. How a regional species pool changes over time is the product of many processes acting at multiple levels of organization. Changes in the functional composition of a regional species pool are changes that occur across all local communities drawn from that species pool. While a species' presence in a local community is due to the availability of the necessary biotic-biotic or biotic-abiotic interactions that enable coexistence, a species' presence in a regional species pool just requires that at least one local community has that set of necessary interactions. The goal of this analysis is to understand when, and possibly for what reasons, mammal ecotypes are enriched or depleted relative to their average diversity. Here, I analyze the diversity history of North American mammals ecotypes for most of the Cenozoic (the last 65 million years). This analysis frames mammal diversity in terms of both their means of interacting with the biotic and abiotic environment (i.e. functional group or ecotype) as well as their regional and global environmental context. Using two hierarchical Bayesian hidden Markov models of diversity, I find that changes to mammal diversity are driven more by the influx of new species than by selective extinction. I also find that the only ecotypes which experience a near constant increase in diversity over time are digitigrade and unguligrade herbivores, while arboreal ecotypes become increasingly rare and in many cases disappear entirely from the species pool over the Cenozoic. Additionally, I find that global temperature is only associated with the origination of some mammal ecotypes but, in almost all cases, does not affect the extinction of mammal ecotypes.

Introduction

Changes to species diversity are the result of evolutionary and ecological processes acting in concert and continually over time. Local communities are shaped by dispersal and local ecological processes such as resource competition and predator-prey relationships. The constituent species of these communities are drawn from a regional species pool, or the set of all species that are present in at least one community within a region (Harrison and Cornell, 2008; Mittelbach and Schemske, 2015; Urban et al., 2008). Species dispersal from the regional species pool to the local communities is a

sorting process shaped by biotic and abiotic environmental filters which are mediated by those
30 species' traits (Cottenie, 2005; Elith and Leathwick, 2009; Harrison and Cornell, 2008; Loeuille and
Leibold, 2008; Shipley et al., 2006; Urban et al., 2008). Regional species pools are shaped by
32 speciation, extinction, migration, and extirpation. The gain or loss of regional diversity is the result
of macroevolutionary dynamics which, in turn, shape the downstream macroecological dynamics of
34 the regional species pool and its constituent local communities (Harrison and Cornell, 2008;
Mittelbach and Schemske, 2015; Urban et al., 2008). In turn, the distribution of species within
36 regional species pool represents the expected distribution of local communities under a purely
diffusive process of community assembly as would be expected neutral theory of biodiversity
38 CITATION.

Fundamentally, all species respond differently to climate and environmental change (Blois and
40 Hadly, 2009). Those species with similar ecological roles within a regional species pool can be
described as belonging to a guild or functional group (Bambach, 1977; Brown and Maurer, 1989;
42 Simberloff and Dayan, 1991; Valentine, 1969; Wilson, 1999). Species within the same functional
group are expected to have more similar macroecological dynamics to each other than to species of
44 a different functional group. By focusing on the relative diversity of functional groups, changes to
diversity are interpretable as changes to the set of ways species within a species pool could interact
46 with the biotic and abiotic environment.

A key question when comparing communities or regional species pools based on their functional
48 composition is whether specific functional groups are enriched or depleted and why; what are the
processes that led to a species pool having the functional composition it does (Blois and Hadly,
50 2009; Brown and Maurer, 1989; McGill et al., 2006; Smith et al., 2008; Weber et al., 2017)?

Comparisons of contemporaneous regional species pools can only determine if a functional group is
52 enriched or depleted in one species pool relative to the other species pools CITATION. These types
of comparison do not take into account if a functional group is enriched or depleted relative to its
54 diversity over time (Blois and Hadly, 2009). While a species pool may be depleted of a functional
group relative to other contemporaneous species pools, that same functional group may be actually
56 be enriched in that species pool relative to its historical diversity. Because the processes which

shape regional species pool diversity (e.g. origination, extinction) operate on much longer time scales than is possible for studies of the Recent, paleontological data provides a unique opportunity to observe and estimate the changes to functional diversity and how species functional traits and environmental context can shape the enrichment or depletion of functional groups within a regional species pool (Blois and Hadly, 2009; Smith et al., 2008). Being able to identify if the diversity of a functional group is depleted relative to their long term average diversity in the species pool is particularly useful in conservation settings; species in depleted groups are most likely more at risk of extinction than species in enriched groups, even if those enriched groups are relatively rare when compared to the functional composition of other contemporaneous species pools.

The paleontological record of North American mammals for the Cenozoic (\sim 66 million years ago to the present) provides one of the best opportunities for understanding how regional species pool functional diversity changes over time. The North American mammal record is a relatively complete temporal sequence for the entire Cenozoic which is primarily, but not exclusively, based on fossil localities from the Western Interior of North America (Alroy, 1996, 2009; Alroy et al., 2000). Additionally, mammal fossils preserve a lot of important physiological information, such as teeth, so that functional traits like the dietary/trophic category of species are easy to estimate (Eronen et al., 2010; Polly et al., 2011, 2015).

The goals of this study are to understand when are unique functional groups enriched or depleted in the North American mammal regional species pool and to estimate the relationship between these changes to regional ecotypic diversity and changes to their environmental context. My contribution is to develop a joint model of observation, origination, survival, as well as the effects of species traits and environmental factors on these processes.

Background

The history of standing diversity for all mammals along with that some individual clades of North American mammals for the Cenozoic has been the focus of considerable study (Alroy, 1996, 2009; Alroy et al., 2000; Badgley and Finarelli, 2013; Blois and Hadly, 2009; Figueirido et al., 2012;

Fraser et al., 2015; Janis, 1993; Janis and Wilhelm, 1993; Pires et al., 2015; Quental and Marshall, 84 2013; Silvestro et al., 2015; Slater, 2015; Smits, 2015). Previous approaches to understanding mammal diversity, both in North America and elsewhere, fall into a number of overlapping 86 categories: total diversity (Alroy, 1996; Alroy et al., 2000; Figueirido et al., 2012; Liow et al., 2008), with/between guild comparisons (Janis et al., 2004; Janis, 2008; Janis et al., 2000; Janis and 88 Wilhelm, 1993; Jernvall and Fortelius, 2004; Pires et al., 2015), within/between clade comparisons (Cantalapiedra et al., 2017; Fraser et al., 2015; Quental and Marshall, 2013; Silvestro et al., 2015; 90 Slater, 2015), and estimating the impact of environmental process on total diversity (Alroy et al., 2000; Badgley and Finarelli, 2013; Badgley et al., 2017; Blois and Hadly, 2009; Eronen et al., 2015; 92 Fraser et al., 2015; Janis, 1993; Janis and Wilhelm, 1993). Each of these studies provide a limited perspective on the macroevolutionary and macroecological processes shaping diversity and 94 diversification. Integration across perspectives is necessary for producing a holistic and internally consistent picture of how the North American mammal species pool has changed through time. One 96 of the goals of this study is to present a framework for approaching hypotheses about diversity and diversification through multiple lenses simultaneously through a joint model so that any inferences 98 are better constrained and the relative importance of species' ecological function, taxonomic affinities, and environmental context may be better elucidated.

100 The narrative of the diversification of North American mammals over the Cenozoic is one of gradual change. There is little convincing evidence that there have been any sudden 102 cross-functional or cross-taxonomic group turnover events in mammal diversity at any point in the Cenozoic record of North America (Alroy, 1996, 2009; Alroy et al., 2000; Eronen et al., 2015; Janis, 104 1993). Instead of being concentrated at specific time intervals, species turnover has been found to be distributed through time. It is then expected then that, for this analysis, turnover events or 106 periods of rapid diversification or depletion should not occur simultaneously for all functional groups under study. Additionally, changes to mammal diversification seem to be primarily driven by 108 changes to origination rate and not to extinction (Alroy, 1996, 2009; Alroy et al., 2000). An unresolved aspect of the general history of mammal diversification is whether that diversity is 110 limited or self-regulating; namely, to what extent is mammal diversification diversity-dependent

(Alroy, 2009; Harmon and Harrison, 2015; Rabosky, 2013; Rabosky and Hurlbert, 2015). Similarity,
112 this question can also be asked of specific functional groups (Jernvall and Fortelius, 2004; Quental
and Marshall, 2013; Silvestro et al., 2015; Van Valkenburgh, 1999).

114 Within the overall narrative of mammal diversity, the histories of some functional and taxonomic
groups are better understood than others. These groups include ungulate herbivores and Neogene
116 carnivores which have particularly good fossil records and have been the focus of previous analyses.

The diversity history of ungulate herbivores has been characterized by more recently originating
118 taxa having longer legs, higher crowned teeth, and a shift from graze-dominated to
browse-dominated diets than their earlier originating counterparts (Cantalapiedra et al., 2017;
120 Fraser et al., 2015; Janis et al., 2004; Janis, 1993, 2008; Janis et al., 2000). The mechanisms which
drive this pattern are theorized to be some combination of tectonic activity driving environmental
122 change such as the drying of the western interior of North America due mountain building and
global temperature and environmental change such as the formation of polar icecaps (Badgley et al.,
124 2017; Blois and Hadly, 2009; Eronen et al., 2015; Janis, 2008).

In contrast, the origination of modern cursorial carnivore forms was not until later in the Cenozoic;
126 this is not to say that carnivore diversity only grew in the late Cenozoic, but that those forms were
late entrants (Janis and Wilhelm, 1993). Instead, the diversity history of carnivores is reflective of
128 density-dependence or some other form of self-regulation (Silvestro et al., 2015; Slater, 2015; Van
Valkenburgh, 1999). Specifically, it has been proposed that different canid clades have replaced each
130 other as the dominate members of their functional group within the species pool (Silvestro et al.,
2015; Van Valkenburgh, 1999). It is then expected that, for this analysis, the diversity of digitigrade
132 and plantigrade carnivores (i.e. the “carnivore” guild of Van Valkenburgh (1999)) should be
relatively constant for the Cenozoic or at least have plateaued by the Neogene.

134 In a relevant study, Smits (2015) found that functional traits such as a species dietary or locomotor
category structure differences in mammal extinction risk. In particular, arboreal taxa were found to
136 have a shorter duration on average than species from other locomotor categories (Smits, 2015). Two
possible scenarios that could yield this pattern were proposed: the extinction risk faced by arboreal

138 species is constant and high for the entire Cenozoic or the Paleogene and Neogene represent
139 different regimes and extinction risk increased in the Neogene, thus driving up the Cenozoic average
140 extinction risk. These two possible explanations have clear and testable predictions with respect to
141 the diversity history of arboreal taxa: 1) if arboreal taxa always have an elevated extinction risk
142 when compared to other taxa, then the diversity history of arboreal taxa is expected to be constant
143 with time, albeit possibly at low diversity; and 2) if the Paleogene and Neogene represent difference
144 selective regimes with the former being associated with lower extinction risk than the latter, then
145 the diversity history of arboreal taxa are expected to be present in the Paleogene but depleted or
146 absent from the species pool during the Neogene.

There is a lack of consensus as to the effect of species body size on mammal diversity and aspects of
148 the diversification processes, specifically extinction (Liow et al., 2008, 2009; Smits, 2015; Tomiya,
149 2013). Species body size is frequently framed as an important biological descriptor because of its
150 correlation with other important and relevant ecological traits such as metabolic rate and home
151 range size (Brown, 1995). It is also relatively easy to estimate for extinct species using proxy
152 measures and regression equations, as was done in this study (see below). However, body size is
153 normally analyzed without simultaneous reference to other relevant species traits (Huang et al.,
154 2017; Liow et al., 2008; Raia et al., 2012; Smith et al., 2004), but see (Smits, 2015); this combined
155 with the high amount of correlation between life history traits and body size limits process-based
156 inference, because the actual causal mechanisms underlying an observed pattern are obscured or
157 missing.

158 The climate history of the Cenozoic can be broadly described as a gradual cooling trend, with polar
159 ice-caps forming in the Neogene (Cramer et al., 2011; Zachos et al., 2008, 2001). There are of course
160 exceptions to this pattern such as the Eocene climatic optimum, the mid-Miocene climatic
161 optimum, and the sudden drop in temperature at the Eocene/Oligocene boundary (Zachos et al.,
162 2008, 2001). In terms of the North American biotic environment, the Cenozoic is additionally
163 characterized by major transition from having closed, partially forested biomes being common in
164 the Paleogene to the landscape being dominated by savannah and grasslands biomes by the
Neogene (Blois and Hadly, 2009; Janis, 1993; Janis et al., 2000; Strömberg, 2005). Additionally, the

¹⁶⁶ landscape structure and topology of North America changed substantially over the Cenozoic with
¹⁶⁷ mountain uplift and other tectonic actives in Western North America (Badgley and Finarelli, 2013;
¹⁶⁸ Blois and Hadly, 2009; Eronen et al., 2015; Janis, 2008). This type of geological activity affects
¹⁶⁹ both local climates as well as continental weather patterns while also mobilizing increased grit into
¹⁷⁰ the environment, something which may be responsible for increasing trend of hyposodony (high
¹⁷¹ crowned teeth) among ungulate and rodent herbivores (Damuth and Janis, 2011; Janis, 1993;
¹⁷² Jardine et al., 2012; Jernvall and Fortelius, 2002) Badgley CITATION.

The Eocene-Oligocene transition has been observed to be associated with extinction of many
¹⁷⁴ ungulate taxa (Janis, 2008). This boundary also marks the transition from the Paleogene to the
Neogene and from herbivores being browsing dominated to grazing dominated, though not
¹⁷⁶ concurrently (Janis, 1993; Strömberg, 2005). Additionally, the Paleogene-Neogene boundary marks
the approximate start of Antarctic ice sheets, which were previously absent (Zachos et al., 2008).
¹⁷⁸ There is an observed stability in estimates of global temperature from the E/O transition till the
end of the Miocene called the Mid-Miocene climatic optimum (Zachos et al., 2008, 2001). The
¹⁸⁰ Mid-Miocene climatic optimum is bookended by periods of temperature decline. We would then
expect that, for the Miocene, turnover and other diversification events would most likely be due to
¹⁸² biological interactions or immigration and not biotic-abiotic interactions because of the constancy
of the climate, and that those groups that are driven primarily by environmental factors, the
¹⁸⁴ Miocene would be a period of marked by an absence of major changes to diversity or the
diversification process.

¹⁸⁶ The effect of climate on mammal diversity and its accompanying diversification process has been
the focus of considerable research with a slight consensus favoring mammal diversification being
¹⁸⁸ more biologically-mediated than climate-mediated when considering the entire Cenozoic (Alroy
et al., 2000; Clyde and Gingerich, 1998; Figueirido et al., 2012). However, differences in temporal
¹⁹⁰ and geographic scale seem to underly the contrast between these two perspectives. For example
when the mammal fossil record analyzed at small temporal and geographic scales a correlation
¹⁹² between diversity and climate is observable (Clyde and Gingerich, 1998). However, when the record
is analyzed at the scale of the continent and most of the Cenozoic this correlation disappears (Alroy

¹⁹⁴ et al., 2000). This result, however, does not go against the idea that there may be short periods of correlation between diversity and climate and that this relationship can change or even reverse
¹⁹⁶ direction over time; this type result means that there is no single direction or longterm correlation between diversity and climate (Figueirido et al., 2012).

¹⁹⁸ In the case of a fluctuating correlation between diversity and climate it is hard to make the argument for an actual causal link between the two without modeling the underlying ecological
²⁰⁰ differences between species; after all, species respond differently based on their individual ecologies (Blois and Hadly, 2009). When analysis is based on diversity or taxonomy alone no mechanisms are
²⁰² possible to infer. Taxonomy, like body size, stands in for many important species traits to the point that mechanistic or process based inference is impossible. While emergent patterns might
²⁰⁴ correspond to taxonomic grouping, this itself is an emergent phenomenon. Instead, by framing hypotheses in terms of species traits and their environmental context, these emergent phenomena
²⁰⁶ can be observed and analyzed rather than assumed.

Foreground

²⁰⁸ The fourth-corner problem is conceptual and statistical framework to explaining the patterns of either species abundance or presence/absence in a community as a product of species traits,
²¹⁰ environmental factors, and the interaction between traits and environment (Brown et al., 2014;
Jamil et al., 2013; Pollock et al., 2012; Warton et al., 2015); effectively uniting climate-based species
²¹² distribution modeling (SDMs) with trait-based community assembly models (CATS, MaxEnt). The fourth-corner problem is normally phrased in modern ecological studies as understanding how traits
²¹⁴ and environment interact to predict the occurrence of species at localities distributed across a region (Jamil et al., 2013; Pollock et al., 2012).

²¹⁶ This conceptual framework can be extended to include time when considering occurrence as a three-dimensional: species at localities in space over time. This extension changes the goal of
²¹⁸ predicting just occurrence to one of predicting species gain and loss at localities. However, the temporal limitations of modern ecological studies limit studying species over their entire durations,

220 where speciation and extinction govern the occurrence of species. By considering the patterns of
fossil occurrences in the geological record the macroevolutionary processes governing species'
222 (macro)ecology can be better understood. One limitation of the fossil record, however, is a lack of
spatial resolution for most taxonomic groups and periods of time. As such, paleontological data
224 "looks" at a different side of the three-dimensional occurrence matrix of the extended fourth-corner
problem than modern ecological data.

226 In this study, the fourth-corner problem is phrased as understanding how mammal functional
groups respond to environmental change in order to predict the origination and survival of species
228 over time (Fig. 1). Additionally, I also consider the incompleteness of the fossil record and the static
effect of other species descriptors not related to functional group on origination and survival.




Figure 1: Conceptual diagram of the analysis at the center of this study. The observed presence matrix (orange) is the empirical presence/absence pattern for all species for all time points; this matrix is an incomplete observation of the "true" presence/absence pattern (purple). This observation process is modeled as a function of both time (red) and a selection of species traits (green). The estimated true presence matrix is modeled as a function of both environmental factors over time (blue) and multiple species traits (green). Additionally, the effects of environmental factors on some of those species traits are also modeled, as traits are expected to mediate the effects of a species environmental context. This diagram is based partially on material presented in Brown et al. (2014) and Warton et al. (2015).

230 My approach to delimiting and assigning mammal functional groups is inspired on the ecocube
heuristic used to classify marine invertebrate species by three functional traits (Bambach et al.,
232 2007; Bush and Bambach, 2011; Bush et al., 2007; Bush and Novack-Gottshall, 2012;
Novack-Gottshall, 2007; Villéger et al., 2011). In this study, the two functional traits used to define
234 a species' functional group are dietary (e.g. herbivore, carnivore, etc.) and locomotor category (e.g.
arboreal, unguligrade, etc.). Species body mass was also included as a species trait in this analysis,
236 but not as a trait for defining a functional group; instead, its inclusion is principally to control for
differences in species dynamics that driven by mass and not functional group.

238 The environmental covariates included in this study are estimates of global temperature as well as
which of three high-level North American plant taxonomic phases corresponds to that temporal
240 unit (Cramer et al., 2011; Graham, 2011). These covariates were chosen because they provide a
characterizations of the environmental context of the entire North American regional species pool
242 for most of the Cenozoic. Importantly, the effects of a species functional group on diversity are
themselves modeled as functions of environmental factors (Fig. 1) allowing for inference as to how a
244 species ecology can mediate selective pressures do to its environmental context.

All observations, paleontological or modern, are made with uncertainty. With presence/absence
246 data this uncertainty comes from not knowing if an absence is a “true” absence or just a failure to
observe (Foote, 2001; Foote and Sepkoski, 1999; Lloyd et al., 2011; Royle and Dorazio, 2008; Royle
248 et al., 2005; Wang and Marshall, 2016). For paleontological data, the incomplete preservation and
sampling of species means that the true times of origination or extinction may not be observed
250 (Foote, 2001; Foote and Sepkoski, 1999; Wang et al., 2016; Wang and Marshall, 2016). The model
used in this analysis is a translation of the conceptual framework described above (Fig. 1) into a
252 statistical model in order to estimate the relative diversity of mammal functional groups over time
and how those functional groups respond to changes to environmental context while taking into
254 account the fundamental incompleteness of the fossil record.

Ultimately, the goals of this analysis are to understand when are different functional groups
256 enriched or depleted in the North American mammal regional species pool and how these changes

in functional diversity are related to changes in species' environmental context. In the analysis
258 performed here, I consider multiple covariates which describe a species' macroecology and
environmental context. In order to analyze this complex, multi-level question and accompanying
260 highly-structured data set, I developed a hierachal Bayesian model combing the fourth-corner
modeling approach with a model of an observation-occurrence or observation-originatation-extinction
262 process.

Materials and Methods

264 Taxon occurrences and species-level information

All fossil occurrence information used in this analysis was downloaded from the Paleobiology
266 Database (PBDB). The initial download restricted occurrences to Mammalia observed in North
America between the Maastrichtian (72-66 Mya) and Gelasian (2.58-1.8 Mya) stages (Cohen et al.,
268 2015). Taxonomic, stratigraphic, and ecological metadata for each occurrence and species was also
downloaded. A new download for a raw, unfiltered PBDB datafile following the same criterion used
270 here is available at <http://goo.gl/2s1geU>. The raw datafile used as a part of this study, along
with all code for filtering and manipulating this download is available at
272 <http://github.com/psmits/copings>.

After being downloaded, the raw occurrence data was then sorted, cleaned, and manipulated
274 programmatically before analysis. Occurrences were restricted to those occurring between 64 and 2
million years ago (Mya); this age restriction was to insure that observation time series lines up with
276 the temperature time series (Cramer et al., 2011). All taxa whose life habit was classified as either
volant (i.e. Chiroptera) or aquatic (e.g. Cetacea) were excluded from this analysis because of their
278 lack of direct applicability to the study of terrestrial species pools.

Many species taxonomic assignments as present in the raw PBDB data were updated for accuracy
280 and consistency. Species present in the PBDB have some taxonomic information, including possible
Family and Order assignments. In order to increase consistency between species and reflect more

recent taxonomic assignments, each species taxonomic assignments updated as follows: 1) species family and order assignemnts as present in the Encyclopedia of life (<http://eol.org>) was downloaded using the *taxize* package for R; 2) for species not present in the EoL or not assigned order, their taxonomic inforation was further updated based on whatever family information was recorded in the PBDB or EoL; 3) for species still missing order assignemnts, their genus information was used to assign either an order or family, which was then used to assign an order.

This procedure is similar to that used in Smits (2015) and is detailed in the code repository associated with this study.

Species functional group is defined as the combination of locomotor and diet categories; the goal is to classify species based on the manner with which they interact with their environment. Mammal species records in the PBDB have life habit (i.e. locomotor category) and dietary category assignments. In order to simplify interpretation, analysis, and per-functional group sample size these classifications were coarsened in a similar manner to Smits (2015) (Table 1). Ground dwelling species locomotor categories were then reassigned based on the ankle posture associated with their taxonomic group, as described in Table 2 (Carrano, 1999). Ankle posture was assumed uniform for all species within a taxonomic group except for those species assigned a non-ground dwelling locomotor category in the PBDB, which retained their non-ground dwelling assignment. All species for which it was possible to assign a locomotor category had one assigned, including species for which post-cranaia are unknown but for which a taxonomic grouping is known. Ground dwelling species which were unable to be reassigned based on ankle posture were excluded from analysis.

Finally, ecotype categories with less than 10 total species were excluded, yielding a total of 18 observed ecotypes out of a possible 24.

Table 2: Ankle posture assignment as based on taxonomy. Assignments are based on (Carrano, 1999). Taxonomic groups are presented alphabetically and without reference for their relatedness.

Family	Stance
Ailuridae	plantigrade
Allomyidae	plantigrade

Continued on next page

Table 2 – continued from previous page

Family	Stance
Amphicyonidae	plantigrade
Amphilemuridae	plantigrade
Anthracotheriidae	digitigrade
Antilocapridae	unguligrade
Apheliscidae	plantigrade
Aplodontidae	plantigrade
Apternodontidae	scansorial
Arctocyonidae	unguligrade
Barbourofelidae	digitigrade
Barylambdidae	plantigrade
Bovidae	unguligrade
Camelidae	unguligrade
Canidae	digitigrade
Cervidae	unguligrade
Cimolodontidae	scansorial
Coryphodontidae	plantigrade
Cricetidae	plantigrade
Cylindrodontidae	plantigrade
Cyriacotheriidae	plantigrade
Dichobunidae	unguligrade
Dinocerata	unguligrade
Dipodidae	digitigrade
Elephantidae	digitigrade
Entelodontidae	unguligrade
Eomyidae	plantigrade

Continued on next page

Table 2 – continued from previous page

Family	Stance
Erethizontidae	plantigrade
Erinaceidae	plantigrade
Esthonychidae	plantigrade
Eutypomyidae	plantigrade
Felidae	digitigrade
Florentiamyidae	plantigrade
Gelocidae	unguligrade
Geolabididae	plantigrade
Glyptodontidae	plantigrade
Gomphotheriidae	unguligrade
Hapalodectidae	plantigrade
Heteromyidae	digitigrade
Hyaenidae	digitigrade
Hyaenodontidae	digitigrade
Hypertragulidae	unguligrade
Ischyromyidae	plantigrade
Jimomyidae	plantigrade
Lagomorpha	digitigrade
Leptictidae	plantigrade
Leptochoeridae	unguligrade
Leptomerycidae	unguligrade
Mammutidae	unguligrade
Megalonychidae	plantigrade
Megatheriidae	plantigrade
Mephitidae	plantigrade

Continued on next page

Table 2 – continued from previous page

Family	Stance
Merycoidodontidae	digitigrade
Mesonychia	unguligrade
Mesonychidae	digitigrade
Micropternodontidae	plantigrade
Mixodectidae	plantigrade
Moschidae	unguligrade
Muridae	plantigrade
Mustelidae	plantigrade
Mylagaulidae	fossorial
Mylodontidae	plantigrade
Nimravidae	digitigrade
Nothrotheriidae	plantigrade
Notoungulata	unguligrade
Oromerycidae	unguligrade
Oxyaenidae	digitigrade
Palaeomerycidae	unguligrade
Palaeoryctidae	plantigrade
Pampatheriidae	plantigrade
Pantolambdidae	plantigrade
Peritychidae	digitigrade
Perissodactyla	unguligrade
Phenacodontidae	unguligrade
Primates	plantigrade
Procyonidae	plantigrade
Proscalopidae	plantigrade

Continued on next page

Table 2 – continued from previous page

Family	Stance
Protoceratidae	unguligrade
Reithroparamyidae	plantigrade
Sciuravidae	plantigrade
Sciuridae	plantigrade
Simimyidae	plantigrade
Soricidae	plantigrade
Suidae	digitigrade
Talpidae	fossorial
Tayassuidae	unguligrade
Tenrecidae	plantigrade
Titanoideidae	plantigrade
Ursidae	plantigrade
Viverravidae	plantigrade
Zapodidae	plantigrade

304

Estimates of species mass used in this study were sourced from multiple databases and papers,
306 especially those focusing on similar macroevolutionary or macroecological questions (Brook and
Bowman, 2004; Freudenthal and Martín-Suárez, 2013; McKenna, 2011; Raia et al., 2012; Smith
308 et al., 2004; Tomiya, 2013); this is similar to Smits (2015). When a species' mass was not available,
proxy measures were used to estimate their mass. For example, given a measurement of a mammal
310 tooth size, it is possible and routine to estimate its mass given some regression equation. The
PBDB has one or more body part measures for many species. These were used as body size proxies
312 for many species, as was the case in Smits (2015). Mass was log-transformed and then rescaled by
first subtracting mean log-mass from all mass estimates, then dividing by two-times its standard
314 deviation; this insures that the magnitude of effects for both continuous and discrete covariates are

Table 1: Species trait assignments in this study are a coarser version of the information available in the PBDB. Information was coarsened to improve per category sample size.

This study		PBDB categories
Diet	Carnivore	Carnivore
	Herbivore	Browser, folivore, granivore, grazer, herbivore.
	Insectivore	Insectivore.
	Omnivore	Frugivore, omnivore.
Locomotor	Arboreal	Arboreal.
	Ground dwelling	Fossorial, ground dwelling, semifossorial, saltatorial.
	Scansorial	Scansorial.

directly comparable (Gelman, 2008; Gelman and Hill, 2007).

- 316 In total, 1400 mammal species occurrence histories were included in this study after applying all of the restrictions above.

Table 3: Regression equations used in this study for estimating body size. Equations are presented with reference to taxonomic grouping, part name, and reference.

Group	Equation	log(Measurement)	Source
General	$\log(m) = 1.827x + 1.81$	lower m1 area	Legendre (1986)
General	$\log(m) = 2.9677x - 5.6712$	mandible length	Foster (2009)
General	$\log(m) = 3.68x - 3.83$	skull length	Luo et al. (2001)
Carnivores	$\log(m) = 2.97x + 1.681$	lower m1 length	Van Valkenburgh (1990)
Insectivores	$\log(m) = 1.628x + 1.726$	lower m1 area	Bloch et al. (1998)
Insectivores	$\log(m) = 1.714x + 0.886$	upper M1 area	Bloch et al. (1998)
Lagomorph	$\log(m) = 2.671x - 2.671$	lower toothrow area	Tomiya (2013)
Lagomorph	$\log(m) = 4.468x - 3.002$	lower m1 length	Tomiya (2013)
Marsupials	$\log(m) = 3.284x + 1.83$	upper M1 length	Gordon (2003)
Marsupials	$\log(m) = 1.733x + 1.571$	upper M1 area	Gordon (2003)
Rodentia	$\log(m) = 1.767x + 2.172$	lower m1 area	Legendre (1986)
Ungulates	$\log(m) = 1.516x + 3.757$	lower m1 area	Mendoza et al. (2006)
Ungulates	$\log(m) = 3.076x + 2.366$	lower m2 length	Mendoza et al. (2006)
Ungulates	$\log(m) = 1.518x + 2.792$	lower m2 area	Mendoza et al. (2006)
Ungulates	$\log(m) = 3.113x - 1.374$	lower toothrow length	Mendoza et al. (2006)

- 318 All fossil occurrences from 64 to 2 million years ago (Mya) were binned into the 19 North American Land Mammal Ages (NALMA) covered by this interval CITATION. The choice of binning by
 320 NALMA reflects the belief that these represent distinct communities or periods of mammal evolution, something that is central to this study. Additionally, because of the inherently discrete
 322 nature of the fossil record it can be hard to re-bin fossils by temporal interval because of the

Table 4: Definitions of the start and stop times of the three plant phases used this study as defined by Graham (2011).

Plant phase	Phase code	Start	Stop
Paleocene-Eocene	Pa-Eo	66	50
Eocene-Miocene	Eo-Mi	50	16
Miocene-Pleistocene	Mi-Pl	16	2

inherent uncertainty in their ages CITATION.

324 Environmental and temporal covariates

The environmental covariates used in this study are collectively referred to as group-level covariates
 326 because they predict the response of a “group” of individual-level observations (i.e. species
 occurrences of an ecotype). Additionally, these covariates are defined for temporal bins and not the
 328 species themselves; as such they predict the individual parts of each species occurrence history. The
 group-level covariates in this study are two global temperature estimates and the Cenozoic “plant
 330 phases” defined by Graham (2011).

Global temperature across most of the Cenozoic was calculated from Mg/Ca isotope record from
 332 deep sea carbonates (Cramer et al., 2011). Mg/Ca based temperature estimates are preferable to
 the frequently used $\delta^{18}\text{O}$ temperature proxy (Alroy et al., 2000; Figueirido et al., 2012; Zachos
 334 et al., 2008, 2001) because Mg/Ca estimates do not conflate temperature with ice sheet volume and
 depth/stratification changes. The former is particularly important to this analysis as the current
 336 polar ice-caps appeared and grew during the second half of the Cenozoic. These properties make
 Mg/Ca based temperature estimates preferable for macroevolutionary and macroecological studies
 338 (Ezard et al., 2016). Temperature was calculated as the mean of all respective estimates for each of
 the NALMA units. The distributions of temperature was then rescaled by subtracting its mean
 340 from all values and then dividing by twice its standard deviation.

The second set of environmental factors included in this study are the Cenozoic plant phases
 342 defined in Graham (2011). Graham’s plant phases are holistic descriptors of the taxonomic
 composition of 12 ecosystem types, which plants are present at a given time, and the relative

		State at $t + 1$		
		0_{never}	1	$0_{extinct}$
State at t	0_{never}	$1 - \pi$	π	0
	1	0	ϕ	$1 - \phi$
	$0_{extinct}$	0	0	1

Table 5: Transition matrix for the birth-death model (Eq. 1). Note that while there are only two state “codes” (0, 1), there are in fact three states: never having originated 0_{never} , present 1, extinct $0_{extinct}$ (Allen, 2011). The two modeled transition probabilities are origination π and survival ϕ .

344 modernity of those plant groups with younger phases representing increasingly modern taxa
 (Graham, 2011). Graham (2011) defines four intervals from the Cretaceous to the Pliocene, though
 346 only three of these intervals take place during the time frame being analyzed. Graham’s plant
 phases was included as a series of “dummy variables” encoding the three phases included in this
 348 analysis (Gelman and Hill, 2007); this means that the Miocene-Pleistocene phase is synonymous
 with the intercept and other phases are defined by their differences from this baseline. The
 350 temporal boundaries of these plant phases, their durations, and abbreviations are defined in Table 4.

Modelling species occurrence

352 At the core of the model used in this study is hidden Markov process where the latent process has
 an absorbing state; also referred to as a discrete-time birth-death model (Allen, 2011) or a
 354 capture-mark-recapture model CITATION. While there are only two state “codes” in a
 presence-absence matrix (i.e. 0/1), there are in fact three states in a birth-death model: not having
 356 originated yet, extant, and extinct. The last of these is the absorbing state, as once a species has
 gone extinct it cannot re-originate (Allen, 2011). Thus, in the transition matrices the probability of
 358 an extinct species changing states is 0 (Table 5). See below for parameter explanations (Tables 6, 7,
 and 8).

360 **Basic model**

I will begin defining the model used in this study by focusing on the basic machinery of the hidden
 362 Markov process at the model’s core. This basic model is similar to the Jolly-Seber

Table 6: Parameters associated with the hidden Markov Model at core of this model (Eq. 1). N is the number of species tracked in this study, and T is the number of time units (NALMAs) covered by this study.

Parameter	dimensions	explanation
y	$N \times T$	observed species presence/absence
z	$N \times T$	“true” species presence/absence
p	$N \times T$	probability of observing a species at time t if it is present
ϕ	$N \times T$	probability of species originating from time t to $t + 1$ if it is not present
π	$N \times (T - 1)$	probability of species surviving at time t , given that it is already originated

capture-mark-recapture model CITATION which has three characteristic probabilities: probability
364 p of observing a species given that it is present, probability π of a species surviving from one time
to another, and probability ϕ of a species first appearing (Royle and Dorazio, 2008) (Table 6). In
366 this formulation, the probability of a species becoming extinct is $1 - \pi$. The inclusion of species and
temporal information means that all three of these probabilities are defined for every species at
368 every time point (Table 6); how this is accomplished is described below. Importantly, only
origination can occur during the first time step as nothing is already present to survive.

$$\begin{aligned}
y_{i,t} &\sim \text{Bernoulli}(p_{i,t} z_{i,t}) \\
z_{i,1} &\sim \text{Bernoulli}(\phi_{i,1}) \\
z_{i,t} &\sim \text{Bernoulli} \left(z_{i,t-1} \pi_{i,t} + \sum_{x=1}^t (1 - z_{i,x}) \phi_{i,t} \right)
\end{aligned} \tag{1}$$

370 The parameters associated with Equation 1 are described in Table 6; this formulation is identical to
that described in Royle and Dorazio (2008). The product term that appears when calculating
372 values of z not at $t = 1$ ensures that once a species goes extinct it does not re-originate. The basic
model described here (Eq. 1) does not include the additional, necessary prior information which is
374 described below.

Expanding on the basics

376 Expanding on the basic model involves modeling the observation, origination and survival
probability as independent multi-level logistic regressions. Origination and survival probabilities

378 share the same covariates and model structure, but observation probability is modeled as a function
of a smaller selection of covariates.

380 The probability of observing a species given that it is present p is modeled as a logistic regression
with a time-varying intercept with an additional zero-centered varying effect for species' functional
382 group, respectively. The effect of species mass was also included through a slope term β^p .

The log-odds of a species originating (logit π) or surviving (logit ϕ) are modeled independently but
384 take the same form: a regression with an intercept that varies by both time and functional group,
an additional taxonomic order varying-intercept term, and the slope term for species mass.

386 Importantly, the time and functional group varying-intercept is itself modeled such that the
intercept for each functional group is a time series with group-level covariates (described below).

388 The expanded model incorporating these regression models is written as

$$\begin{aligned} y_{i,t} &\sim \text{Bernoulli}(p_{i,t} z_{i,t}) \\ p_{i,t} &= \text{logit}^{-1}(u_t + e_{j[i]} + \beta^p m_i) \\ z_{i,1} &\sim \text{Bernoulli}(\phi_{i,1}) \\ z_{i,t} &\sim \text{Bernoulli}\left(z_{i,t-1} \pi_{i,t} + \sum_{x=1}^t (1 - z_{i,x}) \phi_{i,t}\right) \\ \phi_{i,t} &= \text{logit}^{-1}(f_{j[i],t}^\phi + o_{k[i]}^\phi + \beta^\phi m_i). \\ \pi_{i,t} &= \text{logit}^{-1}(f_{j[i],t}^\pi + o_{k[i]}^\pi + \beta^\pi m_i) \end{aligned} \tag{2}$$

How the priors for the varying-effects and coefficients in this expanded model are described below
390 along with the complete model.

Complete model

392 The expanded model (Eq. 2) is still incomplete as it is missing the group-level covariates of interest
such as global temperature, and it is missing all of the necessary priors.

394 Here I describe how the effects of mammal functional group on origination and survival are

Table 7: Parameters for the first expansions

Parameter	dimensions	explanation
u	T	time-varying intercept
e	J	effect of functional group on observation
f^ϕ	$J \times T - 1$	intercept of log-odds ϕ , varies by time and functional group
f^π	$J \times T$	intercept of log-odds π , varies by time and functional group
o^ϕ	K	effect of species' order on log-odds of ϕ
o^π	K	effect of species' order on log-odds of π
β^ϕ	1	effect of species' mass on log-odds of ϕ
β^π	1	effect of species' mass on log-odds of π

modeled. f^ϕ and f^π are modeled as the responses from a multivariate normal distribution, where

396 each functional group's time series is modeled as a regression. The time-series structure of these
 regressions is represented as a random-walk prior for the varying intercept of the group-level
 398 regressions. The effects of these group-level covariates on origination and survival are included for
 each functional group as a vector regression coefficients. The expansion to include this group-level
 400 regression is described in Equation 3, the parameters of which are described in Table 8.

$$\begin{aligned}
 f^\phi &\sim \text{MVN}(\mu^\phi, \Sigma^\phi) \\
 f^\pi &\sim \text{MVN}(\mu^\pi, \Sigma^\pi) \\
 \mu_{j,t}^\phi &= \alpha_{j,t}^\phi + U * \gamma_j^\phi \\
 \mu_{j,t}^\pi &= \alpha_{j,t}^\pi + U * \gamma_j^\pi \\
 \alpha_{j,t}^\phi &\sim \begin{cases} \mathcal{N}(0, 1) & \text{if } t = 1 \\ \mathcal{N}(\alpha_{j,t-1}^\phi, \sigma_j^\phi) & \text{if } t > 1 \end{cases} \\
 \alpha_{j,t}^\pi &\sim \begin{cases} \mathcal{N}(0, 1) & \text{if } t = 1 \\ \mathcal{N}(\alpha_{j,t-1}^\pi, \sigma_j^\pi) & \text{if } t > 1 \end{cases}
 \end{aligned} \tag{3}$$

In hierarchical models like the one described here (Eq. 2, 3) it can be hard to distinguish between

402 the likelihood and prior as data can enter the model through many different parameters CITATION.
 For example, in Equation 2 the model of z can be considered a prior and statements in Equation 3

Table 8: Parameters for the group-level regressions. J is the number of functional groups, and D is the number of group-level covariates.

Parameter	dimensions	explanation
μ^ϕ	$J \times T$	time-series of the mean log-odds of ϕ for each functional group
μ^π	$J \times T$	time-series of the mean log-odds of π for each functional group
Σ^ϕ	$J \times J$	covariance matrix between functional groups for ϕ
Σ^π	$J \times J$	covariance matrix between functional groups for ϕ
α^ϕ	$J \times T$	time-varying intercept of μ^ϕ
α^π	$J \times T$	time-varying intercept of μ^π
σ^ϕ	J	scale of random-walk prior for α^ϕ
σ^π	J	scale of random-walk prior for α^π
γ^ϕ	D	group-level regression coefficients for μ^ϕ
γ^π	D	group-level regression coefficients for μ^π

404 can be considered priors for the parameters which predict ϕ and π . The remaining priors necessary
 to this model, however, are not based on parameter expansion but are prior estimates for the
 406 remaining unmodeled parameters and are sampling statements where no new data enters the model.
 These prior choices are expressed in Equation 4 and are explained below.

408 For the regression coefficients, such as β^ϕ and γ^ϕ , the chosen priors are considered weakly
 informative as they concentrate most of the probability density between -2 and 2. Similarly, the
 410 scale parameters, such as τ^ϕ and σ^ϕ , are also given weakly informative half-Normal priors which
 concentrate most of the probability density between 0 and -2. The covariance matrices, such as Σ^ϕ ,
 412 are decomposed into a vector of scale terms (e.g. τ^ϕ) and correlation matrices (e.g. Ω^ϕ) which were
 then given weakly informative priors. This approach and choice of LKJ priors for the correlation
 414 matrices follows the Stan User Manual CITATION. For parameter vectors which are presented with

only a single prior (e.g. β^ϕ), that prior statement is for each of the elements of that vector.

$$\begin{aligned}
e &\sim \mathcal{N}(0, \sigma^e) \\
\sigma^e &\sim \mathcal{N}^+(1) \\
\beta^p &\sim \mathcal{N}(0, 1) \\
o^\phi &\sim \mathcal{N}(0, v^\phi) \\
o^\pi &\sim \mathcal{N}(0, v^\pi) \\
v^\phi &\sim \mathcal{N}^+(1) \\
v^\pi &\sim \mathcal{N}^+(1) \\
\beta^\phi &\sim \mathcal{N}(0, 1) \\
\beta^\pi &\sim \mathcal{N}(0, 1) \\
\Sigma^\phi &= \text{diag}(\tau^\phi) \Omega^\phi \text{diag}(\tau^\phi) \\
\Sigma^\pi &= \text{diag}(\tau^\pi) \Omega^\pi \text{diag}(\tau^\pi) \\
\tau^\phi &\sim \mathcal{N}^+(1) \\
\tau^\pi &\sim \mathcal{N}^+(1) \\
\Omega^\phi &\sim \text{LKJ}(2) \\
\Omega^\pi &\sim \text{LKJ}(2) \\
\sigma^\phi &\sim \mathcal{N}^+(1) \\
\sigma^\pi &\sim \mathcal{N}^+(1) \\
\gamma^\phi &\sim \mathcal{N}(0, 1) \\
\gamma^\pi &\sim \mathcal{N}(0, 1)
\end{aligned} \tag{4}$$

- ⁴¹⁶ The model used in this study is the complete sampling statement expressed through the combination of equations 2, 3, and 4. These statements taken together form a complete generative
⁴¹⁸ model from which posterior inference of parameter values is possible.

Posterior inference and model adequacy

420 A computer program that implements joint posterior inference the model described above (Eqs. 2,
421 3, and 4) was written in the probabilistic programming language Stan (Stan Development Team,
422 2016). All methods for posterior inference implemented in Stan are derivative-based; this causes
complications for actually implementing the above models, because integers do not have derivatives.
424 In order to infer the values of the matrix of latent discrete parameters z (Tables 6) the log posterior
probabilities of all possible states of the unknown values of z were calculated and summed (i.e.
426 marginalized) (Stan Development Team, 2016).

Species durations at minimum range through from a species first appearance to their last
428 appearance in the fossil record, but the incompleteness of all observations means that the actual
times of origination and extinction are unknown. The marginalization approach used here means
430 that the (log) probabilities of all possible histories for a species are calculated, from the end
members of the species having existed for the entire study interval and the species having only
432 existed between the directly observed first and last appearances to all possible intermediaries (Fig
2) (Stan Development Team, 2016). Marginalization is identical, language-wise, to assuming
434 range-through and then estimating the (log) probability of all possible range extension due to
incomplete sampling.

436 The combined size of the dataset and large number of parameters (Eqs. 2, 3, and 4), in specific the
total number of latent parameters that are the matrix z , means that MCMC based posterior
438 inference is computationally slow. Instead, an approximate Bayesian approach was used: variational
inference. A recently developed automatic variational inference algorithm called “automatic
440 differentiation variational inference” (ADVI) is implemented in Stan and is used here (Kucukelbir
et al., 2015; Stan Development Team, 2016). ADVI assumes that the posterior is Gaussian but still
442 yields a true Bayesian posterior; this assumption is similar to quadratic approximation of the
likelihood function commonly used in maximum likelihood based inference (McElreath, 2016). The
444 principal limitation of assuming the joint posterior is Gaussian is that the true topology of the
log-posterior isn’t estimated; this is a particular burden for scale parameters which are bounded to

	Time Bin							
	1	2	3	4	5	6	7	8
Observed	0	0	0	1	0	1	1	0
-----	-----	-----	-----	-----	-----	-----	-----	-----
Certain	?	?	?	1	1	1	1	?
.....
Potential	0	0	0	1	1	1	1	0
Potential	0	0	1	1	1	1	1	0
Potential	0	1	1	1	1	1	1	0
Potential	1	1	1	1	1	1	1	0
Potential	0	0	0	1	1	1	1	1
Potential	0	0	1	1	1	1	1	1
Potential	0	1	1	1	1	1	1	1
Potential	1	1	1	1	1	1	1	1

Figure 2: Conceptual figure of all possible occurrence histories for an observed species. The first row represents the observed presence/absence pattern for a single species at eight time points. The second row corresponds to the known aspects of the “true” occurrence history of that species. The remaining rows correspond to all possible occurrence histories that are consistent with the observed data. By marginalizing over all possible occurrence histories, the probability of each potential history is estimated. The process of parameter marginalization is described in the text.

446 be positive (e.g. standard deviation).

Of additional concern for posterior inference is the partial identifiability of observation parameters
 448 $p_{t=1}$ and $p_{t=T}$ (Royle and Dorazio, 2008). This issue means that the estimates of sampling
 probabilities at the “edges” of the time series cannot fully be estimated because there are no known
 450 “gaps” in species occurrence histories that are guaranteed to be filled. Instead, the values of the first
 and final columns of the “true” presence-absence matrix z for those observations that do not already
 452 have presences in the observed presence-absence matrix y cannot be estimated (Royle and Dorazio,
 2008). The hierarchical modeling approach used here helps mitigate this problem by pulling the
 454 values of $p_{t=1}$ and $p_{t=T}$ towards the overall mean of p (Gelman et al., 2013), and in fact this
 approach might be more analytically sound than the more ad-hoc approaches that are occasionally
 456 used to overcome this hurdle (Royle and Dorazio, 2008). Additionally, because $p_{t=1}$ and $p_{t=T}$ are
 only partially identifiable, estimates of occurrence θ and origination ϕ at $t = 1$ and estimates of θ , ϕ

458 and survival π at $t = T$ may suffer from similar edge effects. Again, the hierarchical modeling
460 approach used here may help correct for this reality by drawing these estimates towards the overall
460 means of those parameters.

462 After obtaining approximate posterior inference using ADVI, model adequacy and quality of fit
464 were assessed using a posterior predictive check (Gelman et al., 2013). By simulating 100 theoretical
466 data sets from the posterior estimates of the model parameters and the observed covariate
468 information the congruence between predictions made by the model and the observed empirical
470 data can be assessed. These datasets are simulated by starting with the observed states of the
472 presence-absence matrix at $t = 1$; from there, the time series roll forward as stochastic processes
with covariate information given from the empirical observations. Importantly, this is fundamentally
different from observing the posterior estimates of the “true” presence-absence matrix z . The
posterior predictive check used in this study is to compare the observed average number of
observations per species to a distribution of simulated averages; if the empirically observed value
sits in the middle of the distribution then the model can be considered adequate in reproducing the
observed number of occurrences per species.

The ADVI assumption of a purely Gaussian posterior limits the utility and accuracy of the
474 posterior predictive checks because parameter estimates do not reflect the true posterior
distribution and are instead just an approximation (Gelman et al., 2013). Because of this, posterior
476 predictive estimates are themselves only approximate checks of model adequacy. The posterior
predictive check that is used in this study focuses on mean occurrence and not to any scale
478 parameters that might be most affected by the ADVI assumptions.

Given parameter estimates, diversity and diversification rates are estimated through posterior
480 predictive simulations. Given the observed presence-absence matrix y , estimates of the true
presence-absence matrix z can be simulated and the distribution of possible occurrence histories
482 can be analyzed. This is conceptually similar to marginalization where the probability of each
possible occurrence history is estimated (Fig. 2), but now these occurrence histories are generated
484 relative to their estimated probabilities.

The posterior distribution of z gives the estimate of standing diversity N_t^{stand} for all time points as

$$N_t^{stand} = \sum_{i=1}^M z_{i,t}. \quad (5)$$

486 Total regional standing diversity can also be partitioned into the standing diversity of each of the
functional groups.

488 **Results**

The results of the analyses described above take one of two forms: direct inspection of posterior
490 parameter estimates, and downstream estimates of diversity and diversification rates based on
posterior predictive simulations.

492 **Posterior parameter estimates**

The model used here in this study appears to have approximately adequate fit to the data based on
494 the results of the posterior predictive check (Fig. 3). Simulated datasets as estimated from the
models' posterior appears similar in terms of average number of occurrences per species to the
496 observed number of occurrences in the empirical mammal dataset.

Log-odds of observing a species given that it is present varies greatly with time (Fig. 4) with lowest
498 log-odds of observation being during the Gerigian and Harrisonian land-mammal ages. It is
important to note, however, that all land-mammal ages with log-odds of observation greater than 2
500 have very high probabilities of observation, which means that while there may be large differences
in log-odds of observation between land-mammal ages this may not translate to substantial
502 difference in the probability of observation.

There is little variance in the effect of functional group on the log-odds of observing a species that
504 is present (Fig. 5). The only functional group with substantially less than expected log-odds of
observation is scansorial insectivores, indicating that the fossil record of this group is the least




Figure 3: Comparison of the average observed number of occurrences per species (blue line) to the average number of occurrences from 100 posterior predictive datasets using the posterior estimates from the model used in this study. Adequate fit is indicated by the observed value of the test statistic being in the middle of the distribution of those calculated from the simulations.

506 complete of all the functional groups studied. Few functional groups have marginally better than
 expected log-odds of observation, the other insectivorous functional groups have marginally greater
 508 than average log-odds of observation; this is also the case for plantigrade omnivores. These results
 indicate that the observation histories of these functional groups are expected to be complete than
 510 most other functional groups. However, it is important to note that for many functional groups,
 their estimated log-odds of observation are poorly constrained with great uncertainty indicating
 512 little structure in how log-odds of observation varies between functional groups (Fig. 5).

Species mass is found to have a positive effect on probability of observing a species that is present
 514 (Fig. 6). This result indicates that species with greater than average mass are expected to have
 more complete observation histories than species with less than average mass. However, this
 516 estimate does not translate to substantial differences in the estimated probability of observation
 because observation probability is so high for most of the Cenozoic (Fig. 4). In fact, it is only when
 518 land-mammal age observation probability is low that the effect of mass is observable. It is
 important to remember the effect of mass on observation was considered constant over time and
 520 that all differences observation probability between land-mammal ages is driven by variation over
 time. When log-odds of observation is high, differences due to covariate effects translate to very




Figure 4: Ridgeline density plots of the estimates for the log-odds of observation from the time-varying intercept term. Each of the named time units are North American land-mammal ages. The oldest land-mammal age is at the bottom of the stack and the youngest is at the top.

522 small differences in actual probability.

Origination probability varies greatly among functional groups with each functional group
 524 exhibiting a unique time series with a few shared features (Fig. 7). When origination probability is
 below 0.50 this means that few if any new species of that functional group are entering the species
 526 pool, and when origination probability is greater than 0.50 new species of that functional group are
 probably entering the species pool. Finally, if origination probability is approximately 0.50, this
 528 indicates that it is equally likely that a new species is entering the species pool as that it is not.
 The slope of origination probability time-series is also very revealing; when the slope of the time
 530 series is positive then new species are being continually being added to the species pool, and when
 the slope is negative then the number of new species entering the pool is decreasing with time.

532 Most of the functional groups have peak origination probability at the present (Fig. 7); new species
 in these functional groups are continually being added to the species pool. In the case of some
 534 functional groups, such as digitigrade carnivores and fossorial herbivores, this is the culmination of




Figure 5: Ridgeline density plots of the estimates of the effect of functional group on log-odds of observation. Each of the rows correspond to a different functional group as indicated by the dietary and locomotor category combination.

those groups continued growth in the species pool. For other functional groups, such as arboreal

536 herbivores, this peak is a reversal from previously relatively low origination probability; this indicates an expansion of these functional groups following a retraction.

538 Five of the functional groups have peak origination probabilities not at the present: arboreal
carnivores, arboreal insectivores, plantigrade insectivores, scansorial insectivores, and unguiligrade
540 omnivores. The arboreal functional groups reach peak origination probability in the Paleogene,
after which their origination probabilities approach and remain at 0.50, reflecting the loss of these
542 functional groups from the species pool as origination probability never again increases.

Additionally, the uncertainty surrounding in the estimates of origination probability is very large,
544 especially in the Neogene. Large uncertainty in probabilities can reflect complete separation which results from that functional group leaving the species pool and it's (lack of) occurrence is without
546 ambiguity CITATION. The patterns evinced by the other functional groups have similar properties but reach peak origination probability early in the Neogene. Interestingly, origination probability of




Figure 6: Estimates of the effect of species mass on probability of sampling a present species (p). Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units. Probability of observation is presented for each of the NALMAs where the differences in mean probability of observation are due to variation between the time units (Fig. 4).

548 scansorial insectivores has effectively two peaks, once in the late Paleogene and again in the early
549 Neogene. Additionally, as will be discussed later in the context of standing diversity, all five of these
550 functional groups decrease in diversity through the Cenozoic.

551 Origination probability varies greatly amongst mammal orders (Fig. 8). These estimates reflect
552 differences origination probability as well as the relative rarity of that order in the fossil record; if
553 members of that order appear infrequently, they must have lower probability of origination. Orders
554 with greater than average log-odds of origination include Multituberculata, Dinocerata,
555 Didelphimorphia, Creodonta, Condylarthra, Cimolesta, and Acreodi. These orders are major
556 components of the Paleogene fossil record. Orders with lower than average log-odds of origination
557 include Rodentia, Pilosa, Lagomorpha, Eulipotyphyla, Cingulata, Carnivora, and Artiodactyla.
558 These orders are characterized by small body size or primarily Neogene records. Additionally, the
variance between orders is vary large ranging from -3 to 3 log-odds of origination; this large of




Figure 7: Probability of a mammal functional origination at each time point. Each panel depicts 100 time-series sampled from the model’s posterior. The blue line is the mean origination probability as predicted by the group-level predictors. The columns are by locomotor category and rows by dietary category; their intersections are the observed and analyzed ecotypes.

560 variance reflects how species within these orders have very different patterns of origination independent from their origination based on functional ecology (Fig. 7).

562 Species mass is estimated to have a negative relationship with origination probability (Fig. 9). This result means that species with greater than average mass have a lower probability of originating

564 than species with a below average mass. This result is sensible given the left-skewed distribution of

566 mammal species body sizes where large body sizes form the right-hand tail. There are fewer large

568 body-sized mammals which have originated than small body sized mammals. Interestingly, many of

570 the orders with small body sizes (e.g. Rodentia, Lagomorpha) have below average probabilities of

572 originating (Fig. 8); while not completely kosher, when this result is considered together with the effect of mass on origination these effects could be counteracting each other. These results continue

574 to add to the understanding of the heterogeneity and nuance associated with species origination

576 dynamics.




Figure 8: Differences in log-odds of origination based on mammal orders. Positive values correspond to greater log-odds origination than average, while negative values correspond to lower log-odds of origination than average. These estimates reflect the rarity of that order in the fossil record as well as differences in origination.

- 572 The group-level covariates are estimated with high probability (> 0.80) of being different from 0
 (Fig. 10). These results mean that the environmental factors analyzed here are expected to shape
 574 changes in origination probability over time. Importantly, the plant phases and global temperature
 are estimated to affect many of the functional groups
- 576 At least two of the three plant phases are associated with differences in origination probability for
 14 of the 18 functional groups (> 0.85 probability; Table 9). The Paleocene-Eocene phase is found
 578 to be associated with differences in origination probability from the Miocene-Pleistocene for ten of
 the functional groups, all of which are expected to have lower origination probability than the latter
 580 (Table 9). The Eocene-Miocene phase is found to be associated with differences in origination
 probability from the Miocene-Pleistocene for nine of functional groups: eight with a greater
 582 origination probability than the latter, and one with a lower origination probability than the latter
 (Table 9). The Eocene-Miocene phase is expected to be associated with a greater origination
 584 probability than the Paleocene-Eocene for 13 of the functional groups (Table 9).




Figure 9: Mean estimate of the effect of species mass on the probability of a species originating for each of the three plant phases. The effect of mass is considered constant over time and that the only aspect of the model that changes with plant phase is the intercept of the relationship between mass and origination. The three plant phases are indicated by the color of the line. Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units. For clarity, only the mean estimates of the effects of mass and plant phase are plotted.

Temperature is estimated with > 0.85 probability to have an affect origination probability for ten

586 of the 18 functional groups (Table 10). In all cases this relationship is estimated to be negative, meaning that an increase in temperature is associated with a decrease in origination probability.

588 Considering that, on average, temperature decreases through the Cenozoic CITATION, this implies that the origination probability of these ten functional groups may be tracking this long-term trend 590 as opposed to the other functional groups which increases in origination probability independently of temperature.

592 None of the time-series of functional group origination probability are estimated to be either positively or negatively correlated (Fig. 11). This result indicates that functional groups have 594 independent origination histories for the Cenozoic. This result does not preclude the possibility of short term similarities in expansion and decline of origination probability or shared peaks and




Figure 10: Estimated effects of the group-level covariates describing environmental context on log-odds of species origination. These estimates are from the birth-death model. What is plotted is a violin of the distribution of 1000 samples from the approximate posterior.

596 troughs of origination probability. Additionally, if the relationship between two functional groups
 changes over time (e.g. from positive correlation to negative correlation), then it would yield no
 598 overall correlation for the Cenozoic. Finally, it is important to remember that this estimate
 correlation is based on origination probability and not origination rate or diversity.
 600 The survival probability time-series vary greatly between each of the functional groups with each
 exhibiting unique patterns (Fig. 12). Interestingly, unlike origination probability (Fig. 7), survival
 602 probability is frequently estimated with considerable uncertainty. When survival
 probability is below 0.50 then a species that is present is unlikely to survive from one time unit to
 604 the next, while when survival probability is greater than 0.50 species can be expected to survive to
 the next time unit. Finally, when survival probability is approximately 0.50 then survival and
 606 extinction are equally likely. Overall, survival probability is rarely estimated to be greater than 0.50
 with any certainty. This result is consistent with the average occurrence being <1.35 time unit per
 608 species which means that a plurality of species have only a single temporal occurrence (Fig. 3).

Table 9: Posterior probability of the differences in the log-odds of an ecotype originating based on plant phase.

	P(Eo.Mi > 0)	P(Pa.Eo > 0)	P(Eo.Mi > Pa.Eo)
arboreal carnivore	0.575	0.447	0.598
digitigrade carnivore	0.976	0.017	0.998
plantigrade carnivore	0.857	0.780	0.578
scansorial carnivore	0.768	0.154	0.889
arboreal herbivore	0.318	0.357	0.428
digitigrade herbivore	1.000	0.161	0.995
fossorial herbivore	0.999	0.353	0.926
plantigrade herbivore	1.000	0.304	0.998
scansorial herbivore	0.999	0.108	0.998
unguligrade herbivore	0.000	0.000	0.100
arboreal insectivore	0.364	0.003	0.857
fossorial insectivore	0.645	0.341	0.708
plantigrade insectivore	0.794	0.148	0.881
scansorial insectivore	0.916	0.235	0.940
arboreal omnivore	0.590	0.006	0.882
plantigrade omnivore	0.524	0.209	0.762
scansorial omnivore	0.713	0.027	0.938
unguligrade omnivore	0.888	0.127	0.960

The survival probability of many functional groups is frequently approximately 0.50, indicating

610 extinction is frequently random with respect to functional group (Fig. 12). For example, the
 611 survival probability scansorial canirvores is approximately 0.50 for the entire time series which
 612 indicates that there is no best or worst time for this functional groups survival. Similar patterns can
 613 be observed for mean survival probability of arboreal omnivores, fossorial insectivores, and
 614 plantigrade omnivores though all three of these groups have sudden drops in survival probability in
 615 approximately the last 10Mya.

616 Arboreal herbivores are the only functional group for which survival probability is approximately
 617 above 0.50 for the entire Cenozoic (Fig. 12). This result indicates that when an arboreal herbivore
 618 species is present it is expected to survive from one time unit to the next. However, it is important
 619 to note that arboreal herbivores are estimated to have an origination probability below 0.50 for
 620 most of the Cenozoic. Together, these results mean that arboreal herbivore species are rare but are
 621 expected to survive from one time point to the next.

Table 10: Posterior probability that the effects of the two temperature covariates on the log-odds of an ecotype origination are greater than 0. What is estimated is the probability that these estimates are greater than 0; high or low probabilities indicate the “strength” of the covariate in that direction (positive and negative, respectively). These estimates are from the birth-death model.

	$P(\gamma_{temp\ mean} > 0)$
arboreal carnivore	0.355
digitigrade carnivore	0.001
plantigrade carnivore	0.358
scansorial carnivore	0.121
arboreal herbivore	0.219
digitigrade herbivore	0.045
fossilorial herbivore	0.067
plantigrade herbivore	0.000
scansorial herbivore	0.221
unguligrade herbivore	0.339
arboreal insectivore	0.027
fossilorial insectivore	0.219
plantigrade insectivore	0.224
scansorial insectivore	0.192
arboreal omnivore	0.009
plantigrade omnivore	0.087
scansorial omnivore	0.035
unguligrade omnivore	0.129

622 A common feature of multiple functional group’s survival probability time-series is a peak in
 623 survival during the Neogene (Fig. 12). In most cases, these peaks are estimated with little
 624 uncertainty which indicates how apparent this event is. Digitigrade carnivores, digitigrade
 625 herbivores, plantigrade herbivores, scansorial insectivores, unguiligrade herbivores, and unguiligrade
 626 omnivores all peak in survival probability at approximately 25Mya. This peak in survival means
 627 that species of these functional groups which are unlikely to go extinct at this point, potentially
 628 indicating favorable environmental conditions for these groups at the Paleogene-Neogene transition.
 Additionally, this peak does not coincide with the movement from one plant phase to another
 630 (Table 4).

The effect of order on survival probability has much lower variance (Fig. 13) than the effect of order
 632 on origination probability (Fig. 8. Primates, Multituberculata, Eulipotyphla, Dermoptera,
 Creodonta, Condylarthra, Carnivora, and Artiodactyla are estimated to have a lower than average

634 survival probability which implies that species of these orders are expected to be present for a
single time unit. Of these orders, Primates and Multituberculata are expected to have the lowest
636 survival probability of all orders. The orders expected to have greater than average survival
probability are Rodentia, Lagomorpha, and Didelphimorphia.

638 Species mass is estimated to have no relationship or at best a weakly positive relationship with
survival probability (Fig. 14). This result means that differences in mass do not lead to differences
640 in species survival. This result is consistent with previous studies of North American species and
genus survival dynamics CITATION SMITS TOMIYA, and implies that other ecological factors
642 have greater importance on survival than mass alone. GET PROBABILITY ESTIMATE

In contrast to the origination probability, there is little evidence that the group-level covariates
644 have large effects on functional group survival probabilities (Fig. 15). In fact, only the plant phases
are associated with differences in survival probability and only for a relatively small number of
646 functional groups. These results combined with those from the individual-level covariates (Fig. 12,
13, 14) imply that direct interactions (e.g. species–species) are potentially more important to long
648 term species survival than ambient environment (e.g. temperature tolerance). However, because this
estimate of temperature is global in nature, this interpretation is inherently speculative.

650 Functional group survival probability is rarely associated with differences between the three plant
phases (Table 11) with only five pair-wise comparisons having greater than 89% probability of
652 differences in survival between phases. Uniligrade herbivores have an approximately 89%
probability of having lower survival probability during the Paleocene-Eocene than the
654 Miocene-Pleistocene. For digitgrade herbivores, and unguligrade omnivores, the Eocene-Miocene
phase have an approximately 90% probability of having greater survival probability than during the
656 Miocene-Pleistocene phase. In contrast, unguligrade herbivores are estimated to have lower
survival probability in the Eocene-Miocene phase than the Miocene-Pleistocene phase. Finally,
658 unguligrade herbivores have an approximately 99% probability of having a lower survival
probability during the Paleocene-Eocene than the Eocene-Miocene.

660 As stated earlier, temperature is estimated to have no effect on functional group survival

probability (Table 12). This is congruent with previous studies which found no association between
 662 extinction and global temperature CITATION ALROY or no consistent, unidirectional relationship
 between extinction and global temperature CITATION.

Table 11: Posterior probability of the differences in the log-odds of an ecotype surviving based on plant phase.

	P(Eo.Mi > 0)	P(Pa.Eo > 0)	P(Eo.Mi > Pa.Eo)
arboreal carnivore	0.297	0.560	0.328
digitigrade carnivore	0.786	0.367	0.743
plantigrade carnivore	0.411	0.744	0.273
scansorial carnivore	0.428	0.445	0.486
arboreal herbivore	0.256	0.768	0.174
digitigrade herbivore	1.000	0.400	0.942
fossorial herbivore	0.696	0.563	0.565
plantigrade herbivore	0.659	0.508	0.596
scansorial herbivore	0.616	0.539	0.531
unguligrade herbivore	0.000	0.102	0.012
arboreal insectivore	0.289	0.483	0.368
fossorial insectivore	0.532	0.420	0.592
plantigrade insectivore	0.499	0.361	0.605
scansorial insectivore	0.443	0.252	0.634
arboreal omnivore	0.651	0.597	0.591
plantigrade omnivore	0.417	0.549	0.393
scansorial omnivore	0.486	0.525	0.487
unguligrade omnivore	0.929	0.521	0.844

664 None of the time-series of functional group survival probability are estimated to be either positively
 or negatively correlated (Fig. 16); this mirrors the correlations in origination probabiity (Fig. 11).
 666 This result indicates that functional groups probably have independent survival histories for the
 Cenozoic. As with origination probability, this result does not preclude the possibility of short term
 668 similiarities in expansion and decline of orgination probability or shared peaks and troughs of
 survival probability. Additionally, if the relationship between two functional groups changes over
 670 time (e.g. from positive correlation to negative correlation), then it woud yield no overall
 correlation for the Cenozoic. Finally, it is important to remember that this estimate correlation is
 672 based on survival probability and not extinction rate or diversity.

Table 12: Posterior probability that the effects of the two temperature covariates on the log-odds of an ecotype survival are greater than 0. What is estimated is the probability that these estimates are greater than 0; high or low probabilities indicate the “strength” of the covariate in that direction (positive and negative, respectively). These estimates are from the birth-death model.

	$P(\gamma_{temp\ mean} > 0)$
arboreal carnivore	0.665
digitigrade carnivore	0.453
plantigrade carnivore	0.618
scansorial carnivore	0.380
arboreal herbivore	0.761
digitigrade herbivore	0.395
fossorial herbivore	0.429
plantigrade herbivore	0.279
scansorial herbivore	0.345
unguligrade herbivore	0.818
arboreal insectivore	0.489
fossorial insectivore	0.452
plantigrade insectivore	0.435
scansorial insectivore	0.384
arboreal omnivore	0.600
plantigrade omnivore	0.639
scansorial omnivore	0.512
unguligrade omnivore	0.396

Analysis of diversity

674 Standing diversity of the North American mammal species pool estimated from this model exhibits
 an initial increase in diversity followed by a slow decrease till approximately 30Mya, afterwhich
 676 there is a marked increase till approximately 15Mya after which it decreases slightly till it is equal
 to the overall mean diversity of the Cenozoic (Fig. 18). Per-unit standing diveristy is found to be
 678 different from average standing diversity for 12 of 18 time-units (> 85 probability; Table 13).

Diversity is greater than average during the Tiffanian, Wasatchian, Hemingfordian, Barsotvian, and
 680 Clarendonian while diversity is lower than average during the Duchesnean, Chadronian, Orellan,
 Whitneyan, Geringian, Monroecreekian, and Harrisonian. The nadir of diversity is the Orellan while
 682 the apex is the Barstovian (Fig. 18). Interstingly, the rise in diversity among the sampled species
 from the Orellan to the Barstovian is unidirectional and is not estimated to have any temporary
 684 dips in diversification for that entire approximately 15 million year period.

Table 13: Posterior probabilities of diversity N_t^{stand} greater than average standing diversity $\overline{N^{stand}}$ for the whole Cenozoic. The NALMA column corresponds to North American Land Mammal age for that estimate.

NALMA	$P(N_t^{stand} > \overline{N^{stand}})$
Torrejonian	0.79
Tiffanian	0.95
Clarkforkian	0.50
Wasatchian	1.00
Bridgerian	0.69
Uintan	0.75
Duchesnean	0.00
Chadronian	0.01
Orellan	0.00
Whitneyan	0.00
Geringian	0.00
Monroecekian	0.01
Harrisonian	0.11
Hemingfordian	0.96
Barstovian	1.00
Clarendonian	0.93
Hemphillian	0.63
Blancan	0.73

Standing diversity when partitioned by ectype reveals a lot of the complexity behind the pattern

686 of mammal diversity for the Cenozoic (Fig. 19). While each functional group has its own unique
 687 diversity history, there are some broad similarities as is similar to the estimates origination and
 688 survival probability (Fig. 7, 12).

Arboreal ecotypes obtain peak diversity early in the Cenozoic and then decline for the rest of the
 690 time series, becoming increasingly rare or absent as diversity approaches the Recent (Fig. 19).

Arboreal herbivores and omnivores obtain peak diversity at the beginning of the Cenozoic then go
 692 into decline while remaining a small part of the species pool, while arboreal carnivores and
 693 insectivores obtain peak diversity approximately 55 Mya and then quickly decline and become
 694 extremely rare or entirely absent from the species pool. The only arboreal functional group
 695 estimated to not experience a complete disappearance from the species pool are arboreal herbivores.

696 This is consistent with increasing extinction risk in the Neogene compared to the Paleogene as
 697 proposed by Smits (2015).

698 The diversity of plantigrade insectivores, scansorial insectivores, and scansorial omnivores are
estimated to decrease through the Cenozoic (Fig. 19). Plantigrade herbivores and scansorial
700 omnivores have peak diversity at the early Cenozoic and reach low diversity by 30Mya, after which
diversity never increases again. In contrast, scansorial omnivores have nearly constant, above
702 average diversity for the beginning of the Cenozoic till approximately 30Mya, after which diversity
drops and remaining below average diversity for the rest of the Cenozoic.

704 The fossorial functional groups included in this study are estimated to be rare or absent absent for
the first half of the Cenozoic, fossorial herbivores probably having lower diversity than fossorial
706 insectivores (Fig. 19). After fossorial herbivores increase in diversity approximately 30Mya, this
functional group is estimated to quickly reach approximately constant standing diversity for the
708 rest of the Cenozoic. In contrast, fossorial insectivores increase in diversity starting approximately
30Mya and reach max diversity approximately 15Mya, after which this group declines in diversity.

710 Plantigrade carnivores, scansorial herbivores and unguligrade omnivores are estimated to maintain
near constant standing diversity for most of the Cenozoic (Fig. 19). Of these three functional groups,
712 plantigrade carnivores have the greatest variance in standing diversity. Plantigrade carnivores have
greater than average standing diversity from the beginning of the Cenozoic till 50Mya and from
714 approximately 25Mya till approximately 15Mya. This functional group is estimated to be below
average standing diversity from 50Mya to 30Mya, and then from 10Mya till the end of the studied
716 time period. Scansorial herbivores exhibit a similar patterns but with a reversed diversity pattern
for the first 30My of the studied period. Instead of near constant diversity, scansorial herbivores are
718 estimated to have lower than average diversity from the beginning of the Cenozoic till
approximately 50Mya, after which this group has approximately average standing diversity for the
720 rest of the Cenozoic. The unguligrade omnivore functional group has slightly elevated diversity at
the beginning of the Cenozoic and a possible decrease in diversity approximately 15Mya.

722 Scansorial carnivores and plantigrade herbivores have below average standing diversity from the
beginning of the Cenozoic till approximately 20Mya, after which both functional groups increase in
724 diversity till being well above average by the end of the study period (Fig. 19). Plantigrade

omnivores are estimated to be absent or extremely rare in the species pool, only increasing in
726 standing diversity beginning approximately 25Mya. In contrast, scansorial carnivores are estimated
to have been a rare but constant part of the species pool diversity for the entire Cenozoic with an
728 increase in standing diversity starting approximately 25Mya.

Digitigrade carnivores, plantigrade herbivores, and unguligrade herbivores functional groups
730 maintain relatively high standing diversity through out the entire Cenozoic though each exhibits
periods of greater than average and below average standing diversity (Fig. 19). Digitigrade carnivore
732 diversity is estimated to begin the study period below average and then quickly rise to the first peak
in diversity at approximately 55Mya. After this, ditigrade carnivore diversity decreases to below
734 average diversity till approximately 35Mya, after which diversity increases till a second greater peak
in diversity approximately 15Mya. After this second peak in diversity, ditigrade carnivore diversity
736 declines until the end of the study period. Unguligrade herbivores exhibit a similar pattern though
with considerably less uncertainty. In contrast, while plantigrade herbivores have a similar increase
738 and peak in diversity during the first half of the Cenozoic, the functional group does not experience
a second peak in functional diversity till the end of the study period. Additionally, plantigrade
740 herbivores have a longer period of above average standing diversity during the first half of the
Cenozoic, only experiencing a marked decrease in diversity starting approximately 35Mya.

742 The digitigrade herbivore functional group is estimated to be the only group with a near constant
increase in standing diversity through most of the Cenozoic (Fig. 19). There are two periods of
744 decrease in the standing diversity of digitigrade herbivores: the first approximately 10My of the
study period, and a sudden decrease at 15Mya. Beyond these two decreases, this functional group
746 exhibits a remarkable increase in diversity from relative rarity approximately 50-55My till peak
diversity 15-20Mya. Diversity even appears to begin to rebound after the sudden decrease
748 approximately 15Mya.

The waxing and waning of the mammal ecotypes is obvious when comparing changes to estimated
750 relative log-mean of diversity (Fig. 20). While ecotype diversity does appear to change gradually,
there are definite changes to the relative contributions of the ecotypes to the regional species pool.

752 All arboreal ecotypes clearly decrease in relative diversity over the Cenozoic. In contrast the the
digitigrade herbivore, fossorial herbivore, scansorial herbivore, and unguligrade herbivore ecotypes
754 which increase in relative diversity over the Cenozoic. The digitigrade carnivore ecotype increases in
relative diversity until approximately the start of the Neogene, after which it maintains a generally
756 constant relative diversity; this is consistent with previous observations of constant or
density-dependent diversity of the canid guild for the Neogene (Silvestro et al., 2015; Slater, 2015;
758 Van Valkenburgh, 1999), a guild that overlaps with the digitigrade carnivore ecotype. Plantigrade
herbivores remain a constant relative contribution to ecotypic diversity. These results support the
760 hypothesis of a gradual transition from the early Paleogene with a region with more available
habitat for arboreal taxa and less available habitat for many digitigrade and unguligrade taxa, to an
762 environment where arboreal taxa are absent from the species pool and digitigrade and unguligrade
taxa are much more dominant (Fig. 20). It is the relative contributions of digitgrade carnivores,
764 digitigrade herbivores, and unguligrade herbivores which really shape the regional species pool of
the Neogene.

766 Discussion

Both the composition of a species pool and its environmental context change over time, though not
768 necessarily at the same rate or concurrently. Local communities, whose species are drawn from the
regional species pool, have “roles” in their communities defined by their interactions with a host of
770 biotic and abiotic interactors (i.e. a species’ niche). For higher level ecological characterizations like
ecotypes and guilds, these roles are broad and not defined by specific interactions but by the genre
772 of interactions species within that grouping participate in. The diversity of species within an
ecotype or guild can be stable over millions of years despite constant species turnover (Jernvall and
774 Fortelius, 2004; Slater, 2015; Van Valkenburgh, 1999). This implies that the size and scope of the
role of an ecotype or guild in local communities, and the regional species pool as a whole, is
776 preserved even as the individual interactors change. This also implies that the structure of regional
species pools can be constant over time despite a constantly changing set of “players.” There is

778 even evidence that functional groups are at least partially self-organizing and truly emergent
(Scheffer and van Nes, 2006).

780 Comparison of the results from the posterior predictive checks for the pure-presence and birth-death
models supports the conclusion that regional species pool dynamics cannot simply be described by
782 a single occurrence probability and are instead the result of the interplay between the origination
and extinction processes. Additionally, changes to the ecotypic composition and diversification rate
784 of the North American regional species pool are driven primarily by variation in origination and not
extinction (Fig. ??). These aspects of how regional species pool diversity is shaped are not directly
786 observable in studies of the Recent where time scales are short and macroevolutionary dynamics are
inferable solely from phylogeny (Fritz et al., 2013; Price and Schmitz, 2016).

788 Extinction rate for the entire regional species pool through time is highly variable and demonstrates
a saw-toothed pattern with no obvious temporal trends. While a constant mean extinction rate is
790 consistent with previous observation (Alroy, 1996; Alroy et al., 2000), the degree to which mammal
extinction rate is actually variable may not have been equally appreciated as it has been for the
792 marine invertebrate record (Foote, 2000a,b, 2006, 2010). What is most consistent with previous
observations, however, is that diversity seems to be most structured by changes to origination
794 rather than changes to extinction (Alroy, 1996; Alroy et al., 2000).

Comparison of the ecotype specific diversity histories adds a considerable degree of nuance to broad
796 narrative of shifts in functional composition of the North American mammal species pool as being
gradual (Fig. 19). While most ecotypes do not experience sudden shifts in origination or extinction
798 rate (Fig. ??, ??). As with the diversification rate of the entire species pool, the diversification of
individual ecotypes seem principally driven by origination and not extinction. Instead, while species
800 seem to originate in waves (Fig. ??), they appear to leave the regional species pool in an
uncoordinated and individual manner (Fig. ??) which could be considered consistent with the
802 maxim that all species respond differently to environmental change (Blois and Hadly, 2009). Note,
however, this result characterizes the entire North American mammal regional species pool and
804 thus may not reflect the dynamics of individual local communities.

The few large-magnitude, but temporary, increases in ecotype-specific origination rate occur in
806 digitigrade carnivores, digitigrade herbivores, plantigrade herbivores, and unguligrade herbivores.
Importantly, the large peak in diversification and origination rates 16 Mya (Fig. ??) appears driven
808 almost entirely by a massive increase in the origination rate of unguligrade herbivores (Fig. ??).
Additionally, there is some evidence that the origination probabilities of these ecotypes are
810 correlated (Fig. 11, ??). While this result does not mean that there are large and sudden
cross-ecotype changes to the regional species pool, it does suggest that additions to the species pool
812 do not occur in individual ecotypes idiosyncratically.

Arboreal taxa disappear from the regional species pool over the Cenozoic, with long term decline
814 over the Paleogene leading to the disappearance by start of Neogene ~22 Mya. This is partially
consistent with one of the two possible patterns presented here and in Smits (2015) that would
816 result in arboreal taxa having a greater extinction risk than other ecotypes: the Paleogene and
Neogene were different selective regimes and, while the earliest Cenozoic may have been neutral
818 with respect to arboreal taxa, they disappeared quickly over the Cenozoic which may account for
their higher extinction risk. However, these results add some nuance to this scenario as arboreal taxa
820 were declining throughout the Paleogene instead of maintaining a flat diversity as hypothesized
(Smits, 2015). I interpret the decline of arboreal taxa throughout the Paleogene to mean that the
822 shift from closed to open environments began in the Paleogene and led to increasingly hostile
environments for arboreal taxa as opposed to being a sudden change in selective regime between
824 the Paleogene and Neogene. In addition to all arboreal taxa, the diversity of plantigrade and
scansorial insectivores decreases with time (Fig. 19).

826 Digitigrade carnivores have a relatively stable diversity history through the Cenozoic and can be
characterized as varying around a constant mean diversity. This ecotype has a large amount of
828 overlap with the carnivore guild which has been the focus of much research (Janis and Wilhelm,
1993; Pires et al., 2015; Slater, 2015; Van Valkenburgh, 1999). This result is consistent with some
830 form of “control” on the diversity of this ecotype, such as diversity-dependent diversification
(Silvestro et al., 2015; Slater, 2015; Van Valkenburgh, 1999).

832 Both digitigrade and unguligrade herbivores increase in diversity over the Cenozoic. The increase of
these cursorial forms is consistent with the gradual opening up of the North American landscape
834 (Blois and Hadly, 2009; Graham, 2011; Strömberg, 2005) and may indicate a long-term shift in the
interactors associated with those ecotypes leading to increased contribution to the regional species
836 pool. This result may be comparable to the increasing percentage of hypsodont (high-crowned
teeth) mammals in the Neogene of Europe being due to an enrichment of hypsodont taxa and not
838 a depletion of non-hypsodont taxa. Smaller scale increases in fossorial herbivore species, and a lesser
extent plantigrade herbivores, suggests that the increase of interactors may be associated mostly
840 with the herbivore dietary category with locomotor category tempering that relationship. These
results support the conclusion that the increase in digitigrade and unguligrade herbivores is the
842 result of an enrichment of these ecotypes as opposed to being caused by the depletion of other
herbivorous ecotypes; this is further supported by the lack of major changes to the number of
844 extinctions of all herbivore ecotypes (Fig. ??).

An association between plant phase and differences in ecotype occurrence or origination-extinction
846 probabilities is interpreted to mean that an ecotype enrichment or depletion is due to associations
between that ecotype and whatever plants are dominant at that time. Plant phase clearly
848 structures the occurrence and origination probability time series (Fig. ??, 7). These differences in
occurrence or origination translate to the estimates of diversity and diversification rate; the largest
850 spike in diversity, diversification rate, and origination rate all correspond to the onset of the last
plant phase (Fig. ??). The clearest example of the diversity of an ecotype increasing at this
852 particular transition is in scansorial carnivores (Fig. 19); similar shifts in other ecotypes are much
more subtle, as was previously noted for fossorial insectivores.

854 Interestingly, for all of the ecotypes with sudden changes in diversity at this transition the change is
an increase, even if only temporarily. There are two interpretations of these results. A biological
856 interpretation of this result is that, because plant phase associations are only with occurrence or
origination probabilities and not survival, these ecotypes were well suited for the newly available
858 mammal-plant interactions due to the increased modernization of their floral context (Graham,
2011). Alternatively, the increase in diversity associated with the third plant phase may be caused

860 by the edge effect in origination probability that is artificially inflating the number of origination
events (Fig. 7). However, the estimated number of origination events does not have a tremendous
862 spike at this transition, nor is a major increase in the number of origination events sustained (Fig.
??).

864 There are fewer, less obvious shifts in diversity surrounding the transition from the first to second
plant phase, with the following ecotypes having apparent shifts in diversity at 50 My: plantigrade
866 carnivores (down), arboreal omnivores (down), and scansorial omnivores (down). Arboreal
insectivore peak diversity also occurs 50 Mya, and is then followed by a steep decline in diversity
868 till 30 Mya when this ecotype is lost from the species pool. Because plant phase has been found to
structure occurrence/origination (Fig. ??, 7), but not survival (Fig. 12), my interpretation of these
870 results is that new species were not entering the system because there were fewer available
mammal-plant interactions available for those ecotypes. Instead, these ecotypes were poorly suited
872 for the newly available mammal-plant interactions brought upon by the changing environmental
context (Graham, 2011).

874 The temperature covariates are found to have similar effects on occurrence and origination
probabilities (Tables ??, 10). Temperature is found to more often affect ecotype occurrence
876 probabilities than origination probabilities. In most cases, there is a negative association between
temperature and probability of occurring or first originating; this means that if temperature
878 decreases, we would then expect an increase in the probability of occurring or first originating. In
contrast, temperature range is estimated to be a good predictor of survival in only to mammal
880 ecotypes and only marginally for one of those (Table 12). Additionally, both of these cases have
positive relationships, meaning that if temperature decreases it is expected that species survival will
882 also decrease.

The result that temperature does not affect the survival probability of most ecotypes is consistent
884 with previous analysis of mammal diversity (Alroy et al., 2000). The result that temperature affects
origination probability, on the other hand, is in strong contrast to the results Alroy et al. (2000).
886 An important difference between the analyses presented here and that of Alroy et al. (2000) is I am

considering the effect of temperature on the probability of a species originating, assuming it hasn't
888 originated yet while Alroy et al. (2000) analyzes the correlation between the first differences of the
origination and extinction rates with an oxygen isotope curve (Zachos et al., 2001). Origination or
890 extinction rates have very different properties than the origination probabilities estimated here
brought upon by the difference both in definition and units. Origination probability is the expected
892 probability that a species that has never been present and is not present at time t will be present at
time $t + 1$; origination probability is defined for a single species. In contrast, per capita rates are
894 defined (for origination) as the expected number of new species to have originated between time t
and $t + 1$ given the total number of species present at time t ; per capita rates are defined for the
896 standing diversity. It is also important to note that even though there is an edge effect at the last
time interval that causes an increase in the occurrence and origination probabilities of some
898 ecotypes (Fig. ??, 7, the corresponding rates and population level birth/death dynamics do not
share that pattern (Fig. ??, ??, ??). However, it is still possible that the finding that temperature
900 has an effect on origination may simply be because as time approaches the present the number of
species which have originated increases and not because of climatic forcing of origination.

902 All environmental factors are found to affect the occurrence and origination probabilities for most,
but not all, mammal ecotypes (Fig. ??, 10). In contrast, the environmental factors probably do not
904 affect differences in ecotype survival probability (Fig. 15). The focus in previous research on
temperature and major climatic or geological events without other measures of environmental
906 context may have led to confusion in discussions of how the “environment” affects mammal
diversity and diversification (Alroy et al., 2000; Figueirido et al., 2012). The environment or climate
908 are more than just global or regional temperature, it is also the set of all possible biotic and abiotic
interactions that can be experienced by a member of the species pool. By including more
910 descriptors of species’ environmental context than simple an estimate of global temperature a more
complete “picture” of the diversification process is inferred.

912 Analysis of relationship between temperature and origination rate is probably better suited for a
continuous-time birth-death or multilevel stochastic differential equation model instead of a
914 discrete-time model because the both continuous models estimate rates while discrete time models

estimate probabilities (Allen, 2011). The **PyRate** model(s) are based on a continuous-time
916 birth-death process (Silvestro et al., 2015, 2014). Unfortunately, a continuous-time model may be
unsuited for most paleontological data as the fossil record is naturally discrete; fossils are assigned
918 to temporal units, such as stages, which have age ranges. Individual fossils are not assigned
individual numeric ages. This reality was in fact my one of motivations for using discrete-time
920 birth-death model instead of one in continuous-time. There are of course exceptions to this
characterization; the fossil record of graptolites from the Ordovician and Silurian (Crampton et al.,
922 2016) and the fossil record of some mammal orders from Neogene are of high enough resolution that
the application of continuous-time models is appropriate and less fraught.

924 The effect of species mass on either occurrence or origination and extinction was not allowed to
vary by ecotype or environmental context. The primary reason for this modeling choice was that
926 this study focuses on ecotypic based differences in either occurrence, or origination and extinction.
Allowing the effect of body size to vary by ecotype, time, and environmental factors would increase
928 the overall complexity of the model beyond the scope of the study. Instead, body size was included
in order to control for its possible underlying effects (McElreath, 2016). A control means that if
930 there is variation due to body mass, having a term to “absorb” that effect is better than ignoring it.

The only covariate allowed to affect sampling probability was mass and only as a linear predictor.

932 Other covariates, such as the environmental factors considered here, could have affected the
underlying preservation process that limits sampling probability; their exclusion as covariates of
934 sampling/observation was the product of a few key decisions: model complexity, model
interpretability, the scope of this study, and a lack of good hypotheses related to these covariates to
936 warrant their inclusion.

Conclusions

938 These results add a considerable degree of nuance to the narrative of changes to North American
diversity being gradual. My results support the conclusions that ecotypic diversity is shaped more
940 by changes to origination than extinction and that major changes to total diversification rate can

be attributed to increases in origination of only some ecotypes. There are a number of interesting
942 estimated ecotype diversity patterns. While arboreal ecotypes are diverse in the Paleogene, by the
Neogene all arboreal ecotypes dramatically decreased in diversity and became either rare or absent
944 from the regional species pool. The other ecotypes that decrease in diversity over the Cenozoic are
plantigrade and scansorial insectivores and scansorial omnivores. The only ecotypes that
946 demonstrate a sustained pattern of increasing diversity are digitigrade and unguligrade herbivores.
When the environmental covariates analyzed here are inferred to affect the diversification of an
948 ecotype, this effect is virtually always on origination and not survival. This analysis provides a
much more complete picture of North American mammal diversity and diversification, specifically
950 the dynamics of the ecotypic composition of that diversity. By increasing the complexity of analysis
while precisely translating research questions into a statistical model, the context of the results is
952 much better understood. Future studies of diversity and diversification should incorporate as much
information as possible into their analyses in order to better understand or at least contextualize
954 the complex processes underlying that diversity.

Acknowledgements

956 I would like to thank K. Angielczyk, M. Foote, P. D. Polly, R. Ree, and G. Slater for helpful
discussion and advice. This entire study would not have been possible without the Herculean
958 effort of the many contributors to the Paleobiology Database. In particular, I would like to thank J.
Alroy and M. Uhen for curating most of the mammal occurrences recorded in the PBDB. This is
960 Paleobiology Database publication XXX.

References

- 962 Allen, L. J. S. 2011. An introduction to stochastic processes with applications to biology. 2nd ed.
Chapman and Hall/CRC, Boca Raton, FL.

- 964 Alroy, J. 1996. Constant extinction, constrained diversification, and uncoordinated stasis in North
American mammals. *Palaeogeography, Palaeoclimatology, Palaeoecology* 127:285–311.
- 966 ———. 2009. Speciation and extinction in the fossil record of North American mammals. Pages
302–323 *in* R. K. Butlin, J. R. Bridle, and D. Schluter, eds. *Speciation and patterns of diversity*.
968 Cambridge University Press, Cambridge.
- 970 ———. 2010. Fair sampling of taxonomic richness and unbiased estimation of origination and
extinction rates. Pages 55–80 *in* J. Alroy and G. Hunt, eds. *Quantitative Methods in
Paleobiology*. The Paleontological Society.
- 972 Alroy, J., P. L. Koch, and J. C. Zachos. 2000. Global climate change and North American
mammalian evolution. *Paleobiology* 26:259–288.
- 974 Badgley, C., and J. A. Finarelli. 2013. Diversity dynamics of mammals in relation to tectonic and
climatic history: comparison of three Neogene records from North America. *Paleobiology*
976 39:373–399.
- 978 Badgley, C., T. M. Smiley, R. Terry, E. B. Davis, L. R. G. Desantis, D. L. Fox, S. S. B. Hopkins,
T. Jezkova, M. D. Matocq, N. Matzke, J. L. McGuire, A. Mulch, B. R. Riddle, V. L. Roth, J. X.
Samuels, C. A. E. Strömberg, and B. J. Yanites. 2017. Biodiversity and Topographic Complexity:
980 Modern and Geohistorical Perspectives. *Trends in Ecology & Evolution* pages 1–16.
- 982 Bambach, R. K. 1977. Species richness in marine benthic habitats through the Phanerozoic.
Paleobiology 3:152–167.
- 984 Bambach, R. K., A. M. Bush, and D. H. Erwin. 2007. Autecology and the filling of ecospace: Key
metazoan radiations. *Palaeontology* 50:1–22.
- 986 Bloch, J. I., K. D. Rose, and P. D. Gingerich. 1998. New species of Batodonoides (Lipotyphla,
Geolabididae) from the Early Eocene of Wyoming: smallest known mammal? *Journal of
Mammalogy* 79:804–827.

- 988 Blois, J. L., and E. A. Hadly. 2009. Mammalian Response to Cenozoic Climatic Change. *Annual Review of Earth and Planetary Sciences* 37:181–208.
- 990 Brook, B. W., and D. M. J. S. Bowman. 2004. The uncertain blitzkrieg of Pleistocene megafauna. *Journal of Biogeography* 31:517–523.
- 992 Brown, A. M., D. I. Warton, N. R. Andrew, M. Binns, G. Cassis, and H. Gibb. 2014. The fourth-corner solution - using predictive models to understand how species traits interact with
994 the environment. *Methods in Ecology and Evolution* 5:344–352.
- Brown, J. H., and B. A. Maurer. 1989. Macroecology: the division of food and space among species
996 on continents. *Science* 243:1145–1150.
- Brown, J. J. 1995. *Macroecology*. University of Chicago Press, Chicago.
- 998 Bush, A. M., and R. K. Bambach. 2011. Paleoecologic Megatrends in Marine Metazoa. *Annual Review of Earth and Planetary Sciences* 39:241–269.
- 1000 Bush, A. M., R. K. Bambach, and G. M. Daley. 2007. Changes in theoretical ecospace utilization in marine fossil assemblages between the mid-Paleozoic and late Cenozoic. *Paleobiology* 33:76–97.
- 1002 Bush, A. M., and P. M. Novack-Gottshall. 2012. Modelling the ecological-functional diversification of marine Metazoa on geological time scales. *Biology Letters* 8:151–155.
- 1004 Cantalapiedra, J. L., J. L. Prado, and M. T. Alberdi. 2017. Decoupled ecomorphological evolution and diversification in Neogene-Quaternary horses. *Science* 355:627–630.
- 1006 Carrano, M. T. 1999. What, if anything, is a cursor? Categories versus continua for determining locomotor habit in mammals and dinosaurs. *Journal of Zoology* 247:29–42.
- 1008 Clyde, W. C., and P. D. Gingerich. 1998. Mammalian community response to the latest Paleocene thermal maximum: an isotaphonomic study in the northern Bighorn Basin, Wyoming. *Geology*
1010 26:1011–1014.
- Cohen, K. M., S. C. Finney, P. L. Gibbard, and J.-X. Fan. 2015. The ICS International
1012 Chronostratigraphic Chart.

- Cottenie, K. 2005. Integrating environmental and spatial processes in ecological community
1014 dynamics. *Ecology Letters* 8:1175–1182.
- Cramer, B. S., K. Miller, P. Barrett, and J. Wright. 2011. Late Cretaceous-Neogene trends in deep
1016 ocean temperature and continental ice volume: Reconciling records of benthic foraminiferal
geochemistry ($\delta^{18}\text{O}$ and Mg/Ca) with sea level history. *Journal of Geophysical Research: Oceans*
1018 116:1–23.
- Crampton, J. S., R. A. Cooper, P. M. Sadler, and M. Foote. 2016. Greenhouse-icehouse transition
1020 in the Late Ordovician marks a step change in extinction regime in the marine plankton.
Proceedings of the National Academy of Sciences 113:1498–1503.
- Damuth, J., and C. M. Janis. 2011. On the relationship between hypsodonty and feeding ecology in
ungulate mammals, and its utility in palaeoecology. *Biological Reviews* 86:733–758.
- Elith, J., and J. R. Leathwick. 2009. Species distribution models: ecological explanation and
prediction across space and time. *Annual Review of Ecology, Evolution, and Systematics*
1024 40:677–697.
- Eronen, J. T., C. M. Janis, C. P. Chamberlain, and A. Mulch. 2015. Mountain uplift explains
1028 differences in Palaeogene patterns of mammalian evolution and extinction between North
America and Europe. *Proceedings of the Royal Society B: Biological Sciences* 282:20150136.
- Eronen, J. T., P. D. Polly, M. FRED, J. Damuth, D. C. FRANK, V. Mosbrugger,
C. SCHEIDEGGER, N. C. Stenseth, and M. Fortelius. 2010. Ecometrics: The traits that bind
1032 the past and present together. *Integrative Zoology* 5:88–101.
- Ezard, T. H. G., A. Purvis, and H. Morlon. 2016. Environmental changes define ecological limits to
1034 species richness and reveal the mode of macroevolutionary competition. *Ecology Letters*
19:899–906.
- Felsenstein, J. 1985. Phylogenies and the comparative method. *The American Naturalist* 125:1–15.
- Figueirido, B., C. M. Janis, J. A. Pérez-Claros, M. De Renzi, and P. Palmqvist. 2012. Cenozoic

- 1038 climate change influences mammalian evolutionary dynamics. *Proceedings of the National
Academy of Sciences* 109:722–727.
- 1040 Foote, M. 2000a. Origination and extinction components of taxonomic diversity: general problems.
Paleobiology 26:74–102.
- 1042 ———. 2000b. Origination and extinction components of taxonomic diversity: Paleozoic and
post-Paleozoic dynamics. *Paleobiology* 26:578–605.
- 1044 ———. 2001. Inferring temporal patterns of preservation, origination, and extinction from
taxonomic survivorship analysis. *Paleobiology* 27:602–630.
- 1046 ———. 2006. Substrate affinity and diversity dynamics of Paleozoic marine animals. *Paleobiology*
32:345–366.
- 1048 ———. 2010. The geologic history of biodiversity. Pages 479–510 in M. A. Bell, D. J. Futuyma,
W. F. Eanes, and J. S. Levinton, eds. *Evolution since Darwin: the first 150 years*. Sinauer
1050 Associates, Sunderland, MA.
- Foote, M., and J. J. Sepkoski. 1999. Absolute measures of the completeness of the fossil record.
1052 *Nature* 398:415–7.
- Foster, J. R. 2009. Preliminary body mass estimates for mammalian genera of the Morrison
1054 Formation (Upper Jurassic, North America). *PaleoBios* 28:114–122.
- Fraser, D., R. Gorelick, and N. Rybczynski. 2015. Macroevolution and climate change influence
1056 phylogenetic community assembly of North American hoofed mammals. *Biological Journal of the
Linnean Society* 114:485–494.
- Freudenthal, M., and E. Martín-Suárez. 2013. Estimating body mass of fossil rodents. *Scripta
Geologica* 145:1–130.
- Fritz, S. A., J. Schnitzler, J. T. Eronen, C. Hof, K. Böhning-Gaese, and C. H. Graham. 2013.
1060 Diversity in time and space: wanted dead and alive. *Trends in Ecology & Evolution* 28:509–16.

- 1062 Gelman, A. 2008. Scaling regression inputs by dividing by two standard deviations. *Statistics in Medicine* pages 2865–2873.
- 1064 Gelman, A., J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari, and D. B. Rubin. 2013. Bayesian data analysis. 3rd ed. Chapman and Hall, Boca Raton, FL.
- 1066 Gelman, A., and J. Hill. 2007. Data Analysis using Regression and Multilevel/Hierarchical Models. Cambridge University Press, New York, NY.
- 1068 Gordon, C. L. 2003. A First Look at Estimating Body Size in Dentally Conservative Marsupials. *Journal of Mammalian Evolution* page 21.
- 1070 Graham, A. 2011. A natural history of the New World: the ecology and evolution of plants in the Americas. University of Chicago Press, Chicago.
- 1072 Harmon, L. J., and S. Harrison. 2015. Species Diversity Is Dynamic and Unbounded at Local and Continental Scales. *The American Naturalist* 185:000–000.
- 1074 Harrison, S., and H. Cornell. 2008. Toward a better understanding of the regional causes of local community richness. *Ecology Letters* 11:969–979.
- 1076 Huang, S., J. T. Eronen, C. M. Janis, J. J. Saarinen, D. Silvestro, and S. A. Fritz. 2017. Mammal body size evolution in North America and Europe over 20 Myr: similar trends generated by different processes. *Proceedings of the Royal Society B: Biological Sciences* 284:20162361.
- 1080 Jamil, T., W. A. Ozinga, M. Kleyer, and C. J. F. Ter Braak. 2013. Selecting traits that explain species-environment relationships: A generalized linear mixed model approach. *Journal of Vegetation Science* 24:988–1000.
- 1082 Janis, C., J. Damuth, and J. M. Theodor. 2004. The species richness of Miocene browsers, and implications for habitat type and primary productivity in the North American grassland biome. *Palaeogeography, Palaeoclimatology, Palaeoecology* 207:371–398.
- 1086 Janis, C. M. 1993. Tertiary mammal evolution in the context of changing climates, vegetation, and tectonic events. *Annual Review of Ecology and Systematics* 24:467–500.

- . 2008. An evolutionary history of browsing and grazing ungulates. Pages 21–45 in I. J. Gordon and H. H. T. Prins, eds. *The Ecology of Browsing and Grazing*. Springer-Verlag.
- Janis, C. M., J. Damuth, and J. M. Theodor. 2000. Miocene ungulates and terrestrial primary productivity: where have all the browsers gone? *Proceedings of the National Academy of Sciences* 97:7899–904.
- Janis, C. M., and P. B. Wilhelm. 1993. Were there mammalian pursuit predators in the tertiary? Dances with wolf avatars. *Journal of Mammalian Evolution* 1:103–125.
- Jardine, P. E., C. M. Janis, S. Sahney, and M. J. Benton. 2012. Grit not grass: concordant patterns of early origin of hypodonty in Great Plains ungulates and Glires. *Palaeogeography, Palaeoclimatology, Palaeoecology* 365–366:1–10.
- Jernvall, J., and M. Fortelius. 2002. Common mammals drive the evolutionary increase of hypodonty in the Neogene. *Nature* 417:538–40.
- . 2004. Maintenance of trophic structure in fossil mammal communities: site occupancy and taxon resilience. *The American Naturalist* 164:614–624.
- Kucukelbir, A., R. Ranganath, A. Gelman, and D. M. Blei. 2015. Automatic Variational Inference in Stan. Pages 568–576 in NIPS. Vol. 28.
- Legendre, S. 1986. Analysis of mammalian communities from the Late Eocene and Oligocene of Southern France. *Paleovertebrata* 16:191–212.
- Liow, L. H., M. Fortelius, E. Bingham, K. Lintulaakso, H. Mannila, L. Flynn, and N. C. Stenseth. 2008. Higher origination and extinction rates in larger mammals. *Proceedings of the National Academy of Sciences* 105:6097–6102.
- Liow, L. H., M. Fortelius, K. Lintulaakso, H. Mannila, and N. C. Stenseth. 2009. Lower Extinction Risk in SleeporHide Mammals. *The American Naturalist* 173:264–272.
- Lloyd, G. T., J. R. Young, and A. B. Smith. 2011. Taxonomic Structure of the Fossil Record is Shaped by Sampling Bias. *Systematic Biology* 61:80–89.

- 1112 Loeuille, N., and M. a. Leibold. 2008. Evolution in metacommunities: on the relative importance of
species sorting and monopolization in structuring communities. *The American naturalist*
1114 171:788–99.
- 1116 Losos, J. B. 2010. Adaptive radiation, ecological opportunity, and evolutionary determinism. *The
American naturalist* 175:623–39.
- 1118 Losos, J. B., and D. L. Mahler. 2010. Adaptive radiation: the interaction of ecological opportunity,
adaptation, and speciation. Chap. 15, pages 381–420 *in* M. A. Bell, D. J. Futuyma, W. F. Eanes,
and J. S. Levinton, eds. *Evolution since Darwin: the first 150 years*. Sinauer Associates,
1120 Sunderland, MA.
- 1122 Luo, Z.-X., A. W. Crompton, and A.-L. Sun. 2001. A New Mammaliaform from the Early Jurassic
and Evolution of Mammalian Characteristics. *Science* 292:1535–1540.
- 1124 McElreath, R. 2016. *Statistical rethinking: a Bayesian course with examples in R and Stan*. CRC
Press, Boca Raton, FL.
- 1126 McGill, B. J., B. J. Enquist, E. Weiher, and M. Westoby. 2006. Rebuilding community ecology
from functional traits. *TRENDS in Ecology and Evolution* 21:178–185.
- 1128 McKenna, R. T. 2011. Potential for Speciation in Mammals Following Vast , Late Miocene Volcanic
Interruptions in the Pacific Northwest. Masters. Portland State University.
- 1130 Mendoza, M., C. M. Janis, and P. Palmqvist. 2006. Estimating the body mass of extinct ungulates:
a study on the use of multiple regression. *Journal of Zoology* 270:90–101.
- 1132 Mittelbach, G. G., and D. W. Schemske. 2015. Ecological and evolutionary perspectives on
community assembly. *Trends in Ecology and Evolution* 30:241–247.
- 1134 Novack-Gottshall, P. M. 2007. Using a theoretical ecospace to quantify the ecological diversity of
Paleozoic and modern marine biotas Using a theoretical ecospace to quantify the ecological
diversity of Paleozoic and modern marine biotas. *Paleobiology* 33:273–294.
- 1136 Pires, M. M., D. Silvestro, and T. B. Quental. 2015. Continental faunal exchange and the

- asymmetrical radiation of carnivores. *Proceedings of the Royal Society B: Biological Sciences*
1138 282:20151952.
- Pollock, L. J., W. K. Morris, and P. A. Vesk. 2012. The role of functional traits in species
1140 distributions revealed through a hierarchical model. *Ecography* 35:716–725.
- Polly, P., J. Eronen, M. Fred, G. P. Dietl, V. Mosbrugger, C. Scheidegger, D. C. Frank, J. Damuth,
1142 N. C. Stenseth, and M. Fortelius. 2011. History matters: ecometrics and integrative climate
change biology. *Proceedings of the Royal Society B: Biological Sciences* 278:1131–1140.
- 1144 Polly, P. D., A. M. Lawing, J. T. Eronen, and J. Schnitzler. 2015. Processes of ecometric patterning:
modelling functional traits, environments, and clade dynamics in deep time. *Biological Journal of
1146 the Linnean Society* pages n/a–n/a.
- Price, S. A., and L. Schmitz. 2016. A promising future for integrative biodiversity research: an
1148 increased role of scale-dependency and functional biology. *Philosophical Transactions of the
Royal Society B: Biological Sciences* 371:20150228.
- 1150 Quental, T. B., and C. R. Marshall. 2013. How the Red Queen Drives Terrestrial Mammals to
Extinction. *Science* 341:290–292.
- 1152 Rabosky, D. L. 2013. Diversity-Dependence, Ecological Speciation, and the Role of Competition in
Macroevolution. *Annual Review of Ecology, Evolution, and Systematics* 44:1–22.
- 1154 Rabosky, D. L., and A. H. Hurlbert. 2015. Species Richness at Continental Scales Is Dominated by
Ecological Limits. *The American Naturalist* 185:000–000.
- 1156 Raia, P., F. Carotenuto, F. Passaro, D. Fulgione, and M. Fortelius. 2012. Ecological specialization
in fossil mammals explains Cope's rule. *The American Naturalist* 179:328–37.
- 1158 Royle, J. A., and R. M. Dorazio. 2008. Hierarchical modeling and inference in ecology: the analysis
of data from populations, metapopulations and communities. Elsevier, London.
- 1160 Royle, J. A., J. D. Nichols, and M. Kéry. 2005. Modelling occurrence and abundance of species
when detection is imperfect. *Oikos* 110:353–359.

- 1162 Scheffer, M., and E. H. van Nes. 2006. Self-organized similarity, the evolutionary emergence of
groups of similar species. *Proceedings of the National Academy of Sciences* 103:6230–6235.
- 1164 Shipley, B., D. Vile, and E. Garnier. 2006. From plant traits to plant communities: a statistical
mechanistic approach to biodiversity. *Science* 314:812–814.
- 1166 Silvestro, D., A. Antonelli, N. Salamin, and T. B. Quental. 2015. The role of clade competition in
the diversification of North American canids. *Proceedings of the National Academy of Sciences of
the United States of America* 112:8684–9.
- 1168 Silvestro, D., J. Schnitzler, L. H. Liow, A. Antonelli, and N. Salamin. 2014. Bayesian estimation of
speciation and extinction from incomplete fossil occurrence data. *Systematic biology* 63:349–67.
- 1170 Simberloff, D., and T. Dayan. 1991. The Guild Concept and the Structure of Ecological
Communities. *Annual Review of Ecology and Systematics* 22:115–143.
- 1172 Slater, G. J. 2015. Iterative adaptive radiations of fossil canids show no evidence for
diversity-dependent trait evolution. *Proceedings of the National Academy of Sciences*
112:4897–4902.
- 1174 Smith, F. A., J. Brown, J. Haskell, and S. Lyons. 2004. Similarity of mammalian body size across
the taxonomic hierarchy and across space and time. *The American Naturalist* 163:672–691.
- 1176 Smith, F. A., S. K. Lyons, S. Morgan Ernest, and J. H. Brown. 2008. Macroecology: more than the
division of food and space among species on continents. *Progress in Physical Geography*
32:115–138.
- 1178 Smits, P. D. 2015. Expected time-invariant effects of biological traits on mammal species duration.
1180 *Proceedings of the National Academy of Sciences* 112:13015–13020.
- 1182 Stan Development Team. 2016. Stan Modeling Language Users Guide and Reference Manual.
- 1184 Strömberg, C. A. E. 2005. Decoupled taxonomic radiation and ecological expansion of open-habitat
grasses in the Cenozoic of North America. *Proceedings of the National Academy of Sciences of
the United States of America* 102:11980–4.

- Tomiya, S. 2013. Body Size and Extinction Risk in Terrestrial Mammals Above the Species Level.
1188 The American Naturalist 182:196–214.
- Urban, M. C., M. A. Leibold, P. Amarasekare, L. De Meester, R. Gomulkiewicz, M. E. Hochberg,
1190 C. A. Klausmeier, N. Loeuille, C. de Mazancourt, J. Norberg, J. H. Pantel, S. Y. Strauss,
M. Vellend, and M. J. Wade. 2008. The evolutionary ecology of metacommunities. Trends in
1192 Ecology and Evolution 23:311–317.
- Valentine, J. W. 1969. Patterns of taxonomic and ecological structure of the shelf benthos during
1194 Phanerozoic time. Paleontology 12:684–709.
- Van Valkenburgh, B. 1990. Skeletal and dental predictors of body mass in carnivores. Pages
1196 181–205 in J. Damuth and B. J. Macfadden, eds. Body size in mammalian paleobiology:
estimation and biological implications. Cambridge University Press, Cambridge.
- 1198 ———. 1999. Major patterns in the history of carnivorous mammals. Annual Review of Earth and
Planetary Sciences 27:463–493.
- Villéger, S., P. M. Novack-Gottshall, and D. Mouillot. 2011. The multidimensionality of the niche
1200 reveals functional diversity changes in benthic marine biotas across geological time. Ecology
1202 letters 14:561–8.
- Wang, S. C., P. J. Everson, H. J. Zhou, D. Park, and D. J. Chudzicki. 2016. Adaptive credible
1204 intervals on stratigraphic ranges when recovery potential is unknown. Paleobiology 42:240–256.
- Wang, S. C., and C. R. Marshall. 2016. Estimating times of extinction in the fossil record. Biology
1206 Letters 12:20150989.
- Warton, D. I., B. Shipley, and T. Hastie. 2015. CATS regression - a model-based approach to
1208 studying trait-based community assembly. Methods in Ecology and Evolution 6:389–398.
- Weber, M. G., C. E. Wagner, R. J. Best, L. J. Harmon, and B. Matthews. 2017. Evolution in a
1210 Community Context: On Integrating Ecological Interactions and Macroevolution. Trends in
Ecology & Evolution xx:1–14.

- 1212 Wilson, J. B. 1999. Guilds, functional types and ecological groups. *Oikos* 86:507–522.
- Yoder, J. B., E. Clancey, S. Des Riches, J. M. Eastman, L. Gentry, W. Godsoe, T. J. Hagey,
1214 D. Jochimsen, B. P. Oswald, J. Robertson, B. A. J. Sarver, J. J. Schenk, S. F. Spear, and L. J.
Harmon. 2010. Ecological opportunity and the origin of adaptive radiations. *Journal of
1216 Evolutionary Biology* 23:1581–1596.
- Zachos, J. C., G. R. Dickens, and R. E. Zeebe. 2008. An early Cenozoic perspective on greenhouse
1218 warming and carbon-cycle dynamics. *Nature* 451:279–283.
- Zachos, J. C., M. Pagani, L. Sloan, E. Thomas, and K. Billups. 2001. Trends, rhythms, and
1220 aberrations in global climate 65 Ma to present. *Science* 292:686–693.




Figure 11: Posterior mean estimates of the correlations in origination probability between the mammal ecotypes. The lower triangle of the matrix is populated with ellipses corresponding to the level of correlation between the two ecotypes, while the upper triangle of the matrix corresponds to the mean estimated correlation between ecotypes. Darker values correspond to a greater magnitude of correlation with blue values corresponding to a positive correlation and red values a negative correlation.




Figure 12: Probability of a mammal ecotype survival probabilities at each time point as estimated from the birth-death model. Each panel depicts 100 random samples from the model’s posterior. The columns are by locomotor category and rows by dietary category; their intersections are the observed and analyzed ecotypes. Panels with no lines are ecotypes not observed in the dataset.




Figure 13: Differences in log-odds of survival based on mammal orders. Positive values correspond to greater log-odds of survival than average, while negative values correspond to lower log-odds of survival than average.




Figure 14: Mean estimate of the effect of species mass on the probability of a species survival for each of the three plant phases. The effect of mass is considered constant over time and that the only aspect of the model that changes with plant phase is the intercept of the relationship between mass and survival. The three plant phases are indicated by the color of the line. Mass has been log-transformed, centered, and rescaled; this means that a mass of 0 corresponds to the mean of log-mass of all observed species and that mass is in standard deviation units. For clarity, only the mean estimates of the effects of mass and plant plant are plotted.




Figure 15: Estimated effects of the group-level covariates describing environmental context on log-odds of species survival. These estimates are from the birth-death model. What is plotted is a violin of the distribution of 1000 samples from the approximate posterior.




Figure 16: Posterior mean estimates of the correlations in survival probability between the mammal ecotypes. The lower triangle of the matrix is populated with ellipses corresponding to the level of correlation between the two ecotypes, while the upper triangle of the matrix corresponds to the mean estimated correlation between ecotypes. Darker values correspond to a greater magnitude of correlation with blue values corresponding to a positive correlation and red values a negative correlation.




Figure 17: Log diversity

Figure 18: Posterior estimates of the time series of Cenozoic North American mammal diversity; all estimates are from 100 posterior draws. The blue horizontal strip corresponds to the median and 80% credible interval of estimated mean standing diversity.




Figure 19: Posterior of standing log-diversity of North American mammals by ecotype for the Cenozoic as estimated from the birth-death model; 100 posterior draws are plotted to indicate the uncertainty in these estimates and what is technically plotted is log of diversity plus 1.




Figure 20: Mean posterior estimate of relative log standing diversity of 18 North American mammal ecotypes for the Cenozoic. These estimates are calculated from 100 posterior estimates of the true occurrence matrix z as estimated from the birth-death model.