

Cosmopolitan and provincialism dynamics of North American terrestrial mammals across the Cenozoic

Peter D Smits¹ and John Alroy²

¹Committee on Evolutionary Biology, University of Chicago

²Department of Biological Sciences, Macquarie University

September 26, 2013

Abstract

(Keywords: macroevolution, paleoecology, herbivory, trophic structure, biogeography)

1 Introduction

Evolutionary paleoecology is the study of how ecological traits expressed at any level affect the macroevolutionary process, or long term patterns in speciation and extinction (Kitchell, 1985). Community structure plays a fundamental roll in determining which biotic interactions are possible at a given time. The importance of biotic interactions in understanding evolutionary dynamics

In order to investigate how cosmopolitan–endemism dynamics vary across the Cenozoic, the terrestrial mammal records of North America, Europe and South America. The mammalian taxonomic dynamics of each of these three continents have been studied to different degrees (Alroy, 1996; Blois and Hadly, 2009; Eronen et al., 2009; Flynn and Wyss, 1998; Gunnell et al., 1995; Jernvall and Fortelius, 2002, 2004; Macfadden, 1997, 2006; Raia et al., 2006). While all of these continents transitioned from a principally closed and forested environment to a open and savanna like environment over Cenozoic, the timing of the start of this transition was different on each continent (Blois and Hadly, 2009; Eronen et al., 2009; Janis, 1993). Additionally, the mammal lineages present on each continent are divergent especially in the case on the isolated continent of South America (Flynn and Wyss, 1998; Macfadden, 1997, 2006). Because of this it can be expected that there will be a mix of similarities and striking differences between the continents in how taxonomic distributions change over the Cenozoic.

As the environment on each continent moved from a closed, forested environment to an open, savanna-like environment there are a few general predictions of shifts in cosmopolitan–

endemism dynamics. It is expected that as the environment shifted, the ratio of arboreal to land dwelling mammals would shift to include an even greater proportion of land dwelling mammals. Additionally, it is expected there would be an increased similarity between localities, as reflected by decreased endemism in land dwelling taxa. Arboreal taxa, however, are expected to trend towards increasing endemism and decreased site similarity as driven by increasingly fragmented forests.

The shift to grassland environments is frequently connected with the shift from browser-dominated communities to grazer-dominated communities CITATIONS, however in this study the focus is instead on coarser trophic categories such as herbivore and carnivore. There is an expected correlation between herbivore dynamics and land dwelling mammal dynamics because of the expansion of terrestrial herbivores (e.g. horses) CITATIONS.

South America is expected to have experienced rather different cosmopolitan–endemism dynamics than both North America and Europe because of both its long-term isolation and extremely biome provincialism (Flynn and Wyss, 1998; Macfadden, 2006) OTHER CITATIONS. The Amazonian lowland and Andean highlands are radically different biomes that diverged in similarity approximately during the Miocene (Ortiz-Jaureguizar and Cladera, 2006; Pascual and Ortiz-Jaureguizar, 1990) and are now a rainforest and an arid desert, respectively. Because of this, it is expected that there may be increasing and extreme endemism over the Cenozoic. However, the relative paucity of the South American record in comparison to North America and Europe may be a potential confounding factor for interpreting these results.

2 Materials and Methods

Mammalian occurrence information was obtained from the Paleobiology Database (PBDB; <http://www.paleodb.org>). Occurrence information was restricted to terrestrial mammals from the North American fossil record of the Cenozoic. For each occurrence, the locality information was also recorded, most importantly formation name and estimated age. Hierarchical taxonomic information was recorded for each taxon. All partially, ambiguously, or incompletely identified genera or species (e.g. aff., cf., ?) were excluded from analysis.

For each taxon, dietary and life habit information recorded in the PBDB was gathered. Dietary information was then simplified into three categories: herbivore, omnivore, and carnivore. Herbivorous taxa were the amalgam of the PBDB classifications herbivore, grazer, browser, folivore, and granivore. Omnivorous taxa were the amalgam of frugivores and omnivores. Finally, carnivores were the amalgam of all carnivores and insectivores. These three categories were chosen because they represent coarse groups which are identifiable from most mammalian teeth which are the primarily mammalian fossil material. Additionally, these groupings have been used in prior analysis of the effect of ecology on site similarity (Jernvall and Fortelius, 2004).

Temperature information was estimated using the δO^{18} isotope information from the benthic
 foram record for the entire Cenozoic (Zachos et al., 2008). An increase in δO^{18} levels are
 associated with a decrease in atmospheric temperature (Zachos et al., 2008, 2001). Benthic
 foram δO^{18} information has been used previously as climatic information in studies of
 mammalian macroevolutionary patterns (Alroy et al., 2000; Figueirido et al., 2012; Rose
 et al., 2011).

Taxonomic presence-absence was recorded for each formation. Abundance information was
 not included in this analysis, though it is possible (Sidor et al., 2013), because the highly
 variable preservation conditions not just between formations but across the Cenozoic may
 not accurately record abundance information (Damuth, 1982).

Bipartite biogeographic networks were constructed, following Sidor et al. (2013) and Vilhena
 et al. (2013), with species as the taxonomic occurrence and formation as the locality of
 interest. As explained in Sidor et al. (2013), biogeographic networks have many advantages
 over ordination based methods that are frequently used in numerical ecology (Legendre
 and Legendre, 2012). Principally, both taxonomic and locality information are preserved
 in analysis which allows for a more complete understanding of community structure. For
 each temporal bin biogeographic networks were constructed for all taxonomic information,
 individually for each dietary category and each locomotor category.

Four measures of biogeographic network structure were used to assess community change:
 code length as measured via the map equation (Rosvall et al., 2010; Rosvall and Bergstrom,
 2007, 2008), biogeographic connectedness, average number of locality occurrences per taxon,
 and average number of endemics per site. Biogeographic connectedness is defined

$$BC = \frac{O - N}{LN - N} \quad (1)$$

where O is the number of edges or number of occurrences in the biogeographic network,
 N is the number of taxa, and L is the number of localities (Sidor et al., 2013). All four of
 these measures have previously been used to assess cosmopolitan and endemism dynamics
 (Sidor et al., 2013). The four measures of biogeographic network were then calculated for each
 temporal bin for the total network, each of the dietary networks, and each of the locomotor
 category networks.

Network analysis, including calculation of network code length, was done using the **igraph**
 package (Csardi and Nepusz, 2006) for the R language (R Core Team, 2013).

3 Results

4 Discussion

Acknowledgements

Kenneth Angielczyk, David Bapst, Michael Foote, Benjamin Frable, Dallas Krentzel, Carl Simpson. Money? What money?

References

- Alroy, J. 1996. Constant extinction, constrained diversification, and uncoordinated stasis in North American mammals. *Palaeogeography, Palaeoclimatology, Palaeoecology* 127:285–311.
- Alroy, J., P. L. Koch, and J. C. Zachos. 2000. Global climate change and North American mammalian evolution. *Paleobiology* 26:259–288.
- Blois, J. L., and E. A. Hadly. 2009. Mammalian Response to Cenozoic Climatic Change. *Annual Review of Earth and Planetary Sciences* 37:181–208.
- Csardi, G., and T. Nepusz. 2006. The igraph software package for complex network research. *InterJournal Complex Systems*:1695.
- Damuth, J. 1982. Analysis of the preservation of community structure in assemblages of fossil mammals. *Paleobiology* 8:434–446.
- Eronen, J. T., A. R. Evans, M. Fortelius, and J. Jernvall. 2009. The impact of regional climate on the evolution of mammals: a case study using fossil horses. *Evolution* 64:398–408.
- Figueirido, B., C. M. Janis, J. a. Pérez-Claros, M. De Renzi, and P. Palmqvist. 2012. Cenozoic climate change influences mammalian evolutionary dynamics. *Proceedings of the National Academy of Sciences* 109:722–727.
- Flynn, J. J., and a. R. Wyss. 1998. Recent advances in South American mammalian paleontology. *Trends in ecology & evolution* 13:449–54.
- Gunnell, G., M. Morgan, M. C. Mass, and P. D. Gingerich. 1995. Comparative paleoecology of Paleogene and Neogene mammalian faunas: trophic structure and composition. *Palaeogeography, Palaeoclimatology, Palaeoecology* 115:265–286.
- Janis, C. M. 1993. Tertiary mammal evolution in the context of changing climates, vegetation, and tectonic events. *Annual Review of Ecology and Systematics* 24:467–500.

Jernvall, J., and M. Fortelius. 2002. Common mammals drive the evolutionary increase of
120 hypsodonty in the Neogene. *Nature* 417:538–40.

———. 2004. Maintenance of trophic structure in fossil mammal communities: site occupancy
122 and taxon resilience. *American Naturalist* 164:614–624.

Kitchell, J. A. 1985. Evolutionary paleoecology: recent contributions to evolutionary theory.
124 *Paleobiology* 11:91–104.

Legendre, P., and L. Legendre. 2012. *Numerical Ecology*. 3rd ed. Elsevier, Amsterdam.

126 Macfadden, B. J. 1997. Origin and evolution of the grazing guild in New World terrestrial
mammals. *Trends in ecology & evolution* 12:182–187.

128 ———. 2006. Extinct mammalian biodiversity of the ancient New World tropics. *Trends in
ecology & evolution* 21:157–65.

130 Ortiz-Jaureguizar, E., and G. Cladera. 2006. Paleoenvironmental evolution of southern South
America during the Cenozoic. *Journal of Arid Environments* 66:498–532.

132 Pascual, R., and E. Ortiz-Jaureguizar. 1990. Evolving climates and mammal faunas in
Cenozoic South America. *Journal of Human Evolution* 19:23–60.

134 R Core Team. 2013. *R: A Language and Environment for Statistical Computing*. R Foundation
for Statistical Computing, Vienna, Austria.

136 Raia, P., C. Meloro, A. Loy, and C. Barbera. 2006. Species occupancy and its course in the
past: macroecological patterns in extinct communities. *Evolutionary Ecology Research*
138 8:181–194.

Rose, P. J., D. L. Fox, J. Marcot, and C. Badgley. 2011. Flat latitudinal gradient in Paleocene
140 mammal richness suggests decoupling of climate and biodiversity. *Geology* 39:163–166.

Rosvall, M., D. Axelsson, and C. T. Bergstrom. 2010. The map equation. *The European*
142 *Physical Journal Special Topics* 178:13–23.

Rosvall, M., and C. T. Bergstrom. 2007. An information-theoretic framework for resolving
144 community structure in complex networks. *Proceedings of the National Academy of Sciences
of the United States of America* 104:7327–31.

146 ———. 2008. Maps of random walks on complex networks reveal community structure.
Proceedings of the National Academy of Sciences 105:1118–23.

148 Sidor, C. a., D. a. Vilhena, K. D. Angielczyk, A. K. Huttenlocker, S. J. Nesbitt, B. R. Peacock,
J. S. Steyer, R. M. H. Smith, and L. a. Tsuji. 2013. Provincialization of terrestrial faunas
150 following the end-Permian mass extinction. *Proceedings of the National Academy of
Sciences* 110:8129–33.

152 Vilhena, D. a., E. B. Harris, C. T. Bergstrom, M. E. Maliska, P. D. Ward, C. a. Sidor, C. a. E.

- 154 Strömberg, and G. P. Wilson. 2013. Bivalve network reveals latitudinal selectivity gradient
at the end-Cretaceous mass extinction. *Scientific reports* 3:1790.
- 156 Zachos, J. C., G. R. Dickens, and R. E. Zeebe. 2008. An early Cenozoic perspective on
greenhouse warming and carbon-cycle dynamics. *Nature* 451:279–283.
- 158 Zachos, J. C., M. Pagani, L. Sloan, E. Thomas, and K. Billups. 2001. Trends, rhythms, and
aberrations in global climate 65 Ma to present. *Science* 292:686–693.