

# Cosmopolitan and endemism dynamics of terrestrial mammals across the Cenozoic

Peter D Smits

Committee on Evolutionary Biology, University of Chicago

## Abstract

How life history traits, such as diet, are related to the distribution of taxa across the landscape.

## Introduction

Previous work on mammalian site similarity has focused on organismal dietary distributions of terrestrial mammals in the Neogene Old World [1, 2]. Here, I expand that analysis the entire Cenozoic of North America and analyze both dietary and locomotor categories of terrestrial mammals.

## Methods

Mammalian taxonomic occurrence information was obtained from the Paleobiology Database (<http://www.paleodb.org>). Taxonomic occurrence information was restricted to only mammals occurring in North America during the Cenozoic. Ambiguously identified taxa were excluded from all analyses (e.g. aff., cf., ?). Temporal, geologic, dietary and life habit informaiton was also compiled for all taxa.

Because terrestrial assemblages across the Cenozoic do not preserve as complete a record of community structure, taxonomic abundance distributions were not analyzed.

Following Sidor et al. [3] and Vilhena et al. [4]

## Relative abundance




Figure: Relative abundance of mammalian dietary categories.

## Results

## Fancy




Figure: Biogeographic network summary statistics from the 2 My bins.

## Discussion

## Conclusions

## Acknowledgements

John Alroy, Michael Foote, Ken Angielczyk.

## Bibliography

- [1] J. Jernvall and M. Fortelius. Common mammals drive the evolutionary increase of hypsodonty in the Neogene. *Nature*, 417(6888):538–40, May 2002. ISSN 0028-0836. doi: 10.1038/417538a. URL <http://www.ncbi.nlm.nih.gov/pubmed/12037565>.
- [2] J. Jernvall and M. Fortelius. Maintenance of trophic structure in fossil mammal communities: site occupancy and taxon resilience. *American Naturalist*, 164(5):614–624, Nov. 2004. ISSN 1537-5323. doi: 10.1086/424967. URL <http://www.ncbi.nlm.nih.gov/pubmed/15540151>.
- [3] C. a. Sidor, D. a. Vilhena, K. D. Angielczyk, A. K. Huttenlocker, S. J. Nesbitt, B. R. Peacock, J. S. Steyer, R. M. H. Smith, and L. a. Tsuji. Provincialization of terrestrial faunas following the end-Permian mass extinction. *Proceedings of the National Academy of Sciences*, 110(20):8129–33, May 2013. ISSN 1091-6490. doi: 10.1073/pnas.1302323110. URL <http://www.ncbi.nlm.nih.gov/pubmed/23630295>.
- [4] D. a. Vilhena, E. B. Harris, C. T. Bergstrom, M. E. Maliska, P. D. Ward, C. a. Sidor, C. a. E. Strömberg, and G. P. Wilson. Bivalve network reveals latitudinal selectivity gradient at the end-Cretaceous mass extinction. *Scientific reports*, 3:1790, May 2013. ISSN 2045-2322. doi: 10.1038/srep01790. URL <http://www.ncbi.nlm.nih.gov/pubmed/23649088>.