

# **Death and taxa: time invariant differences in mammal species duration**

Peter D Smits<sup>1</sup>

<sup>1</sup>*Committee on Evolutionary Biology, University of Chicago, Chicago, IL, USA.*

**Why species go extinct at different rates remains one of the most fundamental questions in paleobiology <sup>1-5</sup>. Determining which and how biological traits influence extinction risk is vital for understanding the differential diversification of life during the Phanerozoic and for making predictions about species' vulnerability to anthropogenic impacts. Here I present a hierarchical Bayesian survival model of North American Cenozoic mammal species durations as predicted by the species-level ecological factors, time of origination, and phylogenetic relationships. I also explicitly model the effect of species age on extinction risk, relaxing the Law of Constant Extinction which is the critical assumption underlying the Red Queen Hypothesis <sup>2</sup>. This study focuses on time-invariant effects in an effort to characterize aspects of the selective patterns of background extinction in order to identify if the current biodiversity crisis represents a shift selection pressures associated with a shift in "macroevolutionary regime" <sup>6</sup> or arrival at a "tipping point" <sup>7,8</sup>.**

Here I test if extinction is non-random with respect to organismal- and species-level traits during times of background extinction, if and which traits have time-invariant effects on species duration, and if extinction is taxon-age independent among Cenozoic mammals? All of these questions are combined into developing a single model of survival whose parameter estimates directly test these questions. Cenozoic mammals represent an ideal group and time period because their fossil record is well sampled, well resolved both temporally and spatially, and the ecology and taxonomy of individual species are generally understood<sup>1, 4, 10–14</sup>.

This study focuses on identifying the time-invariant effects of organismal and species traits on the expected duration of a species. A time-invariant effect is that when comparing taxa over a long time period there is a consistent effect that is generalizable over the entire period of interest. While the strength of that effect may vary over time, the direction does not change. These consistent effects reveal fundamental differences in selective regime. Periods of background extinction provide a great opportunity to study how traits are related to survival because they represent the majority of geologic time, remain relatively predictable, and change slowly<sup>6, 16</sup>.

The organismal and species traits studied here are dietary and locomotor

categories, bioprovince occupancy, and body mass. Each of these traits describe different aspects of a species' adaptive zone such as energetic cost, population density, expected home range size, set of potential prey items, and dispersal ability<sup>12,18</sup>. It is expected that species with larger geographic ranges have lower extinction rates than species with smaller geographic ranges<sup>6,19</sup>. However, organismal traits directly related to species–environment interactions may play an important role in determining extinction risk. By modeling extinction via traits related to environmental preference, the relative importance of species- and organismal-level properties can be elucidated. All of these traits are analyzed with respect to the individual species origination cohort and phylogenetic position. From there the relative contribution species, cohort, and phylogeny to the unexplained variance can be estimated.

I found that dietary category has a large amount of variation in the pairwise differences of effects on expected duration (Fig. ). Carnivory appears to be associated with a greater expected duration than herbivory or insectivory, while approximately equal to or less than the expected duration of an omnivore. Omnivory is associated with greater expected duration than either herbivory or insectivory. Finally, herbivory and insectivory are associated with approximately equal effects

on expected duration. These results sit nicely beside previous ones<sup>20</sup>. Given that carnivores and omnivores have approximately equal extinction risk, and it has been found previously that carnivores have a greater diversification rate than omnivores, this implies that carnivores have a greater origination rate than omnivores. This comparison implies that herbivores which have the greatest extinction risk must also have a very high origination rate in order to have the greatest diversification rate of these three categories.

For locomotor category, arboreality appears to be associated with a lower expected duration than either scansoriality or a ground dwelling life habit (Fig. ). Scansoriality and a ground dwelling life habit have approximately equal expected durations. As expected, bioprovince occupancy has the largest effect on expected species duration/extinction risk. The other factors all appear to have small effect sizes which is also consistent with expectations (Fig. 5). Body size has a very small, near zero effect on expected duration, similar to the lack of relationship between body size and generic duration<sup>13</sup>. These results consistent with the hypotheses that arboreality is associated with a greater expected extinction risk than either with scansoriality and ground dwelling taxa. Importantly, scansoriality appears to not influence any difference in extinction risk when compared with

ground dwelling taxa. This can be interpreted that arboreal taxa, which require a specific kind of environment, may be more prone to extinction because the lack of permanency of those environments preventing species persistence.

The large difference in time-invariant extinction risk between omnivores and both herbivores and insectivores is most likely related to the concept of “survival of the unspecialized” where less specialized taxa have lower extinction risks <sup>1,21</sup>. Because larger effects are easier to identify, the magnitude of this effect also explains both the early identification and origin of this hypothesis <sup>1</sup>. The very weak if non-existent effect of body size on extinction risk, which can be considered not “significant”, is consistent with some previous results <sup>13</sup>. The direction/sign of the modal estimate of effect is not consistent with the prediction of increase in extinction risk associated with increase in body size <sup>11</sup>. However, the other studies were performed at the generic-level which may or may not involve different processes than at the species-level model.

Of the three sources of variance present in the model, individual species variance accounts for approximately 70% of total variance (Fig. 8). Both cohort and phylogenetic effects account for the other 30% of the observed variance. While both of these effects are the source of approximately 15% of observed vari-

ance individually, the total combined effect factors indicates that neither can be ignored. As  $VPC_{phylo}$ , which is equivalent to phylogenetic heritability, is greater than 0 it is not appropriate to ignore phylogeny when modeling survival<sup>22</sup> as is commonly done in paleontological studies<sup>10,11,15,23–25</sup>.

Because the estimate of the Weibull shape parameter,  $\alpha$ , is greater than 1 extinction risk is expected to increase with taxon age (Table 1). The estimate of  $\alpha$  is also rather tightly constrained, having a small posterior standard deviation.  $\alpha$  is related to the strength of time on extinction risk and is a key parameter in the hazard function  $h(t)$  which can be interpreted as the rate, or approximate probability, of an individual of age  $t$  going extinct. As the value of  $\alpha$  is between 1 and 1.5, extinction risk for a given species only gradually increases with age (Fig. 7). This result has two possible explanations: (1) older taxa being aged out or outcompeted by younger taxa, or (2) as an artifact of the minimum resolution of the fossil record.

The hypothesis that older taxa are being outcompeted or replaced by younger taxa is also consistent with the some recent results<sup>4,5</sup>, both of which require that older taxa have a greater extinction risk than younger taxa.

The other possible explanation for the inferred increase in extinction risk with species age is the minimum resolution which might cause an upward bias in estimates of the Weibull shape parameter  $\alpha$ <sup>27</sup>, an effect which can be observed by the initial plateau in the K-M estimate of  $S(t)$  for the observed (Fig. 2). This plateau is a hallmark of the original paleontological survival analyses<sup>2</sup> which was identified as partially a product of minimum resolution of the fossil records of the different studied groups.

Given this known biasing factor and previous results<sup>4,5</sup>, I hypothesize that the inferred pattern is most likely a combination of these two explanations working on concert. In order to determine the relative importance of these two explanations, more work is required into approaches for directly modeling the minimum resolution of the fossil record.

The estimates for the individual cohort effects show a weak pattern of increased extinction risk in older Cenozoic cohorts and decreased extinction risk in younger cohorts (Fig. 6). However, this pattern is not very strong as there is a large amount of variation, particularly for older cohorts. For example, note the two cohorts between 50 and 55 My that have a much lower extinction risk than other cohorts of similar age. However, it is interesting to note that the appar-

ent shift from older cohorts with a higher extinction risk to younger cohorts with lower extinction risk is approximately 30 Mya or the Paleogene–Neogene boundary. This transition is marked by the opening up of the landscape and the rise of grazers and the decline heavily forested environments. This shift may underly the inferred increased extinction risk associated with arboreal species compared to ground dwelling or scansorial species (Fig. ). However, because the model used here does not allow for change in time-invariant effects, I cannot identify this transition as a tipping point or shift in selective regime <sup>7,8</sup>.

One of the open questions in paleobiology and macroecology is whether the current biodiversity crisis qualified as a mass extinction <sup>7,8,28</sup>. Because change in the magnitude of extinction risk is not necessarily the best indicator of a shift from background to mass extinction <sup>29</sup>, it is more fruitful to look for changes in the direction selection, loss of a selective pressure, or appearance of novel selective pressures. Comparison of the estimated effects of organismal- and species-level traits analyzed here with previous studies demonstrates a mixture of congruence and incongruence.

As expected, large range size is always associated with lower extinction risk in the Recent <sup>30–33</sup>. While I found that body size has no time-invariant effect on



extinction risk, large body size is associated with increased extinction risk in the Recent, though this is variable across environments and clades <sup>30,32,33</sup>. A higher trophic level (e.g. carnivory versus herbivory) is associated with greater extinction risk in Primates and Carnivora <sup>33</sup> which is not congruous with the results found here that carnivores have lower extinction risk than herbivores. Finally, phylogeny has been found to be a factor underlying current mammal species extinction risk, though this effect seems much greater in the Recent than for the whole Cenozoic <sup>31</sup>. Note that the phylogeny of Recent mammals is much better than the primarily taxonomy based phylogeny used here, which may partially account for the difference in effect.

How many of these incongruities are within the standard range of time-variant effects is unknown, though these comparisons across multiple factors do point to our arrival at a tipping point <sup>7,8</sup> and potentially a shift in macroevolutionary regime <sup>6</sup>.

There are a few data quality concerns in this study which are also inherent to almost any paleontological study. Almost all of the body mass estimates were obtained using published regression equations that estimate mass from some other body part (e.g. tooth). These estimates are known with error, which was

not included in the model. If the standard deviation of the residuals from each of these regression equations was known, it would be possible to directly model this as measurement error <sup>34</sup> though this greatly adds to model complexity and decreases some amount of interpretability. Also phylogeny used here is only a coarse, baseline estimate of the actual species relationships. Because of this, the analysis of phylogenetic effect on survival represents a minimum estimate. As it stands though, these results point to the importance of including shared evolutionary history in diversification models.

There are many processes encompassed by background extinction and identifying the exact cause of any one species' reason for extinction is extremely difficult. By focusing on estimating the effects of different ecologies and historical factors on average extinction risk, it is possible to better understand what processes may have driven the resulting pattern of selection (i.e. diversity). Here, I focused on time-invariant factors and their relation to biological selectivity of extinction, possible reasons for the observed time-invariant effects, and the effects of taxon-age on extinction risk. I found that some organismal- and species-level traits such as omnivory and large geographic range size have time-invariant effects on mammal species extinction risk. I also found that there are small but

non-ignorable effects of cohort and phylogeny. Finally, I found putative evidence of increasing extinction risk with species age, though this result may be partially due to the minimum resolution of the fossil record itself <sup>27</sup>.

## **Methods**

**Data and covariate information.** Fossil occurrence information was downloaded from the Paleobiology Database (PBDB). This data set was filtered so that only occurrences identified to the species-level, excluding all “sp.”-s. All aquatic and volant taxa were also excluded. Additionally, all occurrences without latitude and longitude information were excluded.

Species dietary and locomotor category assignments were based on assignments in initial the PBDB which were then reassigned into coarser categories.

All individual fossil occurrences were assigned to 2 My bins ranging through the entire Cenozoic. Species duration was measured as the number of bins from the first occurrence to the last occurrence, inclusive. This bin size was chosen because it approximately reflects the resolution of the North American Cenozoic mammal fossil record <sup>10,14</sup>. The youngest cohort, 0-2 My, was excluded from analysis.

Species body size estimates were sourced from a large selection of primary literature and data base compilations, principally the PBDB, the Neogene Old World Mammal database, and other large scale data collection efforts <sup>35–39</sup>.

For each 2 My bin, a bipartite biogeographic network was created between species occurrences and spatial units, defined as 2x2 latitude–longitude grid cells from an azimuthal equal-area map projection. Emergent bioprovinces within the biogeographic occurrence network were identified using the map equation <sup>40,41</sup>. A bioprovince is a set of species–locality connections that are more interconnected within the group than without.

As there is no single, combined formal phylogenetic hypothesis of all Cenozoic fossils mammals from North America, it was necessary to construct a supertree as a minimum estimate of phylogenetic relations. This was done using matrix representation parsimony to combine multiple large species level phylogenies and the taxonomic information for all the observed species <sup>37,42–44</sup>. Polytomies were resolved in the order of species first appearance and then time-scaled using the “minimum branch length” approach where all branches were a minimum of 0.1 My.

**Survival model.** Species durations were modeled as being drawn from a Weibull distribution with two parameters: shape  $\alpha$  and scale  $\sigma$ .  $\sigma$  was parameterized as a regression model with an exponential link function because it is defined only for values greater than 0. This regression model included eight species-level coefficients for the covariates (dietary and locomotor categories, body size, mean bioprovince occupancy) along with an intercept term. Both continuous covariates were standardized by subtracting the mean value and dividing by 2 times the standard deviation so that the inferred values were comparable with the coefficients from the binary variables <sup>45</sup>. Each of these regression coefficients was given a weakly informative normally distributed prior.

Additionally, a hierarchical term was included for origination cohort along with an individual effect for phylogeny. Cohorts were considered drawn from a shared normal distribution with mean 0 and standard deviation estimated from the data. This standard deviation term was given a weakly informative half-Cauchy prior <sup>34</sup>. Individual phylogenetic effect was considered drawn from a multivariate normal distribution with mean vector of all 0's and covariance matrix being the phylogenetic covariance matrix multiplied by an unknown constant <sup>22,26</sup>.

An important part of survival analysis is censoring observations where the

failure time has not been observed. A right censored is where the point of extinction had not yet been observed in the period of study, such as taxa that are still extant. Left censored observations correspond to observations that went extinct any time between 0 and some known point. In order to account for the minimum resolution of the fossil record encountered here, taxa that occurred in only a single time bin were left censored.

The above model was compared with a nearly identical one where species durations were instead assumed to have been drawn from a exponential distribution which is equivalent to a Weibull distribution with  $\alpha = 1$ .

Parameters were estimated using the Monte Carlo routine implemented in the Stan probabilistic programming language <sup>46</sup>. 4 chains were run in parallel for 20000 samples thinned to every 20th, divided evenly between warm-up and sampling.

Model adequacy was assessed through a multiple posterior predictive checks to determine if the model accurately reflected the observed species durations <sup>34</sup>. The reasoning behind posterior predictive checks being that, if the model accurately captures the observed variation in the data, then simulated data under the fitted model should be similar to the observed data. An example posterior predic-

tive check is comparing an observed summary statistic (e.g. mean) to the distribution of means estimated from 1000 different simulated data sets.

To characterize the contribution to total variance of the different data levels (i.e. individual species, cohort, phylogenetic position), variance partitioning coefficient were calculated using a variation on a randomization approach for models with non-normally distributed errors <sup>47</sup>.

1. Simpson, G. G. *Tempo and Mode in Evolution* (Columbia University Press, New York, 1944).
2. Van Valen, L. A new evolutionary law. *Evolutionary Theory* **1**, 1–30 (1973).
3. Raup, D. M. The role of extinction in evolution. *Proceedings of the National Academy of Sciences* **91**, 6758–6763 (1994).
4. Quental, T. B. & Marshall, C. R. How the Red Queen Drives Terrestrial Mammals to Extinction. *Science* **341**, 290–292 (2013).
5. Wagner, P. J. & Estabrook, G. F. Trait-based diversification shifts reflect differential extinction among fossil taxa. *Proceedings of the National Academy of Sciences of the United States of America* 1–6 (2014).

6. Jablonski, D. Background and mass extinctions: the alternation of macroevolutionary regimes. *Science* **231**, 129–133 (1986).
7. Barnosky, A. D. *et al.* Has the Earth's sixth mass extinction already arrived? *Nature* **470**, 51–57 (2011).
8. Barnosky, A. D. *et al.* Approaching a state shift in Earth's biosphere. *Nature* **486**, 52–8 (2012).
9. Raup, D. M. Cohort Analysis of generic survivorship. *Paleobiology* **4**, 1–15 (1978).
10. Alroy, J. Speciation and extinction in the fossil record of North American mammals. In Butlin, R. K., Bridle, J. R. & Schluter, D. (eds.) *Speciation and patterns of diversity*, 302–323 (Cambridge University Press, Cambridge, 2009).
11. Liow, L. H. *et al.* Higher origination and extinction rates in larger mammals. *Proceedings of the National Academy of Sciences* **105**, 6097 (2008).
12. Smith, F. A., Brown, J., Haskell, J. & Lyons, S. Similarity of mammalian body size across the taxonomic hierarchy and across space and . . . . *The American Naturalist* (2004).



13. Tomiya, S. Body Size and Extinction Risk in Terrestrial Mammals Above the Species Level. *The American Naturalist* E000–E000 (2013).
14. Marcot, J. D. The fossil record and macroevolutionary history of North American ungulate mammals: standardizing variation in intensity and geography of sampling. *Paleobiology* **40**, 237–254 (2014).
15. Payne, J. L. & Finnegan, S. The effect of geographic range on extinction risk during background and mass extinction. *Proceedings of the National Academy of Sciences of the United States of America* **104**, 10506–11 (2007).
16. Raup, D. M. & Boyajian, G. E. Patterns of generic extinction in the fossil record. *Paleobiology* **14**, 109–125 (1988).
17. Finnegan, S., Payne, J. L. & Wang, S. C. The Red Queen revisited: reevaluating the age selectivity of Phanerozoic marine genus extinctions. *Paleobiology* **34**, 318–341 (2008).
18. Jernvall, J. & Fortelius, M. Maintenance of trophic structure in fossil mammal communities: site occupancy and taxon resilience. *American Naturalist* **164**, 614–624 (2004).

19. Roy, K., Hunt, G., Jablonski, D., Krug, A. Z. & Valentine, J. W. A macroevolutionary perspective on species range limits. *Proceedings. Biological sciences / The Royal Society* **276**, 1485–93 (2009).
20. Price, S. A., Hopkins, S. S. B., Smith, K. K. & Roth, V. L. Tempo of trophic evolution and its impact on mammalian diversification. *Proceedings of the National Academy of Sciences of the United States of America* **109**, 7008–12 (2012).
21. Liow, L. H. A test of Simpson’s ”rule of the survival of the relatively unspecialized” using fossil crinoids. *The American naturalist* **164**, 431–43 (2004).
22. Housworth, E. A., Martins, P. & Lynch, M. The Phylogenetic Mixed Model. *The American Naturalist* **163**, 84–96 (2004).
23. Foote, M. & Miller, A. I. Determinants of early survival in marine animal genera. *Paleobiology* **39**, 171–192 (2013).
24. Jablonski, D. & Hunt, G. Larval ecology, geographic range, and species survivorship in Cretaceous mollusks: organismic versus species-level explanations. *The American naturalist* **168**, 556–64 (2006).

25. Hunt, G. The relative importance of directional change, random walks, and stasis in the evolution of fossil lineages. In *Proceedings of the National Academy of...* (2007).
26. Lynch, M. Methods for the analysis of comparative data in evolutionary biology. *Evolution* **45**, 1065–1080 (1991).
27. Sepkoski, J. J. Stratigraphic biases in the analysis of taxonomic survivorship. *Paleobiology* **1**, 343–355 (1975).
28. Alroy, J. The Shifting Balance of Diversity Among Major Marine Animal Groups. *Science* **329**, 1191–1194 (2010).
29. Wang, S. C. On the continuity of background and mass extinction. *Paleobiology* **29**, 455–467 (2003).
30. Fritz, S. a., Bininda-Emonds, O. R. P. & Purvis, A. Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecology letters* **12**, 538–49 (2009).
31. Fritz, S. a. & Purvis, A. Selectivity in mammalian extinction risk and threat types: a new measure of phylogenetic signal strength in binary traits. *Con-*

- servation biology : the journal of the Society for Conservation Biology* **24**, 1042–51 (2010).
32. Liow, L. H., Fortelius, M., Lintulaakso, K., Mannila, H. & Stenseth, N. C. Lower Extinction Risk in SleeporHide Mammals. *The American Naturalist* **173**, 264–272 (2009).
  33. Purvis, a., Gittleman, J. L., Cowlishaw, G. & Mace, G. M. Predicting extinction risk in declining species. *Proceedings. Biological sciences / The Royal Society* **267**, 1947–52 (2000).
  34. Gelman, A. *et al. Bayesian data analysis* (Chapman and Hall, Boca Raton, FL, 2013), 3 edn.
  35. Jones, K. E. *et al. PanTHERIA : a species-level database of life history , ecology , and geography of extant and recently extinct mammals. Ecology* **90**, 2648 (2009).
  36. Smith, F. A. *et al. Similarity of Mammalian Body Size across the Taxonomic Hierarchy and across Space and Time. The American Naturalist* **163**, 672–691 (2004).

37. Raia, P., Carotenuto, F., Passaro, F., Fulgione, D. & Fortelius, M. Ecological specialization in fossil mammals explains Cope's rule. *The American naturalist* **179**, 328–37 (2012).
38. Freudenthal, M. & Martín-suárez, E. Estimating body mass of fossil rodents. *Scripta Geologica* **145**, 1–130 (2013).
39. McKenna, R. T. *Potential for Speciation in Mammals Following Vast , Late Miocene Volcanic Interruptions in the Pacific Northwest*. Masters, Portland State University (2011).
40. Rosvall, M. & Bergstrom, C. T. Maps of random walks on complex networks reveal community structure. *Proceedings of the National Academy of Sciences* **105**, 1118–23 (2008).
41. Vilhena, D. A. *et al.* Bivalve network reveals latitudinal selectivity gradient at the end-Cretaceous mass extinction. *Scientific reports* **3**, 1790 (2013).
42. Janis, C. M., Scott, K. M. & Jacobs, L. L. *Evolution of Tertiary mammals of North America. Vol. 1. Terrestrial carnivores, ungulates, and ungulatelike mammals* (Cambridge University Press, Cambridge, 1998).

43. Janis, C. M., Gunnell, G. F. & Uhen, M. D. *Evolution of Tertiary mammals of North America. Vol. 2. Small mammals, xenarthrans, and marine mammals* (Cambridge University Press, Cambridge, 2008).
44. Bininda-Emonds, O. R. P. *et al.* The delayed rise of present-day mammals. *Nature* **446**, 507–512 (2007).
45. Gelman, A. & Hill, J. *Data Analysis using Regression and Multi-level/Hierarchical Models* (Cambridge University Press, New York, NY, 2007).
46. Stan Development Team. Stan: A c++ library for probability and sampling, version 2.5.0 (2014). URL <http://mc-stan.org/>.
47. Goldstein, H., Browne, W. & Rasbash, J. Partitioning variation in multilevel models. *Understanding Statistics* **1**, 1–12 (2002).

**Acknowledgements** P.D.S would like to thank M. Foote, K. Angielczyk, R. Ree, P.D. Polly for discussion. J. Alroy and Fossilworks/the Paleobiology Database for data gathering, entry, and availability. This is Fossilworks/Paleobiology Database publication number XXX.

**Competing Interests** The authors declare that they have no competing financial interests.

**Correspondence** Correspondence and requests for materials should be addressed to P.D.S. (email: [psmits@uchicago.edu](mailto:psmits@uchicago.edu)).

Table 1: Summaries of 1000 samples drawn from the marginal posteriors for the principle parameters of interest. Because of variable standardization, the intercept can be interpreted as the estimate for the mean observed species. As such, the other values are expected effects of trait values expressed as deviation from the mean. The categorical variables are binary index variables where an observation is of that category or not.  $\hat{R}$  values of less than 1.1 indicate chain convergence for the posterior samples.

	mean	sd	2.5%	25%	50%	75%	97.5%	$\hat{R}$
alpha	1.31	0.03	1.25	1.29	1.31	1.34	1.38	1.01
intercept	-0.82	0.18	-1.17	-0.94	-0.82	-0.70	-0.48	1.00
ground dwelling	-0.12	0.12	-0.34	-0.20	-0.12	-0.04	0.12	1.00
scansorial	-0.13	0.13	-0.38	-0.22	-0.13	-0.05	0.12	1.00
herbivore	0.11	0.12	-0.13	0.03	0.12	0.19	0.35	1.00
insectivore	0.09	0.13	-0.17	0.00	0.09	0.19	0.36	1.00
omnivore	-0.12	0.13	-0.37	-0.20	-0.12	-0.04	0.14	1.00
logit(occupancy)	-0.66	0.08	-0.81	-0.71	-0.65	-0.60	-0.50	1.00
log(size)	-0.05	0.05	-0.15	-0.08	-0.05	-0.01	0.05	1.00
sd cohort	0.33	0.07	0.22	0.29	0.33	0.37	0.48	1.00
sd phylogeny	0.21	0.10	0.07	0.14	0.19	0.26	0.46	1.05



Figure 1: Deviance residuals from the fitted survival model. Each graph depicts the residuals from single draws from the posteriors distributions of all estimated parameters. Positive values indicate an under estimate of the observed duration, while negative values indicate an over estimate of the observed duration. Twelve difference examples are provided here to indicate the lack of individual observation based biases.

Figure 2: Comparison between K-M estimate of survival function (black) from the observed versus K-M estimates from 100 simulated data sets using the fitted model (dark grey). Simulated data sets were generated by drawing parameter values randomly from their estimated posteriors and using the observed covariate information to estimate durations for all the observed species. On the left are the results from the full survival model, while on the right are the results from a simplified model where duration follows an exponential distribution and there is no phylogenetic effect.

Figure 3: The results of additional posterior predictive checks for four summaries of the observed durations, as labeled. Blue vertical indicate the observed value. None of the observed are significantly different from the posterior predictive distributions.

Figure 4: Pairwise differences in effect of the locomotor ( ) and dietary categories ( ) on expected duration from 1000 samples from the posterior distribution. Comparisons of locomotor categories, from top to bottom ( ), are: arboreal versus ground dwelling, arboreal versus scansorial, and ground dwelling versus scansorial. For dietary category, from top to bottom ( ): carnivore versus herbivore, carnivore versus insectivore, carnivore versus omnivore, herbivore versus insectivore, herbivore versus omnivore, and insectivore versus omnivore. Values to the left indicate that the first category is expected to have a greater duration than the second, while values to the right indicate that the first category is expected to have a shorter duration.

Figure 5:

Figure 6: Summaries of estimated individual cohort effect posteriors. Depicted are medians and 80% credible intervals of the estimated posterior distributions. High values correspond to shorter species durations while lower values correspond to greater species durations compared to the mean duration. Lines are placed at the middle of the 2 My origination cohorts.

Figure 7: 100 estimates of the hazard function ( $h(t)$ ) for the observed species mean (grey), along with the median estimated hazard function.  $h(t)$  is an estimate of the rate at which a species of age  $t$  is expected to go extinct. Hazard functions were estimated from random draws from the estimated posterior distributions and evaluated with all covariate information set to 0, which corresponds to the expected duration of the mean species.

Figure 8: Estimates of the variance partitioning coefficients for the three different sources of variance: species, cohort, and phylogeny. Higher values correspond to greater contribution to total observed variance. Each of the estimates is a distribution of 1000 approximating simulations due to the model's non-normality.