

# Evolutionary paleoecology and the biology of extinction

Peter D. Smits  
psmits@uchicago.edu

Dissertation Proposal Hearing  
November 13, 2013  
Committee on Evolutionary Biology  
The University of Chicago

*Committee*

Dr. Michael J. Foote (co-advisor)  
Dr. Kenneth D. Angielczyk (co-advisor)  
Dr. Richard H. Ree  
Dr. P. David Polly

# Introduction and theoretical framework

## 2 Evolutionary paleoecology

Evolutionary paleoecology is defined as the study of the effects of ecological traits and factors on differential rate dynamics, particularly rates of faunal turnover and diversification [35]. Ecological traits and factors are any and all traits expressed by a taxon, at any level, that are involved with biotic–biotic or biotic–abiotic interactions. Diversification is the difference between origination and extinction and is the net pattern of macroevolution. The study of evolutionary paleoecology is then the link between environmental interactions and macroevolution. As a corollary to Kitchell [35]’s definition, Allmon [2] states that in order to correctly link ecological interactions to macroevolution, one must focus on the specific traits and factors that may affect the speciation process. Tacitly included in this is then the study of the biology of extinction and how it relates to ecological properties and interactions [34].

Simpson [68] defined also defined the environment broadly as to include all possible biotic and abiotic interactions as well as the organism itself. This usage of “environment” is frequently misinterpreted or unknown and has lead to a great deal of confusion thus it is important to note here [8, 41, 73]. Related to this, Simpson [68] defined the set of all biotic and abiotic interactions that a lineage experiences as the “adaptive zone.” The adaptive zone represents one of the fundamental metaphors in paleobiology and can be considered the adaptive landscape through time [68, 69].

It is under this framework that I purpose to study how ecological traits associated with range size have affected both the availability of biotic interactions and differential survivorship. I will be studying two very distantly related and biotically different groups: mammals and brachiopods. Both of these groups are considered to have very good fossil records able to reflect massive long term evolutionary patterns [46].

## Paleobiological theory

Extinction, when not during periods of “mass extinction,” is widely considered non-random with respect to biology [26]. Additionally, times of “background extinction” represent the vast majority of geological time compared to periods of “mass extinction.” The exact definition and distinction between periods of background and mass extinction is based on the relative intensity of extinction at a particular time versus times preceding and following. Simply put, the major indicator of a mass extinction is an event that is across taxonomic categories and the biotic interactors before and after the event are dramatically different [26, 27, 37, 38].

Paleobiology is the study of life over time and the processes that generate the observed patterns in diversity and disparity. Intimately related to paleobiology is the concept of macroevolution here defined as the pattern of speciation and extinction over time [28]. The

study of macroevolution is the estimation of the processes underlying these observed patterns. The term origination is frequently used in place of speciation because it includes both speciation and migration because, depending on both the spatial scale and quality of the fossil record, it may be impossible to distinguish between the two.

Macroevolution, as metaphor, subsumes both speciation/extinction (phyletic) and anagenetic/lineage disparity (anagenetic) dynamics [21]. This usage has also been termed the tempo and mode of evolution [68]. This metaphoric usage more closely links paleobiology and macroevolution. This raises an important question: can we have paleobiology without macroevolution?

The contrast to macroevolution is microevolution [21, 68] which is strictly defined as change in allele frequency in a population from one generation to the next. It is important to note that changes in allele frequency affects phenotype frequency and expression and thus, by corollary, describes phenotypic change. Just as Simpson [68] argued, this link can explain rates of evolution and (most) patterns of disparity without invoking any “macroevolutionary” phenomenon.

What makes a definition strong or weak? Why would we prefer one over the other? What is the difference between definition, metaphor and metonymy?

Of major concern with the metaphoric use of macroevolution is that this concept subsumes all aspects of anagenetic change making the difference between microevolution and macroevolution unclear. Interestingly, the link between broad sense macroevolution and Simpson’s tempo and mode of evolution is that Simpson’s statement assigns no hierarchical level to these patterns. The pervasiveness of the use of broad sense macroevolution then is because this usage is explicitly because it is not the actual definition of macroevolution. Effectively, the metaphoric use removes any distinction between the terms and makes the terms useless.

An important theoretical construct in paleobiology is that of emergent traits or properties which are defined as traits that are not reducible to a lower level [22, 28, 42, 76]. An emergent property is effectively a trait that is the product of multiple traits expressed at one or more hierarchical levels in concert and is not reducible [28, 76]. It is extremely important to recognize a fundamental hierarchy in biology in order to allow for emergent traits [76]. In paleobiology, an emergent property is normally one ascribed to the species or genus level as opposed to the organism level and is the root of “species selection” [22, 28, 42, 76]. Range size is considered an emergent property that has continually been demonstrated to play a crucial role in extinction selectivity with species with larger geographic ranges having lower extinction rates than species with smaller geographic ranges [25, 26, 31, 50, 64]. Range size is emergent because no one property of a single organism can explain this trait and instead it is a combination of multiple properties which determines global range size.

Survival can be considered the fundamental measure of fitness or evolution success [16, 52] because ultimately a long-term successful lineage is not one that speciated greatly but one that never went extinct [52]. Because during periods of background extinction patterns of

taxonomic extinction are expected to be non-random with respect to biology [26]. It should  
76 be possible to effectively measure the relative fitness of various ecological traits [34, 35].

Survivorship analysis has a long history in paleobiology [9, 20, 34, 36, 61, 67–69, 74]. The  
78 hazard/extinction rate from a survivorship curve is a statement of the rate at which organisms  
go extinction following origination. Hazard rates are comparable to the diversification rates  
80 estimated via phylogenetic comparative methods [18, 45, 47–49, 56], however hazard rates  
are an estimate of extinction rate which is considered difficult if not impossible to estimate  
82 in this fashion [55].

Here I propose to study the individual and combined effects of organismal traits related to  
84 emergent range size on extinction and, by extension, fitness.

## **Cosmopolitan and endemic mammal dynamics of Ceno- 86 zoic mammals**

*Questions:* How do patterns of community connectedness change over time, specifically in  
88 Cenozoic terrestrial mammals? Do these patterns differ between continents? Are these  
patterns different between groups ecologically different organisms? Are these patterns related  
90 to changes in global temperature?

*Background and Predictions:* During the Cenozoic, there was a global shift from predominately  
92 closed, forested habitats to more open, savanna habitats. This pattern leads to the expectation  
that there would be a relative decrease in arboreal taxa as well as a relative increase in  
94 terrestrial taxa. Additionally, an increase in the relative endemism of arboreal taxa over time  
and an decrease in the relative endemism of terrestrial taxa might have accompanied this.  
96 The timing of this shift was different between continents [70, 71], meaning that the patterns  
reflecting this environmental shift may be globally non-uniform and changes in community  
98 structures could reflect regional scale changes instead of global scale changes. Shifts in  
distribution of taxa according to locomotor category were not necessarily accompanied by  
100 shifts in distribution of dietary categories.

The majority of previous research has focused on mammalian faunal dynamics has focused  
102 on the North American fossil record [3–6, 8, 10, 17, 23, 24, 68, 69]. The major focuses have  
been on the effect of climate change on diversity and distributions between different higher  
104 taxonomic levels. The long term effects of climate change on North American mammalian  
diversity dynamics and community connectedness and similarly remains unresolved and  
106 controversial [5, 8, 10, 17]. The basic predictions are that over the Cenozoic there would be a  
relative increase in endemism in arboreal taxa versus a relative decrease in ground dwelling  
108 endemism. Because of the vast amount of prior work on North American mammalian faunal  
dynamics, this forms the basis for the global predictions made above. The North American  
110 record then inadvertently becomes the baseline comparison for regional differences.

In comparison to North America, the European mammalian fossil is less studied. Importantly, research has focused primarily on faunal dynamics in the Neogene [32, 33, 39, 57–59]. One of the major findings is that, during the Neogene, there was very little shift in relative trophic level abundance [33] while the patterns in dietary shifts were mostly driven by abundance and cosmopolitan herbivores [32]. Because of this, the major predictions for the European record is that occupancy will increase for herbivorous taxa, while increasing or remaining identical in carnivores, and remaining relatively constant for omnivores.

The South American mammalian faunal record appears to reflect two distinct biotic provinces between the North and the South [19, 43, 44, 53]. Because of this, I predict the South American record to have a very different pattern of biogeographic connectedness than either North America or Europe. Namely, the expectation would be a high or progressively increasing degree of endemism along with low connectivity. Also, an expected increase in land-dwelling herbivores relative to arboreal (at least in the south).

A global trend during the Cenozoic was the shift from a “hot house” environment with no polar ice caps to an “ice house” environment with polar ice caps [77, 78]. NEED SOME LITERATURE ON RANGE SIZE SHIFTS AND CLIMATE SEE MACRO NOTES. This transition is predicted to cause major shifts in biomes, causing reorganization of communities CITATIONS. For global mammalian community connectedness and trophic structure, it is predicted that while the environment might be shifting, lineages may adapt in place and overall structure will remain rather constant through time [33].

*Proposed research:* Using methods first proposed by Sidor et al. [66] and Vilhena et al. [75], I propose to construct bipartite biogeographic networks between taxa and localities. A link between a taxon and a locality is formed when that is present at that locality. Here taxa are defined as species and localities are defined as formations. Biogeographic networks will be constructed for every 2 million year bin of the Cenozoic. This bin width is chosen to have minimum 2 formations to be present in the same bin. Additionally, networks will be constructed for each dietary category and each locomotor category.

Network complexity and connectedness is measured using four previously used summary statistics [66]: average number of endemics, average occupancy, biogeographic connectedness, and code length. Biogeographic connectedness is effectively the relationship between the number of endemic taxa and the average occupancy. Code length is a measurement of the complexity and clustering of the graph [62, 63]. A low code length indicates that a graph is compressible into a greater number of subunits without information loss than a graph with a high code length. This means that a low code length indicates lower overall locality similarity than a high code length.

In order to compare whether patterns observed on different continents are similar or different, as well as compare patterns between different categories of ecological traits, HOW DO I DO THIS?

Taxonomic occurrence data will be collected through a combination of the Paleobiological

Database (PBDB; <http://fossilworks.org>), Neogene Old World Database (NOW; <http://www.helsinki.fi/science/now/>), and museum collections. North American fossil mammal data is very well represented and vetted in the PBDB because of the extensive work by John Alroy [3–5]. European fossil mammal data is also well represented between the PBDB and NOW. South American fossil mammal data is available through the PBDB, but is not particularly well vetted and poorly covered. Because of this, South American fossil mammal data will be gathered via various museums such as the Field Museum of Natural History and the American Museum of Natural History.

## **Fitness and extinction risk related to ecology in Cenozoic mammals**

*Questions:* How do ecological characters affect survivorship in mammals? Is any single trait the best predictor of mammalian survivorship, or do multiple traits together predict survivorship better?

*Background and Predictions:* In mammals, three of the arguably most important ecological traits are dietary category, locomotor category, and body size CITATIONS. Additionally, each of these traits are important in determining the emergent species level property of range size. As discussed above, as the Cenozoic progressed on all Continents there was a shift from closed habitat to more open habitat. In the intermediate, one would expect some degree of patchiness in the landscape. Expectedly, there would be a increase followed by a decrease in both speciation rate and extinction rate, with the peak being during the intermediate period.

According to Price et al. [54], herbivores and carnivores should have a greater diversification rate than omnivores. This analysis was global in scope, and purely extant taxa in a comparative phylogenetic context. Diversification rate can increase via either an increase in origination relative to extinction or a decrease in extinction relative to origination. Which of these two processes is occurring is impossible to determine from a phylogeny of only extant organisms [55] which means that only via the analysis of the fossil record is it possible to estimate which process is more likely.

Depending on the continent, body size has been demonstrated to be related to extinction rate [39, 40, 72]. By expanding to include a third continent, South America, I hope to elucidate how differences in taxonomic diversity at a continental level might affect body size mediated extinction rate. Additionally, I will be using alternative methods to better understand the dynamics governing trait based extinction probability.

*Proposed research:* To investigate the effect of ecological traits and climate change on survivorship, I plan to compare different models of survival in order to best understand what are the most important factors in estimating survival probability.

Survivorship analysis is the analysis of time-till-event data. In a paleontological context this

is the time from origination (first appearance date; FAD) till extinction (last appearance date; LAD). Dietary category, locomotor category, and body size will be modeled as time-independent covariates of survival. The climate proxy  $\delta O^{18}$  oxygen curve from Zachos et al. [78] will be modeled as an ancillary time-dependent covariate. Also, constant versus accelerating, decelerating, or time variant extinction rate will be estimated using different fundamental hazard models by comparing the fit various probability distributions to survival.

While many analyses of survivorship are done using generic data [25, 39, 72], there are potential biases in accurately modeling specific level process from generic level data [60, 61, 65, 67, 74]. There are important concerns regarding anagenesis, hierarchical selection, and extant taxa or taxa that did not go extinct in the time frame of interest [60, 61, 67, 74]. Interestingly, the effect of incomplete sampling on estimation of survivorship curves appears rather minimal and uniform [65]. The problems involving extant taxa and taxa that did not go extinct have mostly been dealt with following advances of how to model right-censored data CITATION.

In order to asses potential specific versus generic effects I will estimate differences in estimated survival between specific and generic level survivorship models. Using an approach based on previous work to estimate specific level survival from generic level survival curves [20], or a variant there of, I will measure the deviance between extinction rate estimated from the specific survivorship and the specific level extinction rates estimated from the analysis of the generic survivorship data.

In addition to the above study of mammalian survivorship, I also propose a simulation study to analyze effect of varying sampling probability and/or anagenesis rate on estimating various models of survivorship using `paleotree` [7]. Principally, I am interested in the effect of these paleontological realities on estimation of the hazard function of the survivorship data and in particular departures from a constant, or exponential, hazard function. Alternatives are, for example, models of accelerating or decelerating extinction rate. I intend to revise the previously analyzed effect of sampling on estimation survivorship in this new context [65].

The data necessary to complete the empirical aspectes of this study will be the same as described above for analysis of dynamics of mammalian biogeographic connectedness.

## Extinction risk related to traits affecting habitat selection in Permian brachiopods

*Questions:* How do traits directly related habitat selection and range size relate to extinction risk? Are certain traits more explanatory of extinction risk than others? Does changing environmental and substrate availability affect trait-based extinction risk?

*Background and Predictions:* In brachiopods, three extremely important ecological traits involved in determining possible range size are affixing strategy, substrate preference, and habitat preference. While larval biology is also considered extremely important for determining

range size in marine invertebrates [29, 30]. However, larval ecology does not preserve in brachiopods and thus cannot be used to model survivorship [30].

Affixing strat: pedicle, reclining, cementing. Pedicles are associated with shallow, on-shore environments while reclining on associated with deep, off-shore environments. Note, global Permian pattern is rather heterogeneous in terms of relative abundance between the three affixing strategies CLAPHAM. According to Alexander [1], endemic unattached/reclining taxa have longer durations than all other affixing strategies of other endemic taxa. On the other hand, cosmopolitan pedunculate and cementing taxa have longer durations than unattached/reclining taxa. In terms of regional scales, it will be necessary to correct for relative abundance to actually measure survivability/fitness.

Substrate: carbonate, clastic, mixed. Overall decline in carbonates relative to clastics, decreased fitness for carbonate lovers relative to clastic lovers. Best to be mixed? Brachiopods are mixed slash switchers CARL AND MELANIE'S UNPUBLISHED WORK.

Habitat: on-shore vs off-shore / epicontinental vs oceanic. How possible is this? Very difficult to resolve accurately. Progressive loss of epicontinental seas during the Pharenozoic which would predict over all lower fitness in taxa which occur solely in epicontinental environments. On-shore off-shore dynamics of Sepkoski and Raup.

During the Permian there was a shift from an “ice house” world to a “hot house world” CITATIONS which could be expected to have some major effects on brachiopod survivorship. In particular, taxa in modern Australia would be of particular interest because of the proximity of Australia to the south pole during the Permian and the repeated glacial activity in the region CITATIONS. According to Olszewski and Erwin [51], sea-level and climate change do not wholly explain the ecological dynamics experienced by brachiopods in the Permian of Texas. The prediction then is that the best model of brachiopod survivorship will have to have some biotic component such as affixing strategy or substrate preference. If climate or environmental information, such as habitat preference, is a predictor in the best model of survivorship is less clear cut and necessary to determine empirically.

*Proposed research:* I propose to use a survival analysis approach similar to the one previously described to estimate the differential survivorship of brachiopods leading up to the Permo-Triassic boundary. I restrict the analysis to Australiasia because it represents a relatively continually sampled and well worked area that preserves the majority of the entire Permian CITATIONS. In this case, the time-independent covariates are substrate preference, affixing strategy, and habitat preference. Climate will be modeled as either an ancillary heavyside function or a time-varying covariate depending on the quality of the Permian isotope record. Additionally, as in the mammalian survivorship analysis described above, the time dependence of brachiopod extinction rates will be estimated using different fundamental hazard models by comparing the fit various probability distributions to survival.

Permian brachiopod occurrence information is available via the PBDB and is primarily based on the work of Matthew Clapham [11–15].



## Significance of research

262 Fundamental questions in evolution and paleobiology.

What do certain organisms go extinct while others do not?

264 How does community similarity change over time?

What role does ecology play in survival?

266 Is climate an important factor in survival?

# Timeline

268 Spring/Summer 2014

- Evolution Meeting: preliminary brachiopod survival results

270 • South American fossil mammal data from Field Museum of Natural History collections

Fall 2014/Winter 2015

272 • GSA: survivorship simulation for anagenesis and sampling

Spring/Summer 2015

274 • Evolution Meeting: mammalian survivorship analysis or biogeographic connectedness for North America and Europe

276 • South American fossil mammal data from American Museum of Natural History collections

278 Fall 2015/Winter 2016

280 • SVP or GSA: mammalian survivorship analysis or biogeographic connectedness for North America and Europe

Spring/Summer 2016

282 • Evolution Meeting: brachiopod survival analysis results

Fall 2016/Winter 2017

284 • SVP or GSA: mammalian survivorship analysis or biogeographic connectedness for North America, Europe and South America

286 Spring/Summer 2017

- Evolution Meeting

288 • **Defend**

## References

- [1] R. R. Alexander. Generic longevity of articulate brachiopods in relation to the mode of stabilization on the substrate. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 21: 209–226, 1977.
- [2] W. D. Allmon. Taxic evolutionary paleoecology and the ecological context of macroevolutionary change. *Evolutionary Ecology*, 8(2):95–112, Mar. 1994. ISSN 0269-7653. doi: 10.1007/BF01238244.
- [3] J. Alroy. Constant extinction, constrained diversification, and uncoordinated stasis in North American mammals. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 127: 285–311, 1996.
- [4] J. Alroy. Cope’s rule and the dynamics of body mass evolution in North American fossil mammals. *Science*, 280:731–734, 1998. doi: 10.1126/science.280.5364.731.
- [5] J. Alroy, P. L. Koch, and J. C. Zachos. Global climate change and North American mammalian evolution. *Paleobiology*, 26(1981):259–288, 2000.
- [6] C. Badgley and J. a. Finarelli. Diversity dynamics of mammals in relation to tectonic and climatic history: comparison of three Neogene records from North America. *Paleobiology*, 39(3):373–399, Apr. 2013. ISSN 0094-8373. doi: 10.1666/12024.
- [7] D. W. Bapst. paleotree: an R package for paleontological and phylogenetic analyses of evolution. *Methods in Ecology and Evolution*, 3:803–807, 2012. doi: 10.1111/j.2041-210X.2012.00223.x. URL <http://doi.wiley.com/10.1111/j.2041-210X.2012.00223.x>.
- [8] A. D. Barnosky. Distinguishing the effects of the Red Queen Court Jester on Micene Mammal Evolution in the Northern Rocky Mountains. *Journal of Vertebrate Paleontology*, 21(1):172–185, 2001.
- [9] T. K. Baumiller. Survivorship analysis of Paleozoic Crinoidea: effect of filter morphology on evolutionary rates. *Paleobiology*, 19(3):304–321, 1993.
- [10] J. L. Blois and E. A. Hadly. Mammalian Response to Cenozoic Climatic Change. *Annual Review of Earth and Planetary Sciences*, 37(1):181–208, May 2009. ISSN 0084-6597. doi: 10.1146/annurev.earth.031208.100055.
- [11] M. E. Clapham and D. J. Bottjer. Permian marine paleoecology and its implications for large-scale decoupling of brachiopod and bivalve abundance and diversity during the Lopingian (Late Permian). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 249 (3-4):283–301, June 2007. ISSN 00310182. doi: 10.1016/j.palaeo.2007.02.003. URL <http://linkinghub.elsevier.com/retrieve/pii/S0031018207000600>.
- [12] M. E. Clapham and D. J. Bottjer. Prolonged Permian Triassic ecological crisis recorded by molluscan dominance in Late Permian offshore assemblages. *Proceedings of the National*

*Academy of Sciences of the United States of America*, 104(32):12971–5, Aug. 2007. ISSN 0027-8424. doi: 10.1073/pnas.0705280104. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1941817&tool=pmcentrez&rendertype=abstract>.

- [13] M. E. Clapham and N. P. James. Paleoeecology Of Early-Middle Permian Marine Communities In Eastern Australia: Response To Global Climate Change In the Aftermath Of the Late Paleozoic Ice Age. *Palaaios*, 23(11):738–750, Nov. 2008. ISSN 0883-1351. doi: 10.2110/palo.2008.p08-022r. URL <http://palaaios.sepmonline.org/cgi/doi/10.2110/palo.2008.p08-022r>.
- [14] M. E. Clapham and N. P. James. Regional-scale marine faunal change in Eastern Australia during Permian climate fluctuations and its relationship to local community restructuring. *Palaaios*, 27:627–635, Oct. 2012. ISSN 0883-1351. doi: 10.2110/palo.2012.p12-003r. URL <http://palaaios.sepmonline.org/cgi/doi/10.2110/palo.2012.p12-003r>.
- [15] M. E. Clapham, D. J. Bottjer, C. M. Powers, N. Bonuso, M. L. Fraiser, P. J. Marengo, S. Q. Dornbos, and S. B. Pruss. Assessing the Ecological Dominance of Phanerozoic Marine Invertebrates. *Palaaios*, 21(5):431–441, Oct. 2006. ISSN 0883-1351. doi: 10.2110/palo.2005.P05-017R. URL <http://palaaios.sepmonline.org/cgi/doi/10.2110/palo.2005.P05-017R>.
- [16] W. S. Cooper. Expected time to extinction and the concept of fundamental fitness. *Journal of Theoretical Biology*, 107:603–629, 1984.
- [17] B. Figueirido, C. M. Janis, J. a. Pérez-Claros, M. De Renzi, and P. Palmqvist. Cenozoic climate change influences mammalian evolutionary dynamics. *Proceedings of the National Academy of Sciences*, 109(3):722–727, Jan. 2012. ISSN 1091-6490. doi: 10.1073/pnas.1110246108.
- [18] R. G. Fitzjohn. Quantitative Traits and Diversification. *Systematic Biology*, pages 1–15, 2010. URL [papers2://publication/doi/10.1093/sysbio/syq053](http://publications.sciencemag.org/doi/10.1093/sysbio/syq053).
- [19] J. J. Flynn and a. R. Wyss. Recent advances in South American mammalian paleontology. *Trends in ecology & evolution*, 13(11):449–54, Nov. 1998. ISSN 0169-5347. URL <http://www.ncbi.nlm.nih.gov/pubmed/21238387>.
- [20] M. Foote. Survivorship analysis of Cambrian and Ordovician Trilobites. *Paleobiology*, 14(3):258–271, 1988.
- [21] M. Foote and A. I. Miller. *Principles of Paleontology*. Freeman, New York, third edition, 2007.
- [22] T. A. Grantham. Hierarchical approaches to macroevolution: recent work on species selection and the effect hypothesis. *Annual Review of Ecology and Systematics*, 26:301–321, 1995.
- [23] G. Gunnell, M. Morgan, M. C. Mass, and P. D. Gingerich. Comparative paleoecology of Paleogene and Neogene mammalian faunas: trophic structure and composition.

*Palaeogeography, Palaeoclimatology, Palaeoecology*, 115:265–286, 1995. URL <http://www.sciencedirect.com/science/article/pii/0031018294001150>.

- [24] E. A. Hadly and B. A. Maurer. Spatial and temporal patterns of species diversity in montane mammal communities of western North America. *Evolutionary Ecology Research*, 3:477–486, 2001.

- [25] P. G. Harnik, C. Simpson, and J. L. Payne. Long-term differences in extinction risk among the seven forms of rarity. *Proceedings of the Royal Society B: Biological Sciences*, (October), Oct. 2013. ISSN 0962-8452. doi: 10.1098/rspb.2012.1902. URL <http://rspb.royalsocietypublishing.org/cgi/doi/10.1098/rspb.2012.1902>.

- [26] D. Jablonski. Background and mass extinctions: the alternation of macroevolutionary regimes. *Science*, 231(4734):129–133, 1986.

- [27] D. Jablonski. Mass extinctions and macroevolution. *Paleobiology*, 31(sp5):192–210, 2005. URL [papers2://publication/uuid/F296080D-6609-4C57-9514-C74E5E726340](http://papers2://publication/uuid/F296080D-6609-4C57-9514-C74E5E726340).

- [28] D. Jablonski. Species Selection: Theory and Data. *Annual Review of Ecology, Evolution, and Systematics*, 39(1):501–524, Dec. 2008. ISSN 1543-592X. doi: 10.1146/annurev.ecolsys.39.110707.173510. URL <http://www.annualreviews.org/doi/abs/10.1146/annurev.ecolsys.39.110707.173510>.

- [29] D. Jablonski and G. Hunt. Larval ecology, geographic range, and species survivorship in Cretaceous mollusks: organismic versus species-level explanations. *The American naturalist*, 168(4):556–64, Oct. 2006. ISSN 1537-5323. doi: 10.1086/507994. URL <http://www.ncbi.nlm.nih.gov/pubmed/17004227>.

- [30] D. Jablonski and R. a. Lutz. Larval Ecology of Marine Benthic Invertebrates: Paleobiological Implications. *Biological Reviews*, 58(1):21–89, Feb. 1983. ISSN 1464-7931. doi: 10.1111/j.1469-185X.1983.tb00380.x. URL <http://doi.wiley.com/10.1111/j.1469-185X.1983.tb00380.x>.

- [31] D. Jablonski and K. Roy. Geographical range and speciation in fossil and living molluscs. *Proceedings. Biological sciences / The Royal Society*, 270(1513):401–6, Feb. 2003. ISSN 0962-8452. doi: 10.1098/rspb.2002.2243. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1691247&tool=pmcentrez&rendertype=abstract>.

- [32] J. Jernvall and M. Fortelius. Common mammals drive the evolutionary increase of hypsodonty in the Neogene. *Nature*, 417(6888):538–40, May 2002. ISSN 0028-0836. doi: 10.1038/417538a.

- [33] J. Jernvall and M. Fortelius. Maintenance of trophic structure in fossil mammal communities: site occupancy and taxon resilience. *American Naturalist*, 164(5):614–624, Nov. 2004. ISSN 1537-5323. doi: 10.1086/424967.

- [34] J. Kitchell. Biological selectivity of extinction. In E. G. Kaufman and O. H. Walliser,

editors, *Extinction Events in Earth History*, pages 31–43. Springer-Verlag, Berlin, 1990. URL <http://link.springer.com/chapter/10.1007/BFb0011132>.

[35] J. A. Kitchell. Evolutionary paleoecology: recent contributions to evolutionary theory. *Paleobiology*, 11(1):91–104, 1985.

[36] J. A. Kitchell. The temporal distribution of evolutionary and migrational events in pelagic systems: episodic or continuous. *Paleoceanography*, 2(5):473–487, 1987.

[37] J. A. Kitchell and A. Hoffman. Rates of species-level origination and extinction: functions of age, diversity, and history. *Acta Palaeontologica Polonica*, 36(1):39–67, 1991.

[38] J. A. Kitchell, D. L. Clark, and A. M. Gombos. Biological selectivity of extinction: a link between background and mass extinction. *Palaaios*, 1(5):504–511, 1986.

[39] L. H. Liow, M. Fortelius, E. Bingham, K. Lintulaakso, H. Mannila, L. Flynn, and N. C. Stenseth. Higher origination and extinction rates in larger mammals. *Proceedings of the National Academy of Sciences*, 105(16):6097, 2008. URL [papers2://publication/uuid/5A93DDA3-204F-4D9C-AFAC-A15722A18C61](http://papers2://publication/uuid/5A93DDA3-204F-4D9C-AFAC-A15722A18C61).

[40] L. H. Liow, M. Fortelius, K. Lintulaakso, H. Mannila, and N. C. Stenseth. Lower Extinction Risk in SleeporHide Mammals. *The American Naturalist*, 173(2):264–272, 2009. URL [papers2://publication/doi/10.1086/595756](http://papers2://publication/doi/10.1086/595756).

[41] L. H. Liow, L. Van Valen, and N. C. Stenseth. Red Queen: from populations to taxa and communities. *Trends in ecology & evolution*, 26(7):349–58, July 2011. ISSN 0169-5347. doi: 10.1016/j.tree.2011.03.016. URL <http://www.ncbi.nlm.nih.gov/pubmed/21511358>.

[42] E. A. Lloyd and S. J. Gould. Species selection on variability. *Proceedings of the National Academy of Sciences*, 90:595–599, 1993.

[43] B. J. Macfadden. Origin and evolution of the grazing guild in New World terrestrial mammals. *Trends in ecology & evolution*, 12(5):182–187, 1997.

[44] B. J. Macfadden. Extinct mammalian biodiversity of the ancient New World tropics. *Trends in ecology & evolution*, 21(3):157–65, Mar. 2006. ISSN 0169-5347. doi: 10.1016/j.tree.2005.12.003. URL <http://www.ncbi.nlm.nih.gov/pubmed/16701492>.

[45] W. P. Maddison, P. E. Midford, and S. P. Otto. Estimating a binary character’s effect on speciation and extinction. *Systematic Biology*, 56(5):701, 2007. URL [papers2://publication/uuid/AF092848-7CDC-4A52-A4A4-847F1FF0F020](http://papers2://publication/uuid/AF092848-7CDC-4A52-A4A4-847F1FF0F020).

[46] G. A. Mark and K. W. Flessa. A test for evolutionary equilibria: Phanerozoic brachiopods and Cenozoic mammals. *Paleobiology*, 3(1):17–22, 1977.

[47] S. Nee. Inferring speciation rates from phylogenies. *Evolution*, 55(4):661–8, Apr. 2001. ISSN 0014-3820. URL <http://www.ncbi.nlm.nih.gov/pubmed/11392383>.

- [48] S. Nee, a. O. Mooers, and P. H. Harvey. Tempo and mode of evolution revealed from molecular phylogenies. *Proceedings of the National Academy of Sciences of the United States of America*, 89(17):8322–6, Sept. 1992. ISSN 0027-8424. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2789996&tool=pmcentrez&rendertype=abstract>.
- [49] S. Nee, E. C. Holmes, R. M. May, and P. H. Harvey. Extinction rates can be estimated from molecular phylogenies. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 344(1307):77–82, Apr. 1994. ISSN 0962-8436. doi: 10.1098/rstb.1994.0054. URL <http://www.ncbi.nlm.nih.gov/pubmed/8878259>.
- [50] S. Nürnberg and M. Aberhan. Habitat breadth and geographic range predict diversity dynamics in marine Mesozoic bivalves. *Paleobiology*, 39(3):360–372, Apr. 2013. ISSN 0094-8373. doi: 10.1666/12047. URL <http://www.bioone.org/doi/abs/10.1666/12047>.
- [51] T. Olszewski and D. Erwin. Dynamic response of Permian brachiopod communities to long-term environmental change. *Nature*, 428(April):2–5, 2004. doi: 10.1038/nature02471.1. URL <http://www.nature.com/nature/journal/v428/n6984/abs/nature02464.html>.
- [52] M. E. Palmer and M. W. Feldman. Survivability is more fundamental than evolvability. *PloS one*, 7(6):e38025, Jan. 2012. ISSN 1932-6203. doi: 10.1371/journal.pone.0038025. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3377627&tool=pmcentrez&rendertype=abstract>.
- [53] B. Patterson and R. Pascual. The fossil mammal fauna of South America. *The Quarterly review of biology*, 43(4):409–451, 1968.
- [54] S. a. Price, S. S. B. Hopkins, K. K. Smith, and V. L. Roth. Tempo of trophic evolution and its impact on mammalian diversification. *Proceedings of the National Academy of Sciences of the United States of America*, 109(18):7008–12, May 2012. ISSN 1091-6490. doi: 10.1073/pnas.1117133109. URL <http://www.ncbi.nlm.nih.gov/pubmed/22509033>.
- [55] D. L. Rabosky. Extinction rates should not be estimated from molecular phylogenies. *Evolution*, 64(6):1816–24, June 2010. ISSN 1558-5646. doi: 10.1111/j.1558-5646.2009.00926.x. URL <http://www.ncbi.nlm.nih.gov/pubmed/20030708>.
- [56] D. L. Rabosky, F. Santini, J. Eastman, S. a. Smith, B. Sidlauskas, J. Chang, and M. E. Alfaro. Rates of speciation and morphological evolution are correlated across the largest vertebrate radiation. *Nature Communications*, 4:1–8, June 2013. ISSN 2041-1723. doi: 10.1038/ncomms2958. URL <http://www.nature.com/doifinder/10.1038/ncomms2958>.
- [57] P. Raia, P. Piras, and T. Kotsakis. Turnover pulse or Red Queen? Evidence from the large mammal communities during the Plio-Pleistocene of Italy. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 221(3-4):293–312, June 2005. ISSN 00310182. doi: 10.1016/j.palaeo.2005.02.014. URL <http://linkinghub.elsevier.com/retrieve/pii/S0031018205001033>.

- [58] P. Raia, C. Meloro, A. Loy, and C. Barbera. Species occupancy and its course in the past: macroecological patterns in extinct communities. *Evolutionary Ecology Research*, 8:181–194, 2006.
- [59] P. Raia, F. Carotenuto, J. T. Eronen, and M. Fortelius. Longer in the tooth, shorter in the record? The evolutionary correlates of hypsodonty in Neogene ruminants. *Proceedings. Biological sciences / The Royal Society*, 278(1724):3474–81, Dec. 2011. ISSN 1471-2954. doi: 10.1098/rspb.2011.0273. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3189366&tool=pmcentrez&rendertype=abstract>.
- [60] D. M. Raup. Taxonomic survivorship curves and Van Valen’s Law. *Paleobiology*, 1(1):82–96, Jan. 1975. ISSN 0036-8075. doi: 10.1126/science.49.1254.50. URL <http://www.ncbi.nlm.nih.gov/pubmed/17777225>.
- [61] D. M. Raup. A kill curve for Phanerozoic marine species. *Paleobiology*, 17(1):37–48, 1991.
- [62] M. Rosvall and C. T. Bergstrom. Maps of random walks on complex networks reveal community structure. *Proceedings of the National Academy of Sciences*, 105(4):1118–23, Jan. 2008. ISSN 1091-6490. doi: 10.1073/pnas.0706851105. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2234100&tool=pmcentrez&rendertype=abstract>.
- [63] M. Rosvall, D. Axelsson, and C. T. Bergstrom. The map equation. *The European Physical Journal Special Topics*, 178(1):13–23, Apr. 2010. ISSN 1951-6355. doi: 10.1140/epjst/e2010-01179-1. URL <http://www.springerlink.com/index/10.1140/epjst/e2010-01179-1>.
- [64] K. Roy, G. Hunt, D. Jablonski, A. Z. Krug, and J. W. Valentine. A macroevolutionary perspective on species range limits. *Proceedings. Biological sciences / The Royal Society*, 276(1661):1485–93, Apr. 2009. ISSN 0962-8452. doi: 10.1098/rspb.2008.1232. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2677224&tool=pmcentrez&rendertype=abstract>.
- [65] J. J. Sepkoski. Stratigraphic biases in the analysis of taxonomic survivorship. *Paleobiology*, 1(4):343–355, 1975.
- [66] C. a. Sidor, D. a. Vilhena, K. D. Angielczyk, A. K. Huttenlocker, S. J. Nesbitt, B. R. Peacock, J. S. Steyer, R. M. H. Smith, and L. a. Tsuji. Provincialization of terrestrial faunas following the end-Permian mass extinction. *Proceedings of the National Academy of Sciences*, 110(20):8129–33, May 2013. ISSN 1091-6490. doi: 10.1073/pnas.1302323110.
- [67] C. Simpson. *Levels of selection and large-scale morphological trends*. PhD thesis, University of Chicago, 2006.
- [68] G. G. Simpson. *Tempo and Mode in Evolution*. Columbia University Press, New York, 1944.



- [69] G. G. Simpson. *The Major Features of Evolution*. Columbia University Press, New York, 1953.
- [70] C. a. E. Strömberg. Decoupled taxonomic radiation and ecological expansion of open-habitat grasses in the Cenozoic of North America. *Proceedings of the National Academy of Sciences of the United States of America*, 102(34):11980–4, Aug. 2005. ISSN 0027-8424. doi: 10.1073/pnas.0505700102. URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1189350&tool=pmcentrez&rendertype=abstract>.
- [71] C. a. E. Strömberg, R. E. Dunn, R. H. Madden, M. J. Kohn, and A. a. Carlini. Decoupling the spread of grasslands from the evolution of grazer-type herbivores in South America. *Nature communications*, 4:1478, Jan. 2013. ISSN 2041-1723. doi: 10.1038/ncomms2508. URL <http://www.ncbi.nlm.nih.gov/pubmed/23403579>.
- [72] S. Tomiya. Body Size and Extinction Risk in Terrestrial Mammals Above the Species Level. *The American Naturalist*, pages E000–E000, Sept. 2013. ISSN 00030147. doi: 10.1086/673489. URL <http://www.jstor.org/stable/info/10.1086/673489>.
- [73] L. Van Valen. A new evolutionary law. *Evolutionary Theory*, 1:1–30, 1973. URL <http://ci.nii.ac.jp/naid/10011264287/>.
- [74] L. Van Valen. Taxonomic survivorship curves. *Evolutionary Theory*, 4:129–142, 1979.
- [75] D. a. Vilhena, E. B. Harris, C. T. Bergstrom, M. E. Maliska, P. D. Ward, C. a. Sidor, C. a. E. Strömberg, and G. P. Wilson. Bivalve network reveals latitudinal selectivity gradient at the end-Cretaceous mass extinction. *Scientific reports*, 3:1790, May 2013. ISSN 2045-2322. doi: 10.1038/srep01790.
- [76] E. S. Vrba and N. Eldredge. Individuals, hierarchies and processes: towards a more complete evolutionary theory. *Paleobiology*, 10(2):146–171, 1984.
- [77] J. C. Zachos, M. Pagani, L. Sloan, E. Thomas, and K. Billups. Trends, rhythms, and aberrations in global climate 65 Ma to present. *Science*, 292:686–693, 2001. doi: 10.1126/science.1059412. URL <http://www.sciencemag.org/content/292/5517/686.short>.
- [78] J. C. Zachos, G. R. Dickens, and R. E. Zeebe. An early Cenozoic perspective on greenhouse warming and carbon-cycle dynamics. *Nature*, 451(7176):279–283, Jan. 2008. ISSN 1476-4687. doi: 10.1038/nature06588.