

# Evolutionary paleoecology and the biology of extinction

Peter D Smits

Committee on Evolutionary Biology, University of Chicago

January 7, 2014

Introduction and theory

Brachiopods, environmental preference, and extinction

Ecology and survival in Cenozoic mammals

Community connectedness in Cenozoic mammals

## Introduction and theory

Brachiopods, environmental preference, and extinction

Ecology and survival in Cenozoic mammals

Community connectedness in Cenozoic mammals

## Questions

- ▶ Why do certain taxa go extinct while others do not?
- ▶ How is emergence “formed?” When should it be invoked?
- ▶ Is extinction risk taxon–age independent?
- ▶ When should we expect global, regional, or local processes to dominate?


# Evolutionary paleoecology

*... the consequences of distinct ecological factors on differential rate dynamics, particularly rates of faunal turnover and diversification.*

(Kitchell 1985 *Paleobiology*)

environmental interactions → macroevolution

# Emergent properties




(Grantham 1995 *Ann. Rev. Ecol. Syst.*)

## Species level

Trait that cannot be reduced to organismal level

Product of one or more traits/factors


# Range size



(Jablonski 1987 *Science*)

# Probability of survival

Theoretical  $S(t)$ :




## Survival function

$$S(t) = P(T > t)$$

- ▶  $T$ : survival time ( $\geq 0$ )
- ▶  $t$ : specified time

$\hat{S}(t)$  in practice:




(Kleinbaum and Klein 2012)


# Instantaneous potential of failure (extinction)

Hazard function  $\equiv$  conditional failure rate

$$h(t) = \lim_{\Delta t \rightarrow 0} \frac{P(t \leq T < t + \Delta t | T \geq t)}{\Delta t}$$



# Van Valen's observation of survival



(Liow et al. 2011 *TREE*)

# Law of Constant Extinction

## Definition

Extinction risk in a given adaptive zone is taxon–age independent.

(Van Valen 1973 *Evol. Theory*)

translation: hazard is constant with respect to time  
(**exponential survival**)

$$h(t) = \lambda \iff S(t) = \exp^{-\lambda t}$$

# Brachiopods and mammals: a comparison

## brachiopods

- ▶ Permian (~ 47 My)
- ▶ marine
- ▶ Australasia
- ▶ global warming
- ▶ sessile

## mammals

- ▶ Cenozoic (~ 65 My)
- ▶ terrestrial
- ▶ North America, Europe, South America
- ▶ global cooling
- ▶ motile

# Series of related questions

- ▶ generic level survival in brachiopods
  - ▶ ecological traits re. environmental pref. (emergence)
  - ▶ survival distribution
- ▶ specific level survival in mammals
  - ▶ ecological traits re. range size (emergence)
  - ▶ generic versus specific survival
  - ▶ anagenesis/species:genus simulation
  - ▶ survival distribution
- ▶ community connectedness in mammals
  - ▶ global versus regional versus local scale processes
  - ▶ ecological traits (trophic/locomotion)

Introduction and theory

Brachiopods, environmental preference, and extinction


Ecology and survival in Cenozoic mammals

Community connectedness in Cenozoic mammals


# Traits relating to environment and range size

- ▶ substrate affinity
  - ▶ physical, chemical
  - ▶ availability
- ▶ habitat preference
  - ▶ energetics
  - ▶ availability
- ▶ affixing strategy
  - ▶ energetics
  - ▶ optimality

# Substrate affinity



(Miller and Connolly 2001 *Paleobio.*)



(Foote 2006 *Paleobio.*)

- ▶ carbonates, clastics, mixed
- ▶ lithology/deposition environment
- ▶ Phanerozoic decrease in carbonates:clastics

# Habitat preference

- ▶ on-shore, off-shore, none
- ▶ sea-level and energetics
- ▶ Phanerozoic decrease in on-shore:off-shore

# Affixing strategy

- ▶ pedunculate, reclining, cementing
- ▶ pedunculate:on-shore, reclining:off-shore
- ▶ environmental energetics

# Assigning substrate and habitat

## Probability of assignment

$$P(H_1|E) = \frac{P(E|H_1)P(H_1)}{P(E|H_1)P(H_1) + P(E|H_2)P(H_2)}$$
$$P(E|H) = \binom{n}{k} p^k (1-p)^{n-k}$$


- ▶  $n$ : total # of occ
- ▶  $k$ : # (e.g.) carbonate occ

(Simpson and Harnik 2009 *Paleobiology*)


# Models

# Preliminary results: model comparison

# Preliminary results: best substrate



# Preliminary results: best habitat



Introduction and theory

Brachiopods, environmental preference, and extinction


Ecology and survival in Cenozoic mammals

Community connectedness in Cenozoic mammals

# Ecological traits

- ▶ dietary category
  - ▶ energetics
  - ▶ availability
- ▶ locomotor category
  - ▶ availability
  - ▶ dispersal
- ▶ body size
  - ▶ energetics
  - ▶ home range size

# Predictions: dietary category




- ▶ trophic hierarchy  
(stability → duration)
  - ▶ herb: most stable, longest duration
  - ▶ carni: least stable, shortest duration
  - ▶ omni: avg. stability, avg. duration
- ▶ ↑ diversification
  - ▶ ↑ origination;  $\simeq$  extinction
  - ▶  $\simeq$  origination;  $\downarrow$  extinction


# Predictions: locomotor category

- ▶ Paleogene → Neogene
  - ▶ open → closed environment

# Predictions: body size



(Liow et al. 2008 PNAS)



(Tomiya 2013 Am. Nat.)

- ▶ ↑ body size, ↑ energy req,  
↑ range size, ↓ extinction
- ▶ Europe
  - ▶ lrg body size genera:  
↑ extinction
- ▶ North America
  - ▶ generic body size:  
 $\simeq$  extinction

# Methodology

# Biases to survival: a simulation study

- ▶ bias away from  $h(t) = \lambda$ 
  - ▶ species:genus
  - ▶ anagenesis/cryptic speciation
- ▶ time-homogeneous birth-death model
  - ▶ common phylogenetic model
  - ▶ constant  $p, b$
  - ▶ expected  $S(t) = \exp^{-\lambda t}$
  - ▶ vary (cryptic) anagenesis

Introduction and theory

Brachiopods, environmental preference, and extinction


Ecology and survival in Cenozoic mammals

Community connectedness in Cenozoic mammals

## Definition

The degree to which localities are composed of endemic versus cosmopolitan taxa, and how similar this ratio is across localities.

# Biogeographic networks



(Vilhena *et al.* 2013 *Sci. Reports*)

- ▶ taxa: species
- ▶ locality: 2x2 equal-area map projection grid
- ▶ 2 My intervals
- ▶ PBDB, NOW, museum collections, compilations

## Average relative number of endemics

$$E = \frac{\sum_{i=1}^L \frac{u_i}{n_i}}{L}$$



- ▶  $L$ : number of localities
- ▶  $u$ : number of taxa unique to a locality
- ▶  $n$ : number of taxa at a locality
- ▶  $0 \leq E \leq 1$

## Average relative occupancy per taxon

$$Occ = \frac{\sum_{i=1}^N \frac{l_i}{L}}{N}$$

- ▶  $N$ : total number of taxa
- ▶  $l_i$ : number of localities a taxon occurs at
- ▶  $L$ : number of localities
- ▶  $0 \leq Occ \leq 1$

# Biogeographic connectedness





(Sidor et al. 2013 PNAS)


$$BC = \frac{O - N}{LN - N}$$

- ▶  $O$ : number of occurrences
- ▶  $N$ : total number of taxa
- ▶  $L$ : number of localities
- ▶  $0 \leq BC \leq 1$


# Code length



1111100 110 0110 10011 10000 10011 0110 0011 10011 0001  
000 000 0111 10001 110 011 10001 0111110 00001110 110 0001  
0111 110 0111 0110 11101 11001 11000 10100 00001110 10001 0111  
0000 1010 11010 10111 1001 0100 1001 10111 0100 1001 0100  
0111 010 0111 0110 11001 0110 0011 0100 1001 10111 0011 0100  
0111 10001 1110 11001 0111 0110 0011 0100 1001 10111 1010 0110  
0001




111 0000 11 01 101 100 101 01 0001 0 110 011 00 110 00 111 101 10  
111 000 10 111 000 111 10 011 10 000 111 10 111 10 000 10 011 010  
011 10 000 111 000 1 0001 0 111 010 100 011 00 111 00 011 100 111  
100 111 110 0011 111 01 100 0 100 00 111 00 011 100 111 00 011 100 111  
10 111 000 10 0001 111 0001 0 111 000 110 011 00 011 100 111 00 011 100 111



111 0000 11 01 101 100 101 01 0001 0 110 011 00 110 00 111 101 10  
111 000 10 111 000 111 10 011 10 000 111 10 111 10 000 10 011 010  
011 10 000 111 000 1 0001 0 111 010 100 011 00 111 00 011 100 111  
100 111 110 0011 111 01 100 0 100 00 111 00 011 100 111 00 011 100 111  
10 111 000 10 0001 111 0001 0 111 000 110 011 00 011 100 111 00 011 100 111

(Rosvall and Bergstrom 2008 PNAS)


# Global versus regional versus local scale processes



(Sidor et al. 2013 PNAS)

- ▶ global
  - ▶ corr w/ global climate
  - ▶ multiple regions corr
- ▶ regional
  - ▶  $\downarrow E, \uparrow Occ,$   
 $\uparrow BC, \uparrow code$
- ▶ local
  - ▶  $\uparrow E, \downarrow Occ,$   
 $\downarrow BC, \downarrow code$
- ▶ not mutually exclusive

# Expectations: locomotor category



## ► arboreal

- ▶ ↑  $E$ , ↑ code
- ▶ ↓ BC, ↓ Occ


## ► ground dwelling

- ▶ ↓  $E$ , ↓ code
- ▶ ↑ BC, ↑ Occ

## ► scansorial

- ▶ constant ∨ random

# Expectations: dietary category



(Jernvall and Fortelius 2004 *Am. Nat.*)


- ▶ herbivore
  - ▶ most like all taxa
- ▶ carnivore
  - ▶ constant √ corr w/ herbivores
- ▶ omnivore
  - ▶ constant √ random

# Community connectedness of North America



# Community connectedness of Europe

# Community connectedness of South America



# Preliminary results: NA, Eur



# Preliminary results: locomotor category



# Preliminary results: dietary category



# Questions

## Questions

- ▶ Why do certain taxa go extinct while others do not?
- ▶ How is emergence “formed?” When should it be invoked?
- ▶ Is extinction risk taxon–age independent?
- ▶ When should we expect global, regional, or local processes to dominate?

# Summary of proposed research

## Studies

- ▶ Permian brachiopod trait based survival
- ▶ Cenozoic mammal trait based survival
- ▶ Cenozoic mammal community connectedness

# Acknowledgements

## ► Committee

- ▶ Kenneth D. Angielczyk  
(co-advisor)
- ▶ Michael J. Foote  
(co-advisor)
- ▶ P. David Polly
- ▶ Richard H. Ree

## ► Discussion

- ▶ David Bapst, Megan Boatright, Ben Frable, Colin Kyle, Darcy Ross, Liz Sander
- ▶ John Alroy, Graeme Lloyd, Carl Simpson, Graham Slater

