

Since last meeting

Current projects

Brachiopods

Mammals

Timeline

Since last meeting

Current projects

Brachiopods

Mammals

Timeline

Since last meeting

- ▶ Evolution 2015 talk
- ▶ GSA 2015 talk
- ▶ Chapter 1 published (PNAS)
 - ▶ Effects of biotic traits on mammal species duration
- ▶ Chapter 2 submitted (Evolution)
 - ▶ Interplay between extinction intensity and selectivity in brachiopod extinction
 - ▶ Submitted early October, still in review?
- ▶ Did not submit DDIG

Review possible chapter 1

Review possible chapter 2

Since last meeting

Current projects

Brachiopods

Mammals

Timeline

Regional patterns in the diversification of Paleozoic brachiopods


Question

How does differential taxonomic entrance and loss contribute to regional (e.g. latitudinal) diversity?

Motivation


- ▶ latitudinal diversity gradients
 - ▶ through lense of a diversification process
- ▶ regional as opposed to global
 - ▶ variation within regions may not match global pattern (more biologically relevant?)
 - ▶ partial follow up to brachiopod survival work

Brachiopod latitudinal diversity





(Powell 2007 *G. Eco. Biogeo.*)

Variation in bioersity gradient




(Powell *et al* 2015 *Paleobio.*)

“Modes” of latitudinal diversity




(Powell 2007 *G. Eco. Biogeo.*)

Change in evenness + diversity




(Powell *et al* 2015 *Paleobio.*)

Model structure: Markov model



Model structure: hidden state



Preliminary results

Major assumptions


- ▶ model is only a first-order Markov process
 - ▶ can lead to some taxa existing longer than in actuality
- ▶ any taxon can occur in any geographic unit independent of other units
- ▶ both possibly controlled for by sampling rate through time
 - ▶ further assumes all times and places can be considered similar
- ▶ relaxing all of these assumption increases complexity
 - ▶ expand to whole Phanerozoic and/or other taxonomic groups?

Changes in Cenozoic mammal ecotype composition

Question



How do occurrence ratios of mammalian ecotypes change over time?

Environmental shift



(Stromberg 2005 *PNAS*)

Possible link?



(Smits 2015 *PNAS*)

Multi-logit regression

$$y_i \sim \text{Categorical}(K, \pi)$$

$$\pi_k = \frac{\exp(\beta_{k,j[i]}X_i + \lambda_k)}{\sum_{k=1}^K \exp(\beta_{k,j[i]}X_i + \lambda_{k,i})}$$

$$\text{where } \beta_{K,j[i]}X_i + \lambda_{K,i} = 0$$

$$\lambda_k \sim \text{MVN}(0, \tau_k^2 \Sigma)$$

$$\beta_{k,j} \sim \mathcal{N}(\beta'_k, \sigma_k)$$

Preliminary results

Further developments

- ▶ **NOTE** currently single flat mean; allow trend/multiple?
 - ▶ time order is not currently modeled
- ▶ **NOTE** no phylogenetic effect for $k = K$
 - ▶ every thing is relative **to** it
- ▶ increased categorization (e.g. frugivory), covariates (e.g. climate)?
- ▶ observed taxa represent a proportional sample of reality
 - ▶ how can this be overcome in a **model based** framework?
- ▶ limits to complexity of model due to sample size

Since last meeting

Current projects

Brachiopods

Mammals

Timeline

Things to consider

- ▶ TAing this spring and next year
- ▶ Funding?
 - ▶ FMNH fellow (but I don't spend time at the museum).
- ▶ Estimates for time of completion?
- ▶ Post-doctoral opportunities?