

Title: How macroecology affects macroevolution: the interplay
2 between extinction intensity and trait-dependent extinction in
brachiopods.

4 **Running title:** Trait-dependent extinction in brachiopods

5 **Author:** Peter D Smits, psmits@uchicago.edu, Committee on Evolutionary
6 Biology, University of Chicago, IL, USA.

Keywords: species selection, paleobiology, Bayesian

8 **Word count:** \approx 7000

Table count: 1

10 **Figure count:** 7

Data archival location: Zenodo DOI 10.5281/zenodo.46928.

Abstract

As extinction intensity increases, how do the effects of traits on
 14 taxonomic survival change? Does the extinction rate associated with
 certain traits increase while that of others decreases? Using a hierarchical
 16 Bayesian approach, I develop a model of how the effects of biological
 traits on extinction risk can vary with respect to extinction intensity,
 18 origination cohort (i.e. time of origination), and in relation to each other.
 The emergent traits I analyze in relation to their patterns of
 20 Paleozoic brachiopod genus durations are geographic range, affinity for
 epicontinental seas versus open ocean environments, and body size.
 22 Additionally, I estimate the effects of environmental generalization versus
 specialization on taxonomic survival by allowing environmental preference
 24 to have a nonlinear effect on duration. My analytical framework eschews
 the traditional distinction between background and mass extinction, and
 26 instead considers extinction intensity as a continuum. I find that the
 cohort-specific effects of geographic range and environmental preference
 28 are negatively correlated with baseline extinction intensity. Additionally, I
 find support for greater survival of environmental generalists versus
 specialists in all origination cohorts. These results support the conclusion
 30 that for Paleozoic brachiopods, as extinction intensity increases overall
 extinction selectivity increases.
 32

1 Introduction

34 Extinction is one half of the diversification process (Raup, 1994, Stanley, 1975,
 1979), second only to speciation or origination in shaping diversity; it can also
 36 be the ultimate manifestation of selection as a taxon with a beneficial trait
 should persist for longer on average than a taxon without that beneficial trait
 38 (Jablonski, 2008, Rabosky and McCune, 2010, Raup, 1994, Stanley, 1975).

Species duration is a measure of species fitness CITATION, and trait-associated
40 differences in fitness is the hallmark of (species) selection CITATION.

Jablonski (1986) observed that for bivalves at the end Cretaceous mass
42 extinction event, previous trait-associated differences in survival no longer
mattered except for the case of geographic range. Based on this evidence,
44 Jablonski (1986) proposed the idea of "macroevolutionary modes" and that
mass extinction and background extinction are fundamentally different
46 processes. However, based on estimates of extinction rates over time, there is no
evidence of there being two or more "types" of extinction (Wang, 2003). Instead,
48 extinction rates for marine invertebrates vary continuously as a unimodal
distribution.

50 The apparent disconnect between this theory of macroevolutionary modes and
the observation of continuous variation in extinction rates implies the possibility
52 of a relationship between the strength of selection (extinction **intensity**) and
the association between traits and differences in fitness (extinction
54 **selectivity**) CITATION PAYNE. As extinction intensity increases, what
happens to extinction selectivity? How do trait-associated differences in fitness
56 change as average extinction rate changes over time?

Here I model brachiopod taxon durations as a function of multiple functional
58 taxon traits because trait-dependent differences in extinction risk should be
associated with differences in taxon duration CITATION. Brachiopods are an
60 ideal group for this study as they have an exceptionally complete fossil record
(Foote, 2000b, Foote and Raup, 1996). I focus on the brachiopod record from
62 the post-Cambrian Paleozoic, from the start of the Ordovician (approximately
485 My) through the end Permian (approximately 252 My) as this represents
64 the time of greatest global brachiopod diversity (Alroy, 2010) which results in a
large sample size.

66 The analysis of taxon durations, or time from origination to extinction, falls
under the purview of survival analysis, a field of applied statistics commonly
68 used in health care and engineering (Klein and Moeschberger, 2003) but has a
long history in paleontology (Crampton et al., 2016, Simpson, 1944, 1953, Smits,
70 2015, Van Valen, 1973, 1979). I adopt a hierarchical Bayesian modeling
approach (Gelman et al., 2013, Gelman and Hill, 2007) in order to unify the
72 previously distinct dynamic and cohort paleontological survival approaches
(Baumiller, 1993, Crampton et al., 2016, Ezard et al., 2012, Foote, 1988, Raup,
74 1975, 1978, Simpson, 2006, Van Valen, 1973, 1979).

For analysis of the possible association between extinction intensity and
76 extinction selectivity, average extinction rate and the trait-based differences in
extinction rate need to be estimated. In particular, estimates of how these
78 values vary over time. These requirements necessitate large amounts of data and
parameter rich models. There are two general approaches to this problem:
80 phylogenetic comparative (PCM) and paleobiological. PCM model the
distribution of branch lengths from a time-scaled phylogeny as some kind of
82 birth-death process in order to estimate the speciation and extinction rates
(Fitzjohn, 2010, Goldberg et al., 2011, 2005, Maddison et al., 2007, Rabosky
84 et al., 2013, Stadler, 2013, Stadler and Bokma, 2013, ?) MORE CITATION.
The paleobiological approach to estimating speciation and extinction rates, as
86 well as trait-dependent differences in those rates, is a type of time-series analysis
where the occurrence and sampling histories of fossil taxa are used to estimate
88 the underlying true taxon durations that are fit by a a birth-death process (with
incomplete observation) that fits the observed data.

90 While the capabilities of the PCM approach have increased dramatically from
their first inception, being capable of estimating time-varying rates as well as
92 possible trait-dependent or clade-dependent differences in those rates

CITATION, there are known analytical limitations that hamper analysis of
94 extinction rates and more specifically trait-dependent differences in extinction
in non-ultrametric trees (i.e. those with fossil tips), estimates which are critical
96 to answering the question of how extinction intensity relates to extinction
selectivity. Additionally, many fossil invertebrate groups do not have an inferred
98 phylogeny that covers a significant portion of history of a clade CITATION; this
is the case for post-Cambrian Paleozoic brachiopods. What phylogenies do exist
100 for this group exist at relatively small taxonomic and temporal scales
CITATION, and are unsuited for questions which require large datasets in order
102 to fit parameter rich models. This reality means that a paleobiological approach
to this question is more appropriate than a PCM one.

104 Previous work that has dealt with extinction intensity versus selectivity
approached this problem by independently estimating the extinction intensity
106 and selectivity at different time points or for different origination cohorts and
then comparing those estimates CITATION. I find this approach problematic
108 for a few reasons. First, by treating each time point or cohort as independent
the estimates from each model are not relative to each other and they do not
110 represent the greatest compromise across all available data CITATIONS.
Second, by treating each time point or cohort as independent any and all
112 post-hoc analyses are at risk of false positive results because of multiple
comparisons CITATIONS. Third, post-hoc analysis of correlations between
114 maximum likelihood estimates is much less satisfying or coherent than including
those correlations as estimates of the initial model, something that is not
116 possible when each time point or cohort is considered independent CITATION.

The hierarchical Bayesian approach used here involves developing a single model
118 to estimate all of time points or cohorts simultaneously while also estimating
the group averages for all parameter CITATION. By averaging across groups,

¹²⁰ the analytical advantages of specific prior choices can induce the partial pooling
¹²¹ and regularization of parameter estimates which smooths over groups with small
¹²² sample sizes or weakly estimated effects. Additionally, individual estimates are
¹²³ much more comparable, not only between time points or cohorts, but also
¹²⁴ relative to the groups average CITATION. This approach is conceptually and
¹²⁵ analytically similar to mixed-effects modeling but the Bayesian framework
¹²⁶ allows the use of strongly regularizing priors in order to better constrain
parameter estimates.

¹²⁸ 1.1 Factors affecting brachiopod survival

Conceptually, taxon survival can be considered an aspect of “taxon fitness”
¹³⁰ (Cooper, 1984, Palmer and Feldman, 2012). Traits associated with taxon
survival are thus examples of species (or higher-level) selection, as differences in
¹³² survival are analogous to differences in fitness. The traits analyzed here are all
examples of emergent and aggregate traits (Jablonski, 2008, Rabosky and
¹³⁴ McCune, 2010); specifically I analyze genus-level traits. Emergent traits are
those which are not measurable from individuals of the lower level (e.g.
¹³⁶ organisms within species) such as geographic range, or even fossil sampling rate.
Aggregate traits, like body size or environmental preference, are the average of a
¹³⁸ shared trait across all members of a lower level.

Geographic range is widely considered the most important biological trait for
¹⁴⁰ estimating differences in extinction risk at nearly all times, with large
geographic range associated with low extinction risk (Finnegan et al., 2012,
¹⁴² Harnik et al., 2012, Jablonski, 1986, 1987, 2008, Jablonski and Roy, 2003, Payne
and Finnegan, 2007). This stands to reason even if extinction is completely at
¹⁴⁴ random; a taxon with an unrestricted range is less likely to go extinct at

random than a taxon with a restricted range.

¹⁴⁶ Epicontinental seas are a shallow-marine environment where the ocean has spread over the continental interior or craton with a depth typically less than
¹⁴⁸ 100m. In contrast, open-ocean coastline environments have much greater variance in depth, do not cover the continental craton, and can persist during
¹⁵⁰ periods of low sea level (Miller and Foote, 2009). Because of this, a simple hypothesis that taxa which favor epicontinental seas would be at great risk
¹⁵² during periods of low sea levels, such as during glacial periods, when epicontinental seas are drained. During the Paleozoic (approximately 541–252
¹⁵⁴ My), epicontinental seas were widely spread globally but declined over the Mesozoic (approximately 252–66 My) and have nearly disappeared during the
¹⁵⁶ Cenozoic (approximately 66–0 My) as open-ocean coastlines became the dominant shallow-marine setting (Johnson, 1974, Miller and Foote, 2009, Peters,
¹⁵⁸ 2008, Sheehan, 2001). Taxa in epicontinental environments could also have a greater extinction susceptibility than taxa in open-ocean environments due to
¹⁶⁰ anoxic events due to enhanced water stratification or poor water circulation (Peters, 2007).

¹⁶² Miller and Foote (2009) demonstrated that during several mass extinctions taxa associated with open-ocean environments tend to have a greater extinction risk
¹⁶⁴ than those taxa associated with epicontinental seas. During periods of background extinction, however, they found no consistent difference between
¹⁶⁶ taxa favoring either environment. Miller and Foote (2009) hypothesize that open-ocean taxa may have a greater extinction rate because these environments
¹⁶⁸ would be more strongly affected by waterborne hazards such as fallout from impacts or volcanic events which would propagate more quickly than in
¹⁷⁰ epicontinental environments with sluggish circulation. These two environment types represent the primary identifiable environmental dichotomy observed in

¹⁷² ancient marine systems (Miller and Foote, 2009, Sheehan, 2001). Given these
¹⁷⁴ findings, I would hypothesize that as extinction risk increases, the extinction
risk associated with open-ocean environments should generally increase.

¹⁷⁶ Because environmental preference is defined here as the continuum between
occurring exclusively in open-ocean environments versus epicontinental
¹⁷⁸ environments, intermediate values are considered “generalists” in the sense that
they favor neither end member. A long-standing hypothesis is that generalists or
¹⁸⁰ unspecialized taxa will have greater survival than specialists (Baumiller, 1993,
Liow, 2004, 2007, Nürnberg and Aberhan, 2013, 2015, Simpson, 1944, Smits,
¹⁸² 2015). Because of this, the effect of environmental preference was modeled as a
quadratic function where a concave down relationship between preference and
expected duration indicates that generalists are favored over specialists
¹⁸⁴ end-members.

¹⁸⁶ Body size, measured as shell length, is also considered as a trait that may
potentially influence extinction risk (Harnik, 2011, Payne et al., 2014). Body
size is a proxy for metabolic activity and other correlated life history traits
¹⁸⁸ (Payne et al., 2014). Harnik et al. (2014) analyzed the effect of body size
selectivity in Devonian brachiopods in both a phylogenetic and
¹⁹⁰ non-phylogenetic context; finding that body size was not found to be associated
with differences in taxonomic duration. It has also been found that, at least in
¹⁹² the case of some bivalve subclades, body size can be as important a factor as
geographic range size in determining extinction risk (Harnik, 2011). Given these
¹⁹⁴ results, I expect that if body size has any effect on brachiopod taxonomic
survival it is very small.

¹⁹⁶ It is well known that, given the incompleteness of the fossil record, the observed
duration of a taxon is an underestimate of that taxon’s true duration (Alroy,
¹⁹⁸ 2014, Foote and Raup, 1996, Liow and Nichols, 2010, Solow and Smith, 1997,

Wagner and Marcot, 2013, Wang and Marshall, 2004). Because of this, the
200 concern is that a taxon's observed duration may reflect its relative chance of
being sampled and not any of the effects of the covariates of interest. In this
202 case, for sampling to be a confounding factor there must be consistent
relationship between the quality of sampling of a taxon and its apparent
204 duration (e.g. greater sampling, longer duration). If there is no relationship
between sampling and duration then interpretation can be made clearly; while
206 observed durations are obviously truncated true durations, a lack of a
relationship would indicate that the amount and form of this truncation is not a
208 major determinant of the taxon's apparent duration. By including sampling as a
covariate in the model, this effect can be quantified and can be taken into
210 account when interpreting the estimates of the effects of the other covariates.

2 Materials and Methods

212 2.1 Fossil occurrence information

The brachiopod dataset analyzed here was sourced from the Paleobiology
214 Database (<http://www.paleodb.org>) which was then filtered based on
taxonomic (Rhyconelliformea: Rhynchonellata, Chileata, Obolellida,
216 Kutorginida, Strophomenida, Spiriferida), temporal (post-Cambrian Paleozoic),
stratigraphic, and other occurrence information used in this analysis. Analyzed
218 occurrences were restricted to those with paleolatitude and paleolongitude
coordinates, assignment to either epicontinental or open-ocean environment,
220 and belonging to a genus present in the body size dataset (Payne et al., 2014).
Epicontinental versus open-ocean assignments for each fossil occurrence are
222 partially based on those from Miller and Foote (2009), with additional

occurrences assigned similarly (Miller and Foote, personal communication).

²²⁴ These filtering criteria are very similar to those from Foote and Miller (2013)
with an additional constraint of being present in the body size data set from
²²⁶ Payne et al. (2014). In total, 1130 were genera included in the dataset.

Fossil occurrences were analyzed at the genus level which is common for
²²⁸ paleobiological, macroevolutionary, and macroecological studies of marine
invertebrates (Alroy, 2010, Foote and Miller, 2013, Harnik et al., 2012, Kiessling
²³⁰ and Aberhan, 2007, Miller and Foote, 2009, Nürnberg and Aberhan, 2013, 2015,
Payne and Finnegan, 2007, Simpson and Harnik, 2009, Vilhena et al., 2013).

²³² While species diversity dynamics are frequently of much greater interest than
those of higher taxa (though see Foote 2014, Hoehn et al. 2015), the nature of
²³⁴ the fossil record makes consistent taxonomic assignments at the species level
difficult for all occurrences in space and time. As such, the choice to analyze
²³⁶ genera as opposed to species was in order to assure a minimum level of
confidence and accuracy in the data analyzed here.

²³⁸ Genus duration was calculated as the number of geologic stages from first
appearance to last appearance, inclusive. Durations were based on geologic
²⁴⁰ stages as opposed to millions of years because of the inherently discrete nature
of the fossil record; dates are not assigned to individual fossils themselves but
²⁴² instead fossils are assigned to a geological interval which represents some
temporal range. In this analysis, stages are effectively irreducible temporal
²⁴⁴ intervals in which taxa may occur. Genera with a last occurrence in or after
Changhsingian stage (e.g. the final stage of the study interval) were right
²⁴⁶ censored at the Changhsingian; genera with a duration of only one stage were
left censored (Klein and Moeschberger, 2003). How the likelihood of censored
²⁴⁸ observations is calculated is detailed in section 2.2.

The covariates of duration included in this analysis are geographic range size

²⁵⁰ (r), environmental preference (v, v^2), the statistical interactions between
geographic range size and environmental preference ($r \times v, r \times v^2$), body size
²⁵² (m), and sampling (s). The interaction terms are important to include because
of the known multidirectional relationship between geographic range,
²⁵⁴ environmental preference, and duration CITATION. These interaction terms
help elucidate how the shape of environmental preference may change with
²⁵⁶ increasing geographic range.

Geographic range was calculated as relative occupancy corrected for incomplete
²⁵⁸ sampling. First, the paleolatitude-paleolongitude coordinates for all occurrences
were projected onto an equal-area cylindrical map projection. Each occurrence
²⁶⁰ was then assigned to one of the cells from a 70×34 regular raster grid placed
on the map. Each grid cell represents approximately 250,000 km². The map
²⁶² projection and regular lattice were made using shape files from
<http://www.naturalearthdata.com/> and the **raster** package for R (Hijmans,
²⁶⁴ 2015). For each stage, the total number of occupied grid cells was calculated.
Then, for each temporal bin, the relative occurrence probability of the observed
²⁶⁶ taxa was calculated using the JADE method developed by Chao et al. (2015).
This method accounts for the fact that taxa with an occupancy of 0 cannot be
²⁶⁸ observed which means that occupancy follows a truncated Binomial distribution.
This correction is critical when comparing occupancies from different times with
²⁷⁰ different geographic sampling. Finally, for each genus, the mean relative
occurrence probability was calculated as the average of that genus' occurrence
²⁷² probabilities for all stages it was sampled to yield relative occupancy, my proxy
for geographic range.

²⁷⁴ Environmental preference was defined as probability of observing the ratio of
epicontinental occurrences to total occurrences ($\theta_i = e_i/E_i$) or greater given the
²⁷⁶ background occurrence probability θ'_i as estimated from all other taxa occurring

at the same time (e'_i/E'_i). This measure of environmental preference is
278 expressed.

$$\begin{aligned} p(\theta'_i | e'_i, E'_i) &\propto \text{Beta}(e'_i, E'_i - e'_i) \text{Beta}(1, 1) \\ &= \text{Beta}(e'_i + 1, E'_i - e'_i + 1), \end{aligned} \quad (1)$$

where v is the percent of the distribution defined in equation 1 less than or
280 equal to θ_i . The Beta distribution is used here because it is a continuous
distribution bounded at 0 and 1, which is ideal for modeling percentages.

282 Body size, measured as shell length, was sourced directly from Payne et al.
(2014). These measurements were made from brachiopod taxa figured in the
284 *Treatise on Invertebrate Paleontology* (Williams et al., 2007).

286 The sampling probability for individual taxa was calculated using the standard
gap statistic (Foote, 2000a, Foote and Raup, 1996). The gap statistic is
calculated as the number of stages in which the taxon was sampled minus two
288 divided by the duration of the taxon minus two. Subtracting two from both the
numerator and denominator is because the first and last appearance stages are
290 by definition sampled. Because taxa that were right censored only include a first
appearance, one was subtracted from the numerator and denominator instead of
292 two.

294 The minimum duration for which a gap statistic can be calculated is three
stages, so I chose to impute the gap statistic for all observations with a
duration less than 3. Imputation is the “filling in” of missing observations based
296 on the observed values (Gelman and Hill, 2007, Rubin, 1996). This is fairly
straight forward in a Bayesian framework because both covariates and
298 parameters are considered random variables, meaning that the missing values of
sampling can be modeled as coming from some probability distribution. The
300 model for imputing sampling probability is described in section 2.3.

Prior to analysis, geographic range was logit transformed and body size was
302 natural-log transformed; both of these transformations make these variables
defined for the entire real line. Sampling probability was transformed as
304 $(s(n - 1) + 0.5)/n$ where n is the sample size as recommended by Smithson and
Verkuilen (2006); this serves to slightly shrink the range of the data so that
306 there are no values of 0 or 1. All covariates except for sampling were
standardized by subtracting the mean from all values and dividing by twice its
308 standard deviation, which follows Gelman and Hill (2007). This standardization
means that the associated regression coefficients are comparable as the expected
310 change per 1-unit change in the rescaled covariates. Finally, D is defined as the
total number of covariates, excluding sampling, plus one for the intercept term.

312 2.2 Details of model

Hierarchical modelling is a statistical approach which explicitly takes into
314 account the structure of the observed data in order to model both the within
and between group variance (Gelman et al., 2013, Gelman and Hill, 2007). The
316 units of study (e.g. genera) each belong to a single group (e.g. origination
cohort). Each group is considered a draw from a shared probability distribution
318 (e.g. prior) of all cohorts, observed and unobserved. The group-level parameters,
or the hyperparameters of this shared prior, are themselves given (hyper)prior
320 distributions and are also estimated like the other parameters of interest (e.g.
covariate effects) (Gelman et al., 2013). The subsequent estimates are partially
322 pooled together, where parameters from groups with large samples or effects
remain large while those of groups with small samples or effects are pulled
324 towards the overall group mean. All covariate effects (regression coefficients), as
well as the intercept term (baseline extinction risk), were allowed to vary by
326 group (origination cohort). The covariance between covariate effects was also

modeled.

³²⁸ Genus durations were assumed to follow a Weibull distribution which allows for age-dependent extinction (Klein and Moeschberger, 2003): $y \sim \text{Weibull}(\alpha, \sigma)$.

³³⁰ The Weibull distribution has two parameters: scale σ , and shape α . When $\alpha = 1$, σ is equal to the expected duration of any taxon. α is a measure of the effect of age on extinction risk where values greater than 1 indicate that extinction risk increases with age, and values less than 1 indicate that extinction risk decreases with age. Note that the Weibull distribution is equivalent to the exponential distribution when $\alpha = 1$.

³³⁶ In the case of the right- and left-censored observations mentioned above, the probability of those observations has a different calculation (Klein and Moeschberger, 2003). For right-censored observations, the likelihood is calculated $p(y|\theta) = 1 - F(y) = S(y)$ where $F(y)$ is the cumulative distribution function. In contrast, the likelihood of a left-censored observation is calculated $p(y|\theta) = F(y)$.

³⁴² The scale parameter σ was modeled as a regression following Kleinbaum and Klein (2005) with both varying intercept and varying slopes and the effect of sampling; this is expressed

$$\sigma_i = \exp\left(\frac{-\mathbf{X}_i B_{j[i]} + \delta s_i}{\alpha}\right) \quad (2)$$

³⁴⁶ where i indexes across all observations where $i = 1, \dots, n$ where n is the total number of observations, $j[i]$ is the cohort membership of the i th observation where $j = 1, \dots, J$ where J is the total number of cohorts, X is a $N \times D$ matrix of covariates along with a column of 1's for the intercept term, B is a $J \times D$ matrix of cohort-specific regression coefficients, and δ is the regression coefficient for the effect of sampling s . δ does not vary by cohort.

Each of the rows of matrix B are modeled as realizations from a multivariate
 352 normal distribution with length D location vector μ and $J \times J$ covariance
 matrix Σ : $B_j \sim \text{MVN}(\mu, \Sigma)$. The covariance matrix was then decomposed into a
 354 length J vector of scales τ and a $J \times J$ correlation matrix Ω , defined
 $\Sigma = \text{diag}(\tau)\Omega\text{diag}(\tau)$ where “diag” indicates a diagonal matrix.

356 The elements of μ were given independent normally distributed priors. The
 effects of geographic range size and the breadth of environmental preference
 358 were given informative priors reflecting the previous findings while the others
 were given weakly informative favoring no effect. The correlation matrix Ω was
 360 given an LKJ distributed prior (Lewandowski et al., 2009) that slightly favors
 an identity matrix as recommended by ?. These priors are defined

$$\begin{aligned}\mu^0 &\sim \mathcal{N}(0, 5) \\ \mu^r &\sim \mathcal{N}(-1, 1) \\ \mu^v &\sim \mathcal{N}(0, 1) \\ \mu^{v^2} &\sim \mathcal{N}(1, 1) \\ \mu^{r \times v} &\sim \mathcal{N}(0, 1) \\ \mu^{r \times v^2} &\sim \mathcal{N}(1, 1) \\ \mu^m &\sim \mathcal{N}(0, 1) \\ \tau &\sim \text{C}^+(1) \\ \Omega &\sim \text{LKJ}(2).\end{aligned}\tag{3}$$

362 The log of the shape parameter α was given a weakly informative prior
 $\log(\alpha) \sim \mathcal{N}(0, 1)$ centered at $\alpha = 1$, which corresponds to the Law of Constant
 364 Extinction (Van Valen, 1973).

2.3 Imputation of sampling probability

³⁶⁶ The vector sampling s has two parts: the observed part s^o , and the unobserved
³⁶⁸ part s^u . Of the 1130 total observations, 539 have a duration of 3 or more and
have an observed gap statistic. The gap statistic for the remaining 591
³⁷⁰ observations was imputed. As stated above, the unobserved part is the imputed,
or filled in, based on the observed part s^o . Because sampling varies between 0
and 1, I chose to model it as a Beta regression with matrix W being a
³⁷² $N \times (D - 3)$ matrix of covariates (i.e. geographic range size, environmental
preference, body size; no interactions) as predictors of sampling; this assumes
³⁷⁴ that the sampling value for all taxa come from the same distribution.
Importantly, I make no assumptions of causal structure.

³⁷⁶ Predicting sampling probability using the other covariate that are then included
in the model of duration is acceptable and appropriate in the case of imputation
³⁷⁸ where the sample goal is accurate prediction (Gelman and Hill, 2007, Rubin,
1996). Not including these covariates can lead to biased estimates of the
³⁸⁰ imputed variable; if the covariates themselves are related, not including them
will bias this correlation towards zero which then leads to improper imputation
³⁸² and inference (Rubin, 1996).

The Beta regression is defined

$$s^o \sim \text{Beta}(\phi = \text{logit}^{-1}(X^o\gamma), \lambda), \quad (4)$$

³⁸⁴ where γ is a length D vector of regression coefficients, and X defined as above.

The Beta distribution used in the regression is reparameterized in terms of a
³⁸⁶ mean parameter

$$\phi = \frac{\alpha}{\alpha + \beta} \quad (5)$$

and total count parameter

$$\lambda = \alpha + \beta \quad (6)$$

³⁸⁸ where α and β are the characteristic parameters of the Beta distribution
(Gelman et al., 2013).

³⁹⁰ The next step is to then estimate $s^u|s^o, X^o, X^u, \gamma$, the posterior distribution of
which are folded back into s and used as a covariate of duration (Eq. 2). All the
³⁹² elements of γ , and both δ (Eq. 2) and λ (Eq. 4) were given weakly informative
priors where

$$\begin{aligned} \gamma &\sim \mathcal{N}(0, 1) \\ \delta &\sim \mathcal{N}(0, 1) \\ \lambda &\sim \text{Pareto}(0.1, 1.5). \end{aligned} \quad (7)$$

³⁹⁴ 2.4 Posterior inference and posterior predictive checks

The joint posterior was approximated using a Markov-chain Monte Carlo
³⁹⁶ routine that is a variant of Hamiltonian Monte Carlo called the No-U-Turn
Sampler (Hoffman and Gelman, 2014) as implemented in the probabilistic
³⁹⁸ programming language Stan (Stan Development Team, 2014). The posterior
distribution was approximated from four parallel chains run for 10,000 steps
⁴⁰⁰ each, split half warm-up and half sampling and thinned to every 10th sample for
a total of 4000 posterior samples. Chain convergence was assessed via the scale
⁴⁰² reduction factor \hat{R} where values close to 1 ($\hat{R} < 1.1$) indicate approximate
convergence. Convergence means that the chains are approximately stationary
⁴⁰⁴ and the samples are well mixed (Gelman et al., 2013).

Model adequacy was evaluated using a couple of posterior predictive checks.
⁴⁰⁶ Posterior predictive checks are a means for understanding model fit or adequacy
where the basic idea is that replicated data sets simulated from the fitted model

408 should be similar to the original data and systematic differences between the
409 simulations and observations indicate weaknesses of the model fit (Gelman
410 et al., 2013). For both approaches used here, each posterior predictive datasets
411 were generated from a unique draw from the posterior distribution of each
412 parameter. The two posterior predictive checks used in this analysis are a
413 comparison of a non-parametric estimate of the survival function $S(t)$ from the
414 empirical dataset to the non-parametric estimates of $S(t)$ from the 100 posterior
415 predictive datasets, and comparison of the observed genus durations to the
416 average posterior predictive estimate of $\log(\sigma)$ (Eq. 2). The former is to see if
417 simulated data has a similar survival pattern to the observed, while the latter is
418 to see if the model systematically over- or under- estimates taxon survival.

3 Results

420 Comparison of the posterior predictive estimates of $S(t)$ to the empirical
421 estimate reveal few obvious biases except for the case of values from the far right
422 tail of observed durations (Fig. 1). This result is reinforced by the additional
423 posterior predictive comparison where most estimates are not systematically
424 biased except for a consistent under-estimate of $\log(\sigma)$ for older taxa (Fig. 2).
The results of both posterior predictive checks indicate that, for the majority of
426 observations, model has an adequate fit; of course there is also room for
improvement as I have only approximated a possible data generating process.

428 Body size is estimated to have no effect on taxon duration for the average
429 cohort, with the estimate being nearly 0 (Table 1). The variance between the
430 cohort-specific estimates of the effect of body size τ^m is estimated to be less
431 than the variance of between-cohort estimates of the effect of geographic range
432 size τ^r while having approximately equal variance to the average expected

duration for a cohort.

434 The group-level estimate for the effects of geographic range and environmental
preference on species duration must be considered together because of the
436 complexity of their interaction, as well as their statistical interaction in the
model CITATION. While there are only two covariates, these covariates are
438 associated with five parameters: μ^r , μ^v , μ^{v^2} , μ^{rxv} , and μ^{rxv^2} . I will first
consider the posterior estimates of the individual parameters, then consider
440 them ensemble.

The cohort-level effect of geographic range size on diration indicates large
442 geographic range is associated with long taxon durations (Table 1). Given the
estimates of μ^r and τ^r , there is a less than XXX% ($\pm XXX\%$ SD) probability
444 that this relationships would be reversed ($\Pr(\mathcal{N}(\mu^r, \tau^r) > 0)$). The
between-cohort variance τ^r is tied for approximately lowest variance of all the
446 group-level effects (Table 1).

The estimate of μ^v indicates that, while taxa closer to mean environmental
448 preference are expected to have a greater duration than environmental
specialists, peak expected duration is for species which slightly favor
450 epicontinental over open-ocean facing environments. As explored below, peak
expected duration is also affected by species geographic range. And, as will be
452 discussed, as geographic range increases the peak moves closer to the overall
average geographic range.

454 The estimate of μ^{v^2} indicates that, for the average value of all covariates, the
overall relationship between environmental preference and $\log(\sigma)$ is concave
456 down (Fig. 3, 4, and 5), with only a XXX% ($\pm XXX\%$ SD) probability that
any given cohort is convex up ($\Pr(\mathcal{N}(\mu^{v^2}, \tau^{v^2}) < 0)$).

458 The final important aspect of understanding the effects of geographic range and

environmental preference on taxon duration requires considering their
460 accompanying interaction terms (μ^{rxv}, μ^{rxv^2}). The expected value of these terms
are both positive, indicating a possible positive reenforcement where having
462 both a large geographic range and average environmental preference is expected
to yield a greater duration than having only one of these conditions. However,
464 there is considerable uncertainty around these estimates and considerable
overlap around 0. This result means that for the average cohort, it is only
466 weakly plausible that this interaction matters.

Taken together, the relationship between geographic range, environmental
468 preference, and duration supports the following conclusion: environmental
generalists are always favored over specialists, and this relationship may be
470 reinforced as geographic range increases. However, this reenforcement is not
necessary very plausible as a generality given the estimates of μ^{rxv} and μ^{rxv^2} ,
472 but instead is most likely indicative of considerable between cohort variation.
This last point is supported by the large estimates of τ^{rxv} and τ^{rxv^2} and is
474 further explored below.

The cohort-specific estimates of all the regression coefficients demonstrate a lot
476 of between cohort variance, with no obvious directional trends. As indicated in
Table 1 and detectable visually (Fig. 6), the estimates for β^0 , β^r , and β^m all
478 have much lower variance than the estimates of β^{v^2} , β_{vxr} , β_{vxr^2} .

While most cohort-specific estimates are very similar to the overall cohort-level
480 estimate, there are a few notable excursions away from the overall mean (Fig.
6). There are simultaneous excursions in both β^0 and β^v for cohorts originating
482 in the Givetian (387-382 My) and Frasnian (382-372 My) stages; both of which
directly precede the late Devonian mass extinction event at the
484 Frasnian/Famennian boundary. These cohorts are marked by both a high
extinction intensity and an increase in expected duration for taxa favoring

⁴⁸⁶ epicontinental environments over open-ocean ones; this is consistent with the
⁴⁸⁷ results of Miller and Foote (2009). Interestingly, there is no corresponding
⁴⁸⁸ excursion in the effect of geographic range or for any of the interaction terms
involving environmental preference (Table 1).

⁴⁹⁰ Cohorts originating from the Silurian through the Early Devonian have a
⁴⁹¹ slightly lower extinction intensity than the overall mean; these cohorts are those
⁴⁹² originating in the Llandovery (443-443 My) through the Emsian (407-393 My).
⁴⁹³ This is also a time period is also when there is the lowest overall probability
⁴⁹⁴ that epicontinental favoring taxa are expected to have greater duration than
⁴⁹⁵ open-ocean favoring taxa. Both the Silurian and Devonian periods are notable
⁴⁹⁶ for having been periods with a mostly “hothouse” climate, with no polar icecaps
⁴⁹⁷ and a high sea-level (Edwards and Fanning, 1985, Joachimski et al., 2009,
⁴⁹⁸ Munnecke et al., 2010). A decrease in extinction intensity may indicate that
⁴⁹⁹ brachiopods are better suited to the selective pressures of a hothouse climate as
⁵⁰⁰ well as during periods of high sea-level.

The cohort-specific relationships between environmental preference and $\log(\sigma)$
⁵⁰² were calculated from the estimates of β^0 , β^r , β^v , β^{v^2} , β^{rxv} , and β^{rxv^2} (Fig. ??, ??, ??) and reflect how these two covariates act in concert through the effects of
⁵⁰⁴ these parameters and not just individually (Fig. 6). I've visualized the this
⁵⁰⁵ complex relationship for three different values of taxon mass Because of multiple
⁵⁰⁶ covariates are associated with these parameters, I've visuzlized this relationship
⁵⁰⁷ at three different quantiles of the observed (transformed) body mass values
⁵⁰⁸ (10%, 50%, 90%). While the overall curvature of this relationship does not
⁵⁰⁹ change with either environmental preference or body mass, the y-intercept and
⁵¹⁰ inflection point move leftward towards being centered at average overall
⁵¹¹ environmental preference (Fig. ??, ??, ??). This result supports the conclusion
⁵¹² that larger geographic range and environmental generality are associated with

longer duration than either of those independently. However, as stated above,
514 this result is only somewhat plausible as a generality given the estimates of μ^{rxv}
and μ^{rxv^2} ; note that due to the uncertainty surrounding the position of the
516 inflection point in all three cases it is hard to make strong conclusions.

Beyond results already discussed above in the context of the parameters
518 individually, it is notable that the cohort originating in the Kungurian (279-272
My) is least like the overall expected relationship and has the most sharply
520 curved appearance due to a high estimate β^v^2 (Fig. 6). This cohort has the
biggest difference in extinction risk between environmental generalists and
522 specialists. The cohorts originating during the Emsian (407-393 My) and
Frasnian (382 - 372 My) are tied for second in sharpness of curvature. The least
524 sharply curved cohorts include those originating during Tremadocian (484-477
My), Hirnantian (445-443 My), Llandovery (443-433 My), and Ludlow (427-423
My). Except for the Tremadocian cohort, most of these cohorts originate during
526 the Silurian through the Early Devonian range identified earlier as having lower
528 expected extinction intensity than what is expected from the group-level
estimate.

530 The correlations of the cohort-specific estimates of the regression coefficients are
estimated as the off-diagonal elements of the correlation matrix Ω . Only XXX of
532 the elements of Ω are distinguishable from 0:

(Fig. 10).

534 Comparison of observed values of sampling, as measured by the gap statistic, to
random draws from the posterior estimates of the imputed sampling values
536 indicate that they are very similar (Fig. 11. This result is very encouraging as
this is the ultimate goal of multiple imputation: to fill in missing data with
538 values similar to the observed while taking into account the randomness of that

variable (Gelman and Hill, 2007, Rubin, 1996). The estimates of δ are based on
540 the set of observed values and the entire posterior of imputed values.

Sampling was found to have a negative effect (positive δ) on duration: greater
542 sampling, shorter duration (Table 1). While potentially counter intuitive, this
result is most likely due to some long lived taxa only be sampled in the stages of
544 the first and last appearance. Also, longer lived taxa have more opportunities to
not be sampled than shorter lived taxa. These two factors will lead to this result.

546 While the effect of sampling appears large compared to the other regression
coefficients, this is only because sampling was not standardized like the other
548 covariates. To make the coefficients comparable, δ is multiplied by twice the
posterior mean of the standard deviation of sampling probability; the
550 transformed value of δ is then 0.XXX (± 0.1 SD). This effect is relatively small
compared to the other covariate effects (Table 1). There is then a XXXX%
552 probability that the effect of geographic range μ^r has a greater magnitude than
 δ . Similarly, μ^v has a XXXX% probability of having a greater magnitude of
554 effect than δ .

The Weibull shape parameter α was found to be approximately 1.41 (± 0.05 SD)
556 with a 100% probability of being greater than 1. This result is not consistent
with the Law of Constant Extinction (Van Valen, 1973) and is instead
558 consistent with accelerating extinction risk with taxon age. This may indicate
that older taxa are out-competed by younger taxa, a result consistent with some
560 empirical results (Quental and Marshall, 2013, Smits, 2015, Wagner and
Estabrook, 2014) and (ironically) with a recently proposed Red Queen-like
562 model of evolution (Rosindell et al., 2015). This results, however, is not
consistent with other empirical results (Crampton et al., 2016, Finnegan et al.,
564 2008) and could potentially be caused by the minimum resolution of the fossil
record (Sepkoski, 1975). It is thus unclear if a strong biological inference can be

⁵⁶⁶ made from this result, which means that further work is necessary on the effect
of taxon age on extinction risk.

⁵⁶⁸ 4 Discussion

The generating observation behind this study was that for bivalves at the end
⁵⁷⁰ Cretaceous mass extinction event, the only biological trait that was found to affect extinction risk was geographic range while traits that had previously been
⁵⁷² beneficial had no effect (Jablonski, 1986). This observation raises two linked
questions: how does the effect of geographic range change with changing
⁵⁷⁴ extinction intensity, and how does the effect of other biological traits change
with changing extinction intensity?

⁵⁷⁶ I find that as intensity increases (β^0 decreases), the magnitude of the effect of
geographic range increases. I also find that as intensity increases, the effect of
⁵⁷⁸ favoring epicontinental environments of open-ocean environments is expected to
be increase; this is consistent with the results of Miller and Foote (2009). There
⁵⁸⁰ is no evidence for a correlation between the effect of geographic range and
environmental preference. Additionally, the between-cohort variance in effect of
⁵⁸² geographic range is much less than the between-cohort variance of the effect of
environmental preference which may underlie the lack of correlation between
⁵⁸⁴ these two effects.

Additionally, the lower between-cohort variance in the effect of geographic range
⁵⁸⁶ versus that higher between-cohort variance implies that for cohorts with a
greater than average extinction intensity, the difference in the effect geographic
⁵⁸⁸ range and the group-level effect of geographic range is expected to be smaller
than the difference between the effect of environmental preference and the
⁵⁹⁰ group-level effect of environmental preference.

I find consistent support for the “survival of the unspecialized,” with respect to
592 epicontinental versus open-ocean environmental preference, as a time-invariant
generalization of brachiopod survival; taxa with intermediate environmental
594 preferences are expected to have lower extinction risk than taxa specializing in
either epicontinental or open-ocean environments (Fig. ??), though the
596 curvature of the relationship varies from rather shallow to very peaked (Fig. ??).
However, this relationship is not symmetric about 0, as taxa favoring
598 epicontinental environments are expected to have a greater duration than taxa
favoring open-ocean environments. This description of environment only
600 describes one major aspect of a taxon’s environmental context, with factors
such as bathymetry and temperature being further descriptors of a taxon’s
602 adaptive zone (Harnik, 2011, Harnik et al., 2012, Heim and Peters, 2011,
Nürnberg and Aberhan, 2013); inclusion of these factors in future analyses
604 would potentially improve our understanding of the “survival of the
“unspecialized” hypothesis (Simpson, 1944).

606 Hopkins et al. (2014), in their analysis of niche conservatism and substrate
lithological preference in marine invertebrates, found that brachiopods were
608 among the least “conservative” groups; taxa were found to easily change
substrate preference on short time scales. While substrate preference is not the
610 same as environmental preference (as defined here), a question does arise: are
there three classes of environmental preference instead of two? These classes
612 would be taxa with broad tolerance (“true” generalists), inflexible specialists
 (“true” specialists), and flexible but with a narrow tolerance. A flexible taxon is
614 one with a narrow habitat preference at one time, but with preference that
changes over time with changing environmental availability. My analysis
616 assumes that traits are constant over the duration of the taxon meaning that
this scenario is not detectable; taxa with broad tolerances and flexible taxa with

618 narrow per-stage preference end up being treated the same way. Future work
should explore how environmental preference changes over lineage duration in
620 relation to environmental availability to estimate if the generalists–specialists
continuum is actually ternary relationship.

622 An alternative approach for specifically modeling survival that can take into
account imperfect observation than the method used here is the
624 Cormack-Jolly-Seber (CJS) model (Liow et al., 2008, Liow and Nichols, 2010,
Royle and Dorazio, 2008, Tomiya, 2013). This model is a type of hidden Markov
626 model with an absorbing state (i.e. extinction). In this model, survival is defined
as the probability of surviving from time t to time $t + 1$. Additionally, the effect
628 of preservation and sighting is estimated as probability of observing a taxon that
is present; this can extend the duration of a taxon beyond its last occurrence.
630 This approach is a fundamentally different from the method used in my analysis:
I am estimating the biasing effect of sampling probability on taxon duration to
632 then compare with effects of other covariates, while the CJS model estimates the
pre-sampling fossil record and then estimates per-time unit survival probability.

634 The use of genera as the unit of the study and how to exactly interpret the
effects of the biological traits is an important question. For example, if any of
636 the traits analyzed here are associated with increases in speciation rates, this
might increase the duration of genera through self-renewal (Raup, 1991, 1994),
638 which would be an example of the difference in biological pattern between
species and genera (Jablonski, 1987, 2007, 2008). This could lead to a trait
640 appearing to decrease generic level extinction risk by that trait increasing
species level origination rate instead of decreasing species level extinction risk.

642 The model used here could be improved through either increasing the number of
analyzed traits, expanding the hierarchical structure of the model to include
644 other major taxonomic groups of interest, and the inclusion of explicit

phylogenetic relationships between the taxa in the model as an additional
646 hierarchical effect. An example trait that may be of particular interest is the
affixing strategy or method of interaction with the substrate of the taxon, which
648 has been found to be related to brachiopod survival where, for cosmopolitan
taxa, taxa that are attached to the substrate are expected to have a greater
650 duration than those that are not (Alexander, 1977).

It is theoretically possible to expand this model to allow for comparisons both
652 within and between major taxonomic groups which would better constrain the
brachiopod estimates while also allowing for estimation of similarities and
654 differences in cross-taxonomic patterns. The major issue surrounding this
particular expansion involves finding a similarly well sampled taxonomic group
656 that is present during the Paleozoic. Potential groups include Crinoidea,
Ostracoda, and other members of the “Paleozoic fauna” (Sepkoski, 1981).

658 With significant updates, it would also be possible to compare the brachiopod
record with with Modern groups such as bivalves or gastropods (Sepkoski, 1981),
660 though remembering that the groups may not necessarily share all cohorts with
the brachiopods. This particular model expansion would act as a test of any
662 universal cross-taxonomic patterns in the effects of emergent traits on extinction
such as has been proposed for geographic range (Payne and Finnegan, 2007).
664 Additionally, this expanded model would also act as a test of the distinctness of
the Sepkoski (1981) three-fauna hypothesis in terms of trait-dependent
666 extinction.

Traits like environmental preference or geographic range (Hunt et al., 2005,
668 Jablonski, 1987) are most likely heritable. Without phylogenetic context, this
analysis assumes that differences in extinction risk between taxa are
670 independent of the shared evolutionary history of those taxa (Felsenstein, 1985).
In contrast, the origination cohorts only capture shared temporal context. For

example, if taxon duration is phylogenetically heritable, then closely related taxa may have more similar durations as well as more similar biological traits.

Without taking into account phylogenetic similarity the effects of these biological traits would be inflated solely due to inheritance. The inclusion of phylogenetic context as an additional individual-level hierarchical effect, independent of origination cohort, would allow for determining how much of the observed variability is due to shared evolutionary history versus shared temporal context versus actual differences associated with biological traits (Smits, 2015).

The combination and integration of the phylogenetic comparative and paleontological approaches requires both sources of data, something which is not possible for this analysis because there is no phylogenetic hypothesis for all Paleozoic taxa, something that is frequently the case for marine invertebrates with a good fossil record. When both data sources are available has been possible, however, the analysis can more fully address the questions of interest in macroevolution (Fritz et al., 2013, Harnik et al., 2014, Raia et al., 2012, 2013, Simpson et al., 2011, Slater et al., 2012, Slater, 2013, 2015, Smits, 2015, Tomiya, 2013).

In summary, patterns of Paleozoic brachiopod survival were analyzed using a fully Bayesian hierarchical survival modelling approach while also eschewing the traditional separation between background and mass extinction. I find that cohort extinction intensity is negatively correlated with both the cohort-specific effects of geographic range and environmental preference. These results imply that as extinction intensity increases (β^0) increases, it is expected that both effects will increase in magnitude. However, the change in effect of environmental preference is expected to be greater than the change in the effect of geographic range. Additionally, I find support for greater survival in environmental generalists over specialists in all origination cohorts analyzed;

this is consistent with the long standing “survival of the unspecialized”
700 hypothesis (Baumiller, 1993, Liow, 2004, 2007, Nürnberg and Aberhan, 2013,
2015, Simpson, 1944, 1953, Smits, 2015). The results of this analysis support the
702 conclusion that for Paleozoic brachiopods, as extinction intensity increases
overall extinction selectivity is expected to increase as well.

704 Acknowledgements

I would like to thank K. Angielczyk, M. Foote, P. D. Polly, R. Ree, and G.
706 Slater for helpful discussion during the conception of this study. I'd also like to
thank D. Bapst, N. Pierrehumbert and M. Villarosa Garcia for additional
708 comments. An additional thank you to A. Miller for the epicontinental versus
open-ocean assignments. Finally, thank you to the reviewers for their helpful
710 comments that improved this manuscript. This entire study would not
have been possible without the Herculean effort of the many contributors to the
712 Paleobiology Database. In particular, I would like to thank J. Alroy, M.
Aberhan, D. Bottjer, M. Clapham, F. Fürsich, N. Heim, A. Hendy, S. Holland,
714 L. Ivany, W. Kiessling, B. Kröger, A. McGowan, T. Olszewski, P.
Novack-Gottshall, M. Patzkowsky, M. Uhen, L. Villier, and P. Wager. This work
716 was supported by a NASA Exobiology grant (NNX10AQ446) to A. Miller and
M. Foote. I declare no conflicts of interest. This is Paleobiology Database
718 publication XXX. The code necessary for reproducing these results is available
as a Zenodo archive DOI 10.5281/zenodo.46928.

⁷²⁰ **References**

- Alexander, R. R., 1977. Generic longevity of articulate brachiopods in relation
⁷²² to the mode of stabilization on the substrate. *Palaeogeography, Palaeoclimatology, Palaeoecology* 21:209–226.
- ⁷²⁴ Alroy, J., 2010. The Shifting Balance of Diversity Among Major Marine Animal Groups. *Science* 329:1191–1194.
- ⁷²⁶ ———, 2014. A simple Bayesian method of inferring extinction. *Paleobiology* 40:584–607.
- ⁷²⁸ Baumiller, T. K., 1993. Survivorship analysis of Paleozoic Crinoidea: effect of filter morphology on evolutionary rates. *Paleobiology* 19:304–321.
- ⁷³⁰ Chao, A., T. C. Hsieh, R. L. Chazdon, R. K. Colwell, N. J. Gotelli, and B. D. Inouye, 2015. Unveiling the species-rank abundance distribution by generalizing the Good-Turing sample coverage theory. *Ecology* 96:1189–1201.
- ⁷³² Cooper, W. S., 1984. Expected time to extinction and the concept of fundamental fitness. *Journal of Theoretical Biology* 107:603–629.
- ⁷³⁴ Crampton, J. S., R. A. Cooper, P. M. Sadler, and M. Foote, 2016. Greenhouse–icehouse transition in the Late Ordovician marks a step change in extinction regime in the marine plankton. *Proceedings of the National Academy of Sciences* 113:1498–1503.
- ⁷³⁶ Edwards, D. and U. Fanning, 1985. Evolution and environment in the late Silurian–early Devonian: the rise of pteridophytes. *Philosophical Transactions of the Royal Society B: Biological Sciences* 309:147–165.
- ⁷⁴⁰ Ezard, T. H. G., P. N. Pearson, T. Aze, and A. Purvis, 2012. The meaning of birth and death (in macroevolutionary birth-death models). *Biology Letters* 8:139–42.

- Felsenstein, J., 1985. Phylogenies and the comparative method. *The American Naturalist* 125:1–15.
- 746
- Finnegan, S., N. A. Heim, S. E. Peters, and W. W. Fischer, 2012. Climate change and the selective signature of the Late Ordovician mass extinction. *Proceedings of the National Academy of Sciences* 109:6829–6834.
- 748
- Finnegan, S., J. L. Payne, and S. C. Wang, 2008. The Red Queen revisited: reevaluating the age selectivity of Phanerozoic marine genus extinctions. *Paleobiology* 34:318–341.
- 750
- Fitzjohn, R. G., 2010. Quantitative Traits and Diversification. *Systematic Biology* 59:619–633.
- 754
- Foote, M., 1988. Survivorship analysis of Cambrian and Ordovician Trilobites. *Paleobiology* 14:258–271.
- 756
- , 2000a. Origination and extinction components of taxonomic diversity: general problems. *Paleobiology* 26:74–102.
- 758
- , 2000b. Origination and extinction components of taxonomic diversity: Paleozoic and post-Paleozoic dynamics. *Paleobiology* 26:578–605.
- 760
- , 2014. Environmental controls on geographic range size in marine animal genera. *Paleobiology* 40:440–458.
- 762
- Foote, M. and A. I. Miller, 2013. Determinants of early survival in marine animal genera. *Paleobiology* 39:171–192.
- 764
- Foote, M. and D. Raup, 1996. Fossil preservation and the stratigraphic ranges of taxa. *Paleobiology* 22:121–140.
- 766
- Fritz, S. A., J. Schnitzler, J. T. Eronen, C. Hof, K. Böhning-Gaese, and C. H.

- 768 Graham, 2013. Diversity in time and space: wanted dead and alive. *Trends in
Ecology & Evolution* 28:509–16.
- 770 Gelman, A., J. B. Carlin, H. S. Stern, D. B. Dunson, A. Vehtari, and D. B.
Rubin, 2013. Bayesian data analysis. 3 ed. Chapman and Hall, Boca Raton,
772 FL.
- Gelman, A. and J. Hill, 2007. Data Analysis using Regression and
774 Multilevel/Hierarchical Models. Cambridge University Press, New York, NY.
- Goldberg, E. E., L. T. Lancaster, and R. H. Ree, 2011. Phylogenetic inference
776 of reciprocal effects between geographic range evolution and diversification.
Systematic Biology 60:451–65.
- 778 Goldberg, E. E., K. Roy, R. Lande, and D. Jablonski, 2005. Diversity,
endemism, and age distributions in macroevolutionary sources and sinks. *The
780 American Naturalist* 165:623–33.
- Harnik, P. G., 2011. Direct and indirect effects of biological factors on
782 extinction risk in fossil bivalves. *Proceedings of the National Academy of
Sciences* 108:13594–13599.
- 784 Harnik, P. G., P. C. Fitzgerald, J. L. Payne, and S. J. Carlson, 2014.
Phylogenetic signal in extinction selectivity in Devonian terebratulide
786 brachiopods. *Paleobiology* 40:675–692.
- Harnik, P. G., C. Simpson, and J. L. Payne, 2012. Long-term differences in
788 extinction risk among the seven forms of rarity. *Proceedings of the Royal
Society B: Biological Sciences* 279:4969–4976.
- 790 Heim, N. A. and S. E. Peters, 2011. Regional environmental breadth predicts
geographic range and longevity in fossil marine genera. *PLoS one* 6:e18946.

- 792 Hijmans, R. J., 2015. raster: Geographic data analysis and modeling. URL
<http://CRAN.R-project.org/package=raster>. R package version 2.3-24.
- 794 Hoehn, K. B., P. G. Harnik, and V. L. Roth, 2015. A framework for detecting
natural selection on traits above the species level. Methods in Ecology and
796 Evolution Pp. doi: 10.1111/2041-210X.12461.
- Hoffman, M. D. and A. Gelman, 2014. The No-U-Turn Sampler: Adaptively
798 Setting Path Lengths in Hamiltonian Monte Carlo. Journal of Machine
Learning Research 15:1351–1381.
- 800 Hopkins, M. J., C. Simpson, and W. Kiessling, 2014. Differential niche dynamics
among major marine invertebrate clades. Ecology Letters 17:314–323.
- 802 Hunt, G., K. Roy, and D. Jablonski, 2005. Species-level heritability reaffirmed: a
comment on "On the heritability of geographic range sizes". The American
804 Naturalist 166:129–135.
- Jablonski, D., 1986. Background and mass extinctions: the alternation of
806 macroevolutionary regimes. Science 231:129–133.
- , 1987. Heritability at the species level: analysis of geographic ranges of
808 cretaceous mollusks. Science 238:360–363.
- , 2007. Scale and hierarchy in macroevolution. Palaeontology 50:87–109.
- 810 ———, 2008. Species Selection: Theory and Data. Annual Review of Ecology,
Evolution, and Systematics 39:501–524.
- 812 Jablonski, D. and K. Roy, 2003. Geographical range and speciation in fossil and
living molluscs. Proceedings of the Royal Society B: Biological Sciences
814 270:401–406.
- Joachimski, M. M., S. Breisig, W. Buggisch, J. A. Talent, R. Mawson,

- 816 M. Gereke, J. R. Morrow, J. Day, and K. Weddige, 2009. Devonian climate
and reef evolution: Insights from oxygen isotopes in apatite. *Earth and*
818 *Planetary Science Letters* 284:599–609.
- Johnson, J. G., 1974. Extinction of Perched Faunas. *Geology* 2:479–482.
- 820 Kiessling, W. and M. Aberhan, 2007. Environmental determinants of marine
benthic biodiversity dynamics through Triassic–Jurassic time. *Paleobiology*
822 33:414–434.
- Klein, J. P. and M. L. Moeschberger, 2003. *Survival Analysis: Techniques for*
824 *Censored and Truncated Data*. 2nd ed. Springer, New York.
- Kleinbaum, D. G. and M. Klein, 2005. *Survival analysis: a self-learning text*. 2
826 ed. Springer, New York, NY.
- Lewandowski, D., D. Kurowicka, and H. Joe, 2009. Generating random
828 correlation matrices based on vines and extended onion method. *Journal of*
Multivariate Analysis 100:1989–2001.
- 830 Liow, L. H., 2004. A test of Simpson’s “rule of the survival of the relatively
unspecialized” using fossil crinoids. *The American Naturalist* 164:431–43.
- 832 ———, 2007. Does versatility as measured by geographic range, bathymetric
range and morphological variability contribute to taxon longevity? *Global*
834 *Ecology and Biogeography* 16:117–128.
- Liow, L. H., M. Fortelius, E. Bingham, K. Lintulaakso, H. Mannila, L. Flynn,
836 and N. C. Stenseth, 2008. Higher origination and extinction rates in larger
mammals. *Proceedings of the National Academy of Sciences* 105:6097–6102.
- 838 Liow, L. H. and J. D. Nichols, 2010. Estimating rates and probabilities of
origination and extinction using taxonomic occurrence data:
840 Capture-mark-recapture (CMR) approaches. Pp. 81–94, *in* J. Alroy and

- G. Hunt, eds. Quantitative Methods in Paleobiology. The Paleontological Society.
- Maddison, W. P., P. E. Midford, and S. P. Otto, 2007. Estimating a binary character's effect on speciation and extinction. *Systematic Biology* 56:701.
- Miller, A. I. and M. Foote, 2009. Epicontinental seas versus open-ocean settings: the kinetics of mass extinction and origination. *Science* 326:1106–9.
- Munnecke, A., M. Calner, D. A. T. Harper, and T. Servais, 2010. Ordovician and Silurian sea-water chemistry, sea level, and climate: A synopsis. *Palaeogeography, Palaeoclimatology, Palaeoecology* 296:389–413.
- Nürnberg, S. and M. Aberhan, 2013. Habitat breadth and geographic range predict diversity dynamics in marine Mesozoic bivalves. *Paleobiology* 39:360–372.
- , 2015. Interdependence of specialization and biodiversity in Phanerozoic marine invertebrates. *Nature Communications* 6:6602.
- Palmer, M. E. and M. W. Feldman, 2012. Survivability is more fundamental than evolvability. *PloS one* 7:e38025.
- Payne, J. L. and S. Finnegan, 2007. The effect of geographic range on extinction risk during background and mass extinction. *Proceedings of the National Academy of Sciences* 104:10506–11.
- Payne, J. L., N. A. Heim, M. L. Knope, and C. R. McClain, 2014. Metabolic dominance of bivalves predates brachiopod diversity decline by more than 150 million years. *Proceedings of the Royal Society B: Biological Sciences* 281:20133122.
- Peters, S. E., 2007. The problem with the Paleozoic. *Paleobiology* 33:165–181.

- _____, 2008. Environmental determinants of extinction selectivity in the fossil record. *Nature* 454:626–629.
- Quental, T. B. and C. R. Marshall, 2013. How the Red Queen Drives Terrestrial Mammals to Extinction. *Science* 341:290–292.
- Rabosky, D. L. and A. R. McCune, 2010. Reinventing species selection with molecular phylogenies. *Trends in Ecology & Evolution* 25:68–74.
- Rabosky, D. L., F. Santini, J. Eastman, S. a. Smith, B. Sidlauskas, J. Chang, and M. E. Alfaro, 2013. Rates of speciation and morphological evolution are correlated across the largest vertebrate radiation. *Nature Communications* 4:1–8.
- Raia, P., F. Carotenuto, F. Passaro, D. Fulgione, and M. Fortelius, 2012. Ecological specialization in fossil mammals explains Cope’s rule. *The American Naturalist* 179:328–37.
- Raia, P., F. Carotenuto, F. Passaro, P. Piras, D. Fulgione, L. Werdelin, J. Saarinen, and M. Fortelius, 2013. Rapid action in the Palaeogene, the relationship between phenotypic and taxonomic diversification in Coenozoic mammals. *Proceedings of the Royal Society B: Biological Sciences* 280:20122244.
- Raup, D. M., 1975. Taxonomic survivorship curves and Van Valen’s Law. *Paleobiology* 1:82–96.
- _____, 1978. Cohort Analysis of generic survivorship. *Paleobiology* 4:1–15.
- _____, 1991. *Extinction: Bad Genes or Bad Luck?* Norton, New York.
- _____, 1994. The role of extinction in evolution. *Proceedings of the National Academy of Sciences* 91:6758–6763.

- Rosindell, J., L. J. Harmon, and R. S. Etienne, 2015. Unifying ecology and
890 macroevolution with individual-based theory. *Ecology Letters* 18:472–482.
- Royle, J. A. and R. M. Dorazio, 2008. Hierarchical modeling and inference in
892 ecology: the analysis of data from populations, metapopulations and
communities. Elsevier, London.
- 894 Rubin, D. B., 1996. Multiple imputation after 18+ years. *Journal of the
American Statistical Association* 91:473–489.
- 896 Sepkoski, J. J., 1975. Stratigraphic biases in the analysis of taxonomic
survivorship. *Paleobiology* 1:343–355.
- 898 ———, 1981. A factor analytic description of the Phanerozoic marine fossil
record. *Paleobiology* 7:36–53.
- 900 Sheehan, P., 2001. The late Ordovician mass extinction. *Annual Review of
Earth and Planetary Sciences* 29:331–364.
- 902 Simpson, C., 2006. Levels of selection and large-scale morphological trends.
Ph.D. thesis, University of Chicago.
- 904 Simpson, C. and P. G. Harnik, 2009. Assessing the role of abundance in marine
bivalve extinction over the post-Paleozoic. *Paleobiology* 35:631–647.
- 906 Simpson, C., W. Kiessling, H. Mewis, R. C. Baron-Szabo, and J. Müller, 2011.
Evolutionary diversification of reef corals: a comparison of the molecular and
908 fossil records. *Evolution* 65:3274–3284.
- Simpson, G. G., 1944. *Tempo and Mode in Evolution*. Columbia University
910 Press, New York.
- , 1953. *The Major Features of Evolution*. Columbia University Press,
912 New York.

- Slater, G., L. Harmon, and M. Alfaro, 2012. Integrating fossils with molecular phylogenies improves inference of trait evolution. *Evolution* 66:3931–3944.
- Slater, G. J., 2013. Phylogenetic evidence for a shift in the mode of mammalian body size evolution at the Cretaceous-Palaeogene boundary. *Methods in Ecology and Evolution* 4:734–744.
- , 2015. Iterative adaptive radiations of fossil canids show no evidence for diversity-dependent trait evolution. *Proceedings of the National Academy of Sciences* 112:4897–4902.
- Smithson, M. and J. Verkuilen, 2006. A better lemon squeezer? Maximum-likelihood regression with beta-distributed dependent variables. *Psychological Methods* 11:54–71.
- Smits, P. D., 2015. Expected time-invariant effects of biological traits on mammal species duration. *Proceedings of the National Academy of Sciences* 112:13015–13020.
- Solow, A. and W. Smith, 1997. On fossil preservation and the stratigraphic ranges of taxa. *Paleobiology* 23:271–277.
- Stadler, T., 2013. Recovering speciation and extinction dynamics based on phylogenies. *Journal of Evolutionary Biology* 26:1203–1219.
- Stadler, T. and F. Bokma, 2013. Estimating speciation and extinction rates for phylogenies of higher taxa. *Systematic biology* 62:220–30.
- Stan Development Team, 2014. Stan: A c++ library for probability and sampling, version 2.5.0. URL <http://mc-stan.org/>.
- Stanley, S. M., 1975. A theory of evolution above the species level. *Proceedings of the National Academy of Sciences* 72:646–650.

- _____, 1979. Macroevolution: pattern and process. W. H. Freeman, San
938 Francisco.
- Tomiya, S., 2013. Body Size and Extinction Risk in Terrestrial Mammals Above
940 the Species Level. *The American Naturalist* 182:196–214.
- Van Valen, L., 1973. A new evolutionary law. *Evolutionary Theory* 1:1–30.
- 942 _____, 1979. Taxonomic survivorship curves. *Evolutionary Theory* 4:129–142.
- Vilhena, D. A., E. B. Harris, C. T. Bergstrom, M. E. Maliska, P. D. Ward, C. A.
944 Sidor, C. A. E. Strömberg, and G. P. Wilson, 2013. Bivalve network reveals
latitudinal selectivity gradient at the end-Cretaceous mass extinction.
- 946 Scientific Reports 3:1790.
- Wagner, P. J. and G. F. Estabrook, 2014. Trait-based diversification shifts
948 reflect differential extinction among fossil taxa. *Proceedings of the National
Academy of Sciences* 111:16419–16424.
- 950 Wagner, P. J. and J. D. Marcot, 2013. Modelling distributions of fossil sampling
rates over time, space and taxa: assessment and implications for
952 macroevolutionary studies. *Methods in Ecology and Evolution* 4:703–713.
- Wang, S. C., 2003. On the continuity of background and mass extinction.
954 Paleobiology 29:455–467.
- Wang, S. C. and C. R. Marshall, 2004. Improved confidence intervals for
956 estimating the position of a mass extinction boundary. *Paleobiology* 30:5–18.
- Williams, C., C. H. C. Brunton, and S. J. Carlson, 2007. Treatise on
958 invertebrate paleontology. Part H, Brachiopoda. Geological Society of
America, Boulder, Colorado.

Table 1: Estimates of various parameters in the model used here. These include group-level estimates of the effects of biological traits on brachiopod generic survival, the standard deviation of the between-cohort effects, as well as the estimates of both the effect of sampling δ and the Weibull shape parameter α . The mean, standard deviation (SD), 10th, 50th, and 90th quantiles of the marginal posteriors are presented.

type	parameter	effect of	mean	SD	10%	50%	90%
Mean	μ^i	intercept	-3.05	0.20	-3.30	-3.05	-2.80
	μ_i	-3.18	0.20	-3.45	-3.18	-2.93	
	μ_r	-1.15	0.20	-1.40	-1.14	-0.90	
	μ_v	-0.69	0.22	-0.97	-0.70	-0.41	
	μ_v2	3.10	0.41	2.58	3.12	3.61	
	μ_rxv	0.33	0.43	-0.21	0.32	0.88	
	μ_rxv2	0.17	0.61	-0.62	0.18	0.94	
Standard deviation	μ_m	-0.05	0.13	-0.21	-0.05	0.10	
	τ^i	intercept	0.51	0.11	0.38	0.50	0.65
	τ_i	0.49	0.12	0.35	0.49	0.64	
	τ_r	0.66	0.19	0.42	0.65	0.90	
	τ_v	0.93	0.20	0.69	0.91	1.19	
	τ_v2	1.67	0.47	1.12	1.62	2.31	
	τ_rxv	1.78	0.46	1.23	1.75	2.36	
δ	τ_rxv2	1.90	1.12	0.52	1.79	3.42	
	τ_m	0.48	0.13	0.32	0.47	0.64	
	sampling	0.90	0.15	0.71	0.90	1.09	
Other	δ	0.91	0.16	0.71	0.91	1.11	
	α	1.41	0.05	1.35	1.41	1.47	




Figure 1: Comparison of the empirical estimate of $S(t)$ (highlighted) versus estimates from 100 posterior predictive data sets (black). $S(t)$ corresponds to the probability that the age of a genus t is less than the genus' ultimate duration T . The vertical axis is log10 transformed.




Figure 2: Comparison of all observed genus durations in number of geological stages to the average posterior predictive estimates of $\log(\sigma)$. The dashed, diagonal line corresponds to $x = y$.




Figure 3: The overall expected relationship between environmental affinity v_i and a $\log(\sigma)$ when $r = 0$ and $m = 0$. The 1000 semi-transparent lines corresponds to a single draw from the posterior predictive distribution, while the highlighted line corresponds to the median of the posterior predictive distribution. The overall relationship is concave down with an optimum greater than 0, which means that taxa favoring epicontinental environments are expected to have longer durations than those favoring open-ocean environments. The tick marks along the bottom of the plot correspond to the (rescaled) observed values of environmental preference.




Figure 4: The overall expected relationship between environmental affinity v_i and a $\log(\sigma)$ when $r = 0$ and $m = 0$. The 1000 semi-transparent lines corresponds to a single draw from the posterior predictive distribution, while the highlighted line corresponds to the median of the posterior predictive distribution. The overall relationship is concave down with an optimum greater than 0, which means that taxa favoring epicontinental environments are expected to have longer durations than those favoring open-ocean environments. The tick marks along the bottom of the plot correspond to the (rescaled) observed values of environmental preference.




Figure 5: The overall expected relationship between environmental affinity v_i and a $\log(\sigma)$ when $r = 0$ and $m = 0$. The 1000 semi-transparent lines corresponds to a single draw from the posterior predictive distribution, while the highlighted line corresponds to the median of the posterior predictive distribution. The overall relationship is concave down with an optimum greater than 0, which means that taxa favoring epicontinental environments are expected to have longer durations than those favoring open-ocean environments. The tick marks along the bottom of the plot correspond to the (rescaled) observed values of environmental preference.




Figure 6: Comparison of cohort-specific estimates of β^0 , the effect of geographic range on extinction risk β^r , the effect of environmental preference β^v and β^{v^2} , and body size β^m . Points correspond to the median of the cohort-specific estimate, along with 80% credible intervals. Points are plotted at the midpoint of the cohorts stage of origination in millions of years before present (My). Black, horizontal lines are the overall estimates of covariate effects along with 80% credible intervals (shaded).




Figure 7: Comparison of origination cohort-specific (posterior predictive) estimates of the effect of environmental preference on $\log(\sigma)$ to the mean overall estimate of the effect of environmental preference. Cohort-specific estimates are from 100 posterior predictive simulations across the range of (transformed and rescaled) observed values of environmental preference. The oldest cohort is in the top-right and younger cohorts proceed left to right, with the youngest cohort being the right-most facet of the last row. Panel names correspond to the name of the stage in which that cohort originated.




Figure 8: Comparison of origination cohort-specific (posterior predictive) estimates of the effect of environmental preference on $\log(\sigma)$ to the mean overall estimate of the effect of environmental preference. Cohort-specific estimates are from 100 posterior predictive simulations across the range of (transformed and rescaled) observed values of environmental preference. The oldest cohort is in the top-right and younger cohorts proceed left to right, with the youngest cohort being the right-most facet of the last row. Panel names correspond to the name of the stage in which that cohort originated.




Figure 9: Comparison of origination cohort-specific (posterior predictive) estimates of the effect of environmental preference on $\log(\sigma)$ to the mean overall estimate of the effect of environmental preference. Cohort-specific estimates are from 100 posterior predictive simulations across the range of (transformed and rescaled) observed values of environmental preference. The oldest cohort is in the top-right and younger cohorts proceed left to right, with the youngest cohort being the right-most facet of the last row. Panel names correspond to the name of the stage in which that cohort originated.




Figure 10: Mixed graphical and numerical representation of the correlation matrix Ω of variation in cohort-specific covariate estimates. These correlations are between the estimates of the cohort-level effects of covariates, along with intercept/baseline extinction risk. The median estimates of the correlations are presented numerically (upper-triangle) and as idealized ellipses representing that much correlation (lower-triangle). The darkness of the ellipse corresponds to the magnitude of the correlation.




Figure 11: Histograms of the distribution of gap statistic values from both the observed values and the imputed values from 12 unique posterior realizations. For each panel the observed values are identical but the imputed values are from a single set of their posterior estimates.