

# How cryptic is cryptic diversity? Machine learning approaches to classifying morphological variation in the Pacific Pond Turtle (*Emys marmorata*)

Peter D Smits<sup>1</sup>, Kenneth D Angielczyk<sup>1,2</sup>, and James F Parham<sup>3</sup>

<sup>1</sup>Committee on Evolutionary Biology, University of Chicago

<sup>2</sup>Integrative Research Center, Field Museum of Natural History

<sup>3</sup>Department of Geological Sciences, California State University – Fullerton

September 10, 2015

**Corresponding author:** Peter D Smits, Committee on Evolutionary Biology, University of Chicago, 1025 E. 57th Street, Culver Hall 402, Chicago, IL, 60637, USA; E-mail: psmits@uchicago.edu

## Abstract

We investigate the morphometric identification of cryptic species using machine learning approaches by examining their implications for a recently proposed cryptic turtle species (*Emys pallida*). We collected landmark-based morphometric data from 354 adult *E. marmorata*/*“pallida”* museum specimens. We assigned a classification to each specimen for six different binning schemes based on geographic occurrence data recorded in museum collection archives. We used multiple machine learning methods, both unsupervised and supervised, to compare different classification hypotheses and asked whether it is possible to determine which amongst a set of classification hypotheses is best. In addition, we applied the above approach to a clear-cut example of seven unambiguously distinct species closely related to *E. marmorata*. The results of this study indicate that there is no clear grouping of *E. marmorata*/*“pallida”* based on plastron shape. In contrast, the analysis of the “clear-cut” example demonstrates a near perfect classification, which demonstrates that the methods can recover correct results when an appropriate signal exists. Explanations for the lack of grouping in *E. marmorata* include that possibility that genetic differentiation is not associated with plastron shape variation below the species level and/or that local selective pressures (e.g., from hydrological regime) overwhelm morphological differentiation. A reconsideration of the methods used to delimit *E. “pallida,”* the lack of barriers to gene flow, the strong evidence for widespread admixture between lineages, and the fact that plastron shape can be used to differentiate other emyline species suggest that its lack of diagnosability

most likely reflects the non-distinctiveness of this proposed taxon.

## INTRODUCTION

Molecular systematics has repeatedly demonstrated the existence of cryptic species that can only be diagnosed using genetic data (Stuart et al. 2006; Bickford et al. 2007; Schilck-Steiner et al. 2007; Pfenninger and Schwenk 2007; Clare 2011; Funk et al. 2012). In attempts to streamline the documentation of biodiversity, several methods of species delimitation that rely almost entirely on genetic data have recently been proposed (Pons et al. 2006; Carstens and Dewey 2010; Hausdorf and Hennig 2010; O'Meara 2010; Yang and Rannala 2010; Huelsenbeck et al. 2011). Although strong caveats on the utility of these methods have been raised (Bauer et al. 2000; Carstens et al. 2013), they are nevertheless being used to name species (Leaché and Fujita 2010; Spinks et al. 2014).

The majority of extant taxa, and almost all extinct taxa, are delimited by morphology alone. This disjunction complicates interpretations of variation and diversity in deep time, as apparent morphological stasis may not reflect the true underlying diversity (Eldredge and Gould 1972; Gould and Eldredge 1977; Hunt 2008; Van Bocxlaer and Hunt 2013). Similarly, for many museum specimens of extant taxa (e.g. those preserved in formalin), it is difficult to acquire the genetic data needed for non-morphological species delimitation methods.

These considerations have sparked interest in whether geometric morphometric analyses can capture fine-scale variation that can be used for identifying cryptic species. This would make the task of identifying and maintaining endangered or conserved groups much easier and could contribute to improved classifications of extinct taxa and populations. Most such studies focus on using morphometrics to discover differences between taxa that were identified by other means (Polly 2003; Zelditch et al. 2004; Gaubert et al. 2005; Gündüz et al. 2007; Polly 2007; Demandt and Bergek 2009). Additionally, there has been work on automated taxon identification and classification of taxa into groups (Baylac et al. 2003; Dobigny et al. 2003; MacLeod 2007; van den Brink and Bokma 2011).

Here, we investigate the morphometric identification of cryptic species using machine learning approaches. In particular, we ask whether it is possible to determine which amongst a set of classification hypotheses is best and examine the implications of the results for a recently proposed set of cryptic turtle species.

### *Background and study system*

Machine learning is an extension of known statistical methodology (Hastie et al. 2009) that emphasizes high predictive accuracy and generality at the expense of the interpretability of individual parameters. The basic statistical mechanics are supplemented by randomization, sorting, and partitioning algorithms, along with the maximization or minimization of summary statistics, in order to best estimate a general model for all data, both sampled and unsampled (Hastie et al. 2009). Machine learning approaches have found use in medical research,

epidemiology, economics and automated image identification such as handwritten zip codes (Hastie et al. 2009). The two major classes of machine learning methods are unsupervised and supervised learning. Unsupervised learning methods are used with unlabeled data where the underlying structure is estimated, and they are analogous to clustering and density estimation methods (Kaufman and Rousseeuw 1990). Supervised learning methods are used with labeled data where the final output of data is known and the rules for going from input to output are inferred. These are analogous to classification and regression models (Breiman et al. 1984). Our application of the approaches used in this study illustrates only a sampling of the various methods available for clustering observations and fitting classification models.

Geometric morphometric approaches to identifying differences in morphological variation between different classes, including cryptic species, mostly have used methods like linear discriminate analysis and canonical variates analysis (Polly 2003; Zelditch et al. 2004; Gaubert et al. 2005; Gündüz et al. 2007; Polly 2007; Francoy et al. 2009; Sztencel-Jabonka et al. 2009; Mitrovski-Bogdanovic et al. 2013). Because of their similarity to multivariate approaches like principle components analysis (PCA), these methods are comparatively straightforward ways of understanding the differences in morphology between classes. They also benefit from producing results that can be easily visualized, which aids in the interpretation and presentation of data and results. Most previous morphometric studies did not assess which amongst a set of alternative classification hypotheses was optimal. For example, studies such as those of Caumul and Polly (2005) and Polly (2007) focused on comparing different aspects of morphology and their fidelity to a classification scheme instead of comparing the fidelity of one aspect of morphology to multiple classification schemes. In this context, the study of Cardini et al. (2009) is noteworthy because they compared morphological variation in marmots at the population, regional, and species level and determined the fidelity of shape to divisions at each of these levels.

Here, we used multiple machine learning methods, both unsupervised and supervised, to compare different classification hypotheses. These methods provide different advantages for understanding how to classify taxa, as well as the accuracy of the resulting classifications. Although machine learning methods such as neural networks have been applied to studying shape variation (Baylac et al. 2003; Dobigny et al. 2003; MacLeod 2007; van den Brink and Bokma 2011), including in the context of automated taxon identification and classification of groups, the number of cases remains limited. In the current study, we not only consider pure classification accuracy but also use a statistic of classification strength that reflects the rate at which taxa are both accurately and inaccurately classified.

We analyzed the problem of whether there are distinct subspecies or cryptic species within the western pond turtle, *Emys marmorata* (Baird and Girard 1852) (formerly *Clemmys marmorata*; see Feldman and Parham 2002). *Emys marmorata* is distributed from northern Washington State, USA to Baja California, Mexico. Traditionally, *E. marmorata* was classified into two named subspecies: the northern *E. marmorata marmorata* and the southern *Emys marmorata pallida* (Seeliger 1945), with a central Californian intergrade zone in between. *Emys marmorata marmorata* is differentiated from *E. marmorata pallida* by the presence of

a pair of triangular inguinal scales and darker neck markings. The triangular inguinal plates can sometimes be present in *E. marmorata pallida* although they are considerably smaller. Seeliger (1945) did not formally include the Baja California populations of *E. marmorata* in either taxon, implying the existence of a third distinct but unnamed subspecies.

Previous work on morphological variation in *E. marmorata* has focused primarily on differentiation between populations over a portion of the species' total range (Lubcke and Wilson 2007; Germano and Rathbun 2008; Germano and Bury 2009; Bury et al. 2010); comparatively few studies have included specimens from across the entire range (Holland 1992). Most of these studies considered how local biotic and abiotic factors may contribute to differences in carapace length and found that size can vary greatly between different populations (Lubcke and Wilson 2007; Germano and Rathbun 2008; Germano and Bury 2009). There also has been interest in size-based sexual dimorphism in *E. marmorata* (Holland 1992; Lubcke and Wilson 2007; Germano and Bury 2009), with males being on average larger than females based on total carapace length and other linear measurements. However, the quality of size as a classifier of sex can vary greatly between populations (Holland 1992) because of the magnitude of size differences among populations (Lubcke and Wilson 2007; Germano and Bury 2009). However, the effect of sexual dimorphism on shape, *sensu* Kendall (1977), has not been assessed (Holland 1992; Lubcke and Wilson 2007; Germano and Rathbun 2008).

Of particular importance in the context of cryptic diversity in *E. marmorata* is the morphometric analysis of carapace shape carried out by Holland (1992), who compared populations of *E. marmorata* from three areas of its range. This study concluded that geographic distance was a poor indicator of morphological differentiation, and instead hypothesized that geographic features such as breaks between different drainage basins are probably more important barriers to dispersal and interbreeding. Additionally, (Holland 1992) suggested that morphological differences were more pronounced as the magnitude of barriers and distance increased, but this variation required many variables to adequately capture, implying only very subtle morphological differentiation between putatively distinct populations. Finally, Holland concluded that *E. marmorata* is best classified as three distinct species: a northern species, southern species, and a Columbia Basin species. This classification is similar to that of Seeliger (1945), except elevated to the species level and without recognition of a distinct Baja species.

More recently, the phylogeography of *E. marmorata* and the possibility of cryptic diversity was investigated using molecular data (Spinks and Shaffer 2005; Spinks et al. 2010, 2014). Based on mitochondrial DNA, Spinks and Shaffer (2005) recognized four subclades within *E. marmorata*, a northern clade, a San Joaquin Valley clade, a Santa Barbara clade, and a southern clade. Analyses with nuclear DNA (Spinks et al. 2010) and single-nucleotide polymorphism (SNP) data suggest a primarily north-south division in *E. marmorata*, although the dataset differed from that of Spinks and Shaffer (2005) in the location of this break point. These studies discussed the potential taxonomic implications of their results, with Spinks et al. (2014) going so far as to strongly advocate for the recognition of at least two species (*E. marmorata* and *E. pallida*), and a possible third based on populations in Baja California. However, they did not discuss in detail the morphological characters that would help to

diagnose these species beyond those specified by Seeliger (1945). Given that these characters are somewhat variable within the proposed species, and that Holland (1992) described shell shape variation that might be consistent with this taxonomy, a geometric morphometric analysis of shell shape might provide a reliable way to diagnose groups (whether species or subspecies) within *E. marmorata*.

In this study, we attempt to estimate the best classification scheme of *E. marmorata* based on variation in plastron (ventral shell) shape in order to determine whether this character is consistent with any of the past divisions based on other morphological features or molecular data. We are particularly interested in whether any aspect of plastron shape can be used to reliably diagnose Spinks et al. (2014) proposed species, and if so, the nature of that shape variation.

Because of unclear geographic boundaries between subgroups of *E. marmorata*, we compare multiple hypotheses of morphologically- and molecularly-based classification. We hypothesize that if morphological variation corresponds to class assignment, it should be possible to determine the best classification hypothesis of *E. marmorata* from amongst multiple candidate hypotheses. However, if morphological variation does not correspond to any of the standing hypothesis, then supervised learning model generalization performance will be poor.

## MATERIALS AND METHODS

### *Specimens, sampling, morphometrics*

We collected landmark-based morphometric data from 354 adult *E. marmorata* museum specimens. These specimens are a subset of those included in Angielczyk and Sheets (2007), Angielczyk et al. (2011), and Angielczyk and Feldman (2013), essentially representing the adult individuals included in those papers. We chose to focus on adults because significant changes in plastron shape occur over the course of ontogeny in *E. marmorata* and other emydines (Angielczyk and Feldman 2013).

We assigned a classification to each specimen for the different binning schemes based on geographic occurrence data recorded in museum collection archives. When precise latitude and longitude information were not available we estimated them from locality information. Because Spinks and Shaffer (2005), Spinks et al. (2010), and Spinks et al. (2014) did not use vouchered specimens we were not able to directly sample individuals in their studies. Therefore our specimen classifications were based solely on the geographic information, not explicit assignment using molecular data. Because the exact barriers between different biogeographic regions are unknown and unclear, we represented each hypothesis with two schemes for a total of six different schemes. The schemes differed based on where geographic boundaries were assigned. This changes the classification of certain individuals near the boundaries between groups, providing a test of the robustness of the classification schemes.

Following previous work on plastron variation (Angielczyk and Sheets 2007; Angielczyk et al. 2011; Angielczyk and Feldman 2013), we used TpsDig 2.04 (Rohlf 2005) to digitize 19

landmarks (Fig. 1). Seventeen of the landmarks are at the endpoints or intersection of the keratinous plastral scutes that cover the plastron. Twelve of the landmarks were symmetrical across the axis of symmetry and, in order to prevent issues related to degrees of freedom and other similar concerns (Klingenberg et al. 2002), we reflected these landmarks across the axis of symmetry (i.e. midline) prior to analysis and used the average position of each symmetrical pair. In cases where damage or incompleteness prevented symmetric landmarks from being determined, we used only the single member of the pair. We conducted all subsequent analyses on the resulting “half” plastron. We superimposed the plastral landmark configurations using generalized Procrustes analysis (Dryden and Mardia 1998), after which, we calculated the principal components (PC) of shape using the **shapes** package for R (R Core Team 2013; Dryden 2013).

### *Machine learning analyses*

*Unsupervised learning.*— In order to preserve the relationship between all landmark configurations in shape space, we measured the dissimilarity between observations using Kendall’s Riemannian shape distance or  $\rho$  (Kendall 1984; Dryden and Mardia 1998). We chose this metric because shape space, or the set of all possible shape configurations following Procrustes superimposition, is a Riemannian manifold and thus non-Euclidean (Dryden and Mardia 1998).  $\rho$  varies between 0 and  $\pi/2$  when there is no reflection invariance, which should not be a concern in the case of the half plastral landmark configurations used in the study.

We divisively clustered the shape dissimilarity matrix using partitioning around medoids clustering (PAM), a method similar to  $k$ -means clustering except that instead of minimizing the sum of squared Euclidean distances between observations and centroids, the sum of squared dissimilarities between observations and medoids is minimized (Kaufman and Rousseeuw 1990). Because the optimal number of clusters of shape configurations in the study was unknown, being possibly three, four, or some other value, we estimated clustering solutions in which the number of clusters varied between one and eight. We compared clustering solutions using the gap statistic, which is a measure of goodness of clustering (Tibshirani et al. 2001).

We conducted this analysis using the **cluster** package for R (Maechler et al. 2013).

*Supervised learning.*— We used three different supervised learning, or classification, approaches: linear discriminate analysis, multinomial logistic regression, and random forests. Linear discriminate analysis, also known as canonical variate analysis, is commonly used in studies of geometric morphometric data (Zelditch et al. 2004; Mitteroecker and Bookstein 2011). The other two methods, however, are not. In all cases, the optimal number of PCs used as predictors was chosen via maximum within-sample AUC value, explained below.

Linear discriminate analysis (LDA) attempts to find a linear combination of predictors that best model two or more classes. LDA is very similar to PCA except that instead of finding the linear combination of features that maximize the amount of explained variance in the

data, LDA maximizes the differences between classes. The results of this analysis produces a transformation matrix by which the original features can be transformed to reflect the best discrimination between the classes. In this study, we applied LDA to the eigenscores from a subset of the total number of PCs, ranging from two to six in increasing order of complexity. In total, this produced nine different LDA scaling matrices.

Multinomial logistic regression is an extension of logistic regression, where instead of a binary response there are three or more response classes (Venables and Ripley 2002). Similar to the odds ratios calculated from the coefficients of a logistic regression, the relative risk of a classification can be determined from the coefficients of the model.

Random forest models are an extension of classification and regression trees (CART) (Breiman et al. 1984; Breiman 2001). The goal of CARTs is to use a series of different features (i.e. predictors) to estimate the class of an observation. In top-down induction of decision trees for each member of a given set of predictor variables, attribute value tests are used to estimate the differences between classes. This process, called recursive partitioning, is then repeated on each subset. The recursion continues until the resulting observations all share the same class or no more meaningful partitions are possible. The resulting model is a tree structure by which observations are classified at each intersection via the estimated cutoff points from the attribute tests made during model fitting.

In a random forest model, many CARTs are built from a random subsample of both the features and the observations (specimens). This process is then repeated many times and the parameters of the final model are chosen as the mode of estimates from the distribution of CARTs (Breiman 2001). In addition to classifying the observations, this procedure allows for the features to be ranked in order of importance. This is a generally useful property for studies in which the goal is to describe and model the differences between classes and the relative importance of different predictors.

In this analysis, we used 1000 subtrees to estimate the random forest model parameters. We estimated the best set of predictors necessary for each classification scheme using a recursive feature selection algorithm, and we chose the optimal number of PCs to include based on the AUC of the model. Following the backwards selection algorithm implemented in `caret` (Kuhn 2013), the maximum number of features were included in the initial model, their importance ranked, and the AUC of the model calculated. The lowest ranked feature was then removed, and the AUC of the model recalculated. This was repeated until only one feature, remained. Because PCs were kept in order of importance and not in relation to the amount of variance each PC described, the PCs are not included in the order of ascending eigenvalue.

In classification studies, such as this one, a common metric of performance is area under the receiver operating characteristic curve (AUC). AUC is an estimate of the relationship between the false positive and true positive rates, as opposed to just the true positive rate (accuracy). This relationship is especially useful in cases such as this study where misclassification needs to be minimized just as much as an accurate classification must be obtained. AUC ranges between 0.5 and 1, with 0.5 indicating classification no better than random and 1 indicating

perfect classification (Hastie et al. 2009).

The standard AUC calculation is defined for binary classifications, however in our application there are multiple categories. The alternative calculation that we used follows an all-against-one strategy where the individual AUC values for each class versus all others are averaged to produce a multiclass AUC (Hand and Till 2001). To estimate confidence intervals on the out-of-sample AUC values, we performed a nonparametric bootstrap in which the true and estimated classifications were resampled with replacement. This was done 1000 times.

The ultimate measure of model fit is accurately predicting the values of unobserved samples (Hastie et al. 2009; Kuhn and Johnson 2013). Within-sample performance is inherently biased upwards, so model evaluation requires overcoming this bias. With very large sample sizes, as in this study, part of the sample can be used as the “training set” and the remainder acts as the “testing set.” The former is used for fitting the model where as the later is used for measuring model performance, a process called model generalization. In this analysis, we used 75% of samples as the training set while the remaining 25% were used as the testing set.

It is common for some out of sample observations to be misclassified. This misclassification may be due to the model not accurately representing shape variance, systematic differences between the training and test sets, or systematic differences between the accurately and inaccurately classified samples. Testing and training sets are determined completely at random within each class and with respect to shape. Results were not effected by the individual specimen class assignments to the testing or training sets.

To determine if there were systematic differences in plastron shape between the correctly and incorrectly classified samples, we used a permutation test to estimate if the dissimilarity between the correctly and incorrectly classified individuals was significantly greater than random. The group labels were permuted 1000 times and the distance between the new centroids was calculated. The number of permutations less than the empirical difference divided by 1000 gives a  $p$ -value for the test. Significant results indicate that correctly and incorrectly classified specimens are systematically different. This was done only for classes where there were 10 or more observations.

### *Comparison with clear-cut example*

In addition to the above analysis of classification schemes of *E. marmorata/pallida*, we applied the above approach to a selection of seven morphologically distinct emydine species. This additional analysis was done in order to confirm the efficacy of our supervised learning approach. The seven species analyzed were *Emys blandingii*, *Terrapene coahuila*, *Clemmys guttata*, *Glyptemys insculpta*, *Glyptemys muhlenbergii*, *Emys orbicularis*, and *Terrapene ornata* with a total of 578 specimens analyzed in total. Again, these data were a subset of those used in Angielczyk et al. (2011) and Angielczyk and Feldman (2013).

As with the *E. marmorata/pallida* analysis, we analyzed plastron shape variation. The support for a seven category classification scheme was evaluated using random forest, linear



discriminate analysis, and multinomial logistic regression where the number of PCs used as predictors ranged from two to 11. Data were split into training and testing sets and model performance for both sets was evaluated using the AUC metric.

## RESULTS

### *Unsupervised learning*

Comparison of gap statistic values from PAM clustering show that the optimal, minimal number of clusters is most likely one (Fig. 2). There is some ambiguity in choice because, although it is not statistically different from a solution with only one group, the solution with two groupings does have the greatest mean gap statistic. However, there is no geographical signal in the results of this clustering solution (Fig. 3). Because of this, we assert that there is no means of naturally partitioning plastron shape into distinct subgroups with out reference to external information.

### *Supervised learning*

AUC-based model selection revealed some important patterns of variation and congruence between the classification schemes and the actual data. Generally, the best performing models tended to include as many PCs as possible 4). Note that the best random forest models were determined via recursive feature selection, so PCs were not included in order of percent variance explained. That almost all LDA and multinomial logistic regression models were as complex as possible indicates that the differences between the different groups within each classification scheme are very small.

As part of fitting a random forest model, a ranking of variable importance also is determined. Interestingly, the order of variable importance is not the same as the order of the PCs (5). This means that the variance describing the differences between the classes does not align with the major axes of variance (i.e. the PCs). Such a result is expected if variation between classes was extremely fine grained and not a part of the principal form or function of the plastron. Moreover, this result is consistent with the results from the AUC-based model selection for the multinomial logistic regression and LDA models.

Observed AUC values for all of the optimal models are not exceptionally high (4). In most cases the different proposed classification schemes are generally poor descriptors of the observed variation. It appears that the data set is overwhelmed by noise, making any accurate classifications difficult at best. This observation is cemented with the generalizations of the models to the testing data set (6).

Mean AUC values for the model generalizations, in most cases, are approximately equal to the observed AUC values from the training data set (Table 1). The cases in which the AUC from the generalizations is less than the observed, indicate poor model fit and a poor classification scheme. AUC values from model generalization, or estimating testing data set

membership, does not indicate a clear “best” classification scheme (Fig. 6). Although the scheme with two species has the greatest AUC point estimate for each modeling approach, this scheme is not significantly greater than any other except in some limited cases (e.g. LDA, Table 2). Differences in mean shape between correctly and incorrectly classified observations from test set frequently were statistically significant, though there are exceptions. Again, this test was to determine if the mean shapes were statistically different or not. The frequency of these results, however, is important because it means that the different models are poor predictors of class membership. This may be because differences in plastron shape do not align with the any of the hypothesized classification schemes.

### *Comparison with clear-cut example*

In contrast with the above results, the analysis of the seven morphologically-distinct species demonstrates near perfect classification of the both the in-sample and out-of-sample datasets (Table 4). Additionally, the distribution of ROC scores from 1000 bootstrap replicates are tightly clustered near  $\text{ROC} = 1$  (Fig. 7), which is contrast to the results from the *E. marmorata* case (Fig. 6). These results demonstrate that when there are strong distinctions between the states of the classification schemes, the methods used here can recover them.

## DISCUSSION

The results of this study indicate that there is no clear grouping of *E. marmorata* based on plastron shape.

The unsupervised learning results indicate only a single group of observations being optimal, which is consistent with the results from the generalizations of the supervised learning models. The classification schemes used in the supervised learning models correspond, loosely, to unsupervised learning solutions with multiple groups. Because unsupervised learning solutions with multiple groups are poor descriptors of the observed variation, it is important to see this generally supported by the supervised learning results.

The results from fitting the various supervised learning models to each of the classification schemes generally shows that no one scheme is “best.” Possible explanations include that the genetic differentiation is not associated with plastron shape variation and/or that local selective pressures (e.g. from hydrological regime) overwhelm morphological differentiation. This makes sense given that the plastron is involved in both protection and hydrodynamics, and not necessary mate choice (Rivera 2008; Rivera and Stayton 2011; Stayton 2011; Rivera et al. 2014) and that shell shape in *E. marmorata* is known to vary among populations inhabiting water bodies with different flow regimes (Holland 1992; Lubcke and Wilson 2007; Germano and Bury 2009). Plastron shape does not seem to preserve a strong phylogenetic signal at the interspecific level in emydine turtles (Angielczyk et al. 2011), and our current results suggest that this may be the case for phylogeographic signal within emydine species as well. A final possibility (explored below) is that the proposed classification schemes themselves do not represent significant evolutionary lineages.

Both the low AUC values ( $< 0.9$ ) and the significant difference between the correctly and incorrectly classified observations support the conclusion that none of the hypothesized classification schemes are good descriptors of the observed plastral variation within *E. marmorata*.

Nevertheless, it is important to note that plastron shape is an extremely effective method for differentiating members of the other seven species we investigated. The magnitude of shape differences between the species (measured as Procrustes distance between species' mean shapes) is approximately an order of magnitude greater than the differences between the *E. marmorata* subgroups, and the machine learning methods had no trouble accurately classifying the specimens correctly. These results demonstrate that plastron shape is normally a good marker for species delineation in the closest relatives of *E. marmorata*, and that our lack of results for *E. marmorata* is not simply a shortcoming of the methods we applied. Indeed, they beg the question of what factors have suppressed morphological differentiation of plastron shape in *E. marmorata* and *E. pallida* if they are distinct species. Invoking issues such as the role of the plastron in protection or the need for streamlining are insufficient because the other species are expected to be subject to similar constraints (Stayton 2011; Polly et al.).

### *Is there more than one species of Western Pond Turtle?*

The lack of morphological support for the distinctiveness of *E. pallida* does not, on its own, preclude the recognition of this taxon. However, this apparent lack of congruence does prompt a reexamination of the methods and concepts that led to that taxonomic revision, especially considering that plastron shape is normally a strong indicator of species differences among emydines. In other words, before we can assess the significance of the morphological non-diagnosability, it is essential to evaluate the methods and concepts that led to the initial taxonomic revision.

Spinks et al. (2014) elevated *E. pallida* based on a Bayesian species delimitation analysis of SNP data using BPP (Yang and Rannala 2010). However, Spinks et al. (2014) did not heed the caveats about species delimitation methods raised by Carstens et al. (2013). In addition to specifically addressing the shortcomings of validation methods such as BPP that rely on guide trees and “should be interpreted with caution,” Carstens et al. (2013) also strongly emphasize that “Inferences regarding species boundaries based on genetic data alone are likely inadequate, and species delimitation should be conducted with consideration of the life history, geographical distribution, morphology and behaviour (where applicable) of the focal system. . . .” These caveats evoke the development of the Unified Species Concept (Dayrat 2005; De Queiroz 2007), Integrative Taxonomy (Padial et al. 2010), and other pluralist approaches to species delimitation. None of these considerations were brought to bear on the *E. marmorata* system until now, and in doing so we find the proposal that *E. pallida* is a distinct species to be lacking in a normally robust morphological marker.

The natural history and geographical distribution of *E. marmorata* and *E. pallida* also make

the recognition of these taxa implausible. The data from Spinks et al. (2014) show extensive introgression and admixture in Central California, which is expected because there are no significant barriers to gene flow in this region. Combined with the well-demonstrated ability for testudinoid turtles, including emydids and even *Emys*, to hybridize (e.g. Buskirk et al. 2005; Spinks and Shaffer 2009; Parham et al. 2013) it is hard to imagine how *E. marmorata* and *E. pallida* could maintain their integrity in the face of such admixture. Because the geography, natural history, demonstrated genetic admixture of *E. marmorata*, and comparisons with other morphologically-diagnosable species conflict with the recognition of *E. pallida*, we hypothesize that our inability to classify the morphological data by proposed species is because *E. pallida* is not a distinct species. We agree with Carstens et al. (2013) that “the inferences drawn from species delimitation studies should be conservative, for in most contexts it is better to fail to delimit species than it is to falsely delimit entities that do not represent actual evolutionary lineages.”

We fully agree with Spinks et al. (2014) that *E. marmorata* (*sensu lato*) is a species deserving of strong conservation efforts, and we do not wish to trivialize this need. Moreover, the genetic diversity uncovered by the analysis of Spinks et al. (2014) should be explicitly accounted for in any conservation plan. Given the apparent lack of morphological distinction, however, we consider that this diversity should be considered Evolutionary Significant Units or Distinct Population Segments instead of distinct species.

Finally, it is important to note that the data and analyses we present do not let us definitively say whether the apparent lack of morphological divergence within *E. marmorata* truly reflects the presence of a single species, or if it is an artifact of plastron shape being a poor morphological marker for phylogenetic and phylogeographic divergences, because we could not carry out our morphometric analyses on the specimens from which the genetic data were obtained. The comparisons with the other emydine species, suggests it is because *E. marmorata* is a single species. However, tests of both our preferred conclusion (*E. marmorata* as a single species) and that Spinks et al. (2014) should include morphological and molecular analyses of the same set of voucher specimens, as well as additional tests of species delimitation using alternative methods and corroborating evidence as suggested by Carstens et al. (2013). From a morphological standpoint, support for the validity of “*E. Pallida*” may come from other aspects of morphology, such as carapace shape or other features. Likewise, further investigation of the phylogeographic utility of plastron shape in other turtle species will help to clarify whether the lack of differentiation seen in *E. marmoarata*, and the strong differentiation among the other emydines, is typical or an unusual case.

\*

Acknowledgements Data collection for this project was supported in part by NSF DBI-0306158 (to KDA). G. Miller assisted with data collection and her participation in this research was supported by NSF REU DBI-0353797 (to R. Mooi of CAS). For access to emydine specimens, we thank: J. Vindum and R. Drewes (CAS); A. Resetar (FMNH); R. Feeney (LACM); C.

<sup>446</sup> Austin (LSUMNS); S. Sweet (MSE); J. McGuire and C. Conroy (MVZ); A. Wynn (NMNH);  
P. Collins (SBMNH); B. Hollingsworth (SDMNH); P. Holroyd (UCMP). We are grateful for  
<sup>448</sup> S. Sweet for field assistance and the California Department of Fish and Game for permits.

## BIBLIOGRAPHY

- 450 Angielczyk, K. D. and C. R. Feldman. 2013. Are diminutive turtles miniaturized? The  
ontogeny of plastron shape in emydine turtles. *Biological Journal of the Linnean Society*  
452 108:727–755.
- Angielczyk, K. D., C. R. Feldman, and G. R. Miller. 2011. Adaptive evolution of plastron  
454 shape in emydine turtles. *Evolution* 65:377–394.
- Angielczyk, K. D. and H. D. Sheets. 2007. Investigation of simulated tectonic deformation in  
456 fossils using geometric morphometrics. *Paleobiology* 33:125–148.
- Baird, S. F. and C. Girard. 1852. Descriptions of new species of reptiles collected by the U.S.  
458 Exploring Expedition under the command of Capt. Charles Wilkes. *Proceedings of the  
National Academy of Sciences Philadelphia* 6:174–177.
- 460 Bauer, A. M., J. F. Parham, R. M. Brown, B. L. Stuart, L. Grismer, T. J. Papenfuss,  
W. Bohme, J. M. Savage, S. Carranza, J. L. Grismer, P. Wagner, A. Schmitz, N. B.  
462 Ananjeva, and R. F. Inger. 2000. Availability of new Bayesian-delimited gecko names and  
the importance of character-based species descriptions. *Proceedings of the Royal Society B:  
464 Biological Sciences* 278:490–492.
- Baylac, M., C. Villemant, and G. Simbolotti. 2003. Combining geometric morphometrics  
466 with pattern recognition for the investigation of species complexes. *Biological Journal of  
the Linnean Society* 80:89–98.
- 468 Bickford, D., D. J. Lohman, N. S. Sodhi, P. K. L. Ng, R. Meier, K. Winker, K. K. Ingram,  
and I. Das. 2007. Cryptic species as a window on diversity and conservation. *Trends in  
470 ecology & evolution* 22:148–55.
- Breiman, L. 2001. Random Forests. *Machine Learning* 45:5–32.
- 472 Breiman, L., J. Friedman, C. J. Stone, and R. A. Olshen. 1984. Classification and regression  
trees. Wadsworth International Group, Belmont.
- 474 Bury, R. B., D. J. Germano, and G. W. Bury. 2010. Population Structure and Growth of the  
Turtle *Actinemys marmorata* from the KlamathSiskiyou Ecoregion: Age, Not Size, Matters.  
476 *Copeia* 2010:443–451.
- Buskirk, S. W., J. F. Parham, and C. R. Feldman. 2005. On the hybridisation between two  
478 distantly related Asian turtles (Testudines: *Scalia* x *Mauremys*). *Salamandra* 41:21–26.
- Cardini, A., D. Nagorsen, P. O’Higgins, P. D. Polly, R. W. Thorington Jr, and P. Tongiorgi.  
480 2009. Detecting biological distinctiveness using geometric morphometrics: an example case  
from the Vancouver Island marmot. *Ethology Ecology & Evolution* 21:209–223.
- 482 Carstens, B. C. and T. A. Dewey. 2010. Species Delimitation Using a Combined Coalescent  
and Information-Theoretic Approach: An Example from North American *Myotis* Bats.  
484 *Systematic Biology* 59:400–414.

Carstens, B. C., T. a. Pelletier, N. M. Reid, and J. D. Satler. 2013. How to fail at species  
486 delimitation. *Molecular ecology* 22:4369–83.

Caumul, R. and P. D. Polly. 2005. Phylogenetic and environmental components of mor-  
488 phological variation: skull, mandible, and molar shape in marmots (*Marmota*, Rodentia).  
*Evolution; international journal of organic evolution* 59:2460–72.

490 Clare, E. L. 2011. Cryptic species? Patterns of maternal and paternal gene flow in eight  
neotropical bats. *PloS one* 6:e21460.

492 Dayrat, B. 2005. Towards integrative taxonomy. *Biological Journal of the Linnean Society*  
85:407–415.

494 De Queiroz, K. 2007. Species concepts and species delimitation. *Systematic Biology* 56:879–86.

Demandt, M. H. and S. Bergek. 2009. Identification of cyprinid hybrids by using geometric  
496 morphometrics and microsatellites. *Journal of Applied Ichthyology* 25:695–701.

Dobigny, G., L. Granjon, V. Aniskin, K. Ba, and V. Voloboulev. 2003. A new sigling species  
498 of *Taterillus* (Muridae, Gerbillinae) from West Africa. *Mammalian Biology* 68:299–316.

Dryden, I. L. 2013. shapes: Statistical shape analysis. R package version 1.1-8.

500 Dryden, I. L. and K. Y. Mardia. 1998. Statistical shape analysis. Wiley, New York.

Eldredge, N. and S. J. Gould. 1972. Punctuated equilibria: an alternative to phyletic gradual-  
502 ism. Pages 82–115 *in* *Models in Paleobiology* (T. J. M. Schopf, ed.). Freeman Cooper, San  
Francisco.

504 Feldman, C. R. and J. F. Parham. 2002. Molecular phylogenetics of emydine turtles: taxonomic  
revision and the evolution of shell kinesis. *Molecular Phylogenetics and Evolution* 22:388–98.

506 Francoy, T. M., R. A. O. Silva, P. Nunes-Silva, C. Menezes, and V. L. Imperatriz-Fonseca.  
2009. Gender identification of five genera of stingless bees (Apidae, Meliponini) based on  
508 wing morphology. *Genetics and molecular research* 8:207–214.

Funk, W. C., M. Caminer, and S. R. Ron. 2012. High levels of cryptic species diversity  
510 uncovered in Amazonian frogs. *Proceedings of the Royal Society B: Biological Sciences*  
279:1806–14.

512 Gaubert, P., P. J. Taylor, C. a. Fernandes, M. W. Bruford, and G. Veron. 2005. Patterns  
of cryptic hybridization revealed using an integrative approach: a case study on genets  
514 (*Carnivora*, *Viverridae*, *Genetta* spp.) from the southern African subregion. *Biological*  
*Journal of the Linnean Society* 86:11–33.

516 Germano, D. J. and R. B. Bury. 2009. Variation in body size, growth, and population structure  
of *Actinemys marmorata* from lentic and lotic habitats in Southern Oregon. *Journal of*  
518 *Herpetology* 43:510–520.

Germano, D. J. and G. B. Rathbun. 2008. Growth, population structure, and reproduction of western pond turtles (*Actinemys marmorata*) on the Central Coast of California. *Chelonian Conservation and Biology* 7:188–194.

Gould, S. J. and N. Eldredge. 1977. Punctuated equilibria: the tempo and mode of evolution reconsidered. *Paleobiology* 3:115–151.

Gündüz, I., M. Jaarola, C. Tez, C. Yeniyurt, P. D. Polly, and J. B. Searle. 2007. Multigenic and morphometric differentiation of ground squirrels (*Spermophilus*, *Scuiridae*, *Rodentia*) in Turkey, with a description of a new species. *Molecular phylogenetics and evolution* 43:916–35.

Hand, D. J. and R. J. Till. 2001. A Simple Generalisation of the Area Under the ROC Curve for Multiple Class Classification Problems. *Machine Learning* 45:171–186.

Hastie, T., R. Tibshirani, and J. Friedman. 2009. The elements of statistical learning: data mining, inference, and prediction. 2nd ed. Springer, New York.

Hausdorf, B. and C. Hennig. 2010. Species delimitation using dominant and codominant multilocus markers. *Systematic biology* 59:491–503.

Holland, D. C. 1992. Level and pattern in morphological variation: a phylogeographic study of the western pond turtle (*Clemmys marmorata*). Ph.D. thesis University of Southwestern Louisiana.

Huelsenbeck, J. P., P. Andolfatto, and E. T. Huelsenbeck. 2011. Structurama: bayesian inference of population structure. *Evolutionary bioinformatics online* 7:55–9.

Hunt, G. 2008. Gradual or pulsed evolution: when should punctuational explanations be preferred? *Paleobiology* 34:360–377.

Kaufman, L. and P. J. Rousseeuw. 1990. Finding groups in data : an introduction to cluster analysis. Wiley, New York.

Kendall, D. G. 1977. The diffusion of shape. *Advances in Applied Probability* 9:428–430.

Kendall, D. G. 1984. Shape Manifolds, Procrustean Metrics, and Complex Projective Spaces. *Bulletin of the London Mathematical Society* 16:81–121.

Klingenberg, C. P., M. Barluenga, and A. Meyer. 2002. Shape analysis of symmetric structures: quantifying variation among individuals and asymmetry. *Evolution* 56:1909–1920.

Kuhn, M. 2013. caret: Classification and Regression Training. R package version 5.15-61.

Kuhn, M. and K. Johnson. 2013. Applied predictive modeling. Springer, New York, NY.

Leaché, A. D. and M. K. Fujita. 2010. Bayesian species delimitation in West African forest geckos (*Hemidactylus fasciatus*). *Proceedings. Biological sciences / The Royal Society* 277:3071–7.



- 554 Lubcke, G. M. and D. S. Wilson. 2007. Variation in shell morphology of the Western Pond  
Turtle (*Actinemys marmorata* Baird and Giarard) from three aquatic habitats in Northern  
California. *Journal of Herpetology* 41:107–114.
- 556 MacLeod, N. 2007. Automated taxon identification in systematics: theory, approaches and  
applications. CRC Press, Boca Raton.
- 558 Maechler, M., P. Rousseeuw, A. Struyf, M. Hubert, and K. Hornik. 2013. cluster: Cluster  
Analysis Basics and Extensions. R package version 1.14.4.
- 560 Mitrovski-Bogdanovic, A., A. Petrovic, M. Mitrovic, A. Ivanovic, V. Žikic, P. Starý, C. Vor-  
burger, and v. Tomanovic. 2013. Identification of two cryptic species within the *Praon*  
562 *abjectum* group (Hymenoptera: Braconidae: Aphidiinae) using molecular markers and  
geometric morphometrics. *Annals of the entomological society of America* 106:170–180.
- 564 Mitteroecker, P. and F. Bookstein. 2011. Linear Discrimination, Ordination, and the Visual-  
ization of Selection Gradients in Modern Morphometrics. *Evolutionary Biology* 38:100–114.
- 566 O'Meara, B. C. 2010. New heuristic methods for joint species delimitation and species tree  
inference. *Systematic biology* 59:59–73.
- 568 Padial, J. M., A. Miralles, I. De la Riva, and M. Vences. 2010. The integrative future of  
taxonomy. *Frontiers in Zoology* 7:1–14.
- 570 Parham, J. F., T. J. Papenfuss, P. P. V. Dijk, B. S. Wilson, C. Marte, L. R. Schettino, and  
W. Brian Simison. 2013. Genetic introgression and hybridization in Antillean freshwater  
572 turtles (*Trachemys*) revealed by coalescent analyses of mitochondrial and cloned nuclear  
markers. *Molecular phylogenetics and evolution* 67:176–87.
- 574 Pfenninger, M. and K. Schwenk. 2007. Cryptic animal species are homogeneously distributed  
among taxa and biogeographical regions. *BMC evolutionary biology* 7:121.
- 576 Polly, P. D. 2003. Paleophylogeography of *Sorex araneus*: molar shape as a morphological  
marker for fossil shrews. *Mammalia* 68:233–243.
- 578 Polly, P. D. 2007. Phylogeographic differentiation in *Sorex araneus*: morphology in relation  
to geography and karyotype. *Russian Journal of Theriology* 6:73–84.
- 580 Polly, P. D., C. T. Stayton, E. R. Dumont, S. E. Pierce, E. J. Rayfield, and K. D. Angielczyk.  
???? Combining geometric morphometrics and finite element analysis with evolutionary  
582 modeling: towards a synthesis. *Journal of Vertebrate Paleontology* .
- 584 Pons, J., T. Barraclough, J. Gomez-Zurita, A. Cardoso, D. Duran, S. Hazell, S. Kamoun,  
W. Sumlin, and A. Vogler. 2006. Sequence-Based Species Delimitation for the DNA  
Taxonomy of Undescribed Insects. *Systematic Biology* 55:595–609.
- 586 R Core Team. 2013. R: A Language and Environment for Statistical Computing. R Foundation  
for Statistical Computing Vienna, Austria.

- 588 Rivera, G. 2008. Ecomorphological variation in shell shape of the freshwater turtle *Pseudemys*  
 concinna inhabiting different aquatic flow regimes. *Integrative and comparative biology*  
 590 48:769–87.
- Rivera, G., J. N. Davis, J. C. Godwin, and D. C. Adams. 2014. Repeatability of Habitat-  
 592 Associated Divergence in Shell Shape of Turtles. *Evolutionary Biology* Pages 29–37.
- Rivera, G. and C. T. Stayton. 2011. Finite element modeling of shell shape in the fresh-  
 594 water turtle *Pseudemys concinna* reveals a trade-off between mechanical strength and  
 hydrodynamic efficiency. *Journal of morphology* 272:1192–203.
- 596 Rohlf, F. J. 2005. TpsDig 2.04.
- Schilck-Steiner, B. C., B. Seifert, C. Stauffer, E. Christian, R. H. Crozier, and F. M. Steiner.  
 598 2007. Without morphology, cryptic species stay in taxonomic crypsis following discovery.  
*Trends in ecology & evolution* 22:391–392.
- 600 Seeliger, L. M. 1945. Variation in the Pacific Mud Turtle. *Copeia* 1945:150–159.
- Spinks, P. Q. and H. B. Shaffer. 2005. Range-wide molecular analysis of the western pond  
 602 turtle (*Emys marmorata*): cryptic variation, isolation by distance, and their conservation  
 implications. *Molecular ecology* 14:2047–64.
- 604 Spinks, P. Q. and H. B. Shaffer. 2009. Conflicting mitochondrial and nuclear phylogenies for  
 the widely disjunct *Emys* (Testudines: Emydidae) species complex, and what they tell us  
 606 about biogeography and hybridization. *Systematic biology* 58:1–20.
- Spinks, P. Q., R. C. Thomson, and H. Bradley Shaffer. 2014. The advantages of going large:  
 608 genome wide SNPs clarify the complex population history and systematics of the threatened  
 western pond turtle. *Molecular Ecology* Pages n/a–n/a.
- 610 Spinks, P. Q., R. C. Thomson, and H. B. Shaffer. 2010. Nuclear gene phylogeography reveals  
 the historical legacy of an ancient inland sea on lineages of the western pond turtle, *Emys*  
 612 *marmorata* in California. *Molecular ecology* 19:542–56.
- Stayton, C. T. 2011. Biomechanics on the half shell: functional performance influences patterns  
 614 of morphological variation in the emydid turtle carapace. *Zoology (Jena, Germany)* 114:213–  
 23.
- 616 Stuart, B. L., R. F. Inger, and H. K. Voris. 2006. High level of cryptic species diversity  
 revealed by sympatric lineages of Southeast Asian forest frogs. *Biology letters* 2:470–4.
- 618 Sztencel-Jabonka, A., G. Jones, and W. Bogdanowicz. 2009. Skull Morphology of Two Cryptic  
 Bat Species: *Pipistrellus pipistrellus* and *P. pygmaeus* A 3D Geometric Morphometrics  
 620 Approach with Landmark Reconstruction. *Acta Chiropterologica* 11:113–126.
- Tibshirani, R., G. Walther, and T. Hastie. 2001. Estimating the number of clusters in a  
 622 data set via the gap statistic. *Journal of the Royal Statistical Society: Series B (Statistical  
 Methodology)* 63:411–423.

- 624 Van Bocxlaer, B. and G. Hunt. 2013. Morphological stasis in an ongoing gastropod radiation  
from Lake Malawi. *Proceedings of the National Academy of Sciences* .
- 626 van den Brink, V. and F. Bokma. 2011. Morphometric shape analysis using learning vector  
quantization neural networks an example distinguishing two microtine vole species. *Annales*  
628 *Zoologici Fennici* 48:359–364.
- Venables, W. and B. D. Ripley. 2002. *Modern applied statistics with S*. 4th ed. Springer, New  
630 York.
- Yang, Z. and B. Rannala. 2010. Bayesian species delimitation using multilocus sequence data.  
632 *Proceedings of the National Academy of Sciences* 107:9264–9.
- Zelditch, M. L., D. L. Swiderski, and H. D. Sheets. 2004. *Geometric morphometrics for*  
634 *biologists: a primer*. Elsevier Academic Press, Amsterdam.

Table 1: AUC values for the best model of each classification scheme for both the observed (training) data and the generalized (testing) data. Results from all three different supervised learning approaches are shown here. AUC values range between 0.5 and 1.

Scheme	random forest		multinomial logistic regression		linear discriminate analysis	
	Observed	Generalized	Observed	Generalized	Observed	Generalized
Morph 1	0.63	0.73	0.75	0.79	0.75	0.80
Morph 2	0.61	0.58	0.76	0.77	0.76	0.77
Mito 1	0.63	0.62	0.75	0.63	0.75	0.63
Mito 2	0.77	0.67	0.80	0.64	0.80	0.63
Mito 3	0.56	0.64	0.71	0.74	0.71	0.73
Nuclear	0.56	0.67	0.74	0.62	0.74	0.77

Table 2: Results of bootstrap comparisons between the scheme with the highest mean AUC value and all other schemes, defined  $P(\text{best} - \text{other} > 0)$ . An asterix indicates the best scheme. This was done for each of the three modeling techniques included in this study: random forest (RF), multinomial logistic regression (MLR), and linear discriminate analysis (LDA). Probabilities are the percent of comparisons that are greater than the observed difference in means.

Scheme	RF	MLR	LDA
Morph 1	*	*	1
Morph 2	0.79	0.55	1
Mito 1	0.89	0.94	1
Mito 2	0.82	0.57	*
Mito 3	0.79	0.69	0.73
Nuclear	0.79	0.96	0.96

Table 3: Results of comparisons between correctly and incorrectly classified observations from the testing data set, defined  $P(\text{distance} - \text{simulated} > 0)$ . For each scheme, the classifications with at least 10 observations were tested. This was done for each of the three modeling techniques included in this study: random forest (RF), multinomial logistic regression (MLR), and linear discriminate analysis (LDA).

Scheme	Class	RF	MLR	LDA
Morph 1	CCR	0.77	1.00	1.00
	marm	0.87	0.93	1.00
Morph 2	CCR	0.88	1.00	1.00
	marm	1.00	1.00	0.98
Mito 1	CCR	0.94	0.99	0.98
	marm	0.99	0.98	0.47
Mito 2	marm	0.85	0.97	0.96
	pall	0.94	1.00	1.00
Mito 3	1	0.40	0.92	0.90
	3	0.97	0.99	1.00
Nuclear	marm	1.00	1.00	1.00
	pall	0.99	1.00	1.00

Table 4: Results from classification model estimates of the secondary, multi-species dataset. Models are random forest (RF), linear discriminate analysis (LDA), and multinomial logistic regression (MLR). Comparison of in-sample and out-of-sample AUC of the best performing model, along with the number of predictors. AUC values range between 0.5 (no better than random) and 1 (perfect classification).

	# predictors	In-sample AUC	Out-of-sample AUC
RF	11	1.000	0.999
MLR	6	1.000	0.998
LDA	10	1.000	1.000




Figure 1: Depiction of general plastral shape of *E. marmorata* and position of the 19 landmark used in this study. Anterior is towards the top of the figure.




Figure 2: Results from PAM clustering of the Riemannian shape distance for 8 different number of clusters. Vertical lines are 1 standard deviation of the mean determined from 500 resamples.




Figure 3: Comparison of geographic distribution of clustered observations from the 2 clustering PAM solution. Colour and shape correspond to each of the groups. There is clearly no geographic signal in the data.






Figure 4: Graphical representation of the AUC values from model selection for multinomial logistic regression and linear discriminate analysis, respectively. AUC model selection is based on greatest AUC value. The horizontal axis corresponds to the cumulative number of axes included in the model of interest. A highlighted point corresponds to the AUC best model for that classification scheme.




Figure 5: Variable importance from the random forest models for each of the six classification schemes. Importance is measured as the mean decrease in Gini Index, which is a measure of the strength by which that variable determines CART structure. Indices that are farther to the right indicate greater variable importance.




Figure 6: Bootstrap distributions for generalized AUC values for each of the classification schemes. Each row corresponds to a different modeling approach: multinomial logistic regression (MLR), random forest (RF), linear discriminate analysis (LDA), and LDA using best variables from random forest. Each distribution corresponds to 1000 bootstrap replicates.




Figure 7: Distribution of 1000 bootstrap replicates of the out-of-sample AUC values from the secondary, multi-species dataset. The AUC values are from the random forest (RF), linear discriminate analysis (LDA), and multinomial logistic regression (MLR). AUC values range between 0.5 (no better than random) and 1 (perfect classification).