

Submission 1

Philipp Stärk

April 27, 2021

Problem 1: Self avoiding polymer model:

a) Probability distribution of 4 monomer chain:

Possible values for chain length (found by Drawing): $|\mathbf{R}_3 - \mathbf{R}_0| = 1, \sqrt{5}, 3$.

Probabilities, determined by enumeration and symmetry arguments:

p	Value
$p(\mathbf{R}_3 - \mathbf{R}_0 = 1)$	$\frac{2}{9}$
$p(\mathbf{R}_3 - \mathbf{R}_0 = \sqrt{5})$	$\frac{6}{9}$
$p(\mathbf{R}_3 - \mathbf{R}_0 = 3)$	$\frac{1}{9}$

The probability to find the chain in a straight line is that of $p(|\mathbf{R}_3 - \mathbf{R}_0| = 3) = \frac{1}{9}$.

b) Mean:

$$\langle |\mathbf{R}_3 - \mathbf{R}_0| \rangle = \frac{2}{9} + \sqrt{5} \frac{6}{9} + 3 \frac{1}{9} \approx 2.04$$

Variance:

$$\sigma = \sqrt{\langle |\mathbf{R}_3 - \mathbf{R}_0|^2 \rangle} = \sqrt{\frac{2}{9} + 5 \frac{6}{9} + 9 \frac{1}{9}} \approx 1.91$$

Problem 2: Random walk on a lattice without self-avoidance:

a) Probability distribution $p(X, Y|N)$:

Start with considering random walk in x -direction, assume $N_x = r + l$ steps in this direction, with r the steps in positive and l steps in negative direction, i.e. $X = r - l$.

$$\implies r - l = x, N = r + l \implies r = \frac{N_x + x}{2}, l = \frac{N_x - x}{2}.$$

Thus, we have

$$p(X|N_x) = \binom{N_x}{\frac{N_x + x}{2}} p^{N_x}$$

in one direction (of course analogously in y -direction), with $p = \frac{1}{2}$.

To get the probability $p(X, Y|N)$, we need to sum over all combinations of possible number of path-steps in each direction, where we know:

$$\begin{aligned} N &= N_x + N_y, \\ \forall N_x &\in \{X, \dots, N - Y\}, \\ \forall N_y &\in \{Y, \dots, N - X\}. \end{aligned}$$

Thus, overall, we have:

$$p(X, Y|N) = \sum_{N_x=X}^{N-Y} \binom{N_x}{\frac{N_x+X}{2}} \binom{N-N_x}{\frac{N-N_x+Y}{2}} \frac{1}{4}^N.$$