

Velocity circulation statistics in counterflow turbulence

P. Z. Stasiak

*School of Mathematics, Statistics and Physics, Newcastle University,
Newcastle upon Tyne, NE1 7RU, United Kingdom*

G. Krstulovic

*Université Côte d'Azur, Observatoire de la Côte d'Azur, CNRS, Laboratoire Lagrange,
Boulevard de l'Observatoire CS 34229 - F 06304 NICE Cedex 4, France*

L. Galantucci

Istituto per le Applicazioni del Calcolo "M. Picone" IAC CNR, Via dei Taurini 19, 00185 Roma, Italy

(Dated: February 1, 2026)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

INTRODUCTION

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

METHOD

Superfluid dynamics are modelled by employing the recently developed FOUCAULT model [1]. We follow Schwarz's approach [2], by exploiting the large separation of scales between the ^4He vortex core size a_0 and the average inter-vortex spacing ℓ . We parametrize the superfluid vortex lines as 1D space curves $\mathbf{s}(\xi, t)$ with ξ and t the arclength and time respectively. The corresponding equation of motion is

$$\dot{\mathbf{s}}(\xi, t) = \mathbf{v}_{s\perp} + \beta \mathbf{s}' \times (\mathbf{v}_n - \mathbf{v}_s) + \beta' \mathbf{s}' \times [\mathbf{s}' \times (\mathbf{v}_n - \mathbf{v}_s)]$$

where $\mathbf{s}' = \partial \mathbf{s} / \partial \xi$ is the unit tangent vector at \mathbf{s} , \mathbf{v}_n and \mathbf{v}_s are the normal fluid and superfluid velocities at \mathbf{s} , and β , β' are temperature and Reynolds number dependent mutual friction coefficients. The superfluid velocity \mathbf{v}_s is computed via a de-singularized Biot-Savart integral, accelerated using the tree algorithm [3] (see Supplementary Materials).

RESULTS

-
- [1] L. Galantucci, A. W. Baggaley, C. F. Barenghi, and G. Krstulovic, A new self-consistent approach of quantum turbulence in superfluid helium, *Eur. Phys. J. Plus* **135**, 547 (2020).
 - [2] K.W. Schwarz, Three-dimensional vortex dynamics in superfluid ^4He , *Phys. Rev. B* **38**, 2398 (1988).
 - [3] A. W. Baggaley and C. F. Barenghi, Tree Method for Quantum Vortex Dynamics, *J. Low Temp. Phys.* **166**, 3 (2012).




FIG. 1: Visualisation of turbulent vortex tangles. (a) Counterflow-induced turbulence generated at $T = 2.1$ K with a counterflow velocity of $v_{ns} = \bar{v}_n - \bar{v}_s = 0.94$ cm/s. The blue and red surface rendering shows the normalised kinetic energy density of normal fluid fluctuations E'_n/E'_{max} where $E'_n = |\mathbf{v}_n - \bar{\mathbf{v}}_n|^2$. (b) Turbulence generated by an initial Taylor-Green configuration (see Supplementary Materials for details). In both (a) and (b) superfluid vortex lines are represented as green tubes with exaggerated size.




FIG. 2: Circulation variance $\langle |\Gamma|^2 \rangle$ around square loops of size r . The blue and yellow curves show the normal fluid and super fluid simulations respectively, which are subdivided into the perpendicular (circles) and parallel (diamonds) components. The green squares show the Taylor-Green simulation.




FIG. 3: PDFs of the velocity circulation for different loop sizes $r/\ell > 1$, scaled by the standard deviation σ_Γ . (a) shows the Taylor-Green simulation, while (b) shows both of the normal fluid parallel (diamonds) and perpendicular (circles) components. The inset of (a) shows the flatness $\mathcal{K}(\Gamma) = \langle |\Gamma|^4 \rangle / \sigma_\Gamma^4$, with the same color scheme as in Fig. 2. The black dashed line marks the Gaussian flatness of $\mathcal{K} = 3$.




FIG. 4: Circulation moments $\langle |\Gamma|^p \rangle$ in extended self-similarity (ESS) using $p = 3$. (a) shows the Taylor-Green simulation, while the left and right panels of (b) are the normal fluid perpendicular and parallel component respectively. The insets of each plot give the local slope, where $\lambda_p / \lambda_3 = d \log \langle |\Gamma|^p \rangle / d \log \langle |\Gamma|^3 \rangle$ and the horizontal lines correspond to the K41 scalings $\lambda_p^{K41} = 4p/3$. The shaded regions indicate the classical region where the data used to estimate the exponents.




FIG. 5: Scaling exponents of velocity circulation moments, estimated within the classical range $r > \ell$.

The colour scheme and linestyles is the same as in Fig. 2. The black dashed line represents the self-similar K41 scaling $\lambda_p^{K41} = 4p/3$. The inset shows the proportional deviation from K41 theory.