

Supplementary Materials: Inverse energy transfer in finite-temperature superfluid vortex reconnections

P. Z. Stasiak,¹ C.F. Barenghi,¹ A. Baggaley,¹ L. Galantucci,² and G. Krstulovic³

¹*School of Mathematics, Statistics and Physics, Newcastle University,
Newcastle upon Tyne, NE1 7RU, United Kingdom*

²*Istituto per le Applicazioni del Calcolo “M. Picone” IAC CNR, Via dei Taurini 19, 00185 Roma, Italy*

³*Université Côte d’Azur, Observatoire de la Côte d’Azur, CNRS, Laboratoire Lagrange,
Boulevard de l’Observatoire CS 34229 - F 06304 NICE Cedex 4, France*

NUMERICAL METHOD

Using Schwarz mesoscopic model [1], vortex lines can be described as space curves $\mathbf{s}(\xi, t)$ of infinitesimal thickness, with a single quantum of circulation $\kappa = h/m_4 = 9.97 \times 10^{-8} \text{m}^2/\text{s}$, where h is Planck’s constant, $m_4 = 6.65 \times 10^{-27} \text{kg}$ is the mass of one helium atom, ξ is the natural parameterisation, arclength, and t is time. These conditions are a good approximation, since the vortex core radius of superfluid ^4He ($a_0 = 10^{-10} \text{m}$) is much smaller than any of the length scales of interest in turbulent flows. The equation of motion is

$$\dot{\mathbf{s}}(\xi, t) = \mathbf{v}_s + \frac{\beta}{1 + \beta} [\mathbf{v}_{ns} \cdot \mathbf{s}'] \mathbf{s}' + \beta \mathbf{s}' \times \mathbf{v}_{ns} + \beta' \mathbf{s}' \times [\mathbf{s}' \times \mathbf{v}_{ns}], \quad (1)$$

where $\dot{\mathbf{s}} = \partial \mathbf{s} / \partial t$, $\mathbf{s}' = \partial \mathbf{s} / \partial \xi$ is the unit tangent vector, $\mathbf{v}_{ns} = \mathbf{v}_n - \mathbf{v}_s$, \mathbf{v}_n and \mathbf{v}_s are the normal fluid and superfluid velocities at \mathbf{s} and β, β' are temperature and Reynolds number dependent mutual friction coefficients [2]. The superfluid velocity \mathbf{v}_s at a point \mathbf{x} is determined by the Biot-Savart law

$$\mathbf{v}_s(\mathbf{x}, t) = \frac{\kappa}{4\pi} \oint_{\mathcal{T}} \frac{\mathbf{s}'(\xi, t) \times [\mathbf{x} - \mathbf{s}(\xi, t)]}{|\mathbf{x} - \mathbf{s}(\xi, t)|} d\xi, \quad (2)$$

where \mathcal{T} represents the entire vortex configuration. There is currently a lack of a well-defined theory of vortex reconnections in superfluid helium, like for the Gross-Pitaevskii equation [3–5]. An *ad hoc* vortex reconnection algorithm is employed to resolve the collisions of vortex lines [6].

A *two-way model* is crucial to understand the accurately interpret the back-reaction effect of the normal fluid on the vortex line and vice-versa [7]. We self-consistently evolve the normal fluid \mathbf{v}_n with a modified Navier-Stokes equation

$$\frac{\partial \mathbf{v}_n}{\partial t} + (\mathbf{v}_n \cdot \nabla) \mathbf{v}_n = -\nabla \frac{p}{\rho} + \nu_n \nabla^2 \mathbf{v}_n + \frac{\mathbf{F}_{ns}}{\rho_n}, \quad (3)$$

$$\mathbf{F}_{ns} = \oint_{\mathcal{T}} \mathbf{f}_{ns} \delta(\mathbf{x} - \mathbf{s}) d\xi, \quad \nabla \cdot \mathbf{v}_n = 0, \quad (4)$$




FIG. 1: Schematic diagram of the initial vortex configuration.

where $\rho = \rho_n + \rho_s$ is the total density, ρ_n and ρ_s are the normal fluid and superfluid densities, p is the pressure, ν_n is the kinematic viscosity of the normal fluid and \mathbf{f}_{ns} is the local friction per unit length [8]

$$\mathbf{f}_{ns} = -\mathcal{D}\mathbf{s}' \times [\mathbf{s}' \times (\dot{\mathbf{s}} - \mathbf{v}_n)] - \rho_n \kappa \mathbf{s}' \times (\mathbf{v}_n - \dot{\mathbf{s}}), \quad (5)$$

where \mathcal{D} is a coefficient dependent on the vortex Reynolds number and intrinsic properties of the normal fluid. The regularisation of the mutual friction force onto the normal fluid grid is physically motivated by the strongly localised injection of vorticity during the momentum exchange of point-like particles and viscous flow in classical fluid dynamics [9, 10]. In short, the localised vorticity induced by the relative motion between the vortex lines and the normal fluid is diffused to discretisation of the grid spacing Δx in a time interval ϵ_R . In this way, the delta-forced friction as defined in Eq. 5 is regularised by a Gaussian function, the fundamental solution of the diffusion equation. Further details of the method for classical fluids are contained in [9, 10] and for FOUCAULT in [2].

In this Letter, we report all results using dimensionless units, where the characteristic length scale is $\tilde{\lambda} = D/D_0$, where $D^3 = (1 \times 10^{-3}\text{m})^3$ is the dimensional cube size, $D_0^3 = (2\pi)^3$ is the non-dimensional cubic computational domain. The time scale is given by $\tilde{\tau} = \tilde{\lambda}^2 \nu_n^0 \nu_n$, where the non-dimensional viscosity ν_n^0 resolves the small scales of the normal fluid. In these simulations, these quantities are $\tilde{\lambda} = 1.59 \times 10^{-4}\text{m}$, $\nu_n^0 = 0.32$ and $\tilde{\tau} = 0.366\text{s}$ at $T = 1.9\text{K}$ and $\tilde{\tau} = 0.485\text{s}$ at $T = 2.1\text{K}$. We consider an initial configuration of two pairs of orthogonally vortices, initialised as shown in the schematic of Fig. 1

-
- [1] K.W. Schwarz, Three-dimensional vortex dynamics in superfluid ^4He , *Phys. Rev. B* **38**, 2398 (1988).
 - [2] L. Galantucci, A. W. Baggaley, C. F. Barenghi, and G. Krstulovic, A new self-consistent approach of quantum turbulence in superfluid helium, *Eur. Phys. J. Plus* **135**, 547 (2020).
 - [3] A. Villois, D. Proment, and G. Krstulovic, Irreversible Dynamics of Vortex Reconnections in Quantum Fluids, *Phys. Rev. Lett.* **125**, 164501 (2020).
 - [4] A. Villois, D. Proment, and G. Krstulovic, Universal and nonuniversal aspects of vortex reconnections in superfluids, *Phys. Rev. Fluids* **2**, 044701 (2017).
 - [5] D. Proment and G. Krstulovic, Matching theory to characterize sound emission during vortex reconnection in quantum fluids, *Phys. Rev. Fluids* **5**, 104701 (2020).
 - [6] A. W. Baggaley, The sensitivity of the vortex filament method to different reconnection models, *J. Low Temp. Phys.* **168**, 18 (2012).
 - [7] P. Z. Stasiak, A. W. Baggaley, G. Krstulovic, C. F. Barenghi, and L. Galantucci, Cross-Component Energy Transfer in Superfluid Helium-4, *J Low Temp Phys* 10.1007/s10909-023-03042-5 (2024).
 - [8] L. Galantucci, M. Sciacca, and C.F. Barenghi, Coupled normal fluid and superfluid profiles of turbulent helium II in channels, *Phys Rev B* **92**, 174530 (2015).
 - [9] P. Gualtieri, F. Picano, G. Sardina, and C. M. Casciola, Exact regularized point particle method for multiphase flows in the two-way coupling regime, *Journal of Fluid Mechanics* **773**, 520 (2015).
 - [10] P. Gualtieri, F. Battista, and C. M. Casciola, Turbulence modulation in heavy-loaded suspensions of tiny particles, *Physical Review Fluids* **2**, 034304 (2017).