

UNIVERSITY OF CALIFORNIA

Santa Barbara

Mathematical Approaches to Understanding Mammalian
Circadian Rhythms

by

Peter Corbin St. John

Committee in charge:

Professor Francis J. Doyle III, Chair

Professor Samir Mitragotri

Professor Michelle A. O'Malley

Professor Linda R. Petzold

June 2015

The dissertation of Peter C. St. John is approved.

PROFESSOR SAMIR MITRAGOTRI

DATE

PROFESSOR MICHELLE A. O'MALLEY

DATE

PROFESSOR LINDA R. PETZOLD

DATE

PROFESSOR FRANCIS J. DOYLE III, COMMITTEE CHAIR

DATE

Mathematical Approaches to Understanding Mammalian Circadian Rhythms

Copyright © 2015

by

Peter C. St. John

Acknowledgement

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

This is the second paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

And after the second paragraph follows the third paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest

Acknowledgement

gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

After this fourth paragraph, we start a new paragraph sequence. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Vita of Peter C. St. John

Contact Information

Department of Chemical Engineering
University of California, Santa Barbara
Santa Barbara, CA 93106-5080

Phone: (508) 494-2474
E-mail: pstjohn@engineering.ucsb.edu
Office: Engineering II, Rm. 1508

Education

University of California, Santa Barbara
Ph.D. Candidate, Department of Chemical Engineering

September 2010 - present
Santa Barbara, California

Tufts University
BS, Chemical and Biological Engineering

September 2006 - May 2010
Medford, Massachusetts

Honors and Awards

CAST Student Travel Grant	September 2014
Society for Research on Biological Rhythms (SRBR) Research Merit Award	June 2014
Best Poster, Center for Chronobiology Symposium, UCSD	February 2014
1 st Place, SRBR Logo Competition	January 2014
Mitsubishi Chemical Fellowship Recipient	2012-2015
UCSB Sciencline 2011-2012 Life Science Outstanding Answerer	June 2012
National Science Foundation GRFP Honorable Mention	April 2011
Class of 1947 Victor Prather Prize	May 2010
Max Tischler Prize Scholarship	May 2009
Elected to Tau Beta Pi	September 2008

Publications

St. John, P.C., Taylor, S.R., Abel, J.H., and F.J. Doyle III. Amplitude metrics for cellular circadian bioluminescence reporters (2014) *Biophysical Journal*, 107 (11) pp. 2712-2722

St. John, P.C., Hirota, T., Kay, S.A. and F.J. Doyle III. Spatiotemporal separation of PER and CRY posttranslational regulation in the mammalian circadian clock (2014) *PNAS*, 111 (5) pp. 2040-2045.

Yang, R., Rodriguez-Fernandez, M., **St. John, P.C.**, and F.J. Doyle III. Chapter 8 – Systems Biology (2014) In E. Carson and C. Cobelli (Eds.) *Modelling Methodology for Physiology and Medicine, 2nd Edition*, pp. 159-187.

St. John, P.C., and F.J. Doyle III. Estimating confidence intervals in predicted responses for oscillatory biological models (2013) *BMC Systems Biology* 7:71.

Hirota, T., Lee, J.W., **St. John, P.C.**, Sawa, M., Iwaisako, K., Noguchi, T., Pongsawakul, P.Y., Sonntag, T., Welsh, D.K., Brenner, D.A., Doyle, F.J. III, Schultz, P.G., Kay, S.A., Identification of small molecule activators of cryptochrome (2012) *Science*, 337 (6098) pp. 1094-1097.

Murphy A.R., **St. John P.C.**, Kaplan D.L. Modification of silk fibroin using diazonium coupling chemistry and the effects on hMSC proliferation and differentiation (2008) *Biomaterials*, 29 (19), pp. 2829-2838.

Contributed Talks

St. John, P.C., and F.J. Doyle III. November 2014. Development of Amplitude Response Curves for Single-Cell and Population-Level Circadian Systems. *To be presented at the 2014 AIChE Annual Meeting, Atlanta, GA*

St. John, P.C., and F.J. Doyle III. June 2014. Amplitude metrics for uncoupled cellular circadian bioluminescence reporters. Presented at the Society for Research on Biological Rhythms Meeting, Big Sky, MT.

St. John, P.C., and F.J. Doyle III. October 2012. Cryptochrome balancing for period control: mathematical insights into circadian clock design. Presented at the Model-based Analysis and Control of Cellular Processes Workshop, Purdue University, West Lafayette, IN.

Poster Presentations

St. John, P.C., and F.J. Doyle III. February 2014. Spatiotemporal separation of PER and CRY post-translation regulation. Presented at the UCSD Center for Chronobiology Symposium, La Jolla, CA.

St. John, P.C., T. Hirota, S.A. Kay, and F.J. Doyle III. July 2013. Estimating confidence intervals in model predictions to determine plausible mechanisms for small molecule modifiers. Presented at the Chronobiology Gordon Research Conference, Newport, RI.

St. John, P.C., and F.J. Doyle III. February 2013. Predictive confidence intervals from mathematical circadian models. Presented at the UCSD Center for Chronobiology Symposium, La Jolla, CA.

St. John, P.C., T. Hirota, S. Kay, and F.J. Doyle III. May 2012. Cryptochrome balancing for period control. Presented at the Meeting of the Society for Research on Biological Rhythms, Destin, FL.

St. John, P.C., and F.J. Doyle III. February 2012. Perturbation analysis of circadian clock degradation. Presented at the UCSD Center for Chronobiology Symposium, La Jolla, CA.

St. John, P.C., M. Rodriguez-Fernandez, and F.J. Doyle III. June 2011. Advanced global optimization and sensitivity techniques for analyzing deterministic circadian models. Presented at the 12th Annual UC Systemwide Bioengineering Symposium, Santa Barbara, CA.

Seminars

St. John, P.C. May 2013. Sensitivity analysis in the study of mammalian circadian rhythms. Presented at the UCSB NSF-IGERT systems biology seminar series, Santa Barbara, CA.

Research Experience

University of California, Santa Barbara

Ph.D. Candidate

Computational analysis of the mammalian circadian clock, with a focus on elucidating the functional design consequences of the underlying genetic regulatory network.

Advisor: Francis J. Doyle III

Department: Chemical Engineering

September 2010 - present

Santa Barbara, California

Tufts University <i>Senior Honors Thesis</i> Catalytic Hydrodechlorination of 2-Chlorophenol using Viral-templated Palladium Nanoparticles Advisor: Hyunmin Yi Department: Chemical and Biological Engineering	September 2009 - May 2010 Medford, Massachusetts
University of California, Los Angeles <i>UCLA NanoCER Program (NSF REU)</i> Encapsulation Efficiencies and Release Rates from Water-in-Oil-in-Water Nanoemulsions Advisor: Timothy Deming Department: Bioengineering	June 2009 - August 2009 Los Angeles, California
Tufts University <i>Tufts Summer Scholars</i> Hydrodechlorination of 2-Chlorophenol with Palladium Nanoparticles on Genetically Modified Tobacco Mosaic Virus Scaffolds Advisor: Hyunmin Yi Department: Chemical and Biological Engineering	June 2008 - August 2008 Medford, Massachusetts
Tufts University <i>Undergraduate Research Credit</i> Chemically Modified Silk Fibroin Based Scaffolds for Bone Tissue Engineering Advisor: David Kaplan Department: Biomedical Engineering	October 2007 - May 2008 Medford, Massachusetts

Teaching Experience

University of California, Santa Barbara <i>Teaching Assistant, ChE132c</i> Helped teach undergraduate statistics for two subsequent years: gave three lectures, held office hours and review sessions, and graded homeworks.	January 2013 - December 2013 Santa Barbara, California
University of California, Santa Barbara <i>Teaching Assistant, ChE180a</i> Designed and ran experiments for the junior laboratory course. Also helped in grading student reports.	March 2012 - June 2012 Santa Barbara, California

Community Involvement

Peer Review	January 2014-present
--------------------	-----------------------------

Reviewer for Biophysical Journal; IEEE Control Systems Society Conference; 21st International Symposium on Mathematical Theory of Networks and Systems

Scienceline "Ask A Scientist"

December 2010 - present

Answers science and engineering questions posed by students and teachers from local K-12 schools.

Website: www.sciencealive.ucsbd.edu

UCSB Discover Engineering Weekend

May 2011

Helped organize and run a weekend for local high school students to learn basic engineering principles and apply their knowledge to build miniature alternative energy cars.

Mentoring Experience

Amanda Luan, Undergraduate Student, ICB SSB URAP **June 2014 - December 2014**

Lukas Widmer, Masters Student, ETH Zurich **April 2012 - February 2013**

Andrew Barisser, Rotation Student, BMSE UCSB **September 2012 - December 2012**

Abstract

This is the second paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

And after the second paragraph follows the third paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

After this fourth paragraph, we start a new paragraph sequence. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is

there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

This is the second paragraph. Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Contents

1	Introduction	1
1.1	Biological background	1
1.1.1	Evolutionary history	3
1.1.2	Mechanisms of gene regulation	3
1.1.3	Mammalian genetic circuit	3
1.2	Mathematical modeling	4
1.2.1	Overview of kinetic ODE models	4
1.2.2	Basic terms and assumptions	5
1.2.3	Simulation methods	6
1.2.4	ODE Sensitivity Analysis	8
1.2.5	Previous models of circadian rhythms	9
2	A new model of the core circadian feedback loop	10
2.1	Background	10
2.1.1	Small molecule modulators	10
2.2	Results	13
2.2.1	A new model for period regulation	13
2.2.2	Parameter estimation	21
2.2.3	Model validation and dynamics	26
2.2.4	Prediction of KL001 mechanism	28

2.3	Discussion	30
2.3.1	Insights into circadian network design	30
2.3.2	Identifiability of model parameters	32
3	Identifiability analysis for models of circadian rhythms	33
3.1	Background	33
3.1.1	Identifiability analysis	34
3.2	Methods	36
3.2.1	Collocation methods	36
3.2.2	Generating initial values	39
3.2.3	First order sensitivity analysis	39
3.2.4	Generation of data for bootstrap methods	40
3.2.5	Calculation times	41
3.3	Results	42
3.3.1	Effect of data quality on predictive confidence	43
3.3.2	Application to literature data for model discrimination	46
3.4	Discussion	52
4	Spatiotemporal separation of PER and CRY posttranslational regulation	55
4.1	Background	55
4.1.1	Nuclear entry of PER-CRY complex	56
4.1.2	Importance of posttranslational regulators	57
4.1.3	Functional differences between PER and CRY	58
4.2	Methods	59
4.2.1	Analysis of luminescence profiles	59
4.2.2	Cost function	59
4.2.3	Parameter estimation and bootstrap analysis	63
4.2.4	Selection of parameters for FBXL3 and CKI	63

4.2.5	Numerical experiments	64
4.3	Results	64
4.3.1	Opposite amplitude effects of longdaysin and KL001	64
4.3.2	Main period-determining perturbations	64
4.3.3	Independent mechanisms of PER and CRY regulation	67
4.4	Discussion	75
4.4.1	Phase-dependent activity of small molecule modulators	76
5	Amplitude metrics for cellular bioluminescence reporters	77
5.1	Background	77
5.1.1	Importance of circadian amplitude	78
5.1.2	Amplitude at the single-cell and population level	78
5.1.3	Role of mathematical modeling	79
5.2	Methods	81
5.2.1	Basic definitions	81
5.2.2	Perturbations to limit cycle systems	82
5.2.3	Phase response curves	84
5.2.4	Phase-diffusion model	84
5.2.5	Numerical simulations	86
5.3	Results	86
5.3.1	Definition of an amplitude metric at the single-cell level	86
5.3.2	Single-cell amplitude response curves	87
5.3.3	Population-level response curves	93
5.3.4	Population-level mean expression profiles	94
5.3.5	Application to Experimental Data	105
5.4	Discussion	108
5.4.1	Application to pharmacological control	108
5.4.2	Damping rate is correlated with single-cell noise	109

6 Decay rate as proxy for cellular noise	111
7 Conclusions and future work	112
7.1 Continuation of collaboration with Andrew	112
7.1.1 Experimental validation of model predictions	112
7.1.2 Prediction of fasting perturbations	113
7.1.3 Validity of step vs pulse perturbation	114
7.2 Amplitude maximization through therapeutic treatment	114
7.2.1 Optimal control strategies to maximize peripheral clock amplitude	115
7.2.2 Fewest treatments, combinatorial perturbations, etc.	115
7.3 Iterative optimization of network topology	116
7.3.1 (Idea from Mitsubishi proposal)	116
7.3.2 Design machine-learning algorithm to find optimum model structure	116

List of Figures

1.1	A biological feed forward controller	2
2.1	Connectivity of model 2.1	14
2.2	Repressor degradation profiles	15
2.3	Time course concentration profiles of model 2.1	27
2.4	Model validation against experimental results	28
2.5	Prediction of the KL001 mechanism	29
2.6	Simultaneous knockdown of <i>Cry1</i> and <i>Cry2</i>	31
2.7	First order relative period sensitivities	31
3.1	Parameter estimation and bootstrap methods flowchart	44
3.2	Time-course profiles of the state trajectories for <i>Per</i> mRNA	45
3.3	Parameter and sensitivity identifiability for increasing error	47
3.4	Effect of high-resolution sampling on identifiability	48
3.5	Identifiability comparison of two model structures	50
3.6	Time-series dynamics of fitted models	53
4.1	Small molecule targets	57
4.2	Analysis of circadian reporter luminescence data	60
4.3	Different amplitude effect of small molecule circadian modulators targeting CKI-PER and FBXL3-CRY	65
4.4	Fitted trajectories from bootstrap runs	66

4.5	Bootstrap confidence intervals in relative period sensitivities	68
4.6	Bootstrap predictions of circadian actions of FBXL3-CRY and CKI-PER pathways	69
4.7	Predictions of KL001 and longdaysin results	71
4.8	Comparison of the effects of KL001 and longdaysin	71
4.9	Effect of cytoplasmic PER stabilization on period and E box transcription amplitude	72
4.10	Comparison of two candidate mechanisms for CKI inhibition	73
4.11	Independence of CKI-PER and FBXL3-CRY pathways	74
4.12	Spatiotemporal separation underlies independence of CKI-PER and FBXL3-CRY pathways	75
5.1	Schematic showing trajectories used in the calculation of single-cell phase and amplitude change	83
5.2	Amplitude metrics at the single-cell level measure transient deviations from the limit cycle	88
5.3	Phase and amplitude response curves describe how a single oscillator will respond to a perturbation	89
5.4	Convergence of finite-difference and differential methods	92
5.5	Synchrony affects population-level amplitude	96
5.6	Approximation of an explicit stochastic population by continuous methods	99
5.7	Validation of continuous methods with the Oregonator model	100
5.8	Validation of continuous methods with the model from Hirota <i>et al.</i> , 2012101	
5.9	Response curves describe different types of amplitude change	104
5.10	Method allows direct comparison between model results and experimental bioluminescence profiles	106

List of Models

2.1	A degradation-based model of the core feedback loop	17
2.2	Changed equations for the activation-based model	17
3.1	An expanded parameterization of model 2.1	51
4.1	Model from Relógio <i>et al.</i> 2011 [22]	61
4.2	Model from Leloup & Goldbeter, 2003 [10]	62
5.1	Model adapted from Novák & Tyson, 2008 [97]	96
5.2	The Oregonator model [99]	102

List of Tables

2.1	Parameter values for model 2.1	18
2.2	Summary of cost function entries	22
3.1	Parameter values for model 3.1	49
4.1	Descriptions of the model parameters governing FBXL3-CRY	70
4.2	Descriptions of the model parameters governing CKI-PER	70

Chapter 1

Introduction

1.1 Biological background

Systems biology is a multidisciplinary field, drawing from biology, mathematics, and computer science. In this section, I first provide a background on the biology of circadian rhythms, followed by previous literature on computational modeling.

Ecological competition drives species to optimize their survival and reproduction in a particular environmental niche. Aspects of their environment, including temperature, resource availability, and sunlight are all important factors in determining an individual's success.

In nearly all environments, however, these factors vary greatly from hour to hour and month to month. While food or sunlight might be plentiful during daylight hours, the threat of predators might peak as well. Fortunately, while these critical parameters undergo sharp changes, they do so with a predictable and repeatable schedule. It is not surprising that, in optimizing their survival in an oscillating environment, species have developed a means to predict changes and to ready themselves for the correct behavior.

Biological rhythms are classified as *circadian* if they display a number of defining




Figure 1.1: A biological feed forward controller. Circadian rhythms

characteristics:

- They are *endogenous*, meaning they continue to oscillate even when the organism is isolated from its environment
- They are temperature compensated, maintaining a consistent period for moderate changes in average temperature
- They are *entrainable*, meaning they can adjust their phase and period in response to a changing in environmental signal
- They oscillate with a roughly 24 hour period [1].

1.1.1 Evolutionary history

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.1.2 Mechanisms of gene regulation

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.1.3 Mammalian genetic circuit

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of

the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2 Mathematical modeling

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2.1 Overview of kinetic ODE models

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Spectrum of modeling methods (detail vs. scope)

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2.2 Basic terms and assumptions

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Mass action

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of

the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Michealis-Menten

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Hill-type

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2.3 Simulation methods

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information.

Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Numerical solution of nonlinear ODEs

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Boundary value problem for limit cycle models

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2.4 ODE Sensitivity Analysis

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

ODEs in general

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Period sensitivity

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written

in of the original language. There is no need for special content, but the length of words should match the language.

PRCs

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

1.2.5 Previous models of circadian rhythms

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Chapter 2

A new model of the core circadian feedback loop¹

2.1 Background

2.1.1 Small molecule modulators

Circadian rhythms help to regulate metabolic homeostasis, and therefore compromised circadian oscillations often result in negative health consequences. Since circadian rhythms may be damped by lifestyle factors, many researchers have begun searching for pharmacological agents which might alter circadian rhythmicity. Several proof-of-concept molecules have been found which cause dose-dependent period change in cultured fibroblasts [2]. While chemical biology approaches are able to localize the effects of these molecules to specific targets in the circadian network, it is often difficult to reconcile individual rate changes with the observed systems-level changes in period or amplitude. Mathematical models are therefore essential

¹Portions of this chapter are published in T. Hirota, J. W. Lee, P. C. St. John, M. Sawa, K. Iwaisako, T. Noguchi, P. Y. Pongsawakul, T. Sonntag, D. K. Welsh, D. A. Brenner, F. J. Doyle, P. G. Schultz, and S. A. Kay, "Identification of Small Molecule Activators of Cryptochrome," *Science*, vol. 337, pp. 1094–1097, July 2012.

to understanding the effects of such drugs, as they are able to verify that hypothesized mechanisms of action are mathematically consistent. Furthermore, mathematical models provide a systematic framework by which to examine consequences of various mechanistic assumptions - many of which can be subsequently verified by experimental inquiry. The search for small molecule modulators of circadian rhythms therefore yields the added benefit of strengthening our knowledge of the core clock feedback circuit, as inconsistencies between assumed kinetics and experimental results often lead to a more refined knowledge of the network's dynamics.

In this chapter, I describe a collaborative project in which experimental results inspired the creation of a new model of the core mammalian feedback network. A new small molecule modulator of cryptochrome, KL001, was identified through forward chemical genetic screening [3]. The compound was found to simultaneously increase the half-life of CRY and cause period lengthening. In order to explain these new experimental results while maintaining consistency with published evidence, I constructed a simple mathematical model of the Per/Cry negative feedback loop.

The functional roles of the cryptochrome isoforms (*Cry1* and *Cry2*) in circadian rhythms have long eluded biologists, as although they share a similar structure, perturbations to these genes typically result in opposite trends [4]. Existing genetic evidence and siRNA knockdown studies agree that while suppressing *Cry1* leads to shorter periods, suppressing *Cry2* leads to longer periods [5, 6]. However, some evidence shows the similarity between the two *Crys*. The knockouts $Cry1^{+/-} \ll Cry2^{-/-}$ and $Cry1^{-/-} Cry2^{+/-}$ both show shorter wheel-running activity than their respective WT/double knockouts $Cry1^{+/+} Cry2^{-/-}$ and $Cry1^{-/-} Cry2^{+/+}$ [5]. Using new data from the application of KL001 together with existing experimental results, I developed a hypothesized mechanism to reconcile these seemingly contradictory results:

Network structure

In the core feedback loop, Per and Cry transcription is activated by the CLOCK-BMAL1 complex. The protein products of these genes are transported to the nucleus after a time delay and bind to CLOCK-BMAL1, repressing their own transcription. Evidence indicates that an interaction of PER and CRY proteins is required for the timely nuclear entry of both proteins [7, 8, 9]. I therefore assume that PER and CRY bind and enter the nucleus in a 1:1 stoichiometric ratio, similar to more complicated models of mammalian circadian rhythms [10, 11, 12].

Protein stoichiometry

Quantitative protein assays have revealed that the PER proteins are present at levels nearly ten times less than the CRYs, and are completely consumed at their trough. The CRYs, on the other hand, only fall to about one quarter of their peak value [13]. These results indicate that PER is the rate-limiting component for nuclear entry. This configuration explains why, since /it Cry is in excess, a knockout of either *Cry* does not lead to loss of function at the tissue level. Additionally *Cry1*, the more dominant repressor, is consistently found at higher levels than *Cry2*. However, since the ratio of nuclear to total protein is nearly equivalent for CRY1 and CRY2, PER must import each with equal affinity [13, 14].

Importance of degradation

Studies of the simplest cellular oscillators have illustrated the importance of the degradation of repressor complexes in determining the period of oscillation [15]. In the mammalian circadian clock, FBXL3 has been indicated as a key protein in clearing the nucleus of CRY, and siRNA knockdowns of *Fbxl3* result in period lengthening [6]. While FBXL3 localizes primarily in the nucleus, CRY2 possesses a distinct phosphorylation domain that flags it for cytosolic proteolysis [16]. Knocking down this

domain shortens the period, indicating that degradation rates play a major role in determining the oscillatory period and that stabilization in the cytosol and nucleus are likely not equivalent.

Key time delays

Similar to *Drosophila*, mammalian circadian oscillations are generated by three main time delays in the negative feedback loop: mRNA transcription, nuclear localization of protein, and degradation of the nuclear complexes [17]. In order to capture these dynamics without introducing unnecessary stiffness, I explicitly consider only the concentrations of nuclear mRNA, cytosolic unbound protein, and nuclear protein complexes.

2.2 Results

2.2.1 A new model for period regulation

Since the circadian clock is largely insensitive to the total amount *Cry*, it follows that the core negative loop is comprised of two redundant coupled inhibitory mechanisms. The natural periods of these two loops (running in isolation of the other) can be inferred from their corresponding knockout phenotype, i.e., the CRY1 feedback loop has a long period, while the CRY2 loop has a short period. In the wild-type phenotype, the two isoforms of CRY must compete for the available PER, and thus the nuclear CRY1/CRY2 levels are constrained with a ratio proportional to their relative expression. This suggests that the period of the clock is governed by the nuclear CRY1/CRY2 ratio, where higher amounts of *Cry1* shift the clock closer to the *Cry2*^{-/-} phenotype, and higher amounts of *Cry2* shift the clock closer to the *Cry1*^{-/-} phenotype, described in figure 2.1.




Figure 2.1: Connectivity of model 2.1. Schematic for the core negative feedback loop, consisting of the two redundant *Cry* mechanisms. The wild-type period, consisting of both loops, is a balance between the short and long oscillations.

The CRY1/CRY2 ratio

In silico modeling was used to test the feasibility of the above hypothesis, and to investigate possible mechanisms by which the CRY1/CRY2 ratio might control the period of oscillations. The model uses eight state variables for the three mRNA species (*Per*, *Cry1*, and *Cry2*), three cytosolic proteins, and two nuclear proteins. The differential equations for each state were formulated using standard Hill-type repression, Michaelis-Menten, and mass action kinetics. The 21 unknown kinetic parameters were found by fitting the stoichiometric data from [13], requiring correct periods for the *Cry1*^{-/-} and *Cry2*^{-/-} knockouts.

I investigated two possible mechanisms to explain the long/short period phenotype of the CRY1/CRY2 loops. First, following experimental evidence, I allowed CRY1 and CRY2 to have different inhibitory efficiency [18]. However, optimizations with this structure were unable to find a parameter set with appropriate period sen-




Figure 2.2: Repressor degradation profiles. Time course plots of the simulated total nuclear repressor concentration for the degradation model under various clock conditions. The relative contribution of CRY1 and CRY2 are shaded green and orange, respectively. Longer periods occur when a higher percentage of the nuclear complex is CRY1, taking longer to degrade.

sivities, described in section 2.2.2. Instead, I investigated whether the difference in potency between *Crys* could be explained through difference in degradation rates. To this end, I allowed the degradation rates of nuclear CRY1 and CRY2 to differ, while holding their inhibition constants equal. Using this configuration, the structure was able to fit the experimental sensitivities. In the optimized parameter set, the degradation rate of nuclear CRY2 is higher than that of CRY1, such that for a constant total nuclear CRY, higher fractions of CRY2 cause the repressive complexes to be cleared faster. Since PER is the limiting reagent in nuclear entry, the amount of total CRY that enters the nucleus is largely insensitive to perturbations in cytosolic CRY expression. In effect, the two isoforms of CRY must compete for the available PER, and thus the nuclear CRY1/CRY2 levels are constrained with a ratio proportional to their relative expression. The period of the clock is thus governed by the nuclear CRY1/CRY2 ratio, where higher amounts of *Cry1* shift the clock closer to the *Cry2*^{-/-} phenotype, and higher amounts of *Cry2* shift the clock closer to the *Cry1*^{-/-} phenotype. Figure 2.2 shows the time varying total complex concentration under various perturbations, with the relative contributions from CRY1 and CRY2 highlighted.

Model equations

The model is formulated as set of 8 ordinary differential equations shown in model 2.1 and model 2.2. Additional assumptions made while formulating the model equations are listed below:

- For simplicity and ease of parameter estimation, the model only considers the three genes *Per*, *Cry1*, and *Cry2*, not explicitly considering the three known isoforms of *Per*.
- EBOX activators *Clock* and *Bmal1* are considered constitutively expressed and are represented by the v_{txn} parameters.
- Repression of CLOCK-BMAL1 activity is attained through Hill-type inhibition. The Hill coefficient is fixed at 3, which was found to provide sufficient non-linearity for oscillations. See below for the derivation of the repressor input function used in tables S2A and S2B.
- The degradation of the mRNA species and cytoplasmic proteins is assumed to follow standard Michaelis-Menten kinetics.
- The degradation kinetics of the nuclear proteins are assumed to follow Michaelis-Menten kinetics. Because both $CRY1_n$ and $CRY2_n$ are thought to be degraded by the same pathway in the nucleus, kinetic equations using the *pseudo* steady-state hypothesis were derived for two substrates sharing the same enzyme. As a result, each *Cry* isoform acts as an inhibitor to the other's degradation.

Model 2.1: A degradation-based model of the core feedback loop. Lower case letters (p : *Per*, $c1$: *Cry1*, $c2$: *Cry2*) are mRNA state variables. Upper-case letters (P : PER, $C1$: CRY1, $C2$: CRY2) are the free (cytosolic) proteins. $C1N$: CRY1 and $C2N$: CRY2 are the nuclear proteins.

$$\frac{dp}{dt} = \frac{v_{txn,p}}{k_{txn,p} + (C1N + C2N)^3} - \frac{v_{deg,p} p}{k_{deg,p} + p} \quad (2.1)$$

$$\frac{dc1}{dt} = \frac{v_{txn,c1}}{k_{txn,c} + (C1N + C2N)^3} - \frac{v_{deg,c1} c1}{k_{deg,c} + c1} \quad (2.2)$$

$$\frac{dc2}{dt} = \frac{v_{txn,c2}}{k_{txn,c} + (C1N + C2N)^3} - \frac{v_{deg,c2} c2}{k_{deg,c} + c2} \quad (2.3)$$

$$\begin{aligned} \frac{dP}{dt} = k_{tln,p} p - \frac{v_{deg,P} P}{k_{deg,P} + P} - v_{a,CP} P C1 + v_{d,CP} C1N \\ - v_{a,CP} P C2 + v_{d,CP} C2N \end{aligned} \quad (2.4)$$

$$\frac{dC1}{dt} = c1 - \frac{v_{deg,C1} C1}{k_{deg,C} + C1} - v_{a,CP} P C1 + v_{d,CP} C1N \quad (2.5)$$

$$\frac{dC2}{dt} = c2 - \frac{v_{deg,C2} C2}{k_{deg,C} + C2} - v_{a,CP} P C2 + v_{d,CP} C2N \quad (2.6)$$

$$\frac{dC1N}{dt} = - \frac{v_{deg,CP} C1N}{k_{deg,CP} + C1N + C2N} + v_{a,CP} P C1 - v_{d,CP} C1N \quad (2.7)$$

$$\frac{dC2N}{dt} = - \frac{(v_{deg,CP} m_{C2N}) C2N}{k_{deg,CP} + C2N + C1N} + v_{a,CP} P C2 - v_{d,CP} C2N \quad (2.8)$$

Model 2.2: Changed equations for the activation-based model. The remainder of the model equations (not duplicated below) are found in model 2.1

$$\frac{dp}{dt} = \frac{v_{txn,p}}{k_{txn,p} + (m_{C2N} C1N + C2N)^3} - \frac{v_{deg,p} p}{k_{deg,p} + p} \quad (2.1^*)$$

$$\frac{dc1}{dt} = \frac{v_{txn,c1}}{k_{txn,c} + (m_{C2N} C1N + C2N)^3} - \frac{v_{deg,c1} c1}{k_{deg,c} + c1} \quad (2.2^*)$$

$$\frac{dc2}{dt} = \frac{v_{txn,c2}}{k_{txn,c} + (m_{C2N} C1N + C2N)^3} - \frac{v_{deg,c2} c2}{k_{deg,c} + c2} \quad (2.3^*)$$


$$\frac{dC2N}{dt} = - \frac{v_{deg,CP} C2N}{k_{deg,CP} + C2N + C1N} + v_{a,CP} P C2 - v_{d,CP} C2N \quad (2.8^*)$$

Table 2.1: Parameter values for model 2.1. Parameters for model described in model 2.1.

	Parameter	Description	Degradation	Activation
1	$v_{\text{txn,p}}$	<i>Per</i> Transcription rate	0.195	0.276
2	$v_{\text{txn,c1}}$	<i>Cry1</i> Transcription rate	0.131	0.062
3	$v_{\text{txn,c2}}$	<i>Cry1</i> Transcription rate	0.114	0.053
4	$k_{\text{txn,p}}$	<i>Per</i> Repression constant	0.425	0.425
5	$k_{\text{txn,c}}$	<i>Cry1/2</i> Repression constant	0.259	0.262
6	$v_{\text{deg,p}}$	<i>Per</i> Max degradation rate	0.326	0.472
7	$v_{\text{deg,c1}}$	<i>Cry1</i> Max degradation rate	0.676	0.322
8	$v_{\text{deg,c2}}$	<i>Cry2</i> Max degradation rate	0.608	0.290
9	$k_{\text{deg,p}}$	<i>Per</i> Degradation constant	0.011	0.024
10	$k_{\text{deg,c}}$	<i>Cry1/2</i> Degradation constant	1.149	0.809
11	$v_{\text{deg,P}}$	Max PERc degradation rate	2.970	2.970
12	$k_{\text{deg,P}}$	PERc degradation constant	0.034	0.034
13	$v_{\text{deg,C1}}$	Max CRY1c degradation rate	1.523	1.048
14	$v_{\text{deg,C2}}$	Max CRY2c degradation rate	1.686	1.134
15	$k_{\text{deg,C}}$	CRYc degradation constant	2.017	2.028
16	$v_{\text{deg,CP}}$	CRYn degradation rate	0.101	0.070
17	m_{C2N}	CRY2n degradation multiplier	3.318	3.334
18	$k_{\text{deg,CP}}$	CRYn degradation constant	0.053	0.053
19	$v_{\text{a,CP}}$	CRYn association rate	0.041	0.028
20	$v_{\text{d,CP}}$	CRYn dissociation rate	0.002	0.001
21	$k_{\text{tln,p}}$	PER translation rate	3.000	1.000

Derivation of a shared enzyme degradation rate

To find the rate equations associated with a shared-enzyme degradation mechanism, we derive them from the equilibrium relationships:



The end goal is the degradation rates of the two enzyme complexes:

$$\begin{aligned}
 r_{d,1} &= k_{d,1} [ES1] \\
 r_{d,2} &= k_{d,2} [ES2]
 \end{aligned} \tag{2.10}$$

We invoke the standard pseudo steady-state assumption and set $\frac{d[ES]}{dt} = 0$, and obtain the following production = consumption equalities for [ES1]:

$$k_1[E][S] = k_{-1}[ES1] + k_{d,1}[ES1] \tag{2.11}$$

Solving equation (2.11) for ES1 and substituting in equation (2.9), we obtain

$$\left(\frac{k_{-1} + k_{d,1}}{k_1} \right) [ES1] = ([E_t] - [ES1] - [ES2]) [S1]$$

After defining a useful combined rate constant,

$$K_1 \equiv \frac{k_{-1} + k_{d,1}}{k_1}$$

we can further simplify

$$\begin{aligned} K_1[\text{ES1}] + [\text{S1}][\text{ES1}] &= ([\text{E}_t] - [\text{ES2})][\text{S1}] \\ [\text{ES1}] &= \frac{([\text{E}_t] - [\text{ES2})][\text{S1}]}{K_1 + [\text{S1}]} \end{aligned} \quad (2.12)$$

If we perform the same operations for $[\text{ES2}]$ and plug them into equation (2.12).

$$\begin{aligned} [\text{ES1}] &= \frac{\left(\text{E}_t - \frac{([\text{E}_t] - [\text{ES1}])[\text{S2}]}{K_2 + [\text{S2}]}\right)[\text{S1}]}{K_1 + [\text{S1}]} \\ [\text{ES1}] &= \frac{[\text{E}]_t[\text{S1}]}{K_1 + [\text{S1}]} - \frac{[\text{E}]_t[\text{S1}][\text{S2}]}{(K_1 + [\text{S1}])(K_2 + [\text{S2}])} + \frac{[\text{ES1}][\text{S1}][\text{S2}]}{(K_1 + [\text{S1}])(K_2 + [\text{S2}])} \\ [\text{ES1}] &= \left(\frac{[\text{E}]_t[\text{S1}]}{K_1 + [\text{S1}]}\right) \left(\frac{1 - \frac{[\text{S2}]}{K_2 + [\text{S2}]}}{1 - \frac{[\text{S1}][\text{S2}]}{(K_1 + [\text{S1}])(K_2 + [\text{S2}])}}\right) \end{aligned} \quad (2.13)$$

If we pull the second fraction from (equation (2.13)) into the denominator of the first, we can simplify further.

$$\begin{aligned} [\text{ES1}] &= \frac{[\text{E}]_t[\text{S1}]}{\left(\frac{K_1 + [\text{S1}] - \frac{[\text{S1}][\text{S2}]}{K_2 + [\text{S2}]}}{1 - \frac{[\text{S2}]}{K_2 + [\text{S2}]}}\right)} \\ [\text{ES1}] &= \frac{[\text{E}]_t[\text{S1}]}{\left(\frac{K_2 + [\text{S2}]}{K_2}\right) \left(K_1 + [\text{S1}] - \frac{[\text{S1}][\text{S2}]}{K_2 + [\text{S2}]}\right)} \\ [\text{ES1}] &= \frac{K_2[\text{E}]_t[\text{S1}]}{(K_1 + [\text{S1}])(K_2 + [\text{S2}]) - [\text{S1}][\text{S2}]} \\ [\text{ES1}] &= \frac{K_2[\text{E}]_t[\text{S1}]}{K_1 K_2 + K_2[\text{S1}] + K_1[\text{S2}]} \\ [\text{ES1}] &= \frac{K_2[\text{E}]_t[\text{S1}]}{K_1 K_2 + K_2[\text{S1}] + K_1[\text{S2}]} \end{aligned}$$

Dividing top and bottom by K_2 :

$$[\text{ES1}] = \frac{[\text{E}]_t[\text{S1}]}{K_1 + [\text{S1}] + \frac{K_1}{K_2}[\text{S2}]} \quad (2.14)$$

Substituting (equation (2.14)) into (equation (2.10)) and setting $K_1 = K_2$, $k_{d,1} \neq k_{d,2}$, we obtain the final shared rate laws (with an equivalent analysis for [ES2])

$$r_{d,1} = \frac{V_{\max,1}[S1]}{K_M + [S1] + [S2]}$$

$$r_{d,2} = \frac{V_{\max,2}[S2]}{K_M + [S1] + [S2]}$$

Derivation of Hill-type repression formulas

To derive the equations for the Hill-type inhibition, allowing for different repressive activities (model 2.2), we start with the standard equation for Hill-type regulation:

$$\frac{v_{\max}[A]^n}{K_m^n + [A]^n} \quad (2.15)$$

Allowing for competitive inhibition with two inhibitors, K_m is replaced with the apparent Michaelis-Menten constant

$$K_m^{\text{app}} = K_m \left(1 + \frac{[I_1]}{K_{i,1}} + \frac{[I_2]}{K_{i,2}} \right) \quad (2.16)$$

By assuming constitutive activator concentrations and non-dimensionalizing $[I^*] = \frac{[I]}{K_{i,2}}$, $m = \frac{K_{i,2}}{K_{i,1}}$:

$$\frac{v_{\max^*}}{K_m^* + (m [I_1^*] + [I_2^*])^n}$$

Assuming equal repressive activity ($K_{i,2} = K_{i,1}$), $m = 1$, we obtain the rate equation used in the degradation model.

2.2.2 Parameter estimation

The model equations, specifying the state and parameter dependent time derivatives of each concentration variable, were written in python using the CasADi computer

algebra package [19]. The model was simulated using the SUNDIALS suite of ODE solvers [20]. A parameter-dependent cost function was developed that assigns numerical values reflecting how well a parameter set fits desired features, taken from [13]. The first step in the cost function evaluation is the numerical solution of the limit cycle, described previously [21]. If a limit cycle was unable to be found, the cost function returns a maximum value. Otherwise, the cost function returns a squared difference from the desired value. A priority weight was also attached to each cost entry, such that more important costs would be prioritized. A description of each entry in the cost function, along with the value for experimental and final model is shown in table 2.2 (degradation-based model, model 2.1; activation-based model, model 2.2). To minimize the cost function, we employed a genetic algorithm method used previously for optimizing circadian parameters [11]. In the method, 5000 solutions are calculated at random parameter values within feasible bounds. Those with the best cost function scores are kept and used to generate subsequent solutions. This procedure is iterated for up to 2000 generations, or until convergence criteria are met.

Table 2.2: Summary of cost function entries. The weights for each entry were chosen based on the relative importance of the desired behavior. Entries 1-9 and 12-16 were obtained from the data presented in [13]. SiRNA sensitivities were obtained were taken from [6].

	Description	Weight	Desired	Degradation	Activation
1	<i>per</i> mRNA peak-trough ratio	0.5	> 20	large	large
	$\frac{y_{\max}(Per)}{y_{\min}(Per)}$				

Description		Weight	Desired	Degradation	Activation
2	<i>Cry1</i> mRNA peak-trough ratio	0.5	2.155	8.942	3.748
	$\frac{y_{\max}(Cry1)}{y_{\min}(Cry1)}$				
3	<i>Cry2</i> mRNA peak-trough ratio	0.5	2.236	7.813	3.484
	$\frac{y_{\max}(Cry2)}{y_{\min}(Cry2)}$				
4	PER protein peak to trough ratio	5	> 20	large	large
	$\frac{y_{\max}(\text{PER})}{y_{\min}(\text{PER})}$				
5	CRY1 protein peak-trough ratio	3	3.247	6.385	1.847
	$\frac{y_{\max}(\text{CRY1})}{y_{\min}(\text{CRY1})}$				
6	CRY2 protein peak-trough ratio	3	1.975	8.094	2.347
	$\frac{y_{\max}(\text{CRY2})}{y_{\min}(\text{CRY2})}$				

Description		Weight	Desired	Degradation	Activation
7	Fraction PER of total protein	3	0.105	0.169	0.073
	$\frac{y_{\max}(\text{PER})}{y_{\max}(\text{PER}) + y_{\max}(\text{CRY1}) + y_{\max}(\text{CRY2})}$				
8	Fraction CRY1 of total protein	3	0.555	0.473	0.554
	$\frac{y_{\max}(\text{CRY1})}{y_{\max}(\text{PER}) + y_{\max}(\text{CRY1}) + y_{\max}(\text{CRY2})}$				
9	Fraction CRY2 of total protein	3	0.341	0.358	0.373
	$\frac{y_{\max}(\text{CRY2})}{y_{\max}(\text{PER}) + y_{\max}(\text{CRY1}) + y_{\max}(\text{CRY2})}$				
10	<i>Cry1</i> siRNA period sensitivity	5	< 0	< 0	> 0
	$\frac{\partial T}{\partial v_{\text{deg},\text{c1}}}$				
11	<i>Cry2</i> siRNA period sensitivity	5	> 0	> 0	< 0
	$\frac{\partial T}{\partial v_{\text{deg},\text{c2}}}$				

Description	Weight	Desired	Degradation	Activation
12 <i>Cry1</i> knockout period	5	< 95%	92.1%	193.4%
		$\frac{T(Cry1^{-/-})}{T(WT)}$		
13 <i>Cry2</i> knockout period	5	> 115%	131.8%	99.4%
		$\frac{T(Cry2^{-/-})}{T(WT)}$		
14 Fraction CRY1 entering nucleus	1	0.40	0.195	0.059
		$\frac{y_{\max}(CRY1_n)}{y_{\max}(CRY1 + CRY1_n)}$		
15 Fraction CRY2 entering nucleus	1	0.35	0.139	0.059
		$\frac{y_{\max}(CRY2_n)}{y_{\max}(CRY2 + CRY2_n)}$		
16 Time delay between nuclear re- pressors and mRNA	3	75%	0.850	0.848
		$t_{\max}(\text{nuclear protein}) - t_{\max}(\text{mRNA})$		

	Description	Weight	Desired	Degradation	Activation
17	Time delay between mRNA and cytoplasmic protein	3	25%	0.057	0.095
	$t_{\max}(\text{mRNA}) - t_{\max}(\text{protein})$				
18	Time delay between cytoplasmic protein and nuclear protein	3	0%	0.093	0.057
	$t_{\max}(\text{protein}) - t_{\max}(\text{nuclear protein})$				

2.2.3 Model validation and dynamics

The model was validated by comparing the simulated dynamics to experimental measurements. First, the time course plots of the state variables display reasonable phases and amplitudes, and oscillate with a period of 23.7 hours (figure 2.3). The knockout periods are 21.7 hr (91.4% of WT) for $Cry1^{-/-}$ and 31.5 hr (133% of WT) for $Cry2^{-/-}$, indicating that the two feedback loops are indeed redundant with different free-running periods.

SiRNA knockdowns

SiRNA knockdowns were performed *in silico* by increasing the degradation rate for the corresponding mRNA; shown in figure 2.4, as done previously in [22]. $Cry1$ and $Cry2$ knockdowns in wild type conditions show close agreement to experiment [6],




Figure 2.3: Time course concentration profiles of model 2.1. The three polar plots show the time-varying levels of each clock component for optimal parameter set. In these plots, the amplitude of each variable (always positive) is plotted against its phase, $0 \rightarrow 2\pi$. Since the limit cycle is periodic, this results in a closed curve. Rising mRNA levels caused by low repressor concentrations (CT12) result in accumulating cytosolic protein (left plot). Lower levels of PER prevent all the available CRY from entering the nucleus (middle plot). High levels of nuclear repressors halt transcription until both CRYs have degraded (right plot).

and demonstrate that altering the nuclear CRY1/CRY2 ratio is effective in changing the period of oscillation.

CRY2 cytosolic stabilization

The *Dyrk1a* knockdown in [16] serves as another demonstration of the change in period as a response to a change in nuclear CRY1/CRY2 ratio. To demonstrate the effect in silico, the degradation rate of cytosolic CRY2 was decreased. Matching experimental evidence, the cytosolic CRY2 levels rose, with a corresponding change in the nuclear CRY ratio and decrease in period.

Single cry perturbations

The *Cry1^{+/−} Cry2^{−/−}* and *Cry1^{−/−} Cry2^{+/−}* single/double knockout perturbations ([5]) were approximated with an appropriate knockout and siRNA knockdown, and show the correct period shortening. With one *Cry* knocked out, the levels of cyto-




Figure 2.4: Model validation against experimental results. When comparing against existing experimental results, the model shows the correct response for *Cry1* siRNA, *Cry2* siRNA, and CRY_c stabilization (left, middle), through an adjustment in the nuclear $CRY1/CRY2$ ratio. The model also correctly captures the single/double knockout phenotype of [5], (right)

plasmic PER are no longer stoichiometrically limiting. In this case, less *Cry* leads to less nuclear complex, which is cleared faster.

2.2.4 Prediction of KL001 mechanism

Using the completed model, I investigated possible mechanisms by which KL001 might lengthen the circadian period. Experimental evidence indicates that stabilization of CRY can either lengthen the period (*Fbxl3* knockdown) or shorten the period (*Dyrk1a* knockdown). Assuming equal effect on both isoforms of *Cry*, the model confirms that cytoplasmic stabilization results in period shortening, since excess CRY expedites the nuclear import of the PER:CRY repressive complex. However, stabilizing nuclear CRY (figure 2.2, last column) yields the appropriate period lengthening observed experimentally. We therefore predicted that KL001 acts primarily in the nucleus, potentially through the FBXL3 degradation pathway. Additionally, the model




Figure 2.5: Prediction of the KL001 mechanism. *In silico* response to equal stabilization of both CRYs in the nucleus results in longer periods (middle column), matching observed results. The model correctly predicts that stabilization of individual CRYs in the knockout environments also causes period lengthening.

predicts that the compound's effect on $Cry1^{-/-}$ and $Cry2^{-/-}$ systems would be similar to its effect on the wild type clock, causing dose-dependent period lengthening (figure 2.5, left and right).

Experimental confirmation

Experimental tests of the model's predictions were performed. Nuclear CRY1 and CRY2 levels were up-regulated and almost sustained, respectively, while PER1 level was strongly down-regulated by the compound, supporting stabilization of nuclear CRY, the predicted mechanism. Additionally, continuous treatment with KL001 lengthened the period in both $Cry1$ and $Cry2$ knockout cells in a dose-dependent manner. Similarly, the compound caused period lengthening in both CRY1 and CRY2

knockdown U2OS cells. We further investigated the effect of KL001 in SCN explants, which show robust rhythms even in the absence of *Cry1*, due to intercellular coupling [23]. Both *Cry1* and *Cry2* knockout SCN exhibited dose-dependent period lengthening by the compound treatment. Thus, in a single *Cry* knockout, stabilization of either nuclear CRY1 and CRY2 causes period lengthening, confirming model predictions.

2.3 Discussion

2.3.1 Insights into circadian network design

Mathematical modeling has revealed how the balancing of two redundant feedback loops can provide fine control over the oscillatory period. A contour plot of period vs CRY1 or CRY2 abundance is shown in figure 2.6, which shows that the model's period is largely insensitive to the total amount of CRY (45° line), but highly dependent on the ratio of the two isoforms. Sensitivity analysis of our mathematical model (figure 2.7) reveals that subtle changes in most of the involved rates can have an effect on the clock's free running period, which could be caused by evolutionary noise. Since many studies have linked entrainment to non-natural periods with long-term health problems, a mechanism to align the clock's natural period to that of the environment would be advantageous. It is therefore possible that the period control afforded by CRY1/CRY2 balancing is a deliberately conserved design principle of the circadian clock, which confers period robustness against random mutations of clock components. Indeed, the presence of independent control of the cytosolic CRY2 levels suggests that the biological clock can control its period through simple post-translational modifications [16].




Figure 2.6: Simultaneous knockdown of *Cry1* and *Cry2*. Period contours (% change) resulting from simultaneous *in silico* siRNA knockdowns of *Cry1* (x-axis) and *Cry2* (y-axis). The plot shows that the CRY1/CRY2 ratio determines the period, independent of total CRY, for perturbations up to twice the normal mRNA degradation rate.




Figure 2.7: First order relative period sensitivities. Dimensionless period sensitivities with respect to the kinetic parameters in the degradation-based model. Perturbations to most clock rates result in noticeable changes to the free-running period. Many of the most sensitive parameters are transcriptional rates, which can be easily changed through mutation of DNA promoter regions.

2.3.2 Identifiability of model parameters

Due to the high dimensionality of the model equations and sparsity of the experimental data, it is likely that more than one set of parameters would fit the cost function defined in table 2.2. While the fact that *any* parameter set for the given model equations reproduces experimental results is enough to confirm that our hypothesized mechanism is mathematically consistent, we must be careful in placing too much confidence in predictions based on the particular parameter set generated by the optimization algorithm. While subsequent experimental results confirm many of our model predictions, a more rigorous exploration of parameter space would allow us to determine which predictions are constrained by the available experimental data. However, incorporating error ranges via standard techniques such as bootstrap techniques are infeasible when using a genetic algorithm optimization strategy. In chapter 3, I describe techniques to improve computational efficiency to allow a more systematic evaluation of predictive confidence.

Chapter 3

Identifiability analysis for models of circadian rhythms¹

3.1 Background

A cell's behavior is governed by the dynamic and selective expression of its genes, in which each protein's activity depends on a careful balance between transcription, translation, transport, and degradation rates. These rates, which change with environmental conditions and are often impossible to measure accurately *in vivo* or *in vitro*, determine the function of a regulatory pathway. While studying the roles of individual proteins can often provide some insight on how a particular function is achieved, this approach is limited in explaining complicated cellular phenomena at the scale of dozens to hundreds of interacting genes. With the aid of mathematical models, it is increasingly possible to create *in silico* realizations of genetic regulatory networks to examine their dynamic properties. For example, in the previous chapter, I presented a mathematical model of gene regulation for circadian rhythms,

¹Portions of this chapter are published in P. C. St. John and F. J. Doyle, "Estimating confidence intervals in predicted responses for oscillatory biological models," *BMC Syst. Biol.*, vol. 7, p. 71, July 2013.

using new data from small molecule modulators to gain further insight into clock dynamics.

Essential to understanding how genetic circuits operate is connecting how inputs (i.e., environmental changes, extracellular signals) are processed to give the appropriate outputs (protein expression, cellular response). Models of genetic regulatory networks, often sets of ordinary differential equations (ODEs), contain many unknown parameters that must be estimated from experimental data [24]. Derivatives of the model output with respect to changes in input, known as local sensitivities, are frequently validated experimentally or used to predict potential targets for pharmaceuticals [25]. In chapter 2, I used local sensitivities within the cost function to ensure the model matched previously known experimental results. However, since sensitivities can change drastically with respect to the particular parameter values chosen, the confidence associated with parameter and sensitivity values is an important consideration in model analysis and design.

3.1.1 Identifiability analysis

Practical identifiability analysis is concerned with calculating confidence intervals in parameter estimates resulting from uncertainty in experimental data [26]. Several techniques for such an analysis currently exist, and are commonly used in analyzing biological models [27, 28, 29]. In one method, the inverse of the Fisher information matrix is used to provide estimates of the variance in each parameter. However, since this method assumes a linearized model, the resulting symmetric normal distributions for each parameter do not accurately reflect the mapping of nonlinear models [30]. In the bootstrap method, distributions in parameter estimates are found through optimum fits to repeated physical or *in silico* measurements. While accurate in finding the true nonlinear confidence intervals, this approach requires efficient and robust parameter estimation convergence.

Difficulties imposed by periodic systems

In periodic systems, such as the models examined in this thesis, the behavior (and existence) of limit cycle oscillations is a discontinuous function of the parameters. Optimal values are traditionally found through trial-and-error type approaches [10, 12] or genetic algorithm search strategies [11], both of which are not amenable to bootstrap methods. Additionally, since the solutions are oscillatory, additional care must be taken in the calculation of the first-order sensitivity values. In this chapter, we calculate the sensitivity of the oscillatory period to parameter perturbation, a biologically relevant quantity that is often measured experimentally [21]. Due to these complications, rigorous identifiability analyses of these models are typically not performed.

In this chapter, a bootstrap uncertainty analysis appropriate for oscillatory biological models is developed and applied to the model of circadian rhythms described in chapter 2 [3]. These networks serve as an excellent example of a functional genetic circuit, able to process subtle environmental cues while remaining robust to temperature variations and evolutionary disturbances. Accurate limit cycle models must capture not only the correct time-dependent dynamics, but also the correct input-output response. For circadian rhythms, high-throughput microarrays have provided high-resolution time-series data of gene expression levels [31]. Additionally, knockdown experiments using RNA interference technology (siRNA) and small molecule modulators have resulted in a wealth of data on the dynamic responses to changes in key rates [3, 6, 32, 33]. This data, together with qualitative knowledge of the underlying network structure, permits the use and verification of a suitable uncertainty analysis.

Alternative parameter estimation methods

To enable a bootstrap approach, we employ an efficient parameter estimation routine optimized for limit cycle models. Motivated by the increasing availability of high-resolution time-series measurements, we use an approach similar to multiple shooting, in which a nonlinear and discontinuous parameter estimation problem is transformed into a high-dimensional yet local optimization and solved via nonlinear programming [34]. Since the desired shape of the limit cycle solution is known *a priori*, a relatively accurate initial guess for the parameters and trajectories can be found. By using multiple sets of *in silico* data of varying quality, we illustrate how error in experimental results is propagated to uncertainty in parameter sensitivity. Lower quality data - with either higher error or fewer sampling points - result in wider distributions of limit cycles and less identifiable responses. These results can be used in *a priori* experimental design, finding the minimum sampling points needed for an estimated experimental error to enable accurate modeling. Additionally, we show using literature data how this method can be used to discriminate between candidate model structures, revealing which one yields the highest predictive confidence.

3.2 Methods

3.2.1 Collocation methods

The estimation of the unknown kinetic parameters is accomplished via nonlinear programming [34]. In this method, we divide the limit cycle trajectory $\mathbf{x}(t, \mathbf{p})$ into \mathcal{N} finite elements of length h , and approximate each with a \mathcal{K} degree Lagrange interpolating polynomial, $\mathbf{x}_i^{\mathcal{K}}(t)$, using an internal time $\tau \in [0, 1]$. For finite element

i :

$$\begin{aligned} t &= h(i + \tau) \\ \ell_j(\tau) &= \prod_{k=0, k \neq j}^{\mathcal{K}} \frac{\tau - \tau_k}{\tau_j - \tau_k} \\ \mathbf{x}_i^{\mathcal{K}}(\tau) &= \sum_{j=0}^{\mathcal{K}} \ell_j(\tau) \mathbf{x}_{ij}. \end{aligned} \tag{3.1}$$

We also ensure that the interpolating polynomial matches system dynamics at each collocation point, τ_k , by setting

$$\sum_{j=0}^{\mathcal{K}} \mathbf{x}_{ij} \frac{d\ell_j(\tau_k)}{d\tau} = h\mathbf{f}(\mathbf{x}_{ij}, \mathbf{p}) \tag{3.2}$$

for $k = 1, \dots, \mathcal{K}$.

Additionally, the interpolating polynomials for each finite element must form a continuous function, so the following continuity constraints are imposed:

$$\mathbf{x}_{i+1,0} = \sum_{j=0}^{\mathcal{K}} \ell_j(1) \mathbf{x}_{i,j} \tag{3.3}$$

for $i = 1, \dots, \mathcal{N} - 1$.

Periodic conditions are imposed by setting the beginning of the first element equal to the end of the final element:

$$\mathbf{x}_{0,0} = \sum_{j=0}^{\mathcal{K}} \ell_j(1) \mathbf{x}_{\mathcal{N},j} \tag{3.4}$$

The τ_k values are chosen for optimal accuracy, here we use Gauss-Radau roots so that the resulting method has stiff decay [34]. With $\mathcal{K} = 5$:

$$\tau = \{0.000, 0.057, 0.277, 0.584, 0.860, 1.000\}$$

The interpolating polynomials can now be compared to the experimental data. For each measured value, $\hat{\mathbf{x}}(t_n)$, the corresponding simulated values $\mathbf{x}^{\mathcal{K}}(t_n)$ can be interpolated from \mathbf{x}_{ij} :

$$\mathbf{x}^{\mathcal{K}}(t_n) = \sum_{j=0}^{\mathcal{K}} \ell_j(\tau_n) \mathbf{x}_{ij} \quad (3.5)$$

$$\text{for } n = 1, \dots, \mathcal{M}$$

where i and τ_n are selected for the appropriate finite element and sampling time. The objective function $\Phi(\mathbf{x}, \mathbf{p})$ is thus:

$$\Phi(\mathbf{x}, \mathbf{p}) = \sum_n^{\mathcal{M}} \left(\frac{\mathbf{x}^{\mathcal{K}}(t_n) - \hat{\mathbf{x}}(t_n)}{\mathbf{\sigma}_n} \right)^2 \quad (3.6)$$

where $\mathbf{\sigma}_n$ is the measurement error associated with measurement n . Since \mathbf{x} and $\mathbf{\sigma}$ are vectors, the division in equation (3.6) must be performed element-wise. This cost function was taken from a similar multiple-shooting approach to parameter estimation [35].

Since the cost function (equation (3.6)) and equality constraints (equation (3.2), equation (3.3), and equation (3.4)) now satisfy continuity and differentiability requirements [36], parameter estimation can now be accomplished via constrained nonlinear programming (NLP) instead of a global search strategy. The solution is subject to variable bounds:

$$\begin{aligned} \mathbf{x}_{LB} &\leq \mathbf{x} \leq \mathbf{x}_{UB} \\ \mathbf{p}_{LB} &\leq \mathbf{p} \leq \mathbf{p}_{UB} \end{aligned} \quad (3.7)$$

The numerical implementation is accomplished using IPOPT [37], using the MA57 [38] linear solver. The CasADi computer algebra package [39] was used to provide an interface to the IPOPT numerical libraries and supply derivatives to the cost and equality function calls through automatic differentiation.

3.2.2 Generating initial values

Solution of the NLP described in section 3.2.1 requires a suitable initial guess for the optimal state profiles, \mathbf{x}^* , and kinetic parameters, \mathbf{p}^* . To find approximate values for these variables, a smoothed periodic B-spline, $\tilde{\mathbf{x}}$, is found using experimental data for each state variable using SciPy's interpolate module [40]. Initial values for x_{ij} are obtained by evaluating this spline at each τ_k for each finite element.

$$\begin{aligned} \mathbf{x}_{ij}^* &\approx \tilde{\mathbf{x}}(t_{ij}) \\ \text{where } t_{ij} &= h(i - 1 + \tau_j) \\ \text{for } i &= \{1, \dots, \mathcal{N}\}, j = \{1, \dots, \mathcal{K}\} \end{aligned} \tag{3.8}$$

Since

$$\frac{d\tilde{\mathbf{x}}}{dt} \approx \mathbf{f}(\tilde{\mathbf{x}}, \mathbf{p}), \tag{3.9}$$

approximate values for \mathbf{p} can be obtained by solving the simpler unconstrained NLP,

$$\min_{\mathbf{p}} \sum_i^{\mathcal{N}} \sum_j^{\mathcal{K}} \left(\frac{d\tilde{\mathbf{x}}(t_{ij})}{dt} - \mathbf{f}(\tilde{\mathbf{x}}(t_{ij}), \mathbf{p}) \right)^2 \tag{3.10}$$

in which t_{ij} is the same as in 3.8 and the bounds on p are the same as in equation (3.7).

3.2.3 First order sensitivity analysis

After determining an optimal parameter set for the given experimental data, relevant first order sensitivity coefficients for oscillatory models are found using the procedure from [21], summarized here. First, initial conditions and oscillatory period are verified by solving the boundary value problem (BVP):

$$\min_{\mathbf{x}(0), T} \begin{pmatrix} \mathbf{x}(T) - \mathbf{x}(0) \\ \dot{\mathbf{x}}(0) \end{pmatrix} \tag{3.11}$$

where $\dot{\mathbf{x}}_0(0)$ denotes the time-derivative of the first state variable, evaluated at $t = 0$. This BVP is solved using Newton's method, employing the SUNDIALS packages CVODES for ODE integration and KINSOL for the Newton iterations [20].

Time-dependent parametric sensitivities,

$$\mathbf{S}(t) \equiv \frac{\partial \mathbf{x}(t)}{\partial \mathbf{p}}, \quad (3.12)$$

are obtained by using the staggered-direct method from the CVODES integrator [41]. Sensitivities of the period, $\frac{\partial T}{\partial \mathbf{p}}$, can be obtained directly from sensitivities integrated for one pass of the limit cycle (see [21] or [42] for further details) through a linear solve:

$$\begin{bmatrix} \mathbf{M} - \mathbf{I} & \dot{\mathbf{x}}(T) \\ \frac{\partial \mathbf{f}_0}{\partial \mathbf{x}}(\mathbf{x}(0)) & 0 \end{bmatrix} \begin{bmatrix} \vdots \\ \frac{\partial T}{\partial \mathbf{p}} \end{bmatrix} = \begin{bmatrix} -\mathbf{S}(T) \\ -\frac{\partial \mathbf{f}}{\partial \mathbf{p}}(\mathbf{x}(0)) \end{bmatrix} \quad (3.13)$$

in which \mathbf{M} is the Monodromy matrix, \mathbf{I} is the identity matrix, and the unknown vector contains the relevant period sensitivities.

Since parameter values often span several orders of magnitude, an often more useful measure is the relative period sensitivity, which is independent of the magnitude of the period or parameter value.

$$\frac{\partial \ln T}{\partial \ln \mathbf{p}} = \frac{\mathbf{p}}{T} \frac{\partial T}{\partial \mathbf{p}} = \frac{\partial T}{T} \Big/ \frac{\partial \mathbf{p}}{\mathbf{p}} \quad (3.14)$$

Thus, a relative period sensitivity of 1 indicates that a 1% increase in the parameter value will result in a 1% increase in the period.

3.2.4 Generation of data for bootstrap methods

For each run, two thousand simulated measurements, $\hat{x}_i(t_j)$, were generated from the true data, $\tilde{x}_i(t_j)$, using a normal distribution with $\mu = \tilde{x}_i(t_j)$ and $\sigma_{ij} = \xi \tilde{x}_i(t_j) +$

$\eta \max_j \tilde{x}_i(t_j)$, in which ξ is the relative and η is the absolute error. Each simulated data set was then used to find a unique optimum parameter set, \mathbf{p}^* . Data sets that failed to converge, or reached a steady state solution (in which periodic sensitivities are undefined), were discarded from further analysis.

For the *in silico* data of varying quality used in figures 3.2-3.4, we used the known limit cycle $\mathbf{x}(t)$ to generate data points $\hat{x}_i(t_j)$ at each of \mathcal{M} sampling points. The effect of increasing error and decreasing number sampling points were tested independently:

$$\xi = \{.01, .05, .10, .20, .30\}; \quad \mathcal{M} = 20$$

$$\mathcal{M} = \{30, 20, 15, 10, 5\}; \quad \xi = 0.15$$

Since standard deviations in the data distributions were also used as optimization weights, a small amount of absolute error ($\eta = 0.001$) was added to ensure errors in small values did not dominate the cost function.

3.2.5 Calculation times

Each parameter estimation took approximately 4 seconds on a 2.53GHz processor, with the subsequent limit cycle solution integration and sensitivity calculation taking approximately 0.5 seconds. Due to the parallel nature of the 2000 trials, computation times were alleviated by distributing the tasks onto a cluster of 160 compute nodes.

3.3 Results

Mechanistic models of biological processes are often posed as nonlinear, time-invariant systems of ordinary differential equations (ODEs) [10, 11, 12], of the form:

$$\frac{d\mathbf{x}}{dt} = \mathbf{f}(\mathbf{x}(t), \mathbf{p}) \quad (3.15)$$

in which the vector of state variables $\mathbf{x}(t)$ describe the time-dependent activity of important species (i.e., mRNA, proteins, or metabolites), the parameters \mathbf{p} are the kinetic rate constants, and the vector function $\mathbf{f}(\mathbf{x}(t), \mathbf{p})$ contains the transcription, translation, transport, and degradation rate laws of the gene regulatory network. In modeling rhythmic phenomena, we typically seek models and parameter values that display *limit cycle* oscillations - where for the solution approaches a non-trivial periodic trajectory:

$$\lim_{t \rightarrow \infty} \mathbf{x}(t) = \mathbf{x}(t + T). \quad (3.16)$$

Here the period of oscillation is the smallest $T > 0$ in which equation (3.16) holds. Limit cycle oscillations are independent of the system's initial values $\mathbf{x}(0)$, and are instead determined completely by the parameters \mathbf{p} .

Experimental values for \mathbf{p} are rarely available. Given time-series experimental measurements $\hat{x}_i(t_j)$ for each state variable in a limit cycle system, we find optimal parameters \mathbf{p}^* such that the error between the experimental measurements and the simulated limit cycle is minimized [35]:,

$$\mathbf{p}^* := \arg \min_{\mathbf{p}} \sum_i^{states} \sum_j^{data} \frac{(\hat{x}_i(t_j) - x_i(t_j, \mathbf{p}))^2}{\sigma_{ij}^2}. \quad (3.17)$$

Here σ_{ij} is the standard deviation associated with the measured mean of state i at time j . Using the data points $\hat{x}_i(t_j)$ to generate a suitable initial guess, parameter estimation may proceed via a nonlinear programming approach as described in 3.2.

In this chapter, we assume that all states are measured to demonstrate how initial guesses can be generated directly from the input data. However, for systems with unmeasured states, initial guesses for the trajectory and parameter values can be provided by another approach, such as a global optimization routine. A bootstrap method was implemented by repeatedly sampling input data distributions to calculate a population of optimal parameter fits.

After finding optimal parameter fits, we used the models to predict how perturbations change systems dynamics by performing a first order sensitivity analysis. Since adjustments to periodic systems in response to inputs are often manifested through temporary changes in oscillatory period, relative period sensitivities,

$$\frac{\partial \ln T}{\partial \ln p} \tag{3.18}$$

were calculated due to their independence of parameter magnitude [21, 42, 43]. Relative period sensitivities were integrated into the bootstrap method by calculating appropriate sensitivities for each estimated parameter set.

Of particular importance in determining the reliability of a model prediction is whether an output response maintains a consistent direction despite noise in measurement data. We therefore define a sensitivity value to be practically identifiable for given input data if 95% of the distribution maintains a consistent sign, similar to definitions for parameter identifiability used in previous studies [30, 44]. An overview of the method is shown in figure 3.1.

3.3.1 Effect of data quality on predictive confidence

We first analyze the degree to which uncertainty in input data is propagated to uncertainty in output predictions. To achieve this, we generate *in silico* data from a previously published model of circadian rhythms, using relative error ξ to generate




Figure 3.1: Parameter estimation and bootstrap methods flowchart. The demonstrated method calculates confidence intervals in the sensitivity of limit cycle models. An oscillatory model and experimental (or simulated) data are inputs to the bootstrap method. Unique data sets are then used to calculate optimum limit cycle trajectories. The resulting distribution in sensitivities highlight whether a particular response is identifiable (i.e., consistent across the majority of bootstrap trials.)

normally distributed data ($\sigma_{ij} = \xi \hat{x}_i(t_j)$) at each of \mathcal{M} sampling points. As expected, solution trajectories drifted further from the nominal limit cycle for higher values of error, ξ , or lower sampling density, \mathcal{M} , (figure 3.2). However, the overall shape of the oscillatory profiles remained relatively similar, even for rather high ξ or low \mathcal{M} .

Figure 3.3 shows violin plots of the probability distribution for each parameter set and corresponding sensitivity evaluation for increasing ξ , while figure 3.4 shows similar plots for decreasing \mathcal{M} . Interestingly, there is little correlation between the identifiability of a parameter and its corresponding sensitivity value. For example, v_{dP} , the maximum degradation rate of Per mRNA, shows a very tight clustering about its nominal parameter value, while the sensitivity of this parameter loses identifiability for even small values of ξ . Conversely, K_{dCn} , the Michealis-Menten constant associated with the degradation of nuclear CRY, shows large variations in possible




Figure 3.2: Time-course profiles of the state trajectories for *Per* mRNA. (A) Increasing relative error, ξ , with $\mathcal{M} = 20$. Possible state variable values are shown as shaded regions, obtained by filling between the 5th and 95th percentile for values at each time for 2000 independent parameter estimations. Increasing ξ results in larger deviations from the original model trajectory, shown as a dashed black line. (B) Decreasing number of measurement points, \mathcal{M} , each with $\xi = 0.15$. Higher \mathcal{M} results in trajectories closer to the true trajectory.

parameter values. However, the period sensitivity of KdCn, despite lying close to the x-axis, remains identifiable, indicating a robust prediction. These results reveal which model responses are constrained by the structure and dynamics of the limit cycle oscillations, and which are dependent on the particular parameterization chosen.

Sensitivities that are experimentally distinguishable from zero are the most important for validation. Calculating a typical experimental value for a relative period sensitivity helps to calibrate which sensitivities might be verified experimentally. Referring to a recent RNA interference screen, periods changes of approximately 1 hour (5%) can be reliably measured using luminescence recordings [6]. Assuming an increase in the corresponding mRNA degradation parameter value of 50%, this translates to a relative period sensitivity of 0.1. Thus, many of the identifiable values shown in figures 3.3-3.4 fall within the experimentally measurable range.

3.3.2 Application to literature data for model discrimination

We next apply the method to literature time-course data for core clock components [13]. When modeling a genetic regulatory network, many candidate model equations are often considered. We show that a bootstrap uncertainty analysis can also be useful in discriminating between potential model structures based on predictive confidence. Here two variations of the same model are fit: The first model (model 2.1) was originally optimized using a genetic algorithm approach, and thus contains a minimal number of parameters to reduce optimization complexity. The second model (model 3.1) considered contains independent parameters for each rate expression, increasing the number of parameters from 23 to 35.

The literature data used consisted of 7-8 concentration time points across a 24 hour period. Confidence intervals in the data were not available, so an optimistic 3% relative and 0.5% absolute error was assumed for each data point ($\sigma_{ij} = 0.03 \hat{x}_i(t_j) +$




Figure 3.3: Parameter and sensitivity identifiability for increasing error. Increasing ξ results in a corresponding decrease in the confidence of the parameter and sensitivity estimates. Violin plots of the parameter values (left) and relative period sensitivities (right) show the distribution of values from each parameter estimation. In the plots, a box plot is superimposed above a kernel density plot to convey the distribution of values. The whiskers used extend to the most extreme data point within 1.5x the inner quartile range. Sensitivities in which the 5th and 95th percentile values span the x-axis are deemed non-identifiable (red), as the model's response direction can not be accurately estimated. Higher ξ also results in wider parameter distributions.




Figure 3.4: Effect of high-resolution sampling on identifiability. Lower values of \mathcal{M} result in less constrained parameter and sensitivity values. Similar to figure 3.3, violin plots of the parameters (left) and sensitivities (right) show the distribution from each parameter estimation for decreasing \mathcal{M} . These results highlight the importance of high-resolution time sampling in generating sensitivity information for oscillatory models.

Table 3.1: Parameter values for model 3.1. Parameters were fit to time-series data via nonlinear programming.

	Parameter	Description	Value
1	M_1	<i>Per</i> /CRY2 activity coefficient	3.632
2	Vm_1	<i>Per</i> transcription rate	9.957×10^{-1}
3	Ki_1	<i>Per</i> /CRY inhibition coefficient	1.054×10^{-1}
4	M_2	<i>Cry1</i> /CRY2 activity coefficient	1.000×10^{-3}
5	Vm_2	<i>Cry1</i> transcription rate	2.262×10^{-1}
6	Ki_2	<i>Cry1</i> /CRY inhibition coefficient	2.049×10^{-1}
7	M_3	<i>Cry2</i> /CRY2 activity coefficient	1.000×10^1
8	Vm_3	<i>Cry2</i> transcription rate	1.850×10^{-1}
9	Ki_3	<i>Cry2</i> /CRY inhibition coefficient	1.427×10^{-1}
10	k_1	<i>Per</i> degradation rate	2.920×10^{-1}
11	Km_1	<i>Per</i> degradation self-inhibition	6.609×10^{-1}
12	k_2	<i>Cry1</i> degradation rate	1.000×10^1
13	Km_2	<i>Cry1</i> degradation self-inhibition	1.823×10^{-2}
14	k_3	<i>Cry2</i> degradation rate	4.711×10^{-2}
15	Km_3	<i>Cry2</i> degradation self-inhibition	1.000×10^1
16	k_4	<i>Per</i> translation rate	1.132×10^{-1}
17	k_5	<i>Cry1</i> translation rate	3.409×10^{-1}
18	k_6	<i>Cry2</i> translation rate	1.961×10^{-1}
19	k_7	PER degradation rate	1.000×10^1
20	Km_4	PER degradation self-inhibition	1.911×10^{-3}
21	k_8	CRY1 degradation rate	1.000×10^1
22	Km_5	CRY1 degradation self-inhibition	2.077×10^{-2}
23	k_9	CRY2 degradation rate	1.000×10^1
24	Km_6	CRY2 degradation self-inhibition	1.180×10^{-2}
25	k_{10}	C1N association rate	5.022×10^{-1}
26	k_{11}	C2N association rate	1.035
27	k_{12}	C1N dissociation rate	1.000×10^{-3}
28	k_{13}	C2N dissociation rate	2.070×10^{-1}
29	M_4	CRY1n/CRY2n activity coefficient	1.000×10^1
30	k_{14}	CRY1N degradation rate	1.000×10^1
31	Km_7	CRY1n degradation inhibition	5.206×10^{-3}
32	M_5	CRY2n/CRY2n activity coefficient	1.000×10^1
33	k_{15}	CRY2n degradation rate	1.000×10^1
34	Km_8	CRY2n degradation inhibition	1.829×10^{-2}




Figure 3.5: Identifiability comparison of two model structures. (A) Bootstrap parameter estimations on two model structures using literature time-series data with estimated errors (box plots). Resulting regions of model trajectories are shaded between the 5th and 95th percentile. Per species are shown in purple, *Cry1* in red, and *Cry2* in green. While both models were able to approximately reproduce the same dynamic response, the expanded model was better able to capture differences between the *Cry1* and *Cry2* profiles. (B) Parameter and sensitivity identifiability for the base and expanded models. Violin plots show the parameter and sensitivity distributions, with unidentifiable sensitivities (90% confidence level) highlighted in red. Despite containing more parameters, the expanded model shows better parameter identifiability and higher confidence in its predicted sensitivities. The PER translation rate (1) and PER-CRY association rate (2) sensitivities are consistent across model equations and are highlighted.

Model 3.1: An expanded parameterization of model 2.1. These equations have similar reaction stoichiometry to those in table 2.1, but with more parametric degrees of freedom. This model showed better time-series performance than the more constrained model when fit to time-series data.

$$\begin{aligned}
 \frac{d\mathbf{p}}{dt} &= \frac{Vm_1}{1 + Vm_1 \left(\frac{\mathbf{C1N} + \mathbf{C2N} M_1}{Km_1} \right)^3} - \frac{k_1 \mathbf{p}}{1 + \frac{\mathbf{p}}{Km_1}} \\
 \frac{d\mathbf{c1}}{dt} &= \frac{Vm_2}{1 + Vm_2 \left(\frac{\mathbf{C1N} + \mathbf{C2N} M_2}{Km_2} \right)^3} - \frac{k_2 \mathbf{c1}}{1 + \frac{\mathbf{c1}}{Km_2}} \\
 \frac{d\mathbf{c2}}{dt} &= \frac{Vm_3}{1 + Vm_3 \left(\frac{\mathbf{C1N} + \mathbf{C2N} M_3}{Km_3} \right)^3} - \frac{k_3 \mathbf{c2}}{1 + \frac{\mathbf{c2}}{Km_3}} \\
 \frac{d\mathbf{P}}{dt} &= k_4 \mathbf{p} + k_{12} \mathbf{C1N} + k_{13} \mathbf{C2N} - \frac{k_7 \mathbf{P}}{1 + \frac{\mathbf{P}}{Km_4}} - k_{10} \mathbf{P} \mathbf{C1} - k_{11} \mathbf{P} \mathbf{C2} \\
 \frac{d\mathbf{C1}}{dt} &= k_5 \mathbf{c1} + k_{12} \mathbf{C1N} - \frac{k_8 \mathbf{C1}}{1 + \frac{\mathbf{C1}}{Km_5}} - k_{10} \mathbf{P} \mathbf{C1} \\
 \frac{d\mathbf{C2}}{dt} &= k_6 \mathbf{c2} + k_{13} \mathbf{C2N} - \frac{k_9 \mathbf{C2}}{1 + \frac{\mathbf{C2}}{Km_6}} - k_{11} \mathbf{P} \mathbf{C2} \\
 \frac{d\mathbf{C1N}}{dt} &= k_{10} \mathbf{P} \mathbf{C1} - k_{12} \mathbf{C1N} - \frac{k_{14} \mathbf{C1N}}{1 + \frac{\mathbf{C1N} + \mathbf{C2N} M_4}{Km_7}} \\
 \frac{d\mathbf{C2N}}{dt} &= k_{11} \mathbf{P} \mathbf{C2} - k_{13} \mathbf{C2N} - \frac{k_{15} \mathbf{C2N}}{1 + \frac{\mathbf{C1N} + \mathbf{C2N} M_5}{Km_8}}
 \end{aligned}$$

0.005 $\max(\hat{x}_i)$). Figure 3.5A shows the resulting time-series profiles for bootstrap estimations of each model. While additional kinetic parameters are typically thought to lower the predictive confidence of a model (the ‘curse of dimensionality’), the expanded model is able to better capture the oscillatory profiles with lower variability between solutions. Parameter and sensitivity distributions, figure 3.5B, similarly show how the expanded model parameterization is able to generate more confident predictions in model response. Since the resulting sensitivity identifiability for both models was relatively poor, we highlight sensitivities which pass a 90% confidence level threshold. These results thus indicate higher-resolution data on circadian com-

ponents would help in conferring confidence to model predictions.

Two sensitivities, the PER translation rate (figure 3.5B, 1) and the PER-CRY association rate (2), had high confidence and consistent direction in both the base and expanded parameterization - suggesting that the predicted values are robust to slight changes in both parameter value and model structure. Since a biological system can be modeled using many different combinations of kinetic assumptions, such a technique will likely prove useful in finding consistent predictions which are robust to slight differences in model equations.

Figure 3.6 compares optimal fits for both the base and expanded models to the originally published parameter set [3]. Since the original cost function was concerned mostly with optimizing stoichiometric and knockout data, the refitted models are able to much more accurately represent the time-series dynamics.

3.4 Discussion

Increasingly, mathematical models are being used to study biological systems where traditional experiments would prove infeasible. For example, in the search for drug targets, thousands of possible combinatorial perturbations can be quickly scanned for therapeutic effects using *in silico* modeling. This is especially useful in oscillatory systems with long periods, such as circadian rhythms, where a perturbed *in vitro* or *in vivo* system must be measured for multiple days before changes can be reliably determined.

However, since errors in model responses can arise from either incorrect structure or measurement noise, our confidence in *in silico* predictions is limited. Here we have developed a bootstrap approach suitable for periodic systems, and extended it to include uncertainty in predicted responses. With this method, errors due to local parameter effects can be identified, even in models with complicated dynam-




Figure 3.6: Time-series dynamics of fitted models. Model trajectories for each of the considered models. The original model (purple) shows the time series dynamics for the previously published parameter set, used in Figures 1-4. In these plots, the original model's period and amplitudes were rescaled to best match the experimental data, shown in black (without changing the dynamic profiles). The refitted model (red, dashed) was generated by optimizing the dynamic profile to time series data using the parameter estimation routine described in Supplemental Text 1. The expanded model (green, dashed) consists of a similar network structure, but with independent kinetic parameters for each rate expression (see Supplemental Text 2). These plots show that parameter optimization through nonlinear programming is able to more accurately fit gene and protein expression profiles.

ics. Furthermore, by considering multiple variations in model assumptions, we have demonstrated that more trustworthy model predictions can be found.

Since this method takes advantage of time-series data to generate a strong initial guess for an otherwise difficult parameter estimation, it requires high-resolution data on the concentrations of all species in the model. In many biological systems, such data is only available for the activity levels of certain well-studied species. However, the continued development of high-throughput genomic and proteomic techniques promise to deliver time-series data for a much larger network of components. With expanding datasets, these methods will likely prove useful for the quantitative evaluation of uncertainty in larger biological models.

In the following chapter, I apply this method to several existing circadian models in order find conserved predictions across different model kinetics and parameterizations. Such an approach allows us to further differentiate between the actions of two seemingly similar small molecule perturbations.

Chapter 4

Spatiotemporal separation of PER and CRY posttranslational regulation¹

In the previous chapter, computationally efficient techniques for performing a bootstrap analysis on circadian models was described. In this chapter, we apply this technique to clarify the roles of two posttranslational regulators in the core circadian feedback loop. Since direct experimental evidence of the systems-level effect of these regulators is difficult if not impossible to obtain, we must rely on computational modeling to provide insight into the dynamic consequences of modulating posttranslational activity. For this reason, bootstrap methods are invaluable in generating confident *in silico* results.

4.1 Background

Since circadian and metabolic regulators are tightly integrated, circadian disruptions often manifest in metabolic disease [45]. Recent efforts have therefore sought to gain a mechanistic understanding of these pathways, such that the metabolic burdens

¹Portions of this chapter are published in P. C. St. John, T. Hirota, S. A. Kay, and F. J. Doyle, "Spatiotemporal separation of PER and CRY posttranslational regulation in the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 111, pp. 2040–5, Feb. 2014.

imposed by a 24-hour society might be mitigated. Posttranslational regulators, which play key roles in connecting circadian and metabolic processes, serve as likely targets for future therapeutics – demonstrated by the wealth of available circadian-active small molecules [2].

4.1.1 Nuclear entry of PER-CRY complex

Oscillations in circadian gene transcription are generated through a time-delayed transcription-translation negative-feedback loop. In mammals, transcription factors CLOCK and BMAL1 promote transcription of E box-containing genes *Period* (*Per*) and *Cryptochrome* (*Cry*) (figure 4.1). PER and CRY protein products form heterodimers to accumulate in the nucleus, in which PER is stoichiometrically limiting [13], and subsequently close the negative feedback loop by inhibiting CLOCK-BMAL1-promoted gene expression. While steady-state endpoint assays have shown the possibility of nuclear entry of CRY without PER [9, 46, 47], experiments from *Per1*^{-/-} *Per2*^{-/-} mice demonstrated that PER proteins are required for the timely nuclear accumulation of CRY [13]. Clearance of nuclear repressors reactivates CLOCK-BMAL1, allowing the cycle to begin anew [48].

Experimental evidence on PER/CRY nuclear entry is seemingly contradictory. For nuclear localization of the PER and CRY proteins, experiments from *Per1*^{-/-} *Per2*^{-/-} mice demonstrate that PER proteins are required for timely nuclear accumulation of CRY [13]. While other studies have shown the possibility of nuclear entry of CRY without PER, these results are typically based on steady-state endpoint assays which do not consider the speed of CRY nuclear entry [9, 46, 47]. We therefore consider the formation of the PER-CRY heterodimer as a key step in nuclear entry, which is supported by the fact that to the best of our knowledge all circadian models that consider both PER and CRY employ this kinetic assumption [3, 10, 11, 12, 22].




Figure 4.1: Small molecule targets. Schematic of the core circadian feedback loop, with the targets of small molecule modulators longdaysin (CKI inhibitor) and KL001 (inhibitor of FBXL3-dependent CRY degradation) shown. The size of each molecule is representative of relative concentration

4.1.2 Importance of posttranslational regulators

The stabilities of PER and CRY are tightly regulated: PER proteins are phosphorylated by the casein kinase I family of proteins (CKI δ/ϵ), prompting β -TrCP-mediated degradation [49] and nuclear import [50]. The degradation of CRY proteins is separately regulated by the SCF^{FBXL3} ubiquitin ligase complex [51, 52, 53]. The activities of both CKI-PER and FBXL3-CRY may be further coupled to the cell's metabolic state through AMPK signalling [54]. These posttranslational regulatory mechanisms have a strong effect on period length: the gain-of-function mutant CKI ϵ^{tau} leading to hyperphosphorylation of PER [55] and small molecule CKI inhibitors, such as longdaysin [32], demonstrated that increasing or decreasing CKI-dependent PER phosphorylation shortens or lengthens the period, respectively. In contrast, genetic mutations of FBXL3 [52, 53] and KL001, a small molecule inhibitor of FBXL3-dependent CRY degradation [3], showed that increased CRY stability leads to longer periods.

Since the scale and complexity of the circadian network complicates an intuitive understanding of these relationships, mathematical models have played important roles in understanding how these manipulations affect circadian period [3, 49, 55].

4.1.3 Functional differences between PER and CRY

Given that both CKI and FBXL3 pathways regulate the stability of linked negative factors, it was thought that simultaneous perturbations to both pathways might lead to non-additive effects: i.e., the slowest link would determine the period. However, both small molecule [3] and genetic experiments [56] have demonstrated the independent period effects of these two posttranslational regulations. A recent clarification of the canonical clock feedback circuit has shown that dissociated CRY is the dominant repressor of CLOCK-BMAL1 mediated E box transcription [46]. This distinction helps differentiate between the roles of the otherwise similar PER and CRY proteins, in which the main role of PER in transcriptional repression is likely regulating the timing of nuclear accumulation of CRY. Therefore, while previous mathematical models in which PER acts as a direct repressor have proposed mechanisms for CKI-dependent period lengthening [55, 57], they are likely not suitable for distinguishing between CKI-PER and FBXL3-CRY mediated period change.

In this chapter, we used human cells harboring clock gene reporters together with mathematical modeling to gain insight into the relationship between PER and CRY posttranslational regulation. Consequently, we provide a new mechanism by which CKI-dependent PER phosphorylation controls the circadian period separately from the FBXL3-CRY pathway. The resulting detailed understanding of PER and CRY regulation in the core feedback loop provides a framework on which to interpret metabolic and pharmacological control of circadian rhythms.

4.2 Methods

4.2.1 Analysis of luminescence profiles

Raw luminescence data was first separated into a moving baseline and oscillatory component using a Hodrick-Prescott filter with a smoothing parameter of 1600. Example trajectory decompositions are shown in figure 4.2. Amplitudes (as shown in figure 4.3) were determined by taking the standard deviation in the baseline-subtracted data. Periods were obtained by nonlinear curve fitting, in which a four parameter (initial amplitude, decay, period and phase) damped cosine curve was fit to the baseline-subtracted data. Periods were not shown if the relative amplitude (found by standard deviation) fell below 25%, since noise dominated the periodic trajectory.

4.2.2 Cost function

Models were fit to a cost function of experimental results. *Per*, *Cry*, *Clock*, and *Bmal1* protein and mRNA levels were taken from [13], along with profiles of CRY nuclear localization. For the model from [22], additional activity profiles on *Rev-Erb* and *Ror* were obtained from CircaDB (<http://bioinf.itmat.upenn.edu/circa/>). To score a model trajectory, mRNA state variables were scaled independently to minimize the squared error between model and experiment, since model parameters could be adjusted to give mRNA profiles arbitrary amplitudes. For protein species, where stoichiometric interactions are important, a single scaling parameter was used for all species. Nuclear repressor species, in which only relative measurements were available, were scaled independently. Full model equations for the model from Hirota *et al.*, 2012 are shown in model 2.1, the model from Relógio *et al.* 2011 in model 4.1, and the model from Leloup & Goldbeter, 2003 in model 4.2.




Figure 4.2: Analysis of circadian reporter luminescence data. (A) Raw luminescence data is first cropped by removing the initial transient region (first 12 points). The moving baseline is estimated using a Hodrick-Prescott filter with smoothing parameter 1600 (red dashed line). (B) Data is detrended by subtracting the moving baseline from the raw data. The detrended data is used to calculate the relative amplitude of the oscillations via standard deviation. (C) Periods are estimated by fitting a damped cosine curve (green solid line) to the detrended data.

Model 4.1: Model from Relógio *et al.* 2011 [22].

$$\begin{aligned}
 \frac{d \text{CLKBM1}}{dt} &= \text{BM1n} \cdot \text{kfCLKBM1} - \text{CLKBM1} \cdot \text{dCLKBM1} - \text{CLKBM1} \cdot \text{kdCLKBM1} \\
 \frac{d \text{reverb}}{dt} &= \frac{V3 \max \left(g \cdot \frac{\text{CLKBM1}^v}{k13} + 1 \right)}{\frac{\text{CLKBM1}^v}{k13} \left(\frac{PnCn + PnpCn}{k13} \right)^w + \frac{\text{CLKBM1}^v}{k13} + 1} - \text{dreverb} \cdot \text{reverb} \\
 \frac{d \text{ror}}{dt} &= \frac{V4 \max \left(h \cdot \frac{\text{CLKBM1}^p}{k14} + 1 \right)}{\frac{\text{CLKBM1}^p}{k14} \cdot \left(\frac{PnCn + PnpCn}{k14} \right)^q + \frac{\text{CLKBM1}^p}{k14} + 1} - \text{dror} \cdot \text{ror} \\
 \frac{d \text{REVERBc}}{dt} &= -\text{REVERBc} \cdot \text{dREVERBc} - \text{REVERBc} \cdot \text{kiREVERBc} + \text{kp3} \cdot \text{reverb} \\
 \frac{d \text{RORc}}{dt} &= -\text{RORc} \cdot \text{dRORc} - \text{RORc} \cdot \text{kiRORc} + \text{kp4} \cdot \text{ror} \\
 \frac{d \text{REVERBn}}{dt} &= \text{REVERBc} \cdot \text{kiREVERBc} - \text{REVERBn} \cdot \text{dREVERBn} \\
 \frac{d \text{RORn}}{dt} &= \text{RORc} \cdot \text{kiRORc} - \text{RORn} \cdot \text{dRORn} \\
 \frac{d \text{bm1}}{dt} &= \frac{V5 \max \left(i \cdot \frac{\text{RORn}^n}{k15} + 1 \right)}{\frac{\text{REVERBn}^m}{k15} + \frac{\text{RORn}^n}{k15} + 1} - \text{bm1} \cdot \text{dbm1} \\
 \frac{d \text{BM1c}}{dt} &= -\text{BM1c} \cdot \text{dBM1c} - \text{BM1c} \cdot \text{kiBM1c} + \text{bm1} \cdot \text{kp5} \\
 \frac{d \text{BM1n}}{dt} &= \text{BM1c} \cdot \text{kiBM1c} - \text{BM1n} \cdot \text{dBM1n} - \text{BM1n} \cdot \text{kfCLKBM1} + \text{CLKBM1} \cdot \text{kdCLKBM1} \\
 \frac{d \text{per}}{dt} &= \frac{V1 \max \left(a \cdot \frac{\text{CLKBM1}^b}{k11} + 1 \right)}{\frac{\text{CLKBM1}^b}{k11} \cdot \left(\frac{PnCn + PnpCn}{k11} \right)^c + \frac{\text{CLKBM1}^b}{k11} + 1} - \text{dper} \cdot \text{per} \\
 \frac{d \text{cry}}{dt} &= \frac{V2 \max \left(\frac{\text{CLKBM1}^d}{k12^3} + 1 \right)}{\left(\frac{\text{REVERBn}^f}{k121} + 1 \right) \left(\frac{\text{CLKBM1}^3}{k12^3} + \frac{\text{CLKBM1}^e}{k12} \left(\frac{PnCn + PnpCn}{k12} \right)^f + 1 \right)} - \text{cry} \cdot \text{dcry} \\
 \frac{d \text{Cc}}{dt} &= -\text{Cc} \cdot \text{Pc} \cdot \text{kfPcCc} - \text{Cc} \cdot \text{Pcp} \cdot \text{kfPcpCc} - \text{Cc} \cdot \text{dCc} + \text{PcCc} \cdot \text{kdPcCc} + \text{PcpCc} \cdot \text{kdPcpCc} + \text{cry} \cdot \text{kp2} \\
 \frac{d \text{Pc}}{dt} &= -\text{Cc} \cdot \text{Pc} \cdot \text{kfPcCc} - \text{Pc} \cdot \text{dPc} - \text{Pc} \cdot \text{kphPc} + \text{PcCc} \cdot \text{kdPcCc} + \text{Pcp} \cdot \text{kdphPcp} + \text{kp1} \cdot \text{per} \\
 \frac{d \text{Pcp}}{dt} &= -\text{Cc} \cdot \text{Pcp} \cdot \text{kfPcpCc} + \text{Pc} \cdot \text{kphPc} - \text{Pcp} \cdot \text{dPcp} - \text{Pcp} \cdot \text{kdphPcp} + \text{PcpCc} \cdot \text{kdPcpCc} \\
 \frac{d \text{PcpCc}}{dt} &= \text{Cc} \cdot \text{Pcp} \cdot \text{kfPcpCc} - \text{PcpCc} \cdot \text{dPcpCc} - \text{PcpCc} \cdot \text{kdPcpCc} - \text{PcpCc} \cdot \text{kiPcpCc} + \text{PnpCn} \cdot \text{kePnpCn} \\
 \frac{d \text{PcCc}}{dt} &= \text{Cc} \cdot \text{Pc} \cdot \text{kfPcCc} - \text{PcCc} \cdot \text{dPcCc} - \text{PcCc} \cdot \text{kdPcCc} - \text{PcCc} \cdot \text{kiPcCc} + \text{PnCn} \cdot \text{kePnCn} \\
 \frac{d \text{PnpCn}}{dt} &= \text{PcpCc} \cdot \text{kiPcpCc} - \text{PnpCn} \cdot \text{dPnpCn} - \text{PnpCn} \cdot \text{kePnpCn} \\
 \frac{d \text{PnCn}}{dt} &= \text{PcCc} \cdot \text{kiPcCc} - \text{PnCn} \cdot \text{dPnCn} - \text{PnCn} \cdot \text{kePnCn}
 \end{aligned}$$

Model 4.2: Model from Leloup & Goldbeter, 2003 [10].

$$\begin{aligned}
 \frac{dMP}{dt} &= -MP \cdot kdmP - \frac{MP \cdot vmP}{KmP + MP} + \frac{vsP \cdot BN^n}{BN^n + KAP^n} \\
 \frac{dMC}{dt} &= -MC \cdot kdmC - \frac{MC \cdot vmC}{KmC + MC} + \frac{vsC \cdot BN^n}{BN^n + KAC^n} \\
 \frac{dMB}{dt} &= -MB \cdot kdmB - \frac{MB \cdot vmB}{KmB + MB} + \frac{vsB \cdot KIB^m}{BN^m + KIB^m} \\
 \frac{dPC}{dt} &= -CC \cdot PC \cdot k3 + MP \cdot ksP - \frac{PC \cdot V1P}{Kp + PC} - PC \cdot kdn + PCC \cdot k4 + \frac{PCP \cdot V2P}{Kdp + PCP} \\
 \frac{dCC}{dt} &= -CC \cdot PC \cdot k3 - \frac{CC \cdot V1C}{CC + Kp} - CC \cdot kdn + \frac{CCP \cdot V2C}{CCP + Kdp} + MC \cdot ksC + PCC \cdot k4 \\
 \frac{dPCP}{dt} &= \frac{PC \cdot V1P}{Kp + PC} - \frac{PCP \cdot V2P}{Kdp + PCP} - PCP \cdot kdn - \frac{PCP \cdot vdPC}{Kdp + PCP} \\
 \frac{dCCP}{dt} &= \frac{CC \cdot V1C}{CC + Kp} - \frac{CCP \cdot V2C}{CCP + Kdp} - CCP \cdot kdn - \frac{CCP \cdot vdCC}{CCP + Kd} \\
 \frac{dPCC}{dt} &= CC \cdot PC \cdot k3 - \frac{PCC \cdot V1PC}{Kp + PCC} - PCC \cdot k1 - PCC \cdot k4 - PCC \cdot kdn + \frac{PCCP \cdot V2PC}{Kdp + PCCP} + PCN \cdot k2 \\
 \frac{dPCN}{dt} &= -BN \cdot PCN \cdot k7 + IN \cdot k8 + PCC \cdot k1 - \frac{PCN \cdot V3PC}{Kp + PCN} - PCN \cdot k2 - PCN \cdot kdn + \frac{PCNP \cdot V4PC}{Kdp + PCNP} \\
 \frac{dPCCP}{dt} &= \frac{PCC \cdot V1PC}{Kp + PCC} - \frac{PCCP \cdot V2PC}{Kdp + PCCP} - PCCP \cdot kdn - \frac{PCCP \cdot vdPCC}{Kd + PCCP} \\
 \frac{dPCNP}{dt} &= \frac{PCN \cdot V3PC}{Kp + PCN} - \frac{PCNP \cdot V4PC}{Kdp + PCNP} - PCNP \cdot kdn - \frac{PCNP \cdot vdPCN}{Kd + PCNP} \\
 \frac{dBC}{dt} &= \frac{-BC \cdot V1B}{BC + Kp} - BC \cdot k5 - BC \cdot kdn + \frac{BCP \cdot V2B}{BCP + Kdp} + BN \cdot k6 + MB \cdot ksB \\
 \frac{dBCP}{dt} &= \frac{BC \cdot V1B}{BC + Kp} - \frac{BCP \cdot V2B}{BCP + Kdp} - BCP \cdot kdn - \frac{BCP \cdot vdBC}{BCP + Kd} \\
 \frac{dBN}{dt} &= BC \cdot k5 - BN \cdot PCN \cdot k7 - \frac{BN \cdot V3B}{BN + Kp} - BN \cdot k6 - BN \cdot kdn + \frac{BNP \cdot V4B}{BNP + Kdp} + IN \cdot k8 \\
 \frac{dBNP}{dt} &= \frac{BN \cdot V3B}{BN + Kp} - \frac{BNP \cdot V4B}{BNP + Kdp} - BNP \cdot kdn - \frac{BNP \cdot vdBN}{BNP + Kd} \\
 \frac{dIN}{dt} &= BN \cdot PCN \cdot k7 - IN \cdot k8 - IN \cdot kdn - \frac{IN \cdot vdIN}{IN + Kd}
 \end{aligned}$$

4.2.3 Parameter estimation and bootstrap analysis

Bootstrap parameter estimations were performed as described previously [58], with data from [13] assumed to have a normally distributed 10% relative and 5% absolute error. Since not all states in the models were measured, initial guess values for the trajectory and parameter variables were generated by optimizing the parameter sets first with a genetic algorithm approach, described in [11]. To help ensure bootstrap trials remained in a similar stability region of parameter space (and protect against steady-state solutions), bootstrap parameters were bound between 50% and 150% of their initial value.

4.2.4 Selection of parameters for FBXL3 and CKI

For FBXL3-CRY, parameters that determined the degradation rate of CRY (or CRY containing complexes) were considered to be the most likely candidates. Michealis-Menten degradation parameters were omitted from figure 4.6 since perturbations to such parameters are not easily attributable to changes in FBXL3 binding affinity. In the model presented in [10], CRY is degraded through a series of phosphorylation events, and these parameters were considered as representative of the rate of progression toward ubiquitination of CRY. The forward phosphorylation rates of CRY and nuclear PER-CRY complex were therefore also considered. For CKI-PER, we considered rates that determined the degradation rate and nuclear import rate of PER. Michealis-Menten parameters were not included, similar to FBXL3-CRY. With CRY being the main repressor of E box transcription [46], the degradation rates of PER-CRY complex were not considered as potential mechanisms of CKI. In the models of [10] and [22], the nuclear entry of PER-CRY requires two independent steps: the formation of the PER-CRY complex and the subsequent import of the complex. Therefore, the forward reaction rates of each of these steps were included.

4.2.5 Numerical experiments

Numerical parameter inhibitions were performed by recalculating the limit cycle trajectory for each new parameter set to a tolerance of 10^{-8} , using computational methods described previously [21].

4.3 Results

4.3.1 Opposite amplitude effects of longdaysin and KL001

To gain a more detailed understanding of the roles of CKI-PER and FBXL3-CRY pathways, we applied small molecule compounds longdaysin and KL001, which cause stabilization of PER and CRY, respectively [3, 32] (figure 4.1). We used *Bmal1*- and *Per2-dLuc* as circadian reporters, which represent different loops of the core clock mechanism and show circadian luminescence rhythms with mutually opposite phase. Time-course data on circadian reporter expression under increasing concentrations of longdaysin and KL001 [3] was analyzed for period and amplitude change (figure 4.3). Longdaysin caused dose-dependent increases in period and detrended amplitude to $\approx 50\%$ of control values in both *Bmal1*- and *Per2-dLuc* reporter cells. In contrast, KL001 induced a simultaneous increase in period and strong reduction in amplitude. Modulation of the activity of CKI-PER and FBXL3-CRY is therefore differentiated by an opposite amplitude response.

4.3.2 Main period-determining perturbations

We next used *in silico* modeling to gain mechanistic insight into CKI-PER and FBXL3-CRY mediated circadian regulation. In chapter 2, I described the connection between inhibition of FBXL3-dependent CRY degradation and period change [3]: increasing the stability of nuclear CRY results in longer transcriptional repression and increased




Figure 4.3: Different amplitude effect of small molecule circadian modulators targeting CKI-PER and FBXL3-CRY. (top) Detrended luminescence profiles (first 72 hours, mean of two independent replications) obtained from U2OS reporter cells with increasing concentration of longdaysin and KL001. Black profiles indicate control conditions ($0 \mu\text{M}$), lighter colors indicate higher concentrations of small molecule (from 0.03 to $8 \mu\text{M}$). (bottom) Relative change in period and amplitude of the results in shown on top.

period length. However, while CKI has been linked to modulating PER stability and nuclear entry, it remained unclear which perturbation dominates the period effect, and whether these processes are sufficient to separate the effects of CKI and FBXL3.

To generate predictions that are consistent across slight differences in model assumptions, we chose three mathematical models from the literature based on their moderate size and similar scope [3, 10, 22]. The models included, at a minimum, the expression and nuclear entry mechanisms of PER and CRY. We considered the formation of the PER-CRY heterodimer as a key step in nuclear entry, which is supported by the fact that, to the best of our knowledge, all circadian models that consider both PER and CRY employ this kinetic assumption [3, 10, 11, 12, 22].

Since dynamic models of genetic regulatory networks typically suffer from poor parameter identifiability [59], we demonstrate that our predictions are parameter-independent by employing a bootstrap identifiability analysis [58]. As part of the

bootstrap method, the models were re-fit to experimental data [13] while ensuring appropriate protein stoichiometry. The state trajectories of the resulting 2000 parameter sets for each model are shown in figure 4.4, with reasonable agreement between models and experiment.




Figure 4.4: Fitted trajectories from bootstrap runs. Time series trajectories of the 2000 bootstrap trials for each model. Shaded regions indicate 95% confidence regions. The data were scaled to have a maximum value of 1, except for protein species, where relative values were important for clock stoichiometry.

A first-order period sensitivity analysis, performed on each of the parameter sets, identified which parameters associated with PER and CRY protein activity had the greatest effect on period (figure 4.5). To simplify analysis, we present only those parameters that are associated with experimentally supported mechanisms of CKI and FBXL3 in figure 4.6. We first tested parameters associated with potential FBXL3-CRY activity to evaluate if our method matched the experimentally verified effect of KL001 [3]. Since CRY is the dominant repressor of CLOCK-BMAL1 [46], we attribute degradation rates of the PER-CRY complex to be representative of CRY clearance rates. We found that only parameters governing nuclear CRY degradation show a period lengthening effect upon inhibition, while rates associated with cytoplasmic CRY degradation show period shortening effects. These results match with our

previous assertion that period lengthening occurs via nuclear CRY stabilization [3]. Experimental evidence has also indicated cytoplasmic CRY stabilization may lead to period shortening [16], a result consistent with our mathematical results.

We next describe parameters potentially associated with CKI-dependent regulation of PER localization and stability. Since PER is rate-limiting in the formation of the PER-CRY complex [13], rates associated with complex formation or nuclear import were included in this analysis. Conversely, we did not include degradation rates of PER-CRY nuclear repressive complex, since CRY alone is considered the main repressor. While it was hypothesized in models where PER acts as a direct repressor that the regulation of PER stability would play the dominant role determining the period [55, 57], our new assumptions revealed that parameters governing PER degradation showed only non-identifiable responses. However, inhibition of rates associated with the nuclear entry of the PER-CRY complex showed strong period lengthening effects. These results indicate that under our current understanding of clock kinetics, the regulation of nuclear import likely plays the prominent role in CKI-dependent period regulation.

4.3.3 Independent mechanisms of PER and CRY regulation

Using the model and parameter set of Hirota *et al.*, 2012 [3] and the perturbations identified in figure 4.6, we first confirmed that inhibition of nuclear CRY degradation (vdCn) and PER-CRY nuclear import (vaC1P) reproduced the experimental period and amplitude effects of the small molecules KL001 and longdaysin, respectively (figure 4.7, compare with figure 4.3).

Comparison of the oscillatory profiles of *Per* mRNA and nuclear CRY protein (figure 4.8) revealed that inhibition of FBXL3-dependent CRY degradation caused lingering nuclear CRY to not be completely purged each cycle. This excess repressor during the accumulating phase of *Per* and *Cry* transcripts resulted in lower E box




Figure 4.5: Bootstrap confidence intervals in relative period sensitivities. Violin plots for distributions in relative period sensitivities for parameters associated with PER and CRY proteins. Whiskers extend to the most extreme data point within $1.5x$ the inner quartile range. Distributions in which the 5th and 95th percentile lie on opposite sides of the x -axis are colored red and deemed non-identifiable.




Figure 4.6: Bootstrap predictions of circadian actions of FBXL3-CRY and CKI-PER pathways. Violin plots of the relative period sensitivity of parameters associated with potential mechanisms for FBXL3-CRY (A) and CKI-PER (B) activity. A negative or positive period sensitivity indicate that the period of oscillation will increase or decrease when that rate is inhibited, respectively. Distributions that are not different than 0 with 95% confidence are colored red. Descriptions of the parameters shown are listed in tables 4.1-4.2.

Table 4.1: Descriptions of the model parameters governing FBXL3-CRY

Model	Parameter	Description
[3]	MC2n	CRY2 nuclear multiplicative degradation coefficient
	vdCn	CRY1/2 nuclear degradation rate
	vdC1	CRY1 cytoplasmic degradation rate
	vdC2	CRY2 cytoplasmic degradation rate
[22]	dPnpCn	Phosphorylated PER-CRY complex nuclear degradation rate
	dPnCn	PER-CRY complex nuclear degradation rate
	dPcpCc	Phosphorylated PER-CRY complex cytoplasmic degradation rate
	dPcCc	PER-CRY complex cytoplasmic degradation rate
	dCc	CRY cytoplasmic degradation rate
[10]	V3PC	PER-CRY complex nuclear phosphorylation rate
	vdPCN	Phosphorylated PER-CRY complex nuclear degradation rate
	V1C	CRY cytoplasmic degradation rate
	vdCC	Phosphorylated CRY cytoplasmic degradation rate

Table 4.2: Descriptions of the model parameters governing CKI-PER

Model	Parameter	Description
[3]	vaC1P	PER-CRY complex nuclear entry rate
	vdP	PER cytoplasmic degradation rate
[22]	kiPcpCc	Phosphorylated PER-CRY complex nuclear entry rate
	kfPcpCc	Phosphorylated PER-CRY complex association rate
	kiPcCc	PER-CRY complex nuclear entry rate
	kfPcCc	PER-CRY complex association rate
	dPcp	Phosphorylated PER cytoplasmic degradation rate
	dPc	PER cytoplasmic degradation rate
[10]	k1	PER-CRY complex nuclear entry rate
	k3	PER-CRY complex association rate
	V1P	PER cytoplasmic phosphorylation rate
	vdPC	Phosphorylated PER cytoplasmic degradation rate




Figure 4.7: Predictions of KL001 and longdaysin results. *In silico* reproductions of the circadian reporter experiments in figure 4.3, using the predictions identified in figure 4.6.

amplitudes, providing a likely explanation for the effect of KL001.




Figure 4.8: Comparison of the effects of KL001 and longdaysin. Parameter changes were normalized such that the period change was equal for each pair of perturbations (vdCn: 100% \rightarrow 23%, vaC1P: 100% \rightarrow 51%).

In contrast, stabilization of cytoplasmic PER (lowering vdP) resulted in reduced transcriptional amplitude with minimal period effect (figure 4.9), consistent with experimental findings from the knockdown of β -TrCP, an F box protein responsible for PER degradation [60]. However, other experimental results have shown that down-regulation of β -TrCP leads to longer periods [49], suggesting that further modeling and experimental inquiry is needed on the role of β -TrCP in clock regulation. This period lengthening might be explained through β -TrCP-mediated stabilization of nuclear PER-CRY or by using alternative kinetic assumptions for the rate of PER-CRY binding.




Figure 4.9: Effect of cytoplasmic PER stabilization on period and E box transcription amplitude. Relative period and peak-to-trough amplitude change in *Per* mRNA resulting from a reduction in the vdp parameter in the model from Hirota *et al.*, 2012.

CKI likely regulates nuclear entry rather than PER stability

We further compared the effect of inhibiting PER degradation with inhibiting nuclear import on the oscillatory profile of key clock proteins (figure 4.10) to identify mechanistic differences between the two potential effects of CKI inhibition. Both perturbations increased cytoplasmic PER, suggesting the two mechanisms are difficult to distinguish experimentally. Direct stabilization of PER in the cytoplasm (lowering vdp) lead to two simultaneous trends which shift the period in opposite directions: it shortened the time delay between transcription and inactivation by accelerating the accumulation of cytoplasmic PER and nuclear PER-CRY; and lengthened the repressive phase by increasing the total amount of PER-CRY which enters the nucleus. These perturbations sped and slowed the clock, respectively, and resulted in little period change. In contrast, inhibiting PER-CRY nuclear entry (lowering vaC1P) caused additional free protein to build in the cytoplasm, delaying nuclear accumulation and ultimately increasing the total amount of nuclear PER-CRY. Since both of these trends

work to increase period length, inhibiting PER-CRY nuclear entry resulted in significantly longer cycles. Additionally, the longer cytoplasmic time delay resulted in increased transcription, yielding slightly higher amplitudes (figure 4.8) that closely match the experimental results of the small molecule longdaysin.




Figure 4.10: Comparison of two candidate mechanisms for CKI inhibition. Effects of increasing PER stabilization and nuclear import inhibition on the time profiles of cytoplasmic PER (top) and nuclear CRY (bottom). Parameter values were selected such that the amplitudes of cytoplasmic PER are equal at each level. Lighter colors indicate stronger perturbations (vdP: 100% \rightarrow 22%, vaC1P: 100% \rightarrow 45%), $t = 0$ is set to the onset of PER accumulation.

Since CKI likely regulates both stability and subcellular localization of PER *in vivo*, we considered the effects of simultaneously lowering both PER cytoplasmic degradation and nuclear entry rates, as shown in figure 4.11. The loss of oscillations under extreme reduction of both parameters highlights an interesting role of CKI in conferring robustness to the circadian clock: since oscillations are lost when import of the PER-CRY complex to the nucleus ceases to be rhythmic, CKI ensures lingering PER is purged from the cytoplasm by one pathway or another before E box tran-

scription resumes. This importance has been proven experimentally, as disruption of CKI-mediated regulation leads to compromised circadian oscillations [61].




Figure 4.11: Independence of CKI-PER and FBXL3-CRY pathways. (A) Contour plots of period change (top) and Per mRNA amplitude (bottom) for simultaneous inhibition of both PER degradation (vdP) and nuclear import (vaC1P). The gray shaded region indicates loss of oscillations. (B) Period and amplitude change contour plots for varying both vdCn (CRY nuclear degradation rate) and vaC1P (PER-CRY nuclear import). The consistently straight lines indicate the independence of CKI-PER and FBXL3-CRY mechanisms.

Together, inhibition of CKI by longdaysin may increase the time required before PER-CRY can enter the nucleus to repress transcription, leading to a higher amplitude and longer period. In contrast, KL001 lengthens the period by stabilizing nuclear CRY, resulting in a longer time delay before transcription resumes and lower amplitude from increased E box repression. PER regulation through CKI is therefore partitioned to the accumulating phase, controlling the speed and amount of PER-CRY complex that enters the nucleus. CRY regulation through FBXL3 is par-

titioned independently to the repressive phase, controlling the length of time until CLOCK-BMAL1-dependent transcription resumes (figure 4.12). This independence was reproduced *in silico* by the simultaneous reduction of nuclear CRY degradation and PER-CRY nuclear import (figure 4.11), where nonlinear interactions in amplitude and period between the two perturbations are all but absent.




Figure 4.12: Spatiotemporal separation underlies independence of CKI-PER and FBXL3-CRY pathways. Longdaysin, acting through CKI, lengthens the accumulating phase of the circadian cycle, while KL001, acting through FBXL3, lengthens the declining phase.

4.4 Discussion

An understanding of the interactions between posttranslational regulators is crucial for the further development of circadian pharmacological reagents, as efficient modulation of clock function will assuredly come from simultaneous perturbations to many connected species. In this study, we used circadian reporter cells together with mathematical modeling to provide mechanistic insight into the differences of CKI- and FBXL3- mediated posttranslational regulation of PER and CRY. As a result, we clarified a process by which CKI exerts control over the circadian period, demon-

strated through both the hyperphosphorylating CKI ϵ^{tau} mutant and small molecule CKI inhibitors, such as longdaysin. In developing our predictions, we have used multiple models and parameterizations to ensure our mechanisms are consistent across many *in silico* realizations. These results reinforce the notion that computational modeling is essential in interpreting results in systems with complicated oscillatory feedback. Additionally, *in silico* analyses reveal hidden design principles of biological networks, as this work highlights the importance of the CKI family of kinases in conferring robustness to the circadian cycle.

4.4.1 Phase-dependent activity of small molecule modulators

In this chapter, it was shown that the posttranslational CKI and FBXL3 control separate time regimes of the core clock oscillation. While in the cultured cell experiments the small molecule modulators were continuously present, in pharmacological applications there would likely be dosed at specific times of day. Since their targets play key roles during specific circadian phases, it is likely that there would be a significant difference in small molecule effect depending on the time at which they are given. In the next chapter, I systematically analyze the dynamic consequences associated with *transient* applications of small molecule agonises, rather than dose-dependent continuous applications, as the former will likely have more significant clinical relevance.

Chapter 5

Amplitude metrics for cellular bioluminescence reporters¹

5.1 Background

In the previous chapter, I described used mathematical modeling to explain the systems-level effects of two pharmacological agents. In doing so, it was shown that each compound controlled reactions which primarily occur during specific times of day. Since clinical applications of circadian therapies will likely have a time at which they are administered, it is important to understand the effects of temporary perturbations on circadian amplitude. The effects of temporary perturbations are more complicated, however, since they may effect both the single-cell oscillator as well as the synchrony of cell populations as a whole. In this chapter, I develop the mathematical background required to understand and efficiently predict the response of a population of oscillators to a temporary change.

¹Portions of this chapter are published in P. C. St. John, S. R. Taylor, J. H. Abel, and F. J. Doyle, "Amplitude Metrics for Cellular Circadian Bioluminescence Reporters," *Biophys. J.*, vol. 107, pp. 2712–2722, Dec. 2014.

5.1.1 Importance of circadian amplitude

While the suprachiasmatic nucleus (SCN) in the brain serves as the body's master pacemaker, cells found in peripheral tissues also oscillate in a circadian manner [62]. These peripheral clocks process systemic and SCN-mediated entraining cues, buffering against rhythmic changes in energy availability to maintain metabolic homeostasis [63]. The amplitude of circadian transcription is a relevant factor, and has been shown to play a critical role in phase resetting and entrainment [64, 65]. Recent studies have further highlighted the importance of high peripheral clock amplitudes in maintaining metabolic health. Mice lacking an intact clock have been shown to develop metabolic disease [66], while low amplitude clock oscillations, whether caused by diet [67] or age [68] have also been tied to metabolic disorders. An understanding of how clock amplitudes are regulated is therefore a topic of ongoing research with potential therapeutic applications [69]. Unlike the network of oscillators in the SCN, in which intercellular coupling maintains robust amplitudes even in the absence of external cues, peripheral oscillators are thought to lack a direct mechanism to spontaneously synchronize [70]. As a result, populations of peripheral clocks are likely synchronized by common external cues, with stochastic effects and cell heterogeneity driving entrained populations gradually toward desynchrony.

5.1.2 Amplitude at the single-cell and population level

The development of immortalized peripheral oscillator cell lines with bioluminescent reporters has allowed high-throughput analysis of the responses of circadian rhythms to genetic and pharmacological manipulation [32, 71]. In addition to providing experimental tractability, these systems provide detailed information on the amplitude of oscillations in gene transcription, a measure often lacking from earlier experiments using wheel-running activity. As a result, these cell lines have proven useful

in studying core clock connectivity and stoichiometry [72]. However, since *in vitro* experiments typically measure entire cultures of cells, data collected at the population-level can obscure the response of the clock at the scale of the gene-regulatory network. Even when individual cells can be recorded, stochastic noise hinders accurate amplitude determination. As a result, two studies using similar perturbations to understand the mechanism of light-induced amplitude reduction reached differing conclusions, in which either single-cell amplitude reduction or population-level desynchrony was identified as the dominant factor [73, 74].

5.1.3 Role of mathematical modeling

Mathematical models have long been used to understand the results of circadian experiments [10, 75], aided by definitions and computational techniques designed to match modeling predictions to experimental data. One such definition is the response function, a general technique that maps a change in an output variable to a temporary change in parameters [76]. For instance, the phase response curve (PRC) has been used to characterize the entrainment behavior of both experimental and mathematical systems [77, 78, 79], and in analyzing the synchrony of populations of oscillators [80]. Accurate and efficient numerical routines for finding infinitesimal PRCs have therefore been developed [42, 59, 81]. In addition to changes in phase, phase-dependent changes in amplitude are an important factor in understanding circadian rhythms. Amplitude response curves (ARCs) were first used in the clock literature to understand the effects of light pulses on simple phase-amplitude models, and were useful in predicting phase singularity behavior [82]. ARCs have also been used to characterize perturbations to groups of oscillators through desynchrony [74, 83], at the level of the single oscillator [84, 85], and even in studying the effect of entrainment phase on SCN rhythm amplitude [86]. However, previous definitions of the ARC are inconsistent between studies and do not simultaneously consider

amplitude effects at the single-cell and population level.

The choice of modeling framework dictates the type of amplitude response which can be predicted. Ordinary differential equation (ODE) models of gene regulation are capable of describing the amplitude and phase-resetting behavior of single cells, but fail to capture the collective dynamics of a population of oscillators. Likewise, phase-only models correctly capture the change in synchrony of a population, but do not capture fluctuations in amplitude of individual oscillators. Explicit stochastic simulations of populations of cells are capable of realistically capturing both single-cell and population-level effects, and have been successfully used to understand the response of coupled oscillators to external VIP perturbation [87]. However, these methods are computationally expensive, and cannot be used for an analytical understanding of amplitude response.

In this chapter, I describe approaches to quantify amplitude change in a population of non-interacting oscillators. By exploiting the independence of each oscillator, we derive computationally efficient methods to approximate the mean dynamics of full stochastic simulations. Additionally, our method allows the calculation of ARCs at both the single-cell and population level, allowing the behavior of the system to be quickly profiled. Specifically, we use ODE models to describe the transient amplitude response at the single-cell level, coupled with a phase probability density function to describe population-level dynamics. Following a perturbation with a finite duration, a limit cycle oscillator undergoes a transient change in amplitude. When the mean expression level of a population of oscillators is also considered, an additional change in amplitude is incurred due to the change in synchrony of the population, which persists until synchrony is changed by subsequent perturbations. We therefore observe a separation in timescales between the effects on clock output mediated at the single-cell level and those mediated by population synchrony, allowing the source of an amplitude change to be qualitatively inferred by inspection of bioluminescence

data. Characterizing the mechanisms and consequences of both types of amplitude regulation will be important in understanding how peripheral amplitudes are maintained, and may lead to the design of pharmacological or behavioral strategies to boost circadian amplitudes.

5.2 Methods

5.2.1 Basic definitions

Ordinary differential equation (ODE) models take the general form

$$\frac{dx}{dt} = f(x(t), p), \quad (5.1)$$

in which $x(t)$ represents the concentrations of the state variables, such as mRNA and protein concentrations, f contains information on the production, degradation, and reactivity of the states, and p are the kinetic parameters which govern reaction kinetics. Limit cycle models are ODE models in which the solution approaches a steady state oscillatory trajectory, satisfying the equation:

$$\lim_{t \rightarrow \infty} [x(t + T) - x(t)] = 0. \quad (5.2)$$

The period is the smallest $T > 0$ for which equation (5.2) holds. The points on the stable limit cycle are denoted by $x^\gamma(\theta)$ with each point assigned to a value of a phase variable $\theta \in [0, 2\pi)$. For convenience, time in equation (5.1) can be rescaled such that the period is 2π :

$$\tilde{t} = \frac{2\pi}{T}t; \quad \tilde{f} = \frac{T}{2\pi}f; \quad \frac{dx}{d\tilde{t}} = \tilde{f}(x(\tilde{t}), p). \quad (5.3)$$

The phase variable θ is therefore defined on the limit cycle as $\theta = \tilde{t} \bmod 2\pi$, with $\theta = 0$ assigned to a unique and identifiable point.

5.2.2 Perturbations to limit cycle systems

In this chapter we restrict our analysis to temporary perturbations capable of entraining an oscillatory system, excluding permanent parameter changes arising, for instance, from genetic knockout experiments. The most simple entraining perturbation involves adding or removing components to a limit cycle system, resulting in a perturbed trajectory $x(t)$. Here the initial conditions are determined by the strength of the perturbation and the phase at which it is applied:

$$x(0) := x^\gamma(\theta_0) + \Delta x(0). \quad (5.4)$$

This trajectory evolves according to equation (5.1), eventually returning to x^γ . It is useful to express this trajectory by the deviation from the limit cycle:

$$\Delta x(\tilde{t}) = x(\tilde{t}) - x^\gamma(\tilde{t} + \theta_0). \quad (5.5)$$

In addition to perturbations directly to the state of the system, oscillators can also be perturbed by temporary changes to the parameters. For a parameter pulse Δp of duration \tilde{d} (in radians) that ends at θ_0 , the oscillator deviates from the limit cycle trajectory according to:

$$x_{\tilde{d}}(-\tilde{d}) = x^\gamma(\theta_0 - \tilde{d}) \quad (5.6)$$

$$\frac{dx_{\tilde{d}}}{d\tilde{t}} = \tilde{f}(x_{\tilde{d}}(\tilde{t}), p + \Delta p). \quad (5.7)$$

The pulse trajectory $x_{\tilde{d}}(\tilde{t})$ is then integrated from $\tilde{t} = -\tilde{d} \rightarrow 0$, at which point the pulse is removed. A temporary parameter pulse is therefore equivalent to a state perturbation, but with the perturbation at $t = 0$ defined by:

$$\Delta x(0) = x_{\tilde{d}}(0) - x^\gamma(\theta_0). \quad (5.8)$$




Figure 5.1: Schematic showing trajectories used in the calculation of single-cell phase and amplitude change. A perturbation $x(\tilde{t}, p)$, blue, from the limit cycle solution $x^\gamma(\tilde{t}, p)$, red, ultimately returns to the limit cycle with a phase shift $x^\gamma(\tilde{t} + \Delta\theta, p)$, green.

A schematic depicting an example perturbed and reference trajectory is shown in figure 5.1.

The response of the system to very small perturbations is more efficiently calculated using ODE sensitivity analysis [88]. The sensitivity matrix,

$$S(t) = \frac{dx(t)}{dx(0)} = \lim_{\Delta x(0) \rightarrow 0} \frac{\Delta x(t)}{\Delta x(0)}, \quad (5.9)$$

can be calculated directly by integrating

$$\frac{d}{dt} S(t) = \frac{df(x(t), p)}{dx} S(t) \quad (5.10)$$

with $S(0) = I$.

5.2.3 Phase response curves

Quantifying phase changes following a perturbation has been well studied and is particularly relevant for models of circadian rhythms [42, 81]. A phase response curve (PRC) maps the change in phase resulting from the same perturbation applied at each initial phase. Infinitesimal PRCs, the derivative of the phase change with respect to the perturbation, can be defined for state and parameter-impulse perturbations [81]. Methods for efficiently calculating these quantities using ODE sensitivity analysis have been developed [81], with the important result that the parameter- and state-impulse PRCs can be related by the Jacobian matrix:

$$\frac{d}{d\tilde{t}} \frac{d\theta}{dp} = \frac{d\theta}{dx} \frac{d\tilde{f}}{dp}. \quad (5.11)$$

This result follows from the fact that in the limit of an infinitely short and small parameter pulse, $\Delta x(0) \rightarrow d\tilde{f}/dp \tilde{d} \Delta p$.

5.2.4 Phase-diffusion model

Large populations of oscillators are typically described using phase-only models [89], in which the state of each oscillator is represented only by its phase, θ . The synchrony of the population can be modeled using a probability density function $p(\theta, \tilde{t})$ that describes the probability of finding an oscillator at each phase [90]. The usefulness of probability density functions in describing the phase and amplitude responses of populations of circadian cells has been previously shown [74]. As with all probability density functions,

$$\int_0^{2\pi} p(\theta, \tilde{t}) d\theta = 1. \quad (5.12)$$

The shape of $p(\theta, \tilde{t})$ changes as the cells advance in time. Stochastic effects cause the population to gradually desynchronize as slight cycle-to-cycle deviations are propa-

gated throughout the population [91]. For an infinite population of oscillators, these effects are well-described by a Fokker-Plank equation [92]:

$$\frac{\partial p}{\partial \tilde{t}} = \frac{\partial p}{\partial \theta} + d \frac{\partial^2 p}{\partial \theta^2}. \quad (5.13)$$

Due to the rescaling of t , the mean period of the population is 2π . Here, the $\partial p / \partial \theta$ term, analogous to convection, describes the mean oscillatory period, while the $\partial^2 p / \partial \theta^2$ term describes the diffusion of phases across $[0, 2\pi)$. The phase diffusivity parameter d (in units of inverse radians) describes the speed with which the population desynchronizes and can be fit to experimental data [93]. Equation (5.13) has periodic boundary conditions, with initial condition $\phi(\theta)$ as the phase population at $t = 0$:

$$\text{BCs: } p(0, \tilde{t}) = p(2\pi, \tilde{t}) \quad (5.14)$$

$$\frac{\partial p}{\partial \theta}(0, \tilde{t}) = \frac{\partial p}{\partial \theta}(2\pi, \tilde{t}) \quad (5.15)$$

$$\text{IC: } p(\theta, 0) = \phi(\theta). \quad (5.16)$$

The solution of equations (5.13-5.16) is well-characterized, with $p(\theta, \tilde{t})$ evolving in time as the convolution of the initial conditions with a wrapped normal distribution with mean \tilde{t} and standard deviation $\sqrt{2d\tilde{t}}$ [94]:

$$p(\theta, \tilde{t}) = \phi(\theta) * \mathcal{WN}(\theta; \tilde{t}, \sqrt{2d\tilde{t}}), \quad (5.17)$$

in which the wrapped normal distribution [95] is defined as

$$\mathcal{WN}(\theta; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} \sum_{k=-\infty}^{\infty} \exp \left[\frac{-(\theta - \mu + 2\pi k)^2}{2\sigma^2} \right]. \quad (5.18)$$

Since the convolution of two normal distributions is also a normal distribution, it is efficient when possible to describe $\phi(\theta)$ as a normal distribution with mean μ_0 and

standard deviation σ_0 , such that $p(\theta, \tilde{t})$ can be found analytically through

$$p(\theta, \tilde{t}) = \mathcal{WN}(\theta; \mu_0 + \tilde{t}, \sqrt{\sigma_0^2 + 2d^2\tilde{t}^2}). \quad (5.19)$$

5.2.5 Numerical simulations

ODE models were simulated in Python, using the computer algebra package CasADI [39]. Stochastic simulations were performed using the StochKit2 package [96]. The codes used to generate the figures are included as a supplemental file.

5.3 Results

A signal to a population of limit cycle oscillators can affect amplitude in two ways. First, individual cells are perturbed from their limit cycles, and exhibit transient dynamics before settling back to steady state amplitudes. Secondly, the phases of the population are changed, resulting in a permanent change in population synchrony. Therefore, in order to derive continuous approximations to the dynamics of a large population of non-interacting oscillators, we first demonstrate how ARCs can be found at both the single-cell and population level.

5.3.1 Definition of an amplitude metric at the single-cell level

Following a perturbation to a single cell, the path that the perturbed trajectory $x(t)$ takes in returning to the limit cycle will have a different amplitude than the unperturbed trajectory. While such an amplitude change will only have a finite duration, it plays an important role when perturbations are repeatedly received by the clock, such as a peripheral oscillator entrained to daily metabolic stimuli. To define an amplitude change metric for such a case, we compare a perturbed trajectory $x(t)$ to a phase-shifted limit cycle reference $y(t)$, for which $x(t) \rightarrow y(t)$ for sufficiently long

times. Since $x(t)$ approaches the reference as $t \rightarrow \infty$, the means of both trajectories are equal and can be calculated by

$$\mu := \int_0^{2\pi} \frac{x^\gamma(\theta)}{2\pi} d\theta. \quad (5.20)$$

The amplitude change metric is defined as

$$\begin{aligned} \Delta A(x(t), y(t)) &:= \int_0^\infty (x(t) - \mu)^2 - (y(t) - \mu)^2 dt \\ &= \int_0^\infty h(t) dt. \end{aligned} \quad (5.21)$$

This amplitude metric was chosen over alternatives, such as peak-trough distance, for its analytical tractability. The integrand in equation (5.21), abbreviated by $h(t)$, compares the variances of the reference and perturbed trajectories. When $h(t) > 0$, the trajectory is further from the mean than the reference, and similarly when $h(t) < 0$ the trajectory is closer to the mean than the reference. Thus the overall amplitude change can be calculated by integrating $h(t)$ until the two trajectories converge, returning an amplitude value for each state variable. Amplitude change for a two-dimensional oscillator is easy to visualize graphically. In figure 5.2, the same state perturbation $\Delta x(0)$ applied at two different phases results in opposite amplitude changes, depending on whether the perturbation shifts the trajectory to the interior of the limit cycle (reduced amplitudes) or to the outside the limit cycle (increased amplitudes). Trajectories for the second state variable, $Y(\tilde{t})$, and the corresponding integrand for the amplitude change equation, $h(\tilde{t})$, demonstrate how this transient change is quantified.

5.3.2 Single-cell amplitude response curves

Similar to the PRC, we denote the phase-dependent amplitude change following a perturbation as an amplitude response curve (ARC), using the metric presented in




Figure 5.2: Amplitude metrics at the single-cell level measure transient deviations from the limit cycle. The same perturbation applied at two different phases can result in opposite amplitude effects. The state-space representation (**B**) reveals the path both perturbations take to return to the limit cycle. The time-series representation (**C**) shows how the perturbation in blue results in an amplitude decrease, while the one in red results in an amplitude increase. (**D**) This amplitude change is quantified by integrating $h(\tilde{t})$, the difference in variance from each solution to the limit cycle as defined in equation (5.21), from $t = 0 \rightarrow \infty$. Here the shaded region indicates the area under the curve, ΔA , for the perturbations shown in red and blue. Model adapted from [97].

equation (5.21). In figure 5.3, we calculate the PRC and ARC for a state perturbation of three different strengths. The weakest perturbation results in type 1 (weak) resetting, in which the phase response curve is continuous, while the strongest perturbation results in type 0 (strong) resetting, in which the PRC is discontinuous. This transition occurs once the state perturbation is strong enough to push the trajectory over the unstable fixed point located at the middle of the limit cycle. For a perturbation of intermediate strength, at a critical phase the trajectory will be pushed close to the fixed point and take a long time to recover the steady state amplitude. The behavior at this singularity point is indicated by a sharp dip in the ARC for this perturbation strength.

Infinitesimal versions of response curves are often more general and easier to compute than those that track a specific perturbation strength [76]. We therefore




Figure 5.3: Phase and amplitude response curves describe how a single oscillator will respond to a perturbation. (A-B) The phase and amplitude response curves for three perturbations of increasing strengths demonstrate the transition from type 1 phase resetting for a weak stimulus (blue) to type 0 phase resetting from a strong stimulus (red). (C) The state space representation for the perturbations of varying strengths demonstrate how the trajectories return to the limit cycle. For the particular perturbation shown in green, the trajectory starts very close to the singularity, and therefore takes a long time to recover. (D) For an oscillator perturbed to the singularity, a strong amplitude reduction ensues. Here, the oscillator with an initial phase $\theta \approx 3\pi/2$ is perturbed by $\Delta x = 1.1$ at $\tilde{t} = 0$. Several cycles are required for normal amplitudes to be restored, corresponding to a dip in the ARC (C, green).

derive an expression for the infinitesimal ARC, defined as

$$\frac{dA}{dx} := \lim_{\Delta x(0) \rightarrow 0} \frac{\Delta A(x(\tilde{t}), x^\gamma(\tilde{t} + \theta_0 + \Delta\theta))}{\Delta x(0)} = \int_0^\infty \lim_{\Delta x(0) \rightarrow 0} \frac{h(t)}{\Delta x(0)} dt. \quad (5.22)$$

While this quantity could be calculating by using a very small $\Delta x(0)$, it is more accurate and efficient to derive a direct method to calculate equation (5.22) using the ODE sensitivities defined in equation (5.10). For simplicity, we define $t_\theta = \tilde{t} + \theta_0$, which allows the time variable for the perturbation to vary from $0 \rightarrow \infty$ while tracking the appropriate phase on the limit cycle. Since $\Delta\theta \rightarrow 0$ as $\Delta x(0) \rightarrow 0$, we Taylor expand the limit cycle trajectory around t_θ :

$$x^\gamma(t_\theta + \Delta\theta) = x^\gamma(t_\theta) + \frac{dx^\gamma(t_\theta)}{d\theta} \Delta\theta + O(\Delta\theta^2) \quad (5.23)$$

$$= x^\gamma(t_\theta) + \tilde{f}(x^\gamma(t_\theta)) \Delta\theta + O(\Delta\theta^2). \quad (5.24)$$

Simplifying the integrand in equation (5.22),

$$\frac{h(t)}{\Delta x(0)} = \frac{1}{\Delta x(0)} \left[(x^\gamma(t_\theta) + \Delta x(\tilde{t}) - \mu)^2 - (x^\gamma(t_\theta + \Delta\theta) - \mu)^2 \right] \quad (5.25)$$

$$= \frac{1}{\Delta x(0)} \left[(x^\gamma(t_\theta) + \Delta x(\tilde{t}) - \mu)^2 - (x^\gamma(t_\theta) + \tilde{f}(x^\gamma(t_\theta)) \Delta\theta - \mu)^2 \right] \quad (5.26)$$

$$= \frac{1}{\Delta x(0)} \left[(\Delta x(\tilde{t}) - \tilde{f}(x^\gamma(t_\theta)) \Delta\theta) (\Delta x(\tilde{t}) + \tilde{f}(x^\gamma(t_\theta)) \Delta\theta + 2(x^\gamma(t_\theta) - \mu)) \right]. \quad (5.27)$$

Taking the limit of this integrand as $\Delta x(0) \rightarrow 0$ cancels several differential terms, and allows the remainder to be substituted with quantities that can be calculated using ODE sensitivity analysis:

$$\lim_{\Delta x(0) \rightarrow 0} \frac{h(t)}{\Delta x(0)} = 2 \left(\lim_{\Delta x(0) \rightarrow 0} \frac{\Delta x(\tilde{t})}{\Delta x(0)} - \tilde{f}(x^\gamma(t_\theta)) \lim_{\Delta x(0) \rightarrow 0} \frac{\Delta\theta}{\Delta x(0)} \right) (x^\gamma(t_\theta) - \mu) \quad (5.28)$$

$$= 2 \left(S(\tilde{t}) - \tilde{f}(x^\gamma(t_\theta)) \frac{d\theta}{dx} \right) (x^\gamma(t_\theta) - \mu). \quad (5.29)$$

Note that in equation (5.29), $d\theta/dx$ represents the derivative of $\Delta\theta$ with respect to the perturbation, and is therefore a scalar quantity. The infinitesimal state-impulse ARC may therefore be calculated directly from the ODE sensitivity matrix and the PRC, allowing the amplitude change for an infinitesimal perturbation to be calculated exactly and efficiently:

$$\frac{dA}{dx} = \int_0^\infty 2 \left(S(\tilde{t}) - \tilde{f}(x^\gamma(t_\theta)) \frac{d\theta}{dx} \right) (x^\gamma(t_\theta) - \mu) dt. \quad (5.30)$$

Here, the first term of the integrand ($S - \tilde{f}\dot{\theta}$) tracks the distance from the perturbed trajectory to the limit cycle, which decays to zero as $t \rightarrow \infty$. The second term ($x^\gamma - \mu$) weights this distance by whether or not the deviation occurs above or below the oscillatory mean, yielding negative amplitude changes when the trajectory is perturbed closer to the mean. Just as with the parameter-impulse PRC, the infinitesimal parameter-impulse ARC is defined as

$$\frac{d}{d\tilde{t}} \frac{dA}{dp} := \lim_{\tilde{d}, \Delta p \rightarrow 0} \frac{\Delta A}{\tilde{d} \Delta p}, \quad (5.31)$$

and may be calculated from the state-impulse version with the following relationship:

$$\frac{d}{d\tilde{t}} \frac{dA}{dp} = \frac{dA}{dx} \frac{df}{dp}. \quad (5.32)$$

As with equation (5.11), this equivalency reflects the fact that for a pulse of infinitely short duration, a parameter change is equivalent to changing the state of the system along the direction specified by the Jacobian. Convergence between the methods in equations (5.30-5.32) and finite-difference approaches is shown in figure 5.4, which demonstrates that the numerically-efficient differential ARCs remain representative even for moderate-strength perturbations.




Figure 5.4: Convergence of finite-difference and differential methods. (A) Perturbations of decreasing strength to the state of the oscillator result in phase and amplitude response curves that match the differential limit. (B) Temporary perturbations of decreasing strength to a kinetic parameter. The duration of each perturbation is fixed to 0.2 radians. Finite difference approximations closely match the differential method, which is offset in phase by 0.1 radians to account for the nonzero pulse duration.

5.3.3 Population-level response curves

We next briefly describe how to calculate the PRC and ARC for a phase-density model. These approaches are commonly used in understanding phase models [74, 90], and are presented here to match previous definitions for single cells. A phase transition curve, $g(\theta) = \theta + \Delta\theta$, maps the phase of an oscillator before perturbation to its phase after the perturbation. Since individual oscillators are neither created nor destroyed during the perturbation, it is possible - yet numerically difficult - to directly calculate the phase probability distribution following perturbation using the standard change of variables relation:

$$\hat{p}(\theta, \tilde{t}) \, dg(\theta) = p(\theta, \tilde{t}) \, d\theta. \quad (5.33)$$

However, it is easier to estimate the mean and standard deviation of the perturbed population $\hat{p}(\theta, \tilde{t})$ using directional statistics [95]. A population defined on the unit circle can be described by a complex variable $z = \rho e^{i\bar{\theta}}$, where $\bar{\theta}$ is the mean phase and ρ , the synchronization index, is related to the standard deviation of the population. For $\rho = 1$, the population is clustered about one mean phase, while for $\rho = 0$ the population is evenly balanced across the unit circle. Complex variables for the population before and after perturbation can be calculated via

$$z := \int_0^{2\pi} e^{i\theta} p(\theta, \tilde{t}) \, d\theta \quad (5.34)$$

$$\hat{z} := \int_0^{2\pi} e^{i\theta} \hat{p}(\theta, \tilde{t}) \, d\theta = \int_0^{2\pi} e^{ig(\theta)} p(\theta, \tilde{t}) \, d\theta. \quad (5.35)$$

In equation (5.35), we avoid calculating the perturbed population explicitly by instead integrating over the new phases at the prior population density function. Population-

level amplitude and phase responses can therefore be calculated by

$$\Delta\bar{\theta} = \angle z - \angle \hat{z} \quad (5.36)$$

$$\Delta\rho = |z| - |\hat{z}|. \quad (5.37)$$

Population-level PRCs and ARCs can be tabulated by solving equations (5.34-5.37) for populations $p(\theta, \tilde{t})$ with different mean phases. It is important to note that the ARC at the population level strongly depends on the slope of the PRC, as has been shown previously [74].

5.3.4 Population-level mean expression profiles

To efficiently capture the population-level effects of bioluminescence experiments, we couple the detailed single-cell ODE model to a phase-density model. Previous work has used limit cycle models to estimate population-level parameters, such as desynchronization rate, for phase-only models [93]. In this chapter, we use an ODE model to calculate the response to a perturbation at each phase, and subsequently take the weighted average of these responses according to the phases of the cells in the population.

Assuming each oscillator in the population follows the dynamics described by $x^\gamma(\theta)$, the unperturbed mean population-level expression, $\bar{x}(\tilde{t})$, can be found by taking the weighted average of the expression level over the current population:

$$\bar{x}(\tilde{t}) = \int_0^{2\pi} x^\gamma(\theta) p(\theta, \tilde{t}) d\theta. \quad (5.38)$$

Phase-diffusion models can explain why the gradual damping from experimental population-level data closely resembles an exponentially damped sinusoid, a result that has been shown experimentally [70] and computationally [93]. Our method

similarly demonstrates exponential decay: for the idealized system $x^\gamma(\theta) = \cos(\theta)$ starting from a synchronized state:

$$\bar{x}(\tilde{t}) = \frac{1}{\sqrt{4\pi d\tilde{t}}} \int_{-\infty}^{\infty} \cos(x) \exp\left(-\frac{(x - \tilde{t})^2}{4d\tilde{t}}\right) dx \quad (5.39)$$

$$= \Re \left[\frac{1}{\sqrt{4\pi d\tilde{t}}} \int_{-\infty}^{\infty} \exp(ix) \exp\left(-\frac{(x - \tilde{t})^2}{4d\tilde{t}}\right) dx \right] \quad (5.40)$$

$$= \Re \left[e^{(i-d)\tilde{t}} \right] \quad (5.41)$$

$$= e^{-d\tilde{t}} \cos(\tilde{t}). \quad (5.42)$$

Due to the smoothing effect of phase diffusion, higher frequency sinusoidal components of the limit cycle are damped faster than lower frequency components, resulting in an exponentially decaying sinusoids even for limit cycles that are not sinusoidal in sufficiently disperse populations. The analytical result in equation (5.42) allows us to easily estimate the phase diffusivity parameter in equation (5.13) by simply fitting an exponentially damped sinusoid to detrended bioluminescence data.

Using our continuous approximation to population-level dynamics, we demonstrate an example unperturbed trajectory using a model adapted from [97], see model 5.1. In figure 5.5, an initially jagged population density smooths and widens over time as cells desynchronize, similar to the effects of diffusion. While each cell's expression level follows the limit cycle, the population amplitude gradually damps with time as cells with diverse phases are averaged together.

Next we describe how to calculate population-level mean expression following a perturbation. The perturbed trajectory $\hat{x}(\tilde{t})$ can be decomposed into contributions from two sources. First, for long times after the perturbation, each individual oscillator will have returned to the limit cycle $x^\gamma(\theta)$, but with a new population density

Model 5.1: Model adapted from Novák & Tyson, 2008 [97]. Used for ARC demonstrations in figures 5.2-5.5 ($P = 4$) and figure 5.6 ($P = 2$). It should be noted that the use of the stochastic simulation algorithm for reactions with non-elementary propensities may inaccurately represent the noise of the full system. In this case, we simply fit the noise characteristics of the approximated model (using the volume parameter) to yield physiologically realistic desynchronization rates. The volume of the stochastic simulations was chosen as $\Omega = 250$.

$$\begin{aligned}\frac{dX}{dt} &= \frac{1}{1+Y} - X \\ \frac{dY}{dt} &= k_t X - k_d Y - \frac{Y}{\alpha_0 + \alpha_1 Y + \alpha_2 Y^2}\end{aligned}$$

Parameter	Value	Parameter	Value
k_t	20	α_0	0.005
k_d	1	α_1	0.05
P	4 (or 2)	α_2	0.1




Figure 5.5: Synchrony affects population-level amplitude. (A) Stochastic fluctuations cause a population of cells to gradually desynchronize with time. The phase probability density, $p(\theta, \tilde{t})$, gradually widens as it advances in phase according to the mean period. (B) The mean amplitude from a population of oscillators is determined by the probability density function $p(\theta, \tilde{t})$ and the oscillator's limit cycle $x^\gamma(\theta)$. As time passes, population-level rhythms resemble an exponentially damped sinusoid.

$\hat{p}(\theta, \tilde{t})$. This steady-state perturbed trajectory $\hat{x}_{ss}(\tilde{t})$ can be found by

$$\hat{x}_{ss}(\tilde{t}) = \int_0^{2\pi} x^\gamma(\theta) \hat{p}(\theta, \tilde{t}) d\theta. \quad (5.43)$$

For very long times following perturbation, the new phase probability density could be approximated using the initial mean and standard deviation found through equation (5.35), as jagged profiles in the population density will eventually smooth to a normal distribution. However, for shorter times following perturbation, $\hat{p}(\theta, \tilde{t})$ must be calculated numerically.

The second contribution to the perturbed population trajectory comes from deviations from limit cycle oscillations in each cell. We calculate the population-level effect of these deviations by averaging over the deviations that occur at each phase. We define the deviation trajectory $\delta x(\theta, \tilde{t})$ for each phase as the distance between the perturbed trajectory and the phase-adjusted reference:

$$\delta x(\theta_0, \tilde{t}) := x(\tilde{t}) - x^\gamma(\tilde{t} + \theta_0 + \Delta\theta) \quad (5.44)$$

$$\therefore \lim_{\tilde{t} \rightarrow \infty} \delta x(\theta, \tilde{t}) = 0. \quad (5.45)$$

Since the perturbed trajectory ultimately converges with the phase-adjusted reference, deviations will converge to zero. Since the phase change, $\Delta\theta$, associated with a perturbation at each phase is likely not known prior to calculating the perturbed trajectory, it is difficult to tabulate deviation trajectories associated with each final phase, $\theta_0 + \Delta\theta$. It is therefore more straightforward to find the average effect of single-cell perturbations at the population level by weighting the deviations by the phase density function prior to perturbation. The population-level response to a perturbation, $\hat{x}(\tilde{t})$, is therefore defined as

$$\hat{x}(\tilde{t}) = \int_0^{2\pi} x^\gamma(\theta) \hat{p}(\theta, \tilde{t}) + \delta x(\theta, \tilde{t}) p(\theta, \tilde{t}) d\theta. \quad (5.46)$$

Here the first term is equivalent to $\hat{x}_{ss}(\tilde{t})$, the steady-state perturbed trajectory, while the second term decays to zero as individual oscillators return to their steady-state amplitude. The accuracy of the continuous phase-diffusion model was tested by explicitly simulating a perturbation to a population of 225 uncoupled oscillators using a stochastic simulation algorithm [96, 98]. Good agreement between the stochastic population and continuous approximation was shown in phase shift, single-cell level amplitude change, and alteration of population synchrony (figure 5.6). These results indicate the use of a continuous probability function is justified in the case of cultured cellular reporter systems, which may contain up to 10^5 individual cells per culture [70]. In addition, the method was verified by similar simulations using the Oregonator [99], a model containing only mass-action terms, and a detailed mechanistic model of circadian rhythms [3] at a variety of phase timings (figure 5.7 and figure 5.8). Good agreement is seen between the continuous and stochastic simulations, indicating the proposed method is suitable for predicting the population-level responses for a variety of model types. The slightly reduced accuracy seen in the larger model 2.1 demonstrates a limitation of the method, which should be considered before applying the method to extreme cases. As equation (5.45) tabulates the response of the system from the limit cycle, systems which are perturbed from states far from the deterministic limit cycle will not be captured appropriately. Therefore, the difference in accuracy between model 2.1 and model 5.2 likely comes from increased deviation about the deterministic limit cycle in a greater number of spatial dimensions. The continuous approximation is therefore able to capture the population-level dynamics of a wide variety of limit cycle models and perturbations.

We next demonstrate the usefulness of single-cell and population level ARCs in predicting the mean response of a population. Using a detailed single-cell model (model 2.1), we calculate the response of the system, specifically the average CRY2 protein expression level, to a temporary increase in *Per* transcription rate. In part A




Figure 5.6: Approximation of an explicit stochastic population by continuous methods. (A) A population of 225 stochastic oscillators was simulated using the Gillespie stochastic simulation algorithm (SSA). The mean protein expression, Y , is plotted as a function of time. A 50% reduction in the protein translation rate to each of the oscillators is applied from $\tilde{t} = -\pi/4 \rightarrow 0$, resulting in both single-cell and population-level amplitude change (red solid line). This population is approximated by the continuous methods described in this manuscript, in which the decay parameter $d = 0.025$ is estimated to match the stochastic-induced desynchrony of the control population (black solid line). The initial standard deviation $\sigma_0 = 0.48$ is similarly matched to the stochastic population. The resulting predicted unperturbed and perturbed trajectories, $\bar{x}(\tilde{t})$ and $\hat{x}(\tilde{t})$ respectively, closely match the stochastically modeled values. (B) Phase histograms for the stochastic population are shown at several phases, both before and after the desynchronizing perturbation. Phase probability-density functions for the continuous approximation are also shown, with close agreement between stochastic and continuous simulations. This close approximation validates the use of ODE models and phase-diffusion populations in deriving amplitude and phase-response behavior for networks of uncoupled cells.




Figure 5.7: Validation of continuous methods with the Oregonator model. (A) The Oregonator model (model 5.2) was tested as an example of a mass-action limit cycle oscillator. A population of 2000 oscillators was perturbed at size different mean phases, μ , by increasing the c_2 parameter by 50% for $d = \pi$ (highlighted region). Only the average Y_2 variable is plotted.





Figure 5.8: Validation of continuous methods with the model from Hirota et al., 2012. (A) (B) The model described in [3] (model 2.1) was similarly tested at a variety of mean phases by reducing the vdp parameter by 28.5% and plotting the resulting $c2$ state variable (as in figure 5.10). Parameters for the continuous approximations were found by estimating the phase, initial standard deviation, period, and phase diffusivity of the control population.

Model 5.2: The Oregonator model [99]. The parameters used are as presented in [98]. Used in figure 5.7 as an example of a limit cycle oscillator using only mass-action kinetics.

The individual chemical reactions (used for stochastic simulation) are:



These reactions can be converted to a standard ODE model, in which only the intermediate variables Y are considered, as:

$$\begin{aligned}
 \frac{dY_1}{dt} &= c_1x_1 \cdot Y_2 - c_2 \cdot Y_1 \cdot Y_2 + c_3x_2 \cdot Y_1 - c_4 \cdot Y_1^2 \\
 \frac{dY_2}{dt} &= -c_1x_1 \cdot Y_2 - c_2 \cdot Y_1 \cdot Y_2 + c_5x_3 \cdot Y_3 \\
 \frac{dY_3}{dt} &= c_3x_2 \cdot Y_1 - c_5x_3 \cdot Y_3
 \end{aligned}$$

Parameter	Value
c_1x_1	2
c_2	0.1
c_3x_2	104
c_4	0.016
c_5x_3	26

Parameter	Value

of figure 5.9, we plot the resulting single-cell and population-level response curves. In this case, the population-level phase change is a slightly smoothed version of the single-cell PRC, since the population has an averaging effect on incoming perturbations with each cell receiving the input at a slightly different internal phase. This smoothing follows from the consistent definition of phase between the single-cell and population level. In the ARC plot, however, the shape of the single-cell and population-based ARC are different, since they describe different types of changes (finite vs. sustained) in the output trajectories.

Using these response curves, we demonstrate characteristic features of amplitude change mediated at the single-cell and population-level. In part B of figure 5.9, the perturbation at θ_1 does not cause a significant change in phase or population synchrony, yet strongly reduces single-cell amplitudes. The resulting population-level trajectory appears damped for the first cycle, until the oscillators within the population return to the limit cycle and normal amplitudes are restored. In part C, the perturbation at θ_2 reduces population synchrony while increasing single-cell amplitudes, yielding amplitudes that are transiently increased before the population settles to a lower-amplitude, desynchronized trajectory.

These examples demonstrate the qualitative differences between amplitude change mediated at the single-cell and population level. Their characteristic features allow us to distinguish between these two sources of amplitude change without explicitly recording single-cell amplitudes: amplitude change induced at the population-level will be sustained, but may be masked in the short term by changes to single-cell level amplitudes. Similarly, a change induced at the single-cell level will be evident only at short time-scales. These results underscore the importance in considering both single-cell and population-mediated amplitude change in predicting the effects of daily stimuli on clock amplitudes.




Figure 5.9: Response curves describe different types of amplitude change. A population of cells is simulated using a detailed model of circadian rhythms (model 2.1). Each cell in the population is subjected to a 20% increase in the vtp parameter (*Per* transcription rate) for $d = \pi/2$, inducing both a phase and amplitude change. (A) The phase response curves are shown for both the single limit cycle oscillator and population average, left. The population-level ARC is shown together with the single-cell amplitude response, with both curves normalized to $\sigma = 1$. (B-C) Population-level rhythms, $\hat{x}(\tilde{t})$ resulting from perturbations given at the indicated phases in A (red and cyan lines, respectively). The population transitions from the unperturbed population, $\bar{x}(\tilde{t})$, to the steady-state perturbed population, $\hat{x}_{ss}(\tilde{t})$. (B) A perturbation at this phase yields a transient amplitude reduction resulting from perturbations at the single-cell level, however, there is little change in population synchrony. Population-level rhythms are therefore transiently damped before regaining normal amplitudes. (C) Oscillators are desynchronized, but with a transient increase in limit cycle amplitude. It therefore takes some time before the amplitude reduction from desynchrony is realized.

5.3.5 Application to Experimental Data

We conclude by using the proposed modeling framework to reconcile two experimental studies that measured the effect of light on the circadian amplitude of photosensitive fibroblast cells [73, 74]. Both studies sought to determine the main factor of amplitude reduction following a transient light pulse, in which either desynchrony alone [74] or a combination of single-cell amplitude reduction and desynchrony [73] was identified as the dominant factor.

Data on the response of melanopsin-responsive and control NIH3T3 mouse fibroblasts cells is shown in figure 5.10 [74]. The experimental data is de-noised using a discrete wavelet transform [100]. In the experiment, a light pulse is given to desynchronize a colony of cells, resulting in drastically reduced amplitudes. A second (longer) light pulse subsequently re-synchronizes the population, resulting in increased amplitudes. To capture this phenomenon *in silico*, we used a recent model of the core circadian feedback circuit [3]. In order to find parameters for the phase density function, we assumed an initial phase population with $\sigma = 0.1$ and calculated a phase diffusivity parameter $d = 0.104$ from the experimental control trajectory.

The next step in capturing the experimental data was finding a perturbation capable of desynchronizing the system. To find such a perturbation, we calculated population-level ARCs at several different knockdown strengths for each parameter in the model. We selected the degradation rate of *Per* mRNA from the feasible parameters due to PER's known induction by CREB following photo-perturbation [101]. While not an exact match of the experimental system, reducing degradation rate yields a similar response to increasing transcription. The parameter was reduced by 28.5% during the light pulses, a value fit to maximize light-induced desynchrony. The output variable, PER2-luciferase in the experimental system, was chosen to be represented by *Cry1* mRNA - an E box activated gene that is buffered from the direct effects of the parameter perturbation. The single-cell and population-level response




Figure 5.10: Method allows direct comparison between model results and experimental bioluminescence profiles. (A) Response curves for the parameter perturbation used to model the first light pulse in (B). The initial phase of the system is chosen to coincide with the strong dip in the population-level ARC (left). (B) De-noised bioluminescence data from Ukai *et al.* 2007 (top) demonstrates population-level amplitude change resulting from two light pulses [74]. Circadian oscillations are suppressed by the first light pulse and rescued by the second. These perturbations were reproduced *in silico* (bottom) using the model from Hirota *et al.* 2012. The experimental results are qualitatively captured by the model, demonstrating the suitability of the method in capturing bioluminescence experiments. The relative amplitude of the model equations has been scaled to match that of the normalized bioluminescence profiles. (C) Example probability density functions for the model trajectory shown in (A). The initial population (red) is effectively desynchronized (orange) following the first light pulse.

curves for this parameter choice are shown in figure 5.10, which demonstrates the strong desynchronization that occurs at $\theta \approx \pi/4$. Since the reporter used in the experimental system likely has a phase lag from the corresponding mRNA, we did not try to match the initial phase of the simulation to experiment. Instead, the initial phase of simulation was chosen such that the first pulse occurred when the system was at the phase that corresponded with the minimum of the population-level ARC.

Population-level trajectories were found by solving for the resulting limit cycle trajectories at every phase, and subsequently finding the weighted average of these trajectories according the phase density (as described in equations (5.38 and 5.46)). The simulated control and perturbed trajectories, Fig. 5B, closely match the experimental results, in which the model captures both the phase shifts and amplitude modulation of the light pulses. The phase probability density function for the light-sensitive model trajectory is shown in Fig. 5C for several representative time points. Changes to the synchronization of the population from each perturbation are readily apparent, with both perturbations inducing a bimodal distribution in the phase density. While experimental data on individual cell phases was unavailable for this data set, bimodal distributions in SCN neuron firing following a phase shift have previously been seen experimentally [102]. As higher-frequency features, these bimodal distributions dissipate as the phases diffuse. Amplitude change in this case is mediated both at the population and single-cell level, as indicated by their representative ARCs. The contributions from each source are summarized in Fig. S4.

Several quantitative differences between the model and experiment do appear. Most importantly, the acute amplitude change following the light pulse is not correctly captured by the single-cell model. From the experimental system, it appears the light pulse temporarily reduces PER2-luciferase amplitudes immediately following perturbation, separate from the overall change in population synchrony. This result is most obvious after the second light pulse, where rhythms require some

time before their maximum amplitude is reached. Such a delay is indicative of perturbations to the limit cycle system, suggesting a contribution from single-cells in determining the population amplitude. Light-induced amplitude suppression in fibroblasts is therefore likely mediated at the single-cell level at short time-scales, and at the population level for longer time-scales. To improve the fit of the model to the data, the response curves could be fit to match experimentally predicted values. Infinitesimal PRCs and ARCs calculated at the single-cell level are numerically efficient to evaluate, and could be more readily incorporated into a parameter estimation algorithm than explicit stochastic simulations of a population of oscillators.

5.4 Discussion

In this chapter, we have described new tools for understanding amplitude change in independent cells based on analytically tractable methods for simulating a large population of oscillators. These tools synthesize single-cell level ODE sensitivity analysis metrics with population-level measures of synchrony, allowing the quantification of clock amplitude at multiple levels of biological organization. Using these tools, we have demonstrated how ODE models can be directly compared to experimental bioluminescence profiles, which currently serve as a standard system to investigate clock perturbations. More generally, we have demonstrated the differences between acute and prolonged changes in amplitude following temporary perturbations, and characterized the mechanisms that give rise to each.

5.4.1 Application to pharmacological control

While our examples have focused on light-mediated perturbations, these approaches could also be applied to pharmacological perturbations. Some difficulty arises in obtaining such data in cultured cells, however, as entraining perturbations would

require that pharmacological agents be introduced for only a finite duration. As medium or temperature changes are often enough to resynchronize cultured cells, a transient application is difficult to achieve experimentally. The search for clock-enhancing molecules has therefore tended to focus on constant drug concentrations: for instance, dose-dependent period or amplitude change following inhibition of CK1 δ or similar targets [2]. For *in vivo* systems, however, pharmacokinetics dictates a finite duration of action for both naturally secreted hormones and pharmacological therapies. More effective treatments might therefore be designed by explicitly accounting for such a transient response, perturbing peripheral clocks at the right phase to induce resynchronization and an increase in single-cell level amplitudes.

Maintaining robust rhythmicity in peripheral oscillators has been linked to protection against metabolic disease. It is therefore interesting that liver cells have not developed a stronger mechanism of intercellular coupling, such as in the SCN, to maintain robust amplitudes in the absence of external cues. Perhaps the ability to quickly re-entrain to a phase shift, which is often slowed by coupling [65], has historically been more advantageous than protection against highly variable food intake. Regardless, the necessities of modern society often dictate irregular circadian-metabolic regimes that damp peripheral clock oscillations. For such instances, we hope that the mathematical theory presented here will permit the development of optimized treatment regimes.

5.4.2 Damping rate is correlated with single-cell noise

One result from this chapter, which has previously been discussed in the circadian literature [93], is that in noninteracting populations, stochastic noise at the single-cell level affects the damping rate of a population of oscillators. Stochastic noise is often a relevant circadian parameter, as it describes the precision of the underlying oscillators. In the following chapter, I show how population-level recordings, even those

from high-throughput circadian screens, can be used to gain estimates of stochastic noise at the single-cell level. Using such an approach, it is possible to gain genome-wide insight into how noise is regulated in the cell-autonomous feedback loop.

Chapter 6

Decay rate as proxy for cellular noise

Chapter 7

Conclusions and future work

7.1 Continuation of collaboration with Andrew

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.1.1 Experimental validation of model predictions

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of

the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.1.2 Prediction of fasting perturbations

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

small-molecule activators of SIRT1

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.1.3 Validity of step vs pulse perturbation

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Characterization of insulin effect

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.2 Amplitude maximization through therapeutic treatment

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some

nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.2.1 Optimal control strategies to maximize peripheral clock amplitude

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.2.2 Fewest treatments, combinatorial perturbations, etc.

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.3 Iterative optimization of network topology

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.3.1 (Idea from Mitsubishi proposal)

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

7.3.2 Design machine-learning algorithm to find optimum model structure

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like “Huardest gefburn”? Kjift – not at all! A blind text like this gives you

information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

Bibliography

- [1] J. C. Dunlap, J. J. Loros, and P. J. DeCoursey, *Chronobiology: Biological Timekeeping*. Sinauer Associates, Inc., 2009.
- [2] Z. Chen, S.-H. Yoo, and J. S. Takahashi, "Small molecule modifiers of circadian clocks," *Cell. Mol. Life Sci.*, vol. 70, pp. 2985–98, Aug. 2013.
- [3] T. Hirota, J. W. Lee, P. C. St. John, M. Sawa, K. Iwaisako, T. Noguchi, P. Y. Pongsawakul, T. Sonntag, D. K. Welsh, D. A. Brenner, F. J. Doyle, P. G. Schultz, and S. A. Kay, "Identification of Small Molecule Activators of Cryptochrome," *Science*, vol. 337, pp. 1094–1097, July 2012.
- [4] E. V. McCarthy, J. E. Baggs, J. M. Geskes, J. B. Hogenesch, and C. B. Green, "Generation of a novel allelic series of cryptochrome mutants via mutagenesis reveals residues involved in protein-protein interaction and CRY2-specific repression," *Mol. Cell. Biol.*, vol. 29, pp. 5465–76, Oct. 2009.
- [5] G. T. van der Horst, M. Muijtjens, K. Kobayashi, R. Takano, S. Kanno, M. Takao, J. de Wit, A. Verkerk, A. P. Eker, D. van Leenen, R. Buijs, D. Bootsma, J. H. Hoeijmakers, and A. Yasui, "Mammalian Cry1 and Cry2 are essential for maintenance of circadian rhythms," *Nature*, vol. 398, pp. 627–30, Apr. 1999.
- [6] E. E. Zhang, A. C. Liu, T. Hirota, L. J. Miraglia, G. Welch, P. Y. Pongsawakul, X. Liu, A. Atwood, J. W. Huss, J. Janes, A. I. Su, J. B. Hogenesch, and S. A. Kay, "A genome-wide RNAi screen for modifiers of the circadian clock in human cells," *Cell*, vol. 139, no. 1, pp. 199–210, 2009.
- [7] K. Miyazaki, M. Mesaki, and N. Ishida, "Nuclear entry mechanism of rat PER2 (rPER2): role of rPER2 in nuclear localization of CRY protein," *Mol. Cell. Biol.*, vol. 21, pp. 6651–9, Oct. 2001.
- [8] C. H. Ko and J. S. Takahashi, "Molecular components of the mammalian circadian clock," *Hum. Mol. Genet.*, vol. 15, pp. R271–7, Oct. 2006.
- [9] K. Kume, M. J. Zylka, S. Sriram, L. P. Shearman, D. R. Weaver, X. Jin, E. S. Maywood, M. H. Hastings, and S. M. Reppert, "mCRY1 and mCRY2 are essential components of the negative limb of the circadian clock feedback loop," *Cell*, vol. 98, no. 2, pp. 193–205, 1999.

- [10] J.-C. Leloup and A. Goldbeter, "Toward a detailed computational model for the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 100, pp. 7051–6, June 2003.
- [11] H. P. Mirsky, A. C. Liu, D. K. Welsh, S. A. Kay, and F. J. Doyle, "A model of the cell-autonomous mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 106, pp. 11107–12, July 2009.
- [12] D. B. Forger and C. S. Peskin, "A detailed predictive model of the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 100, pp. 14806–11, Dec. 2003.
- [13] C. Lee, J.-P. Etchegaray, F. R. Cagampang, A. S. I. Loudon, and S. M. Reppert, "Posttranslational mechanisms regulate the mammalian circadian clock," *Cell*, vol. 107, pp. 855–67, Dec. 2001.
- [14] Y. Lee, R. Chen, H.-m. Lee, and C. Lee, "Stoichiometric relationship among clock proteins determines robustness of circadian rhythms," *J. Biol. Chem.*, vol. 286, pp. 7033–42, Mar. 2011.
- [15] N. A. Cookson, L. S. Tsimring, and J. Hasty, "The pedestrian watchmaker: genetic clocks from engineered oscillators," *FEBS Lett.*, vol. 583, pp. 3931–7, Dec. 2009.
- [16] N. Kurabayashi, T. Hirota, M. Sakai, K. Sanada, and Y. Fukada, "DYRK1A and glycogen synthase kinase 3beta, a dual-kinase mechanism directing proteosomal degradation of CRY2 for circadian timekeeping," *Mol. Cell. Biol.*, vol. 30, pp. 1757–68, Apr. 2010.
- [17] E. Vielhaber, E. Eide, A. Rivers, Z.-H. Gao, and D. M. Virshup, "Nuclear Entry of the Circadian Regulator mPER1 Is Controlled by Mammalian Casein Kinase Ie," *Mol. Cell. Biol.*, vol. 20, pp. 4888–4899, July 2000.
- [18] E. A. Griffin Jr., D. Staknis, and C. J. Weitz, "Light-Independent Role of CRY1 and CRY2 in the Mammalian Circadian Clock," *Science*, vol. 286, pp. 768–771, Oct. 1999.
- [19] J. Andersson, B. Houska, and M. Diehl, "Towards a Computer Algebra System with Automatic Differentiation for use with Object-Oriented modelling languages," *3rd Int. Work. Equation-Based Object-Oriented Model. Lang. Tools*, pp. 99–105, 2010.
- [20] A. C. Hindmarsh, P. N. Brown, K. E. Grant, S. L. Lee, R. Serban, D. A. N. E. Shumaker, and C. S. Woodward, "SUNDIALS : Suite of Nonlinear and Differential / Algebraic Equation Solvers," *ACM Trans. Math. Softw.*, vol. 31, no. 3, pp. 363–396, 2005.
- [21] A. K. Wilkins, B. Tidor, J. White, and P. I. Barton, "Sensitivity Analysis for Oscillating Dynamical Systems," *SIAM J. Sci. Comput.*, vol. 31, no. 4, p. 2706, 2009.

- [22] A. Relógio, P. O. Westermark, T. Wallach, K. Schellenberg, A. Kramer, and H. Herz, "Tuning the Mammalian Circadian Clock: Robust Synergy of Two Loops," *PLoS Comput. Biol.*, vol. 7, p. e1002309, Dec. 2011.
- [23] A. C. Liu, D. K. Welsh, C. H. Ko, H. G. Tran, E. E. Zhang, A. A. Priest, E. D. Buhr, O. Singer, K. Meeker, I. M. Verma, F. J. Doyle, J. S. Takahashi, and S. A. Kay, "Intercellular coupling confers robustness against mutations in the SCN circadian clock network," *Cell*, vol. 129, pp. 605–16, May 2007.
- [24] R. N. Gutenkunst, J. J. Waterfall, F. P. Casey, K. S. Brown, C. R. Myers, and J. P. Sethna, "Universally sloppy parameter sensitivities in systems biology models," *PLoS Comput. Biol.*, vol. 3, pp. 1871–78, Oct. 2007.
- [25] D. B. Kell and J. D. Knowles, "The role of modeling in systems biology," in *Syst. Model. Cell. Biol. from concepts to nuts bolts* (Z. Szallasi, J. Stelling, and V. Periwal, eds.), pp. 3–18, Cambridge: MIT Press, 2006.
- [26] A. Raue, C. Kreutz, T. Maiwald, J. Bachmann, M. Schilling, U. Klingmüller, and J. Timmer, "Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood," *Bioinformatics*, vol. 25, pp. 1923–9, Aug. 2009.
- [27] M. Nihtilä and J. Virkkunen, "Practical identifiability of growth and substrate consumption models," *Biotechnol. Bioeng.*, vol. 19, pp. 1831–50, Dec. 1977.
- [28] J. E. Jiménez-Hornero, I. M. Santos-Dueñas, and I. García-García, "Optimization of biotechnological processes. The acetic acid fermentation. Part II: Practical identifiability analysis and parameter estimation," *Biochem. Eng. J.*, vol. 45, pp. 7–21, June 2009.
- [29] A. Holmberg, "On the practical identifiability of microbial growth models incorporating Michaelis-Menten type nonlinearities," *Math. Biosci.*, vol. 62, pp. 23–43, Nov. 1982.
- [30] M. Joshi, A. Seidel-Morgenstern, and A. Kremling, "Exploiting the bootstrap method for quantifying parameter confidence intervals in dynamical systems," *Metab. Eng.*, vol. 8, pp. 447–55, Sept. 2006.
- [31] M. E. Hughes, L. DiTacchio, K. R. Hayes, C. Vollmers, S. Pulivarthy, J. E. Baggs, S. Panda, and J. B. Hogenesch, "Harmonics of circadian gene transcription in mammals," *PLoS Genet.*, vol. 5, p. e1000442, Apr. 2009.
- [32] T. Hirota, J. W. Lee, W. G. Lewis, E. E. Zhang, G. Breton, X. Liu, M. Garcia, E. C. Peters, J.-P. Etchegaray, D. Traver, P. G. Schultz, and S. A. Kay, "High-Throughput Chemical Screen Identifies a Novel Potent Modulator of Cellular Circadian Rhythms and Reveals CKI α as a Clock Regulatory Kinase," *PLoS Biol.*, vol. 8, p. e1000559, Dec. 2010.

- [33] T. Hirota, W. G. Lewis, A. C. Liu, J. W. Lee, P. G. Schultz, and S. A. Kay, "A chemical biology approach reveals period shortening of the mammalian circadian clock by specific inhibition of GSK-3beta.," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 105, pp. 20746–51, Dec. 2008.
- [34] L. T. Biegler, *Nonlinear Programming: Concepts, Algorithms, and Applications to Chemical Processes*. SIAM, 2010.
- [35] H. G. Bock, E. Kostina, and J. P. Schloder, "Numerical methods for parameter estimation in nonlinear differential algebraic equations," *GAMM Mitteilungen*, vol. 408, no. 2, pp. 376 – 408, 2007.
- [36] C. A. Floudas, *Nonlinear and Mixed-Integer Optimization*. New York, New York, USA: Oxford University Press, 1995.
- [37] A. Wachter and L. T. Biegler, "On the implementation of an interior-point filter line-search algorithm for large-scale nonlinear programming," *Math. Program.*, vol. 106, pp. 25–57, Apr. 2006.
- [38] HSL, "A collection of Fortran codes for large scale scientific computation," 2011.
- [39] J. Andersson, *A General-Purpose Software Framework for Dynamic Optimization*. Phd thesis, Arenberg Doctoral School, KU Leuven, Department of Electrical Engineering (ESAT/SCD) and Optimization in Engineering Center, Kasteelpark Arenberg 10, 3001-Heverlee, Belgium, Oct. 2013.
- [40] E. Jones, T. Oliphant, and P. Peterson, "SciPy: Open Source Scientific Tools for Python," 2001.
- [41] R. Serban and A. C. Hindmarsh, "CVODES: the Sensitivity-Enabled ODE Solver in SUNDIALS," in *Proc. IDETC/CIE 2005*, (Long Beach, CA), 2005.
- [42] M. A. Kramer, H. Rabitz, and J. M. Calo, "Sensitivity analysis of oscillatory systems," *Appl. Math. Model.*, vol. 8, pp. 328–340, Oct. 1984.
- [43] E. G. Bure and E. Rosenwasser, "The study of the sensitivity of oscillatory systems," *Autom. Remote Control*, vol. 7, pp. 1045–1052, 1974.
- [44] D. E. Zak, G. E. Gonye, J. S. Schwaber, and F. J. Doyle, "Importance of input perturbations and stochastic gene expression in the reverse engineering of genetic regulatory networks: insights from an identifiability analysis of an in silico network.," *Genome Res.*, vol. 13, pp. 2396–405, Nov. 2003.
- [45] J. Bass, "Circadian topology of metabolism.," *Nature*, vol. 491, pp. 348–56, Nov. 2012.
- [46] R. Ye, C. P. Selby, N. Ozturk, Y. Annayev, and A. Sancar, "Biochemical analysis of the canonical model for the mammalian circadian clock.," *J. Biol. Chem.*, vol. 286, pp. 25891–902, July 2011.

- [47] K. Yagita, F. Tamanini, M. Yasuda, J. H. J. Hoeijmakers, G. T. J. van der Horst, and H. Okamura, "Nucleocytoplasmic shuttling and mCRY-dependent inhibition of ubiquitylation of the mPER2 clock protein," *EMBO J.*, vol. 21, pp. 1301–14, Mar. 2002.
- [48] J. S. Takahashi, H.-K. Hong, C. H. Ko, and E. L. McDearmon, "The genetics of mammalian circadian order and disorder: implications for physiology and disease," *Nat. Rev. Genet.*, vol. 9, pp. 764–75, Oct. 2008.
- [49] S. Reischl, K. Vanselow, P. I. O. Westermark, N. Thierfelder, B. Maier, H. Herz, and A. Kramer, "Beta-TrCP1-mediated degradation of PERIOD2 is essential for circadian dynamics," *J. Biol. Rhythms*, vol. 22, pp. 375–86, Oct. 2007.
- [50] A. Takano, Y. Isojima, and K. Nagai, "Identification of mPer1 phosphorylation sites responsible for the nuclear entry," *J. Biol. Chem.*, vol. 279, pp. 32578–85, July 2004.
- [51] L. Busino, F. Bassermann, A. Maiolica, C. Lee, P. M. Nolan, S. I. H. Godinho, G. F. Draetta, and M. Pagano, "SCFFbxl3 controls the oscillation of the circadian clock by directing the degradation of cryptochrome proteins," *Science*, vol. 316, pp. 900–4, May 2007.
- [52] S. I. H. Godinho, E. S. Maywood, L. Shaw, V. Tucci, A. R. Barnard, L. Busino, M. Pagano, R. Kendall, M. M. Quwailid, M. R. Romero, J. O'neill, J. E. Chesham, D. Brooker, Z. Lalanne, M. H. Hastings, and P. M. Nolan, "The after-hours mutant reveals a role for Fbxl3 in determining mammalian circadian period," *Science*, vol. 316, pp. 897–900, May 2007.
- [53] S. M. Siepka, S.-H. Yoo, J. Park, W. Song, V. Kumar, Y. Hu, C. Lee, and J. S. Takahashi, "Circadian mutant Overtime reveals F-box protein FBXL3 regulation of cryptochrome and period gene expression," *Cell*, vol. 129, pp. 1011–23, June 2007.
- [54] Y. Lee and E.-K. Kim, "AMP-activated protein kinase as a key molecular link between metabolism and clockwork," *Exp. Mol. Med.*, vol. 45, p. e33, Jan. 2013.
- [55] M. Gallego, E. J. Eide, M. F. Woolf, D. M. Virshup, and D. B. Forger, "An opposite role for tau in circadian rhythms revealed by mathematical modeling," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 103, pp. 10618–23, July 2006.
- [56] E. S. Maywood, J. E. Chesham, Q.-J. Meng, P. M. Nolan, A. S. I. Loudon, and M. H. Hastings, "Tuning the period of the mammalian circadian clock: additive and independent effects of CK1εTau and Fbxl3Afh mutations on mouse circadian behavior and molecular pacemaking," *J. Neurosci.*, vol. 31, pp. 1539–44, Jan. 2011.
- [57] K. Vanselow, J. T. Vanselow, P. I. O. Westermark, S. Reischl, B. Maier, T. Korte, A. Herrmann, H. Herz, A. Schlosser, and A. Kramer, "Differential effects of

- PER2 phosphorylation: molecular basis for the human familial advanced sleep phase syndrome (FASPS)," *Genes Dev.*, vol. 20, pp. 2660–72, Oct. 2006.
- [58] P. C. St. John and F. J. Doyle, "Estimating confidence intervals in predicted responses for oscillatory biological models," *BMC Syst. Biol.*, vol. 7, p. 71, July 2013.
- [59] R. Gunawan and F. J. Doyle, "Isochron-based phase response analysis of circadian rhythms," *Biophys. J.*, vol. 91, pp. 2131–41, Sept. 2006.
- [60] K. Ohsaki, K. Oishi, Y. Kozono, K. Nakayama, K. I. Nakayama, and N. Ishida, "The role of beta-TrCP1 and beta-TrCP2 in circadian rhythm generation by mediating degradation of clock protein PER2," *J. Biochem.*, vol. 144, pp. 609–18, Nov. 2008.
- [61] H. Lee, R. Chen, Y. Lee, S. Yoo, and C. Lee, "Essential roles of CKIdelta and CKIepsilon in the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 106, pp. 21359–64, Dec. 2009.
- [62] K. A. Lamia, K.-F. Storch, and C. J. Weitz, "Physiological significance of a peripheral tissue circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 105, pp. 15172–7, Sept. 2008.
- [63] B. Kornmann, O. Schaad, H. Bujard, J. S. Takahashi, and U. Schibler, "System-driven and oscillator-dependent circadian transcription in mice with a conditionally active liver clock," *PLoS Biol.*, vol. 5, p. e34, Feb. 2007.
- [64] C. S. Pittendrigh, W. T. Kyner, and T. Takamura, "The Amplitude of Circadian Oscillations: Temperature Dependence, Latitudinal Clines, and the Photoperiodic Time Measurement," *J. Biol. Rhythms*, vol. 6, pp. 299–313, Dec. 1991.
- [65] U. Abraham, A. E. Granada, P. I. O. Westermark, M. Heine, A. Kramer, and H. Herzel, "Coupling governs entrainment range of circadian clocks," *Mol. Syst. Biol.*, vol. 6, p. 438, Nov. 2010.
- [66] B. Marcheva, K. M. Ramsey, E. D. Buhr, Y. Kobayashi, H. Su, C. H. Ko, G. Ivanova, C. Omura, S. Mo, M. H. Vitaterna, J. P. Lopez, L. H. Philipson, C. a. Bradfield, S. D. Crosby, L. JeBailey, X. Wang, J. S. Takahashi, and J. Bass, "Disruption of the clock components CLOCK and BMAL1 leads to hypoinsulinaemia and diabetes," *Nature*, vol. 466, pp. 627–31, July 2010.
- [67] M. Hatori, C. Vollmers, A. Zarrinpar, L. DiTacchio, E. A. Bushong, S. Gill, M. Leblanc, A. Chaix, M. Joens, J. A. J. Fitzpatrick, M. H. Ellisman, and S. Panda, "Time-restricted feeding without reducing caloric intake prevents metabolic diseases in mice fed a high-fat diet," *Cell Metab.*, vol. 15, pp. 848–60, June 2012.

- [68] H.-C. Chang and L. Guarente, "SIRT1 Mediates Central Circadian Control in the SCN by a Mechanism that Decays with Aging," *Cell*, vol. 153, pp. 1448–1460, June 2013.
- [69] P. C. St. John, T. Hirota, S. A. Kay, and F. J. Doyle, "Spatiotemporal separation of PER and CRY posttranslational regulation in the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 111, pp. 2040–5, Feb. 2014.
- [70] D. K. Welsh, S.-H. Yoo, A. C. Liu, J. S. Takahashi, and S. A. Kay, "Bioluminescence imaging of individual fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression," *Curr. Biol.*, vol. 14, pp. 2289–95, Dec. 2004.
- [71] C. Ramanathan, H. Xu, S. K. Khan, Y. Shen, P. J. Gitis, D. K. Welsh, J. B. Hogenesch, and A. C. Liu, "Cell type-specific functions of period genes revealed by novel adipocyte and hepatocyte circadian clock models," *PLoS Genet.*, vol. 10, p. e1004244, Apr. 2014.
- [72] J. E. Baggs, T. S. Price, L. DiTacchio, S. Panda, G. A. Fitzgerald, and J. B. Hogenesch, "Network features of the mammalian circadian clock," *PLoS Biol.*, vol. 7, p. e52, Mar. 2009.
- [73] S. R. Pulivarthy, N. Tanaka, D. K. Welsh, L. De Haro, I. M. Verma, and S. Panda, "Reciprocity between phase shifts and amplitude changes in the mammalian circadian clock," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 104, pp. 20356–61, Dec. 2007.
- [74] H. Ukai, T. J. Kobayashi, M. Nagano, K.-h. Masumoto, M. Sujino, T. Kondo, K. Yagita, Y. Shigeyoshi, and H. R. Ueda, "Melanopsin-dependent photoperturbation reveals desynchronization underlying the singularity of mammalian circadian clocks," *Nat. Cell Biol.*, vol. 9, pp. 1327–34, Nov. 2007.
- [75] S. Becker-Weimann, J. Wolf, H. Herz, and A. Kramer, "Modeling feedback loops of the Mammalian circadian oscillator," *Biophys. J.*, vol. 87, pp. 3023–34, Nov. 2004.
- [76] D. A. Rand, B. V. Shulgin, D. Salazar, and A. J. Millar, "Design principles underlying circadian clocks," *J. R. Soc. Interface*, vol. 1, pp. 119–30, Nov. 2004.
- [77] S. Daan and C. S. Pittendrigh, "A Functional analysis of circadian pacemakers in nocturnal rodents," *J. Comp. Physiol. A*, vol. 106, pp. 253–266, Oct. 1976.
- [78] S. R. Taylor, F. J. Doyle, and L. R. Petzold, "Oscillator model reduction preserving the phase response: application to the circadian clock," *Biophys. J.*, vol. 95, pp. 1658–73, Aug. 2008.
- [79] B. Pfeuty, Q. Thommen, and M. Lefranc, "Robust entrainment of circadian oscillators requires specific phase response curves," *Biophys. J.*, vol. 100, pp. 2557–65, June 2011.

- [80] J. K. Kim, Z. P. Kilpatrick, M. R. Bennett, and K. Josić, "Molecular Mechanisms that Regulate the Coupled Period of the Mammalian Circadian Clock," *Biophys. J.*, vol. 106, pp. 2071–2081, May 2014.
- [81] S. R. Taylor, R. Gunawan, L. R. Petzold, and F. J. Doyle, "Sensitivity Measures for Oscillating Systems: Application to Mammalian Circadian Gene Network," *IEEE Trans. Automat. Contr.*, vol. 53, pp. 177–188, Jan. 2008.
- [82] M. E. Jewett and R. E. Kronauer, "Refinement of a limit cycle oscillator model of the effects of light on the human circadian pacemaker," *J. Theor. Biol.*, vol. 192, pp. 455–65, June 1998.
- [83] P. Achermann and H. Kunz, "Modeling Circadian Rhythm Generation in the Suprachiasmatic Nucleus with Locally Coupled Self-Sustained Oscillators: Phase Shifts and Phase Response Curves," *J. Biol. Rhythms*, vol. 14, pp. 460–468, Dec. 1999.
- [84] D. R. van der Veen, J. Shao, Y. Xi, L. Li, and G. E. Duffield, "Cardiac atrial circadian rhythms in PERIOD2::LUCIFERASE and per1:luc mice: amplitude and phase responses to glucocorticoid signaling and medium treatment," *PLoS One*, vol. 7, p. e47692, Jan. 2012.
- [85] O. Castejón, A. Guillamon, and G. Huguet, "Phase-amplitude response functions for transient-state stimuli," *J. Math. Neurosci.*, vol. 3, p. 13, Jan. 2013.
- [86] F. van Oosterhout, E. a. Lucassen, T. Houben, H. T. VanderLeest, M. C. Antle, and J. H. Meijer, "Amplitude of the SCN clock enhanced by the behavioral activity rhythm," *PLoS One*, vol. 7, p. e39693, Jan. 2012.
- [87] S. An, R. Harang, K. Meeker, D. Granados-Fuentes, C. a. Tsai, C. Mazuski, J. Kim, F. J. Doyle, L. R. Petzold, and E. D. Herzog, "A neuropeptide speeds circadian entrainment by reducing intercellular synchrony," *Proc. Natl. Acad. Sci. U. S. A.*, vol. 110, pp. E4355–61, Nov. 2013.
- [88] H. Rabitz, M. Kramer, and D. Dacol, "Sensitivity Analysis in Chemical Kinetics," *Annu. Rev. Phys. Chem.*, vol. 34, pp. 419–461, Oct. 1983.
- [89] J. Rougemont and F. Naef, "Collective synchronization in populations of globally coupled phase oscillators with drifting frequencies," *Phys. Rev. E*, vol. 73, p. 011104, Jan. 2006.
- [90] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence*. Springer-Verlag Berlin Heidelberg, 1984.
- [91] J.-n. Teramae and D. Tanaka, "Robustness of the noise-induced phase synchronization in a general class of limit cycle oscillators," *Phys. Rev. Lett.*, vol. 93, p. 204103, Nov. 2004.

Bibliography

- [92] R. B. Stein, "A theoretical analysis of neuronal variability," *Biophys. J.*, vol. 5, pp. 173–94, Mar. 1965.
- [93] J. Rougemont and F. Naef, "Dynamical signatures of cellular fluctuations and oscillator stability in peripheral circadian clocks," *Mol. Syst. Biol.*, vol. 3, p. 93, Jan. 2007.
- [94] G. S. Chirikjian, "Gaussian Distributions and the Heat Equation," in *Stoch. Model. Inf. Theory, Lie Groups, Vol. 1*, vol. 1, ch. 2, pp. 31–61, Boston: Birkhäuser Boston, 2009.
- [95] K. V. Mardia and P. E. Jupp, *Directional Statistics*. John Wiley & Sons, 2009.
- [96] K. R. Sanft, S. Wu, M. Roh, J. Fu, R. K. Lim, and L. R. Petzold, "StochKit2: software for discrete stochastic simulation of biochemical systems with events," *Bioinformatics*, vol. 27, pp. 2457–8, Sept. 2011.
- [97] B. Novák and J. J. Tyson, "Design principles of biochemical oscillators," *Nat. Rev. Mol. Cell Biol.*, vol. 9, pp. 981–91, Dec. 2008.
- [98] D. T. Gillespie, "Exact Stochastic Simulation of Coupled Chemical Reactions," *J. Phys. Chem.*, vol. 84, no. 25, pp. 2340–2361, 1977.
- [99] R. J. Field and R. M. Noyes, "Oscillations in chemical systems. IV. Limit cycle behavior in a model of a real chemical reaction," *J. Chem. Phys.*, vol. 60, no. 5, p. 1877, 1974.
- [100] T. L. Leise and M. E. Harrington, "Wavelet-based time series analysis of circadian rhythms," *J. Biol. Rhythms*, vol. 26, pp. 454–63, Oct. 2011.
- [101] S. a. Tischkau, J. W. Mitchell, S.-H. Tyan, G. F. Buchanan, and M. U. Gillette, "Ca²⁺/cAMP response element-binding protein (CREB)-dependent activation of Per1 is required for light-induced signaling in the suprachiasmatic nucleus circadian clock," *J. Biol. Chem.*, vol. 278, pp. 718–23, Jan. 2003.
- [102] J. H. T. Rohling, H. T. VanderLeest, S. Michel, M. J. Vansteensel, and J. H. Meijer, "Phase resetting of the mammalian circadian clock relies on a rapid shift of a small population of pacemaker neurons," *PLoS One*, vol. 6, p. e25437, Jan. 2011.