

THE HIGGS BOSON FOR MATHEMATICIANS. LECTURE NOTES ON GAUGE THEORY AND SYMMETRY BREAKING

M. J. D. HAMILTON

ABSTRACT. These notes form part of a lecture course on gauge theory. The material covered is standard in the physics literature, but perhaps less well-known to mathematicians. The purpose of these notes is to make spontaneous symmetry breaking and the Higgs mechanism of mass generation for elementary particles more easily accessible to mathematicians interested in theoretical physics. We treat the general case with an arbitrary compact gauge group G and an arbitrary number of Higgs bosons and explain the situation in the classic case of the electroweak interaction where $G = SU(2) \times U(1)$. Prerequisites are only a basic knowledge of Lie groups and manifolds. No prior knowledge of gauge theory or bundle theory is assumed.

1. INTRODUCTION

Spontaneous symmetry breaking and the Higgs mechanism of mass generation constitute an important part of the Standard Model of elementary particles. Just as gauge theories play a significant role in pure mathematics, like the Donaldson and Seiberg-Witten theories of smooth 4-manifolds, it seems worthwhile to study the Higgs mechanism also from a mathematical and, in particular, geometric point of view.

It is useful to consider the general case of an arbitrary compact Lie group G and an arbitrary number of Higgs bosons from the very beginning, because it will then become clearer what is special in the situation of the electroweak $SU(2) \times U(1)$ -theory that appears in the Standard Model of particle physics. Regarding abstraction we chose to stay midway between mathematics and physics: We treat the general case of a compact gauge group G , but we want to keep the physical intuition of fields as maps on spacetime with values in a vector space rather than sections in a vector bundle over the spacetime manifold. We therefore essentially treat the case of trivial bundles over an arbitrary spacetime manifold M . Many notions can be generalized to arbitrary bundles. Furthermore, since we are mainly interested in the generation of the mass of elementary particles, which appears in the Lagrangian in terms of

Date: 24 Jul 2020 .

Key words and phrases. gauge theory, spontaneous symmetry breaking, Higgs boson.

order two, we do not describe the interaction between the Higgs boson and other particles, corresponding to terms of order three and higher in the fields.

Our main references are: For Sections 2, 4, 8 and 9 the lecture notes [Louis 2011] and the book [Mosel 1999], for Section 3 the lecture notes [Ziller 2010] and for Sections 5, 6 and 7 the book [Bleecker 1981]. Good references with background on the Standard Model and symmetry breaking are [Robinson 2011] and [Ryder 1996] (to name but two). For the general definition of Yukawa couplings in Section 9 we did not find a suitable reference. The definition here is a (provisional) generalization of the example appearing in the electroweak gauge theory.

We use the Einstein summation convention throughout and always assume that gauge groups are compact, connected Lie groups.

2. GAUGE THEORIES

Many interesting theories in physics, like the Standard Model of elementary particles, are gauge theories. The idea of gauge theories is that *symmetry generates interaction*. What does that mean?

Gauge theories are certain **field theories**; in particular, matter is described by fields. These field theories have a certain symmetry, called **gauge symmetry**. (One could argue that the major step is actually to describe matter by fields and to collect several such fields into multiplets; see below. Once this is done, the introduction of gauge symmetry is quite natural.) Like every symmetry, gauge symmetry consists of two parts:

- Action or transformation
- Invariance

2.1. Transformation. In the case of gauge symmetry the action consists of *local gauge transformations*. Suppose M is our spacetime manifold and

$$\psi_i: M \longrightarrow S \quad \text{for } i = 1, \dots, n,$$

are matter fields defined on M . Here S denotes a (usually complex) vector space. The fields ψ_i have a certain behaviour under Lorentz transformations, for example, they could be scalars, spinors, etc.¹

We now collect these matter fields into what could be called a **multiplet** Ψ , i.e. a vector of fields:

$$\Psi = \begin{pmatrix} \psi_1 \\ \vdots \\ \psi_n \end{pmatrix}.$$

¹We assume that M has a Riemannian or Lorentzian metric if we want to define spinors or if we contract spacetime indices, but some parts of the theory of gauge symmetry and symmetry breaking do not involve any metric on the manifold M .

Mathematically a multiplet is a field

$$\Psi: M \longrightarrow S \otimes V,$$

where V is a complex vector space of dimension n . The space $S \otimes V$ is also called **twisted**, in particular in the case of spinors.

We also have a compact, connected Lie group G , called the **gauge group**, that acts as a linear representation on the multiplet space:

$$G \times V \longrightarrow V, \quad (g, v) \longmapsto g \cdot v.$$

The Lie group G then also acts on the space $S \otimes V$ (trivially on the first factor). The action of G therefore mixes the components ψ_i of the multiplet Ψ according to the representation on V , in the sense that under a group element $g \in G$ every component ψ_i of the multiplet gets mapped to some linear combination of ψ_1, \dots, ψ_n .

Definition 2.1. A (local) gauge transformation is a map

$$\sigma: M \longrightarrow G$$

of the spacetime manifold M into the Lie group G that acts in every point $x \in M$ on the multiplet $\Psi(x)$ via the representation on V :

$$\Psi \longmapsto \sigma \cdot \Psi, \quad (\sigma \cdot \Psi)(x) = \sigma(x) \cdot \Psi(x).$$

The word "local" indicates that the group element $\sigma(x) \in G$ is allowed to depend on the point x in spacetime.

The set of all gauge transformations forms an infinite dimensional group

$$\mathcal{G} = C^\infty(M, G)$$

of smooth maps from spacetime M into the Lie group G . Gauge transformations thus result in an action of \mathcal{G} on the (infinite dimensional) vector space

$$C^\infty(M, S \otimes V)$$

of all possible fields.

2.2. Invariance. Every field theory has field equations, i.e. certain PDEs, that describe the evolution of the field in spacetime. The invariance in the case of gauge theories means that *the field equations are invariant under gauge transformations*: If Ψ is a solution to the field equations, then $\sigma \cdot \Psi$ should also be a solution of the equations for every local gauge transformation $\sigma: M \rightarrow G$.

Instead of invariance of the field equations one often demands that the **Lagrangian** \mathcal{L} of the field theory, from which the field equations are derived (as Euler-Lagrange equations), is invariant under gauge transformations. In this case the field equations will be gauge invariant as well: If the Lagrangian

\mathcal{L} is gauge invariant and a multiplet field Ψ is a critical point of the **action functional**

$$S[\Psi] = \int \mathcal{L}[\Psi] dx,$$

then $\sigma \cdot \Psi$ is a critical point for every gauge transformation $\sigma: M \rightarrow G$.

Invariance of the Lagrangian is stronger than invariance of the field equations, because the latter only means invariance of the set of critical points of the action. A gauge invariant Lagrangian is relevant in particular in quantum field theories where the full action (and not just its critical points) appears in path integrals.

If we demand gauge invariance of the field equations (or Lagrangian) we are almost inevitably led to the introduction of a gauge field A_μ that has values in the Lie algebra \mathfrak{g} of G . The reason is that in general a PDE will be invariant under gauge transformations only if the partial derivatives of the field in the directions of spacetime are **covariant derivatives**: If ∇_μ is the covariant derivative of the multiplet Ψ in the direction e_μ , we demand that

$$\nabla_\mu(\sigma \cdot \Psi) = \sigma \cdot (\nabla_\mu \Psi) \quad \forall \Psi \in \mathcal{C},$$

i.e.

$$\sigma \nabla_\mu \sigma^{-1} = \nabla_\mu$$

for all gauge transformations $\sigma: M \rightarrow G$ (here the inverse $\sigma^{-1}(x)$ means taking the inverse of the group element $\sigma(x) \in G$). The standard derivative ∂_μ does not have this behaviour because there is a second term due to the Leibniz rule (σ is a function on M). To a covariant derivative ∇_μ the gauge transformation σ looks like a constant.

The simplest way to construct a covariant derivative is to write

$$\nabla_\mu = \nabla_\mu^A = \partial_\mu + A_\mu,$$

where the **gauge field** A is a 1-form on M with values in the Lie algebra \mathfrak{g} . The set of all gauge fields is $\Omega^1(M, \mathfrak{g})$. If we insert a basis vector field e_μ on the spacetime manifold M into A , we get a field

$$A_\mu: M \longrightarrow \mathfrak{g}.$$

Every representation of a Lie group induces a representation of its Lie algebra, hence we get an action of A_μ on the multiplet space V and thus an action on the multiplets $\Psi: M \rightarrow S \otimes V$. The action of the covariant derivative on multiplets

$$(\nabla_\mu^A \Psi)(x) = (\partial_\mu \Psi)(x) + A_\mu(x) \cdot \Psi(x)$$

is given by the standard derivative of each component of Ψ in the point $x \in M$ plus the action of the Lie algebra element $A_\mu(x)$ on the multiplet $\Psi(x)$.

We need to find the correct behaviour of the gauge field under gauge transformations: This can be written down most easily in the case that G is a group

of matrices, so that we can differentiate in the standard way a gauge transformation in a spacetime direction (there is a formula involving the adjoint representation of the group G and the Maurer-Cartan form in the general case, see [Baum 2014, p. 79] and [Naber 2011, p. 303]).

Lemma 2.2. *If we demand that under a local gauge transformation σ we have*

$$A_\mu \mapsto A'_\mu = \sigma A_\mu \sigma^{-1} - (\partial_\mu \sigma) \sigma^{-1},$$

then we have

$$\nabla_\mu^{A'} = \sigma \nabla_\mu^A \sigma^{-1},$$

so that ∇_μ^A is a covariant derivative.

Proof. We calculate

$$\begin{aligned} \nabla_\mu^{A'}(\sigma \cdot \Psi) &= (\partial_\mu + A'_\mu)(\sigma \cdot \Psi) \\ &= (\partial_\mu \sigma) \cdot \Psi + \sigma \cdot (\partial_\mu \Psi) + \sigma A_\mu \sigma^{-1} \sigma \cdot \Psi - (\partial_\mu \sigma) \sigma^{-1} \sigma \cdot \Psi \\ &= \sigma \cdot \nabla_\mu^A \Psi. \end{aligned}$$

We see that the gauge field A_μ with its transformation behaviour exactly cancels the term from the Leibniz rule applied to ∂_μ . \square

To ensure gauge symmetry of the field equations for the matter multiplet Ψ we therefore have to introduce another field A_μ that compensates or absorbs with its behaviour under gauge transformations the transformation of the field Ψ . Only the combined transformations of the gauge field and the matter multiplet leave the field equations invariant.

Note that if the space of multiplets is $S \otimes V$, where S is a space of spinors, we can define a **twisted Dirac operator** that acts on the multiplets via

$$D^A \Psi = i\gamma^\mu \nabla_\mu^A \Psi = i\gamma^\mu (\partial_\mu + A_\mu) \Psi,$$

where the gamma matrices γ^μ act on all of the spinor components ψ_i separately (mathematically, the action of $i\gamma_\mu$ on a spinor is the Clifford multiplication of the basis vector e_μ with the spinor). This Dirac operator appears in the Lagrangian and the field equations for fermion multiplets Ψ .

It is sometimes useful to choose a basis $\alpha_1, \dots, \alpha_r$ of the Lie algebra \mathfrak{g} (here r is the dimension of G). Then we can write

$$A_\mu = \sum_{i=1}^r A_\mu^i \alpha_i.$$

The basis elements α_i are called **gauge symmetry generators** and the components $A_\mu^i : M \rightarrow \mathbb{R}$ are called **gauge boson fields**.

2.3. Interactions. Since the covariant derivative of the matter field Ψ involves the field A_μ , the field equations will now become a certain coupled system of PDEs for the matter multiplets Ψ and the gauge field A_μ . The existence of a coupled, non-linear system of equations means that there is an **interaction** between the matter fields and the gauge field.

In general, non-linear terms in field equations correspond to interactions (terms of higher than quadratic order in the Lagrangian). One can introduce direct interactions between particles of the same field ϕ by considering terms like ϕ^3, ϕ^4 , etc. in the Lagrangian. In gauge theories the interactions between particles of the field Ψ happen via the gauge field A_μ . We want to define the relevant Lagrangians in the case of a gauge theory.

We first choose a **Hermitian scalar product** on the complex vector space V such that the action of the gauge group G on V is **unitary** (this is always possible since G is compact). Without loss of generality we can assume that $V = \mathbb{C}^n$ and the scalar product is given by

$$\langle v, w \rangle = v^\dagger w,$$

where

$$v^\dagger = (v_1^*, \dots, v_n^*), \quad w = \begin{pmatrix} w_1 \\ \vdots \\ w_n \end{pmatrix}.$$

Definition 2.3. If the components ψ_i are scalar fields so that $S = \mathbb{C}$, then we define the **Klein-Gordon Lagrangian** by

$$\mathcal{L} = (\nabla^\mu \Psi)^\dagger (\nabla_\mu \Psi) - m^2 \Psi^\dagger \Psi,$$

where m is the mass of the field.

If the components ψ_i are spinor fields so that S is some complex spinor space, we first set

$$\bar{\psi}_i = \psi_i^\dagger \gamma^0$$

and

$$\bar{\Psi} = (\bar{\psi}_1, \dots, \bar{\psi}_n).$$

Note that $\bar{\psi}_i \psi_i$ and $\bar{\Psi} \Psi$ are Lorentz scalars.

Definition 2.4. The **Dirac Lagrangian** for a spinor field is

$$\mathcal{L} = \bar{\Psi} D^A \Psi - m \bar{\Psi} \Psi.$$

We also have to introduce another set of dynamic field equations for the gauge field A_μ itself (in the Lagrangian and hence the field equations for Ψ , the gauge field appears without derivatives). This involves the **field strength** (or **curvature**)

$$F_{\mu\nu} = F_{\mu\nu}^A = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

The field strength F is a 2-form with values in \mathfrak{g} , i.e. an element in $\Omega^2(M, \mathfrak{g})$. The field strength transforms under (matrix) gauge transformations σ as

$$F_{\mu\nu}^A \longmapsto F_{\mu\nu}^{A'} = \sigma F_{\mu\nu}^A \sigma^{-1},$$

(in general the transformation is given by the adjoint action of the Lie group G). In a basis $\alpha_1, \dots, \alpha_r$ for the Lie algebra we can write

$$F_{\mu\nu} = \sum_{i=1}^r F_{\mu\nu}^i \alpha_i.$$

Since G is compact, it has a bi-invariant Riemannian metric that corresponds to an Ad-invariant positive definite scalar product on the Lie algebra \mathfrak{g} .

Definition 2.5. Choosing an Ad-invariant positive definite scalar product k on \mathfrak{g} and an associated orthonormal basis α_i , then the **Yang-Mills Lagrangian** is given by

$$\mathcal{L} = -\frac{1}{4}k(F^{\mu\nu}, F_{\mu\nu}) = -\frac{1}{4}F_i^{\mu\nu}F_{\mu\nu}^i,$$

where summation over Lie algebra indices i is understood.

If the group G is non-abelian, there are non-linear terms in the field equations derived from the Yang-Mills Lagrangian. This means that there is a direct interaction between gauge bosons.

Remark 2.6. One can easily check that each of the three Lagrangians defined above is invariant under gauge transformations.

The total, gauge invariant Lagrangian for a matter field and a gauge field is the sum of the Klein-Gordon or Dirac Lagrangian and the Yang-Mills Lagrangian.

2.4. Moduli space. Mathematically we have in a gauge theory an action of the group \mathcal{G} of gauge transformations on the infinite dimensional space

$$\mathcal{C} = C^\infty(M, S \otimes V) \times \Omega^1(M, \mathfrak{g})$$

of pairs of multiplet fields and gauge fields (configurations):

$$\mathcal{G} \times \mathcal{C} \longrightarrow \mathcal{C}$$

$$\sigma \cdot (\Psi, A_\mu) = (\sigma \cdot \Psi, \sigma A_\mu \sigma^{-1} - (\partial_\mu \sigma) \sigma^{-1}).$$

Gauge invariance means that the action of \mathcal{G} preserves the space of solutions $\mathcal{S} \subset \mathcal{C}$ to the (coupled) field equations for Ψ and A_μ . The quotient space \mathcal{S}/\mathcal{G} could be called the **moduli space** of the field equations (see e.g. [Morgan 1996] for the case of the moduli space of the gauge invariant Seiberg-Witten equations).

2.5. **Summary.** The steps in the construction of a gauge theory are therefore:

- Collect the matter fields into a multiplet Ψ on which a compact Lie group G acts.
- Demand gauge invariance of the field equations under local gauge transformations.
- Introduce the gauge field A_μ to get a covariant derivative in order to formulate gauge invariant field equations.
- The field equations become a coupled, non-linear system of equations for Ψ and A_μ . This means that there is an interaction between the matter fields and the gauge fields and thus, via the gauge field, between matter fields. If the group G is non-abelian there is also a direct interaction between the gauge fields, because of the commutator in the definition of the field strength $F_{\mu\nu}^A$.

Example 2.7. As an example we consider **QCD**, the gauge theory that describes the interaction between **quarks**. Each **flavour** f of quarks (u, d, c, s, t, b) is described by a triplet q_f of complex 4-component Dirac spinors q_f^r, q_f^s, q_f^b on spacetime $M = \mathbb{R}^4$ of the form

$$q_f = \begin{pmatrix} q_f^r \\ q_f^s \\ q_f^b \end{pmatrix}.$$

The three components of q_f correspond to the three **colours** that every flavour of quarks can have. A triplet q_f is therefore an element of $S \otimes V$, where S denotes the vector space of 4-spinors and $V = \mathbb{C}^3$ is a 3-dimensional complex vector space. The group $SU(3)$ acts via the standard representation on V and thus on the triplets q_f . Since the group $SU(3)$ is non-abelian there is an interaction between the gauge bosons, that are called **gluons**.

3. COMPACT LIE GROUPS

We recall some facts about Lie groups and Lie algebras. For more details see, for example, [Ziller 2010].

3.1. **Ad-invariant scalar products on Lie algebras.** A Lie algebra \mathfrak{g} is called **simple** if the dimension of \mathfrak{g} is greater than 1 and \mathfrak{g} has no non-trivial ideals. A Lie algebra \mathfrak{g} is called **compact** if it is the Lie algebra of a compact Lie group G .

Theorem 3.1. *Let \mathfrak{g} be a compact Lie algebra. Then \mathfrak{g} is the direct sum of an abelian ideal and compact simple ideals.*

Since ideals in the Lie algebra correspond to normal subgroups, this means that up to a finite quotient any compact, connected Lie group G is of the form

$$G = U(1) \times \cdots \times U(1) \times G_1 \times \cdots \times G_s,$$

where G_1, \dots, G_s are compact simple Lie groups.

Theorem 3.2. *Every compact simple Lie algebra \mathfrak{g} has up to a positive constant a unique Ad-invariant positive definite scalar product.*

The existence follows because (by averaging) the associated compact Lie group G has a bi-invariant Riemannian metric. Uniqueness follows by Schur's Lemma, because the adjoint action of a simple Lie algebra is irreducible (an Ad-invariant subspace of \mathfrak{g} is an ideal). We get:

Corollary 3.3. *Let G be a compact, connected Lie group of the form*

$$G = U(1) \times \cdots \times U(1) \times G_1 \times \cdots \times G_s,$$

*(up to a finite quotient) and k an Ad_G -invariant positive definite scalar product on the Lie algebra \mathfrak{g} . Then k is the orthogonal direct sum of a positive definite scalar product k_0 on $u(1) \oplus \cdots \oplus u(1)$ and Ad_{G_i} -invariant positive definite scalar products k_i on the Lie algebras \mathfrak{g}_i . These scalar products (relative to some fixed scalar products) are determined by real constants, called **coupling constants** in physics.*

In gauge theory we have to choose an Ad-invariant scalar product on the Lie algebra \mathfrak{g} because only then the Yang-Mills Lagrangian in Definition 2.5 will be gauge invariant. The Yang-Mills Lagrangian thus involves (in our notation implicitly) the choice of coupling constants.

3.2. Conventions. We compare the conventions used in physics and mathematics for scalar products and coupling constants. We fix a compact Lie algebra \mathfrak{g} which is either simple or isomorphic to $u(1)$. Let k_0 be a fixed Ad-invariant positive definite inner product on \mathfrak{g} and β_1, \dots, β_r an associated orthonormal basis. Let g be the coupling constant.

(a) In **mathematics** we choose the scalar product

$$k = \frac{1}{g} k_0$$

with orthonormal basis

$$\alpha_i = g\beta_i, \quad i = 1, \dots, r.$$

We expand the gauge field A_μ with values in \mathfrak{g} as

$$A_\mu = \sum_{l=1}^r A_\mu^l \alpha_l.$$

The covariant derivative is

$$\nabla_\mu^A = \partial_\mu + A_\mu.$$

The curvature is

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu], \quad F_{\mu\nu} = \sum_{l=1}^r F_{\mu\nu}^l \alpha_l.$$

The Yang-Mills Lagrangian is

$$\mathcal{L} = -\frac{1}{4}k(F^{\mu\nu}, F_{\mu\nu}) = -\frac{1}{4}F_{\mu\nu}^l F_l^{\mu\nu}.$$

- (b) In **physics** we choose the scalar product k_0 and the orthonormal basis β_1, \dots, β_r . We expand the gauge field B_μ with values in $ig \subset \mathfrak{g} \otimes \mathbb{C}$ as

$$B_\mu = \sum_{l=1}^r B_\mu^l \beta_l.$$

There are two conventions for the covariant derivative:

$$\nabla_\mu^B = \partial_\mu \pm ig B_\mu.$$

The curvature is

$$G_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu \pm ig[B_\mu, B_\nu], \quad G_{\mu\nu} = \sum_{l=1}^r G_{\mu\nu}^l \beta_l.$$

The Yang-Mills Lagrangian is

$$\mathcal{L} = -\frac{1}{4}k'_0(G^{\mu\nu}, G_{\mu\nu}) = -\frac{1}{4}G_{\mu\nu}^l G_l^{\mu\nu}$$

where k'_0 denotes the Hermitian inner product on $\mathfrak{g} \otimes \mathbb{C}$ associated to k_0 .

- (c) The correspondence between both sides is given by mapping

$$A_\mu \rightarrow \pm ig B_\mu, \quad F_{\mu\nu} \rightarrow \pm ig G_{\mu\nu}.$$

If the representation of the Lie group G on the multiplet space V is unitary so that the action of an element of the Lie algebra \mathfrak{g} is skew-Hermitian, the field B_μ will then act as a Hermitian operator. We have $\nabla_\mu^A = \nabla_\mu^B$ and

$$A_\mu^l = \pm i B_\mu^l, \quad F_{\mu\nu}^l = \pm i G_{\mu\nu}^l.$$

4. MASSIVE VECTOR BOSONS AND CHIRAL FERMIONS

It is clearly observed in experiments that there are elementary particles in nature that have a mass. We have seen in Definitions 2.3 and 2.4 ways how to formulate a gauge invariant Lagrangian for massive particles. However, there are certain situations where a mass cannot be introduced in such a direct way into a gauge theory without spoiling gauge invariance. It is known that these situations, that we want to describe now, occur in nature. The Higgs field is a solution to this problem because it opens up the possibility of introducing a mass for elementary particles in an indirect and gauge invariant way.

4.1. Massive vector bosons. The Yang-Mills Lagrangian

$$\mathcal{L} = -\frac{1}{4}F_i^{\mu\nu}F_{\mu\nu}^i,$$

describes a massless gauge boson. It is observed in experiments that certain gauge bosons have a mass. To introduce a mass directly means to add a term of the form

$$\frac{1}{2}m^2 A_i^\mu A_\mu^i$$

to the Yang-Mills Lagrangian. However, this term is clearly not invariant under the gauge transformations in Lemma 2.2. We will later see in Theorem 7.8 that such a term can be introduced in a gauge invariant way because of the interaction between the gauge field and the Higgs field.

4.2. Chiral fermions. Recall (see e.g. [Morgan 1996]) that if the dimension of the manifold M is $2m$, then the dimension of the target space S of complex Dirac spinors on M is 2^m . Under the spin group (Lorentz transformations on spinors) this space splits into two irreducible representations S_L and S_R , called left-handed and right-handed Weyl spinors, of complex dimension 2^{m-1} . A Dirac multiplet

$$\Psi \in S \otimes V$$

thus decomposes into

$$\Psi = \begin{pmatrix} \Psi_L \\ \Psi_R \end{pmatrix} \in (S_L \otimes V) \oplus (S_R \otimes V).$$

We restrict to dimension $2m = 4$. We have

$$\bar{\Psi} = \Psi^\dagger \gamma^0 = (\bar{\Psi}_R, \bar{\Psi}_L) = (\Psi_R^\dagger, \Psi_L^\dagger)$$

and, since multiplication with $i\gamma^\mu$ (Clifford multiplication with a basis vector e_μ) turns a left-handed into a right-handed spinor and vice versa, the part

$$\bar{\Psi} D^A \Psi$$

of the Dirac Lagrangian splits as

$$\bar{\Psi}_L i\gamma^\mu (\partial_\mu + A_\mu) \Psi_L + \bar{\Psi}_R i\gamma^\mu (\partial_\mu + A_\mu) \Psi_R.$$

This term will stay gauge invariant even if the corresponding unitary representations of the gauge group G on V are chosen *independently* for left-handed and right-handed spinors. In fact, we can choose different multiplet spaces V_L and V_R that twist the left-handed and right-handed spinors, so that the multiplet field takes value in

$$(S_L \otimes V_L) \oplus (S_R \otimes V_R).$$

It is known from experiments that a realistic gauge theory of elementary particles must be **chiral** in this sense. The twisted Dirac operator D^A is thus a first order differential operator from

$$C^\infty(M, (S_L \otimes V_L) \oplus (S_R \otimes V_R))$$

into

$$C^\infty(M, (S_R \otimes V_L) \oplus (S_L \otimes V_R)).$$

In the case of vector bundles this is precisely the situation studied in the **Atiyah-Singer index theorem** for Dirac operators, which from a physics point of view is a theorem about chiral multiplets of fermions on even dimensional manifolds; for more details see [\[Berline et al. 2004\]](#).

The mass term $m\bar{\Psi}\Psi$ in the Dirac Lagrangian, however, is of the form

$$m (\bar{\Psi}_L \Psi_R + \bar{\Psi}_R \Psi_L) = 2m \operatorname{Re} (\bar{\Psi}_L \Psi_R).$$

This term clearly does not stay invariant if the left-handed and right-handed representations on V are not identical (the term as such is not defined if $V_L \neq V_R$). We consider a slightly more general situation:

Definition 4.1. Let V_R and V_L be unitary representations of a Lie group G . We define a **mass form** as a G -invariant form

$$\kappa: V_L \times V_R \longrightarrow \mathbb{C}$$

which is complex antilinear in the first argument and complex linear in the second.

We then have:

Theorem 4.2. Suppose that V_L and V_R are irreducible, unitary, non-isomorphic representations of G . Then every mass form is identically zero.

Proof. We define \bar{V}_L^* as

$$\bar{V}_L^* = \{\alpha: V_L \rightarrow \mathbb{C} \mid \alpha \text{ is } \mathbb{C}\text{-antilinear}\}.$$

This complex vector space has a G -representation defined by

$$(g \cdot \alpha)(v_L) = \alpha(g^{-1} \cdot v_L)$$

for $g \in G, v_L \in V_L$. The map

$$V_L \longrightarrow \overline{V}_L^*, \quad v_L \longmapsto \langle \cdot, v_L \rangle,$$

where $\langle \cdot, \cdot \rangle$ is the Hermitian form on V_L , defines a complex linear G -equivariant isomorphism.

Suppose a mass form $\kappa \neq 0$ exists. Then

$$V_R \longrightarrow \overline{V}_L^*, \quad v_R \longmapsto \kappa(\cdot, v_R)$$

is a complex linear G -equivariant map. In total we get a complex linear G -equivariant map

$$V_R \longrightarrow V_L$$

which is non-zero, because $\kappa \neq 0$. By Schur's Lemma this map has to be an isomorphism of the representations V_R and V_L (because the kernel and image of the map are G -invariant), contradicting our assumption. \square

We will see in Section 9 how a mass for fermions in a chiral gauge theory can be introduced in a gauge invariant way via Yukawa couplings of the fermions to the Higgs field, involving certain *trilinear* forms.

5. THE HIGGS FIELD

We want to consider gauge theories in which the **symmetry is spontaneously broken**, leading to the existence of one or several Higgs bosons. For example, the electroweak interaction in the Standard Model is a spontaneously broken gauge theory.

We assume that we have a gauge theory with a compact, connected Lie group G . To describe symmetry breaking we need a **Higgs field** $\Phi: M \rightarrow E$. Here $E = \mathbb{C}^n$ and we have a unitary representation of the gauge group G on E . The Higgs field is a scalar under Lorentz transformations (more precisely, a multiplet of n complex scalars under the action of the gauge group G). We can arrange that the action of the Lie group G on the space E is unitary with respect to the standard Hermitian scalar product $\langle v, w \rangle = v^\dagger w$. The real part $\text{Re}\langle \cdot, \cdot \rangle$ is a positive definite scalar product on E .

We also assume that there is a **potential**

$$V: E \longrightarrow \mathbb{R}$$

which is invariant under the action of G . The potential appears in the Lagrangian for the Higgs field Φ :

$$\mathcal{L} = \left(\nabla^{A\mu} \Phi \right)^\dagger \left(\nabla_\mu^A \Phi \right) - V(\Phi),$$

where

$$V(\Phi)(x) = V(\Phi(x))$$

for a spacetime point $x \in M$. This is a Lagrangian of Klein-Gordon type plus the potential which describes a certain self-interaction of the Higgs field.

5.1. Spontaneously broken gauge theories. In a gauge theory we demand that the Lagrangian is invariant under gauge transformations. Solutions to the field equations will then map to solutions under arbitrary gauge transformations $\sigma: M \rightarrow G$. *This is still true in a spontaneously broken gauge theory.*

Definition 5.1. A vector $v_0 \in E$ is called **vacuum vector** if it is a local minimum of the potential V . A Higgs field Φ is called **vacuum** if its value is constant and equal to v_0 on all of spacetime. The stabilizer (isotropy) subgroup $H \subset G$ of all elements $g \in G$ with $g \cdot v_0 = v_0$ is called the **unbroken subgroup** of G . A gauge theory containing a Higgs field is called **spontaneously broken** if $H \neq G$.

In a spontaneously broken gauge theory the Lagrangian and hence the field equations are invariant under the full gauge group G . However, the vacuum value of the Higgs field is only invariant under a smaller subgroup H . Note that the symmetry can only be broken if $v_0 \neq 0$. *The non-zero vacuum value of the Higgs field is precisely the reason why elementary particles have a mass.* We consider an example:

Example 5.2. In the case of the electroweak interaction we have $E = \mathbb{C}^2$ and $G = SU(2) \times U(1)$. The potential of the Higgs field is

$$V(v) = -\frac{1}{2}\mu v^\dagger v + \frac{1}{2}\lambda (v^\dagger v)^2$$

with certain constants $\mu, \lambda > 0$. A vector v_0 is a vacuum vector if and only if

$$||v_0|| = \sqrt{\frac{\mu}{2\lambda}}.$$

The set of vacuum vectors (minima of the potential V) thus forms a 3-sphere in \mathbb{C}^2 around the origin of radius $||v_0||$.

5.2. The Hessian of the Higgs potential. Let v_0 be a vacuum vector. We consider Higgs fields Φ whose values are in the vicinity of v_0 :

$$\Phi = v_0 + \Delta\Phi,$$

where $\Delta\Phi$ is the **shifted Higgs field**. We want to derive an approximation to $V(\Phi)$ for small values of $\Delta\Phi$ using the Taylor expansion of the potential V .

The vector space E (more precisely, the tangent space to E in the point v_0) can be split as

$$E = T_{v_0}E = W_{v_0} \oplus W_{v_0}^\perp,$$

where W_{v_0} is the tangent space to the **orbit** of the gauge group G through the vacuum vector v_0 :

$$W_{v_0} = T_{v_0}(G \cdot v_0).$$

Note that the orbit $G \cdot v_0$ (since G is compact) is diffeomorphic to the quotient space G/H , where H is the unbroken subgroup.

Definition 5.3. We set d for the dimension of W_{v_0} . This is equal to the codimension of H in G .

Let $\text{Hess}(V)$ denote the **Hessian** of the potential V . The Hessian is a symmetric linear map

$$\text{Hess}(V): T_v E \longrightarrow T_v E$$

on the tangent space to E in a point v and can be defined on any Riemannian manifold as

$$\text{Hess}(V)(X) = \nabla_X \text{grad } V,$$

where in our situation X is a vector tangent to E and ∇ denotes the (flat) Levi-Civita connection induced by the positive definite inner product on E . This linear map is symmetric in the sense that

$$\langle \text{Hess}(V)(X), Y \rangle = \langle X, \text{Hess}(V)Y \rangle.$$

It corresponds to the matrix of second derivatives

$$\text{Hess}(V) = \left(\frac{\partial^2 V(x)}{\partial x_i \partial x_j} \right)$$

in standard coordinates x on the vector space E .

We want to diagonalize this symmetric linear map in the vacuum point v_0 . Since V has a minimum along the whole orbit of G through v_0 (as the potential is invariant under the action of G) it follows that $\text{grad } V$ vanishes along the orbit and hence

$$\text{Hess}(V)_{v_0}(X) = 0$$

for vectors $X \in W_{v_0}$ tangent to the orbit. This implies in particular that W_{v_0} and $W_{v_0}^\perp$ are invariant subspaces of the Hessian.

Lemma 5.4. *We can thus find (real) orthonormal bases consisting of eigenvectors*

$$e_1, \dots, e_d \in W_{v_0}$$

$$f_1, \dots, f_{2n-d} \in W_{v_0}^\perp$$

of the Hessian in v_0 where the e_i have eigenvalue 0 and the f_j have non-negative eigenvalues (because v_0 is a local minimum). We set $2m_{f_j}^2$ for the eigenvalue of f_j (with $m_{f_j} \geq 0$).

5.3. The Goldstone and Higgs bosons.

Definition 5.5. We expand the shifted Higgs field $\Delta\Phi$ in the orthonormal eigenbasis of the Hessian:

$$\Delta\Phi = \frac{1}{\sqrt{2}} \sum_{i=1}^d \xi_i e_i + \frac{1}{\sqrt{2}} \sum_{j=1}^{2n-d} \eta_j f_j,$$

where ξ_i and η_j are real scalar fields on the spacetime manifold M . The ξ_i are called **Goldstone bosons**, the η_j are called **Higgs bosons**. The number of Goldstone bosons is equal to the dimension of G/H . The number of Higgs bosons is equal to the real dimension of the multiplet space E minus the dimension of G/H . The Goldstone bosons correspond to perturbations of the Higgs field Φ along the orbit of G through the vacuum vector v_0 , while the Higgs bosons correspond to perturbations orthogonally to the orbit. As particles in the associated quantum field theory the Goldstone and Higgs bosons are thus minimal excitations of the Higgs field from its vacuum state.

Theorem 5.6. *Up to second order in the shifted Higgs field we have by the Taylor formula (the first order term vanishes, because v_0 is a critical point)*

$$V(\Phi) \approx V(v_0) + \frac{1}{2} \sum_{j=1}^{2n-d} m_{f_j}^2 \eta_j^2.$$

This is (up to the irrelevant constant $V(v_0)$) the standard form of a the Klein-Gordon mass term for real scalar fields η_j (see Definition 2.3). The **Goldstone bosons have zero mass**, while the **Higgs bosons have mass** $m_{f_j} \geq 0$. The Lagrangian \mathcal{L} for the Higgs field Φ expressed in terms of the shifted Higgs field then contains Klein-Gordon summands of the form

$$\frac{1}{2} \left((\partial^\mu \eta_j)(\partial_\mu \eta_j) - m_{f_j}^2 \eta_j^2 \right),$$

plus other terms, some of which we will determine later.

Example 5.7. In the case of electroweak theory we have $G = SU(2) \times U(1)$ and $H = U(1)$, embedded in G as a diagonal subgroup, not as the second factor (we will define the precise representation of G on the vector space E later in Section 8 and determine the subgroup H). Therefore we have three Goldstone bosons and one Higgs boson, because E has complex dimension 2.

One can check that the **mass of the Higgs boson** is $\sqrt{\mu}$: We take

$$v_0 = \begin{pmatrix} 0 \\ \sqrt{\frac{\mu}{2\lambda}} \end{pmatrix}.$$

Then W_{v_0} is the real span of

$$\begin{pmatrix} 1 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} i \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 0 \\ i \end{pmatrix}$$

and $W_{v_0}^\perp$ is the real span of

$$\begin{pmatrix} 0 \\ 1 \end{pmatrix}.$$

In the standard coordinates $x_1 + ix_2, x_3 + ix_4$ for \mathbb{C}^2 the only non-vanishing entry of the Hessian $\text{Hess}(V)_{v_0}$ at v_0 is the 33-component, which is 2μ . This implies the claim.

6. UNITARY GAUGE

In this section we want to discuss the notion of unitary gauge in the mechanism of symmetry breaking. We use the same notation as before: We consider a Higgs field $\Phi: M \rightarrow E$ on the spacetime manifold M with values in a vector space $E = \mathbb{C}^n$. On the vector space E we have a representation of the compact gauge group G , unitary with respect to the standard Hermitian scalar product $\langle \cdot, \cdot \rangle$. We choose a vacuum vector $v_0 \in E$.

We then saw that we can write $\Phi = v_0 + \Delta\Phi$, where the shifted Higgs field $\Delta\Phi$ can be decomposed as

$$\Delta\Phi = \frac{1}{\sqrt{2}} \sum_{i=1}^d \xi_i e_i + \frac{1}{\sqrt{2}} \sum_{j=1}^{2n-d} \eta_j f_j,$$

with real valued scalar fields ξ_i and η_j on spacetime M , called Goldstone bosons and Higgs bosons. Here the vectors e_i are tangential to the orbit of the gauge group G through v_0 and the vectors f_j are orthogonal to the orbit.

Note that the orbit through v_0 is orthogonal to the vector v_0 because of our assumption that the action of the gauge group G is unitary. Therefore the space spanned by the vectors f_j will contain a direction parallel to v_0 . In the case of the electroweak theory with $G = SU(2) \times U(1)$ this is actually the only orthogonal direction. Hence in this situation there is only one Higgs boson.

The scalar fields ξ_i and η_j depend both on the scalar field Φ and the vacuum vector v_0 . We can therefore say that for a given Higgs field Φ and a given vacuum vector v_0 we have certain Goldstone bosons ξ_i and Higgs bosons η_j . We always consider v_0 and the vectors e_i and f_j fixed.

6.1. Unitary gauge and the Goldstone bosons. The intuition behind the Goldstone bosons is that they do not correspond to physical fields (or particles), because they can be *gauged away* (the Higgs bosons, on the contrary, cannot be gauged away). We set:

Definition 6.1. We call a smooth gauge transformation $\sigma: M \rightarrow G$ **unitary gauge** for a given Higgs field Φ with respect to a vacuum vector v_0 if all Goldstone bosons of the transformed field $\sigma \cdot \Phi$ with respect to the vacuum vector v_0 vanish identically on M . We say that a Higgs field is in unitary gauge if all its Goldstone bosons vanish identically.

We want to find an equivalent condition for a gauge to be unitary. Since the Lie group G acts on E there is an induced action of the Lie algebra \mathfrak{g} on E . We write $A \cdot v$ for $A \in \mathfrak{g}$ and $v \in E$. Since the action of G is through unitary isomorphisms, the action of \mathfrak{g} is through skew-Hermitian endomorphisms, i.e.

$$\langle A \cdot v, w \rangle + \langle v, A \cdot w \rangle = 0 \quad \forall A \in \mathfrak{g} \quad \forall v, w \in E.$$

Definition 6.2. For a point $x \in M$ we consider the linear map

$$s_x^\Phi: \mathfrak{g} \longrightarrow \mathbb{R}, \quad A \longmapsto \operatorname{Re} \langle \Phi(x), A \cdot v_0 \rangle.$$

Bleecker calls this map **fiber derivative** (see [Bleecker 1981]).

We can then show:

Lemma 6.3. *Let Φ be a given Higgs field and v_0 a given vacuum vector. Then all Goldstone bosons of Φ with respect to v_0 vanish in the point $x \in M$ if and only if $s_x^\Phi \equiv 0$.*

Proof. Since $A \in \mathfrak{g}$ acts through skew-Hermitian endomorphisms, we have

$$\operatorname{Re} \langle \Phi(x), A \cdot v_0 \rangle = \operatorname{Re} \langle \Delta \Phi(x), A \cdot v_0 \rangle.$$

Note that $\mathfrak{g} \cdot v_0 = W_{v_0}$, the tangent space to the orbit. Hence

$$\operatorname{Re} \langle \Delta \Phi(x), A \cdot v_0 \rangle = \frac{1}{\sqrt{2}} \sum_{i=1}^d \operatorname{Re} \langle \xi_i(x) e_i, A \cdot v_0 \rangle.$$

The claim follows because the vectors e_i are also elements of W_{v_0} . □

Definition 6.4. The **unitary set** of a Higgs field Φ with respect to the vacuum vector v_0 is defined as

$$U(\Phi, v_0) = \{(x, g) \in M \times G \mid \operatorname{Re} \langle g \cdot \Phi(x), A \cdot v_0 \rangle = 0 \quad \forall A \in \mathfrak{g}\}.$$

We then get:

Theorem 6.5. *A smooth gauge transformation $\sigma: M \rightarrow G$ is a unitary gauge for Φ with respect to v_0 if and only if the map*

$$M \longrightarrow M \times G, \quad x \longmapsto (x, \sigma(x))$$

has image in the unitary set $U(\Phi, v_0)$.

To find a unitary gauge σ we thus have to rotate the vectors $\Phi(x) \in E$ for each $x \in M$ through the action of the gauge group in such a way that the shifted vectors $\Delta(\sigma(x) \cdot \Phi(x))$ of the transformed field are perpendicular to the orbit of G through v_0 .

Example 6.6. In the case of electroweak theory we have $E = \mathbb{C}^2$ and $G = SU(2) \times U(1)$. The vacuum vector is some non-zero $v_0 \in \mathbb{C}^2$. Its orbit under the gauge group G is the 3-sphere of radius $\|v_0\|$ around the origin. The only direction perpendicular to the orbit through v_0 is the direction of v_0 itself. If Φ is a given Higgs field, a unitary gauge σ therefore has to ensure that all transformed field values $\sigma(x) \cdot \Phi(x)$, as x ranges over spacetime M , are multiples of the vector v_0 .

More generally we have:

Corollary 6.7. *A gauge transformation $\sigma: M \rightarrow G$ is a unitary gauge for a Higgs field Φ with respect to a vacuum vector v_0 if and only if all transformed field values $\sigma(x) \cdot \Phi(x)$, for $x \in M$, lie in the vector subspace of E that contains the vector v_0 and which is the orthogonal complement to the tangent space of the orbit under G through v_0 .*

6.2. Existence of unitary gauges. The existence problem for unitary gauges seems to be non-trivial even in the local case. Bleecker has a criterion when at least locally unitary gauges exist. Recall that H is the so-called unbroken subgroup of G fixing the vacuum vector v_0 .

Definition 6.8. Let \mathfrak{h}^\perp denote some orthogonal complement of the Lie algebra \mathfrak{h} of H inside \mathfrak{g} . For a point $(x, g) \in U(\Phi, v_0)$ we set

$$B_{(x,g)}^\Phi: \mathfrak{h}^\perp \times \mathfrak{h}^\perp \longrightarrow \mathbb{R}, \quad (A, B) \longmapsto \operatorname{Re}\langle g \cdot \Phi(x), A \cdot B \cdot v_0 \rangle.$$

Bleecker calls this map the **broken Hessian**. It is a bilinear, symmetric form because

$$\operatorname{Re}\langle g \cdot \Phi(x), [A, B] \cdot v_0 \rangle = 0$$

for $(x, g) \in U(\Phi, v_0)$.

Bleecker then proves the following theorem:

Theorem 6.9. *If $(x, g) \in U(\Phi, v_0)$ is a point such that $B_{(x,g)}^\Phi$ is non-degenerate, then there exists a unitary gauge $\sigma: W \rightarrow G$ in a neighbourhood $W \subset M$ of x such that $\sigma(x) = g$.*

The proof uses the regular value theorem.

7. MASS GENERATION FOR GAUGE BOSONS

The purpose of this section is to describe how masses of gauge bosons are generated through the Higgs field.

7.1. Broken und unbroken gauge bosons. We assume that the gauge group G is a compact, connected Lie group of dimension r . The Lie algebra \mathfrak{g} has an Ad-invariant positive definite scalar product k , determined by the coupling constants.

Let $H \subset G$ denote the stabilizer subgroup of the vacuum vector $v_0 \in E$. We denote by $\mathfrak{h} \subset \mathfrak{g}$ the Lie algebra of H and by \mathfrak{h}^\perp its orthogonal complement with respect to the scalar product k . We denote the dimension of \mathfrak{h}^\perp as before by d .

Definition 7.1. We define a positive semidefinite, bilinear symmetric form m on \mathfrak{g} by

$$m: \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathbb{R}, \quad (A, B) \longmapsto \operatorname{Re} \langle A \cdot v_0, B \cdot v_0 \rangle.$$

Here $\langle \cdot, \cdot \rangle$ denotes the Hermitian inner product on the vector space E .

Note that $A \cdot v_0 = 0$ for all $A \in \mathfrak{h}$, while the map $A \mapsto A \cdot v_0$ is injective on the orthogonal complement \mathfrak{h}^\perp . Hence the restriction of m onto the Lie algebra \mathfrak{h} vanishes identically and the restriction onto the orthogonal complement \mathfrak{h}^\perp is positive definite. Since m is a symmetric form we can diagonalize it and get:

Proposition 7.2. *We can find orthonormal bases (with respect to k) $\alpha_1, \dots, \alpha_d$ of the subspace \mathfrak{h}^\perp (called **broken generators**) and $\alpha_{d+1}, \dots, \alpha_r$ of the subspace \mathfrak{h} (called **unbroken generators**) such that the symmetric form m is diagonal in this basis. We have*

$$m(\alpha_i, \alpha_i) = \frac{1}{2} M_i^2 \geq 0,$$

where $M_i > 0$ for the broken generators and $M_i = 0$ for the unbroken generators.

If $A_\mu: M \rightarrow \mathfrak{g}$ is a gauge field, then we can write

$$A_\mu = \sum_{i=1}^r A_\mu^i \alpha_i,$$

where A_μ^i are the **broken and unbroken gauge bosons**, respectively. The numbers M_i are the **masses of the gauge bosons**. The masses M_i are proportional to the norm $\|v_0\|$ of the vacuum value of the Higgs field. The masses M_i also depend on the choice of the scalar product on \mathfrak{g} via the choice of orthonormal basis $\{\alpha_i\}$ and thus on the coupling constants.

7.2. The combined Lagrangian. For a gauge field A_μ we define as before the covariant derivative

$$\nabla_\mu^A = \partial_\mu + A_\mu$$

and the curvature (or field strength)

$$F_{\mu\nu} = dA_{\mu\nu} + [A_\mu, A_\nu] = \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu].$$

We can also express the field strength in our chosen orthonormal basis for the Lie algebra \mathfrak{g} :

$$F_{\mu\nu} = \sum_{i=1}^r F_{\mu\nu}^i \alpha_i.$$

The total Lagrangian for the Higgs field Φ and the gauge field A_μ is now:

$$\mathcal{L} = \left(\nabla^{A_\mu} \Phi \right)^\dagger \left(\nabla_\mu^A \Phi \right) - V(\Phi) - \frac{1}{4} F_i^{\mu\nu} F_{\mu\nu}^i.$$

Here we assumed the standard inner product $\langle v, w \rangle = v^\dagger w$ on $E = \mathbb{C}^n$. There is a summation over Lie algebra indices i in the last term.

We write $\Phi = v_0 + \Delta\Phi$ with the shifted Higgs field $\Delta\Phi$ as before. We want to determine the terms up to second order in the fields $\Delta\Phi$ and A_μ in the Lagrangian \mathcal{L} , i.e. the terms corresponding to "free" fields. Higher order terms contain interactions between these fields. A direct calculation shows:

Theorem 7.3. *Up to terms of second order the Lagrangian \mathcal{L} is given by*

$$\begin{aligned} \mathcal{L} \approx & (\partial^\mu \Delta\Phi)^\dagger (\partial_\mu \Delta\Phi) + 2\text{Re} (\partial^\mu \Delta\Phi)^\dagger (A_\mu \cdot v_0) + (A^\mu \cdot v_0)^\dagger (A_\mu \cdot v_0) \\ & - V(\Phi) - \frac{1}{4} (dA_i^{\mu\nu}) (dA_{\mu\nu}^i). \end{aligned}$$

7.3. Simplifying the Lagrangian. We want to simplify each of the summands. *It is very useful now to assume that the Higgs field Φ is given in unitary gauge.* By this we mean that all Goldstone bosons of Φ vanish on all of space-time M identically. We can always do a transformation of the Higgs field into unitary gauge (if such a gauge exists) because the Lagrangian \mathcal{L} is gauge invariant.

Lemma 7.4. *If Φ is in unitary gauge, then*

$$\text{Re} (\partial^\mu \Delta\Phi)^\dagger (A_\mu \cdot v_0) = 0.$$

Proof. Note that $A_\mu \cdot v_0$ is tangential to the orbit of v_0 under G , while the shifted Higgs field $\Delta\Phi$ and therefore its derivative is everywhere on M orthogonal to the orbit. This implies the claim. \square

Lemma 7.5. *If Φ is in unitary gauge, then*

$$(\partial^\mu \Delta\Phi)^\dagger (\partial_\mu \Delta\Phi) = \frac{1}{2} \sum_{j=1}^{2n-d} (\partial^\mu \eta_j) (\partial_\mu \eta_j),$$

where η_j are the Higgs bosons.

Proof. Follows immediately, because under our assumption that the Goldstone bosons vanish we have

$$\Delta\Phi = \frac{1}{\sqrt{2}} \sum_{j=1}^{2n-d} \eta_j f_j.$$

□

Lemma 7.6. *Up to second order in the Higgs bosons we have*

$$V(\Phi) \approx V(v_0) + \frac{1}{2} \sum_{j=1}^{2n-d} m_{f_j}^2 \eta_j^2,$$

where m_{f_j} is the mass of the j -th Higgs boson.

Proof. The formula can be found in Theorem 5.6

□

Lemma 7.7. *We have*

$$(A^\mu \cdot v_0)^\dagger (A_\mu \cdot v_0) = \frac{1}{2} \sum_{i=1}^d M_i^2 A_i^\mu A_\mu^i,$$

where M_i are the masses defined above.

Proof. We have

$$(A^\mu \cdot v_0)^\dagger (A_\mu \cdot v_0) = m(A^\mu, A_\mu)$$

for the bilinear form m on \mathfrak{g} introduced in the beginning. This implies the claim by our choice of basis α_i . □

We now collect all terms and get:

Theorem 7.8. *If the Higgs field Φ is in unitary gauge, then up to second order in the shifted Higgs field and the gauge field the Lagrangian is given by*

$$\begin{aligned} \mathcal{L} \approx & \frac{1}{2} \sum_{j=1}^{2n-d} (\partial^\mu \eta_j) (\partial_\mu \eta_j) - \frac{1}{2} \sum_{j=1}^{2n-d} m_{f_j}^2 \eta_j^2 \\ & - \frac{1}{4} \sum_{i=1}^d (dA_i^{\mu\nu}) (dA_{\mu\nu}^i) + \frac{1}{2} \sum_{i=1}^d M_i^2 A_i^\mu A_\mu^i \\ & - \frac{1}{4} \sum_{i=d+1}^r (dA_i^{\mu\nu}) (dA_{\mu\nu}^i). \end{aligned}$$

Here we removed the irrelevant constant $V(v_0)$. This Lagrangian has the following interpretation:

- The two terms in the first line are the Klein-Gordon Lagrangian for $2n - d$ real scalar Higgs bosons η_j of mass m_{f_j} .

- The two terms in the second line are the Lagrangian for d broken gauge bosons A_μ^1, \dots, A_μ^d of mass M_i .
- The term in the third line is the Lagrangian for $r - d$ unbroken massless gauge bosons $A_\mu^{d+1}, \dots, A_\mu^r$.

This is the **Brout-Englert-Higgs mechanism** of creating in a gauge invariant way masses for gauge bosons.

8. THE $SU(2) \times U(1)$ -THEORY OF ELECTROWEAK INTERACTIONS

In this section we want to discuss the Higgs mechanism in the special case of the **electroweak interaction**. Our gauge group is $G = SU(2) \times U(1)$. We choose the scalar product on the Lie algebra $\mathfrak{g} = \mathfrak{su}(2) \oplus \mathfrak{u}(1)$ in such a way that the following elements form an orthonormal basis:

$$\begin{aligned}\beta_l &= g \frac{i\sigma_l}{2} \in \mathfrak{su}(2) \quad (l = 1, 2, 3) \\ \beta_4 &= g' \frac{i}{2} \in \mathfrak{u}(1),\end{aligned}$$

where σ_l are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

and the positive real numbers g and g' are the coupling constants corresponding to $SU(2)$ and $U(1)$.

The vector space E in which the Higgs field takes value is $E = \mathbb{C}^2$. We have an action of G on E where the generators $\beta_i \in \mathfrak{g}$ act on a vector

$$v = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \in \mathbb{C}^2$$

as

$$\begin{aligned}\beta_l \cdot v &= g \frac{i\sigma_l}{2} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \quad (l = 1, 2, 3) \\ \beta_4 \cdot v &= g' \frac{i}{2} \begin{pmatrix} v_1 \\ v_2 \end{pmatrix}.\end{aligned}$$

Hence the action of $SU(2)$ on E is the standard 2-dimensional representation. The vacuum vector is given by

$$v_0 = \begin{pmatrix} 0 \\ \sqrt{\frac{\mu}{2\lambda}} \end{pmatrix} = \begin{pmatrix} 0 \\ ||v_0|| \end{pmatrix} \in \mathbb{C}^2,$$

where μ and λ are the parameters of the Higgs field potential that can be found in Example 5.2.

8.1. **The gauge bosons.** A direct calculation shows that the symmetric, bilinear form

$$m: \mathfrak{g} \times \mathfrak{g} \longrightarrow \mathbb{R}$$

(see Definition 7.1) is given in the basis β_i by

$$m(\beta_i, \beta_j) = \frac{||v_0||^2}{4} \begin{pmatrix} g^2 & 0 & 0 & 0 \\ 0 & g^2 & 0 & 0 \\ 0 & 0 & g^2 & -gg' \\ 0 & 0 & -gg' & g'^2 \end{pmatrix}.$$

If we define a new orthonormal basis

$$\alpha_1 = \beta_1$$

$$\alpha_2 = \beta_2$$

$$\alpha_3 = \frac{1}{\sqrt{g^2 + g'^2}}(g\beta_3 - g'\beta_4)$$

$$\alpha_4 = \frac{1}{\sqrt{g^2 + g'^2}}(g'\beta_3 + g\beta_4),$$

then the bilinear form m becomes diagonal:

$$m(\alpha_i, \alpha_j) = \frac{||v_0||^2}{4} \begin{pmatrix} g^2 & 0 & 0 & 0 \\ 0 & g^2 & 0 & 0 \\ 0 & 0 & g^2 + g'^2 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

We see that the subalgebra of the **stabilizer group** is given by

$$\mathfrak{h} = \text{span}(\alpha_4).$$

Indeed, α_4 acts on the vacuum vector as

$$\alpha_4 \cdot v_0 = \frac{1}{\sqrt{g^2 + g'^2}} \left(g'g \frac{i\sigma_3}{2} + gg' \frac{i}{2} \right) \begin{pmatrix} 0 \\ ||v_0|| \end{pmatrix} = 0.$$

The subspace of **broken generators** is given by

$$\mathfrak{h}^\perp = \text{span}(\alpha_1, \alpha_2, \alpha_3).$$

We have

- **three massive gauge bosons**, two gauge bosons of mass $\frac{1}{\sqrt{2}}||v_0||g$ and one gauge boson of mass $\frac{1}{\sqrt{2}}||v_0||\sqrt{g^2 + g'^2}$.
- We also have **one massless gauge boson**.

8.2. The physics notation. In physics the following notation is used: We set

$$\tan \theta_W = \frac{g'}{g},$$

where θ_W is the **Weinberg angle**. Then

$$\alpha_3 = \cos \theta_W \beta_3 - \sin \theta_W \beta_4$$

$$\alpha_4 = \sin \theta_W \beta_3 + \cos \theta_W \beta_4.$$

Hence the basis (α_3, α_4) is rotated by an angle θ_W with respect to (β_3, β_4) (clockwise in our situation). We can then decompose our gauge field

$$A_\mu = \sum_{i=1}^4 A_\mu^i \beta_i$$

as

$$A_\mu = W_\mu^+ \frac{1}{\sqrt{2}}(\alpha_1 - i\alpha_2) + W_\mu^- \frac{1}{\sqrt{2}}(\alpha_1 + i\alpha_2) + Z_\mu^0 \alpha_3 + \gamma_\mu \alpha_4,$$

where

- $W_\mu^\pm = \frac{1}{\sqrt{2}}(A_\mu^1 \pm iA_\mu^2)$ are gauge bosons of mass

$$m_W = \frac{1}{\sqrt{2}} \|v_0\| g$$

- $Z_\mu^0 = \cos \theta_W A_\mu^3 - \sin \theta_W A_\mu^4$ is a gauge boson of mass

$$m_Z = \frac{1}{\sqrt{2}} \|v_0\| \sqrt{g^2 + g'^2}$$

- $\gamma_\mu = \sin \theta_W A_\mu^3 + \cos \theta_W A_\mu^4$ is the massless photon.

We have

$$\cos \theta_W = \frac{m_W}{m_Z}.$$

We already calculated the mass of the Higgs boson in Example 5.7.

8.3. Charges.

Definition 8.1. Let Ψ be some multiplet of matter fields in the electroweak theory, taking value in a space $S \otimes V$, where the complex vector space V of dimension r is the multiplet space and we have some unitary representation of the gauge group $G = SU(2) \times U(1)$ on V . The generators β_l of $su(2)$ act as

$$\beta_l \rightarrow igT_l \quad (l = 1, 2, 3)$$

and the generator β_4 of $u(1)$ acts as

$$\beta_4 \rightarrow ig' \frac{Y}{2},$$

where T_l and Y are certain *Hermitian* operators on V . The eigenvalues of T_3 are called **weak isospin** and of Y **weak hypercharge**. Since T_3 and Y commute in any representation of $SU(2) \times U(1)$ we can find an orthonormal basis of V of common eigenvectors for both operators. If we identify V with \mathbb{C}^r via this basis, then both T_3 and Y act as diagonal matrices whose entries are the charges of the corresponding multiplet component fields ψ_i .

Remark 8.2. The eigenvalues of the weak isospin operator T_3 and thus the charges are determined by the **weights** of the representation. The weights are elements in the dual space of the **Cartan subalgebra** in $su(2) \otimes \mathbb{C}$, spanned by $i\sigma_3$. See [Ziller 2010] for more details.

If we set

$$\begin{aligned} T_+ &= (T_1 - iT_2) \\ T_- &= (T_1 + iT_2) \\ Q &= T_3 + \frac{Y}{2}, \end{aligned}$$

then a general gauge field $A_\mu: M \rightarrow \mathfrak{g}$ acts on the multiplets with value in $S \otimes V$ as

$$\begin{aligned} A_\mu \rightarrow & \frac{ig}{\sqrt{2}}(W_\mu^+ T_+ + W_\mu^- T_-) \\ & + Z_\mu^0 \frac{ig^2 T_3 - ig'^2 \frac{Y}{2}}{\sqrt{g^2 + g'^2}} \\ & + \gamma_\mu \frac{igg'}{\sqrt{g^2 + g'^2}} Q. \end{aligned}$$

It follows that the **elementary electric charge** is given by

$$e = \frac{gg'}{\sqrt{g^2 + g'^2}} = g \sin \theta_W.$$

For example, on the Higgs field

$$\Phi = \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix}$$

with values in E the charge operators act as

$$\begin{aligned} T_3 &= \begin{pmatrix} \frac{1}{2} & 0 \\ 0 & -\frac{1}{2} \end{pmatrix} \\ Y &= \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ Q &= \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}. \end{aligned}$$

We can directly read off the charges.

9. YUKAWA COUPLINGS

Using Yukawa couplings we can generate masses for chiral fermions via the Higgs field in a gauge invariant way. We consider two definitions:

Definition 9.1. Suppose that V_A, V_B, V_C are unitary representation spaces of a compact Lie group G . Then we define a **multiplet triple product** as a \mathbb{C} -linear map

$$\tau: V_A \otimes V_B \otimes V_C \longrightarrow \mathbb{C}$$

which is invariant under the action of G . Suppose similarly that S_A, S_B, S_C are representations of the Lorentz spin group (for example, scalars or spinors). Then we define a **Lorentz triple product** as a \mathbb{C} -linear map

$$\mu: S_A \otimes S_B \otimes S_C \longrightarrow \mathbb{C}$$

which is invariant under Lorentz transformations, i.e. a scalar.

We then have:

Proposition 9.2. Suppose τ is a multiplet triple product and μ is a Lorentz triple product. Then

$$\mathcal{L} = g_Y \text{Re}(\mu \otimes \tau): (S_A \otimes V_A) \otimes (S_B \otimes V_B) \otimes (S_C \otimes V_C) \longrightarrow \mathbb{R}$$

is for every constant $g_Y \in \mathbb{R}$ a G -invariant scalar. It defines a gauge invariant, cubic Lagrangian called **Yukawa coupling** (the constant g_Y is also called Yukawa coupling).

As a non-quadratic term the Yukawa Lagrangian describes an interaction between certain fields. We want to consider this notion in the case of the electroweak interaction of leptons.

9.1. Action of the gauge group. Let

$$\begin{aligned} V_A &= V_B = \mathbb{C}^2 \\ V_C &= \mathbb{C} \end{aligned}$$

and $G = SU(2) \times U(1)$. We consider the following representations of G on these three complex vector spaces, defined on the generators β_i by:

$$\beta_l \cdot (v_A, v_B, v_C) = \left(g \frac{i\sigma_l}{2} v_A, g \frac{i\sigma_l}{2} v_B, 0 \right)$$

for $l = 1, 2, 3$, and

$$\beta_4 \cdot (v_A, v_B, v_C) = \left(-g' \frac{i}{2} v_A, +g' \frac{i}{2} v_B, -g' i v_C \right).$$

Here g and g' are the coupling constants as before. The action of $SU(2)$ on V_A and V_B is the standard representation and on V_C the trivial representation.

9.2. Triple products. We define \bar{V}_A as the complex conjugate vector space of V_A . It consists of the elements

$$\bar{v}_A = v_A^\dagger$$

for $v_A \in V_A$ and has complex structure defined by

$$i\bar{v}_A = (-iv_A)^\dagger.$$

We have an induced action of G on \bar{V}_A .

Proposition 9.3. *The linear map*

$$\begin{aligned} \tau: \bar{V}_A \otimes V_B \otimes V_C &\longrightarrow \mathbb{C} \\ (\bar{v}_A, v_B, v_C) &\longmapsto \bar{v}_A v_B v_C \end{aligned}$$

is invariant under the action of G and defines a multiplet triple product.

Proof. It is easy to check that

$$\tau(\beta_i \cdot \bar{v}_A, v_B, v_C) + \tau(\bar{v}_A, \beta_i \cdot v_B, v_C) + \tau(\bar{v}_A, v_B, \beta_i \cdot v_C) \equiv 0$$

for all $i = 1, 2, 3, 4$. This implies the claim. \square

We let

$$S_A = S_L \quad (\text{the space of left-handed Weyl spinors})$$

$$S_B = \mathbb{C}$$

$$S_C = S_R \quad (\text{the space of right-handed Weyl spinors}).$$

We define \bar{S}_A as the conjugate space of left-handed Weyl spinors. It consists of the spinors $\bar{\psi}_L = \psi_L^\dagger$ for $\psi_L \in S_A$ and has complex structure defined by

$$i\bar{\psi}_L = \overline{(-i\psi_L)}.$$

Proposition 9.4. *The linear map*

$$\begin{aligned} \mu: \bar{S}_A \otimes S_B \otimes S_C &\longrightarrow \mathbb{C} \\ (\bar{\psi}_A, \phi_B, \psi_C) &\longmapsto \phi_B \bar{\psi}_A \psi_C \end{aligned}$$

defines a Lorentz triple product.

Proof. The proof follows because $\bar{\psi}_A \psi_C$ is a Lorentz scalar. \square

9.3. Yukawa coupling in the electroweak theory. We set

$$E_L = \begin{pmatrix} E_L^1 \\ E_L^2 \end{pmatrix} = \begin{pmatrix} \nu_{eL} \\ e_L \end{pmatrix} \in S_A \otimes V_A$$

$$\Phi = \begin{pmatrix} \Phi_1 \\ \Phi_2 \end{pmatrix} \in S_B \otimes V_B$$

$$e_R \in S_C \otimes V_C.$$

Here e_L and e_R correspond to the left-handed and right-handed **electron** (which are apriori unrelated fields; see below for an explanation of the common name "electron") and ν_{eL} is the left-handed **electron neutrino**. There are another two analogous generations in the Standard Model: the muon with the muon neutrino and the tau with the tau neutrino. The field Φ is the Higgs field.

In the Standard Model all right handed fermions, including the quarks, are singlets under the weak isospin group $SU(2)$ and transform in the trivial representation, while the left handed fermions form doublets (the three generations mentioned above as well as three generations of quark doublets) that transform in the standard representation of $SU(2)$. Left-handed and right-handed fermions therefore also transform in different representations of the weak hypercharge group $U(1)$, because each pair has the same electric charge. So far only left-handed neutrinos have been observed [Drewes 2013]. The electroweak interaction is thus a gauge theory with chiral matter.

Combining Propositions 9.2, 9.3 and 9.4 we have

Corollary 9.5. *The term*

$$\begin{aligned}\mathcal{L} &= 2g_Y \text{Re}(\bar{E}_L \Phi e_R) \\ &= 2g_Y \text{Re}(\Phi_1 \bar{E}_L^1 e_R + \Phi_2 \bar{E}_L^2 e_R),\end{aligned}$$

for a constant $g_Y > 0$, is a gauge invariant Lorentz scalar and defines a Yukawa coupling.

Here we introduced a factor of 2 for convenience. This is the classic Yukawa coupling that appears in the electroweak gauge theory of leptons. If we use the expansion

$$\Phi = \begin{pmatrix} 0 \\ ||v_0|| \end{pmatrix} + \Delta\Phi$$

for the Higgs field and keep only terms up to second order in the fields (without interactions), we get

$$\begin{aligned}\mathcal{L} &\approx 2||v_0||g_Y \text{Re}(\bar{e}_L e_R) \\ &= ||v_0||g_Y(\bar{e}_L e_R + \bar{e}_R e_L).\end{aligned}$$

This is precisely the mass term for a Dirac electron

$$e = \begin{pmatrix} e_L \\ e_R \end{pmatrix}$$

of mass $m_e = ||v_0||g_Y$. The previously unrelated Weyl spinors e_L and e_R thus combine after symmetry breaking into one Dirac spinor e (see [Robinson 2011] and [Srednicki 2007]). We also see that the neutrino remains massless. The determination of the electron mass (and similarly the mass of other fermions, like the quarks) is thus reduced to the determination of the parameter g_Y in

the Yukawa coupling and the norm $||v_0||$ of the vacuum value of the Higgs field.

9.4. Summary. In the Standard Model we have the following types of interactions (terms in the Lagrangian of order three or higher in the fields):

- Interactions between the gauge bosons and the fermions and between the gauge bosons and the Higgs field (via the covariant derivative)
- Interactions between gauge bosons (only in non-abelian gauge theories)
- Self-interactions of the Higgs field (via the Higgs potential)
- Interactions between the fermions and the Higgs field (via Yukawa couplings)

The famous 18 parameters of the classical Standard Model (with massless neutrinos), that so far have to be determined by experiments, are therefore:

- Two coupling constants of the electroweak interaction (or one coupling constant and the Weinberg angle) and the coupling constant of the strong interaction.
- The Yukawa couplings between the Higgs field and the nine massive fermions (three leptons, six quarks).
- The parameters λ and μ of the Higgs field (or the mass of the Higgs boson and the vacuum value of the Higgs field).
- There are another four parameters coming from the CKM matrix which is related to the weak interaction of quarks.

REFERENCES

- [Baum 2014] Helga Baum, *Eichfeldtheorie. Eine Einführung in die Differentialgeometrie auf Faserbündeln*, Springer Verlag, Berlin Heidelberg (2014).
- [Berline et al. 2004] Nicole Berline, Ezra Getzler, Michèle Vergne, *Heat Kernels and Dirac Operators*, Springer-Verlag, Berlin Heidelberg (2004).
- [Bleecker 1981] David Bleecker, *Gauge theory and variational principles*, Addison-Wesley Publishing Company, Reading, Massachusetts (1981).
- [Drewes 2013] Marco Drewes, *The phenomenology of right handed neutrinos*, Int. J. Mod. Phys. E **22** (2013).
- [Louis 2011] Jan Louis, *Quantum field theory II*, Lecture notes, University of Hamburg 2011.
- [Morgan 1996] John W. Morgan, *The Seiberg-Witten equations and applications to the topology of smooth four-manifolds*, Princeton University Press, Princeton, New Jersey (1996)
- [Mosel 1999] Ulrich Mosel, *Fields, symmetries, and quarks*, Springer Verlag, Berlin Heidelberg (1999).
- [Naber 2011] Gregory L. Naber, *Topology, geometry and gauge fields*, Springer, New York Dordrecht Heidelberg London (2011).
- [Robinson 2011] Matthew Robinson, *Symmetry and the Standard Model. Mathematics and particle physics*, Springer, New York Dordrecht Heidelberg London (2011).
- [Ryder 1996] Lewis H. Ryder, *Quantum field theory*, Cambridge University Press, Cambridge (1996).

- [Srednicki 2007] Mark Srednicki, *Quantum field theory*, Cambridge University Press, Cambridge
- [Ziller 2010] Wolfgang Ziller, *Lie groups. Representation theory and symmetric spaces*, Lecture notes, University of Pennsylvania 2010.

INSTITUTE FOR GEOMETRY AND TOPOLOGY, UNIVERSITY OF STUTTGART, PFAFFENWALDRING
57, 70569 STUTTGART, GERMANY
Email address: mark.hamilton@math.lmu.de