

Quantum Processes Near Black Holes*

G. W. Gibbons

University of Cambridge, D.A.M.T.P.

Silver Street, Cambridge, England

Abstract

A general review is given of quantum processes near black holes with a special emphasis on the Hawking Thermodynamic Emission Process. Astrophysical applications are not discussed.

I wish in this talk to summarize recent work on quantum effects near black holes. In doing so I wish to confine myself to giving an outline of what principles go into the calculations and the results. I shall not discuss any astrophysical applications (for which see e.g. [40]). I have tried to put the various results in some sort of perspective and I hope that in doing so I have not given insufficient weight to anyone's contribution or incorrectly judged it. If I have done so I apologize in advance. In my talk I hope to indicate what parts of the theory look satisfactory and which require more work and also I shall try to indicate parallels which other parts of physics — especially the theory of quantum processes in strong external electromagnetic fields.

The first indication of potentially interesting effects arose when Penrose pointed out the existence of what has come to be known as the “Penrose Effect” [1]. This arises because of the existence of negative energy orbits in-side the “ergosphere” of a rotating black hole (the region where the Killing vector which is timelike at infinity becomes spacelike). Given a region of negative energy orbits it is possible to extract energy — in this case the rotational energy of the black hole. One simply drops in a particle with positive energy E_1 , and lets it split (inside the region) into 2 particles one with positive energy E_2 , which emerges and the other with negative energy E_3 which remains inside. Since $E_1 = E_2 + E_3$

*This is a remake of the paper *Quantum Processes Near Black Holes* originally published in the first Marcel Grossmann Meeting on the Recent Progress of the Fundamentals of General Relativity, 1975, pages 449-458. The new version was typed out by Pete Su on November 18, 2024.

we have $E_2 > E_1$. This situation also occurs in electromagnetism near a point charge, in special relativity or indeed in any deep enough potential well [2]. It also occurs near charged black holes [3].

In fact in any electromagnetic background which is stationary, axisymmetric and invariant under simultaneous inversion of time and angle coordinates one finds that the energy E and angular momentum L of a particle of mass m and charge e must satisfy

$$(E dt + L d\phi + eA)^2 > m^2 \quad (1)$$

where A is the vector potential which falls to zero at infinity. This expression (or a simple generalization of it if there is a third constant of the motion) determines two surfaces $E^\pm(r, \theta)$ in the (E, r, θ) space between which a classical particle cannot exist. If the surface E^+ can fall below $-m$ we have just the required situation referred to sometimes as “level crossing”. The region r , is referred to as a “generalized ergosphere” for the mode in question. It is easy to check the existence of such a region in “superheavy” atoms. If

$$A = \Phi dt + B d\phi; \quad \Omega = g_{\phi t}/(g_{tt}) \quad (2)$$

the rate of rotation of inertial frames and $\sigma^2 = (g_{\phi t})^2 - g_{\phi\phi}g_{tt}$ we have

$$E^\pm = e\Phi + (L + eB)\Omega \pm \sigma^2 \sqrt{m^2 + (L + eB)^2}. \quad (3)$$

On a horizon $\sigma \rightarrow 0$ and $\Omega \rightarrow \Omega_H$, $\Phi + \Omega B \rightarrow \Omega_H$, and $E^\pm \rightarrow e\Phi_H + L\Omega_H = \mu_H$ which may be thought of as a chemical potential for the mode in question.

From the duality between waves and particles one expects a similar phenomenon to occur for waves and indeed this turns out to be the case (Misner [4] and Zeldovich [5]), and one has here the phenomenon of “super radiance.” For a classical scalar field this arises because the conserved flux vector

$$J_\mu = \frac{(\bar{\phi} \nabla_\mu \phi - \phi \nabla_\mu \bar{\phi})}{2i} \quad (4)$$

need not necessarily be future directed timelike. An incident wave carrying positive flux can send negative flux down the hole and the reflected positive flux can be greater than the incident flux. All of this is very reminiscent of the well known “Klein Paradox” situation [6] and indeed in the most general case of a charged, rotating black hole we have a rather close analogy to the Klein Paradox.

In our previous notation we find the ϕ can be written as $\phi = e^{iEt} e^{iL\phi} \chi$ and χ obeys

$$\frac{1}{\sigma} (\nabla_A \sigma \nabla^A \chi) + [(E dt + L d\phi + eA)^2 - m^2] \chi = 0 \quad (5)$$

where ∇_A denotes covariant differentiation in the r, θ plane. The conserved flux is

$$J = (E dt + L d\phi + A) |\chi|^2 + \frac{(\bar{\chi} d\chi - \chi d\bar{\chi})}{2i} \quad (6)$$

The null generator of the horizon is $\ell = \partial/\partial t - \Omega_H \partial/\partial \phi$. The flux through the horizon is $\propto \langle J, \ell \rangle \propto E - \mu_H$. Thus $E < \mu_H$ but $E^2 < m^2$ so we have superradiance. This is of course just the previous criterion.

For classical spin $\frac{1}{2}$ fields the situation is different the conserved flux vector

$$J = \bar{\psi} \Gamma_\mu \psi \quad (7)$$

is always future directed timelike and so simple super radiance is not possible [7]. However it is still possible for negative energy to fall down the hole since the stress tensor of a spin $\frac{1}{2}$ field does not obey the positive energy condition. Note that in both these cases a "hole" is necessary. Super radiance cannot occur unless a particle or energy can be trapped inside a certain region. Having seen how super radiance is possible, the analogy with "stimulated emission" is very close. On rather general grounds — Dirac [8], Feynman [9], Einstein [10] one expects — at least for bosons a related "spontaneous emission." Further each mode should be emitted with a coefficient just given by the super radiant coefficient (Starobinsky [11]). Note that while these physical arguments seem quite compelling one possible objection is that they seem to imply that a black hole can be some sort of thermal equilibrium with a surrounding heat bath. This as we shall see will turn out to be the case but at the early stages of this subject this seemed rather puzzling. Before I go on, it seems worthwhile here to point out that interesting as these speculations seem, the motivation for following them up would have been rather low had it not been in one's mind that rather small black holes (masses $>$ planck mass $\sim 10^{-5}$ g) had been postulated earlier by Hawking [12] as possibly arising in the early stages of a chaotic big bang universe, although the idea of black holes smaller than the Chandrasekhar limit had been suggested earlier by Zeldovich [13]. In this connection these early speculations brought to light an amusing coincidence Starobinsky [11] pointed out that the order of magnitude for the time for spontaneous loss of all of its angular momentum by a black hole of mass must be (in units such that $G = c = \hbar = k = 1$)

$$t \sim M^3 \quad (8)$$

Thus a hole would lose all of its angular momentum in less than 10^{10} years if its mass were less than 10^{-13} cm — a number not without significance in other contexts. The extension of these ideas to the charge of a black hole [14] showed similarly that unless the hole had a mass of this order, e^2/m_e , it would be energetically favorable for it to discharge itself even if it possessed a single

electron charge. The rate was expected to depend on the field strength and in a Schwinger [15] type way. Thus unless the electric field is less than the critical field mile the rate is very fast. This implied that to have a charge comparable with its mass the black hole mass must exceed e^2/m_e (which is coincidentally the least mass of a “classical geon” [16]. Essentially the same ideas seem to have occurred to Zaumen [17] independently. The story has also been taken up by Ruffini and Deruelle [19] and Ruffini and Damour [20]. These estimates made it very unlikely that mini black holes possessed charge.

Having seen the physical ideas which enter it remained to give them a more rigorous expression. The first person to tackle this problem was Unruh [21]. Since there is at present no well worked out candidate for a quantum theory of gravity Unruh adopted an approach in which the gravitational field was treated as a classical background—the so called external field approach. Thus one takes the equations describing a free quantum field in flat space and minimally couples them to the external field by the replacement $\partial_\mu \rightarrow \nabla_\mu - ieA_\mu$. This does not always yield a sensible theory [41] but in the case of spin 0, $\frac{1}{2}$ and 1 a workable theory results.

The next problem one encounters is the definition of particle states or a vacuum state. This may be summarized as follows:: the basic strategy of the quantum theory of fields is to resolve a field into normal modes. The coefficients of these normal modes obey the familiar bose einstein/ fermi-dirac commutation/anticommutation relations. This gives field commutation/anticommutation relations which are independent of the choice of normal modes—provided they are properly normalized with the natural sesquilinear form available:

$$\frac{1}{2i} \int (\bar{\phi} \nabla_\mu \phi - \phi \nabla_\mu \bar{\phi}) d\Sigma^\mu \quad \text{for spin 0} \quad (9)$$

$$\int \bar{\psi} \Gamma^\mu \psi d\Sigma^\mu \quad \text{for spin } \frac{1}{2} \quad (10)$$

What is *not* independent of the choice is the vacuum state. Any transformation of the normal modes (Bogoliubov transformation) which mixes up particle and antiparticle modes (or positive and negative frequencies to use a conventional expression) will give an inequivalent definition of the vacuum state. Indeed, as seems to occur in most practical examples, the number of “created particles” diverges and the two different state vectors may not even be connected by a unitary transformation [22]. Unruh made a particular choice — essentially that the particle modes be positive frequency with respect to the Killing vector $\partial/\partial t$ in the Kerr solution. He then computed the stress tensor expectation value $\langle T_{\mu\nu} \rangle$ in this state and found that $\langle T_{00} \rangle$ was infinite, $\langle T_{0r} \rangle$ was finite and corresponded to an outward flux of super radiant modes at the expected rate. Similar results were subsequently found by Ford [23]. It should be mentioned that the gravitational

background used was the maximally extended Kerr solution. We see that in general we meet three generic types of problem:

1. choice of vacuum state
2. infinities in T
3. breakdown for higher spins.

All of these problems occur and are familiar in the corresponding electromagnetic case. The next advance came with the work of Hawking [24]. He realized that

1. One can only satisfactorily define particle states at infinity
2. One must for a satisfactory treatment take collapse into account.

To take the first point; Hawking decided in the spirit of the matrix approach to define two vacua, the in vacuum and the out vacuum $|0_{-}\rangle$, $|0_{+}\rangle$. Provided past infinity constitutes a Cauchy surface (thus excluding the mixed white hole/black hole situation considered by Unruh) one may define an initial no particle state by the usual prescription of associating positive frequencies with particle states and conversely negative frequencies with anti-particle states. Since the idea of positive frequency is invariant under the asymptotic symmetry group, the B.M.S. group. This remains valid even in the presence of gravitational radiation. Indeed one can show that the gravitational field of a plane wave (like a plane electromagnetic wave) is incapable of producing particles [25]. Since at infinity any ingoing radiation will be effectively plane it is clear that as far as past infinity is concerned we have a reasonable definition of what it means to say that there are initially no ingoing particles. Similarly we can identify outgoing particles at future infinity. The task of identifying particle states in the interaction region is much less clear. This is perhaps not unreasonable. Physically a particle is really a certain sort of normal mode with high symmetry. Mathematically it is connected with irreducible representations of the Poincaré Group [26]. Neither of these concepts is applicable near the black hole. These remarks do *not* apply to the strong electromagnetic field around a large black hole since here the typical wavelength of the created particles is much smaller than the horizon size.

Now to turn to the next point. The problem is to count the number of out-going particles in state which initially contained no ingoing particles. We are working in the Heisenberg picture.

$$N_i^{out} = \langle 0_{-} | (a_i^{out})^{\dagger} (a_i^{out}) | 0_{-} \rangle \quad (11)$$

Taking the collapse into account and the very high redshifts associated with the formation of the event horizon Hawking was able to show that this number

diverges which he also showed corresponded to a steady emission at a rate

$$R_i = \Gamma_i \left(\exp \left(\frac{E_i - N_i \Omega_H - e_i \Phi_H}{T} \right) \mp 1 \right)^{-1} \quad \begin{array}{l} - \text{ for bosons} \\ + \text{ for fermions} \end{array} \quad (12)$$

where E_i is the energy of the outgoing particles; N_i the angular momentum; Ω_H the angular velocity of the hole; e_i the charge of the outgoing particle; Φ_H the electrical potential of the hole. Γ_i is the absorbtion coefficient of the hole for classical waves of energy E_i , etc, and T is related to a constant κ (“the surface gravity”) which plays an important role in black hole physics. For a black hole of mass M , charge Q these constants are

$$\begin{aligned} \Omega_H &= J/M^2 \\ \kappa &= r_+ - r_-/(2r^2) \\ r_0^2 &= r_+^2 + \frac{J^2}{M^2} \\ \Phi_H &= Qr_+/r_0^2 \\ r_{\pm} &= M \pm \sqrt{M^2 - \frac{J^2}{M^2} - Q^2} \\ T &= \kappa/2\pi. \end{aligned}$$

In terms of an “irreducible mass” $M_0 = r_0$ and the event horizon area A_H we have

$$A_H = 16\pi M_0^2 = 4\pi r_0^2 \quad (13)$$

$$M^2 = \left(M_0 + \frac{Q^2}{4M_0^2} \right)^2 + \frac{J^2}{4M_0^2} \quad (14)$$

There are several remarkable features of this result. Firstly, observe that while it was necessary to include the collapse in order to give meaning to the calculation the details of the collapse do not enter at all into the answer. If one does not consider the collapse then the most natural way of doing the calculations (i.e. in the fully extended Kruskal manifold) give no production [27]. It should be pointed out that the result one gets in the Kruskal manifold depends crucially on what boundary conditions one sets on the past horizon. A suitable choice could in principle yield any desired result. Unruh has pointed out that one may obtain Hawking’s result by choosing as part of one’s set of normal modes a set entering the external region of the black hole from the past horizon and behaving like a complex exponential of the affine parameter on the past horizon (see his article). Secondly in order to obtain the result it is necessary to take into account normal modes of arbitrarily high energy (way above the Planck frequency) essentially

because of the red shifting effect of the horizon. Indeed, if one imposes a cutoff frequency of ω_c the emission will stop after a time $t \sim (1/\kappa) \log(\omega_c/\kappa)$. This in fact is likely to be true in any calculation since it comes from considering how close to the horizon (in terms of an affine parameter) you have to be in order that you can send signals which reach infinity at a retarded time.

Thirdly we have here just the thermal emission which was lacking in our previous physical arguments — although the analogy between atomic levels and a black hole is not absolutely precise. It should be mentioned here that the idea that a black hole has associated with it a temperature and an entropy (which is of course what these results imply) had been suggested previously by Beckenstein [28] on the basis of certain analogies between black hole physics and thermodynamics (cf. [29]) — the non-decreasing event horizon area playing the part of entropy. The precise relation is now seen to be $S_H = A_H/4$.

One could of course spend a great deal of time describing the implications of these spectacular results. In what follows I shall bring out just a few points. Before proceeding, however, I should note that Hawking’s work encompasses all the previous results of Unruh and Starobinsky as limiting cases. The way in which this comes about for charged black holes and the relation of all this to the well known formulae due to Schwinger [15] for particle creation in uniform electric fields has been described by myself [30] cf also [20]. There are of course obvious analogies with thermionic emission from metal surfaces, $e \Phi_H$ acting as a “work function”.

It is sometimes said that the super-radiant processes like particle creation in a uniform static electric field do not require a time dependent field in contradistinction to the Hawking process. This is not really correct. The stationary (and in general non physical) backgrounds can only be used in conjunction with boundary conditions. These boundary conditions relate to what particles enter the space from the past horizon or from the infinite past respectively. This decision is made (via the choice of boundary conditions for Feynman propagators or a choice of a complete set of normal modes) in a way appropriate for a situation in which the interaction is “turned off” in the infinite past. From this point of view the “method of level crossing” amounts to:

1. a particular choice of normal modes (proportional to $e^{i\omega t}$).
2. a method of deciding whether the ingoing antiparticles overlap with outgoing particles (i.e. whether the Bogoliubov coefficients vanish) or not
3. an elegant representation of how large the effects will be by computing transmission coefficients in a certain effective potential. This is explained in more detail by Dr. Damour in his article and in [30] but the essential

point is that what is a particle or anti-particle is determined by the flux it carries through a Cauchy surface. As I explained earlier this depends, for horizon modes, on $E - \mu_H$. Thus roughly speaking, a wave can appear as antiparticle near the horizon ($E - \mu_H < 0$) but as a particle at infinity ($E > 0$).

From what has been said above it is clear that for all its spectacular successes, Hawking's derivation of his result is not entirely satisfactory. For an alternative attempt see [31], for skeptical remarks [32], [33]. This it seems to me not due to any fault on Hawking's part but due to the intrinsic problems with the whole external field theory method. For instance we still have the divergences in $\langle T_{\mu\nu} \rangle$ (and in the total number of particles) which need to be taken into account before we begin to feed back this expectation value into a Hartree-Fock version of Einstein's equations, if indeed that is appropriate. Some of these infinities can be absorbed as infinite renormalizations of the cosmological constant and Newton's constant G . Some, however, are of an essentially new nature, as has been pointed out by DeWitt [34]. In terms of an effective Lagrangian they correspond to terms of the form $R_{ab}R^{ab}$ and $(R_{ab}g^{ab})^2$. No detailed calculations of these have been given in the black case but there seems little ground for doubting their existence. Thus even the external field case external field problem is unrenormalizable which is closely related to the fact that the only quantum theory of gravity which is amenable to calculations suffers from the same defect [35]. One might also worry that particle interactions have not been taken into account. This is presumably valid for large holes but for holes of size 10^{-13}cm this considerable flux of particles in such a small volume should surely require the use of interacting field theory.

All of these problems are deep and difficult, what I want to do in the final part of the talk is to cut through the Gordian Knot as it were, disregard the original field theory derivation and hang everything on the thermodynamic idea.

Before doing so it is perhaps worthwhile trying to see why a black hole should give a thermal spectrum at all. The most natural interpretation is that the Hawking process consists of very many individual events consisting of the creation of a virtual pair near the hole and the tunnelling of a member (presumably carrying negative energy) through the horizon, the other reaching infinity.

What is seen is a statistical ensemble of such independent events each happening with a probability proportional to the phase space available. The factor Γ_i arises because the particles are created (in some sense) near the horizon and have to tunnel out through a combined curvature and centrifugal barrier. In all such calculations we need a constraint and in fact if we constrain the rate of dissipation of entropy or irreducible mass we obtain the Hawking formulae —

with of course the temperature undetermined. Thus a black hole emits thermally because it is the most likely thing for it to do. Support for the idea comes from the work of Wald [22] and Parker [36] who have shown that the statistics of outgoing particles are those of a black body.

Given that we have a small body whose temperature decreases the more energy it loses. This is, of course, a manifestation of the well known fact that gravitating systems can have negative specific heats. The evolution of such a system will be towards a hotter and hotter state —the black hole presumably disappearing altogether, and presumably with it the baryons that make it up. To my knowledge now one has mechanism whereby black holes can emit more baryons than antibaryon without making use of long range fields carrying baryonic charge [37]. If we adopt the thermodynamic viewpoint even when particle interactions are taken into account, then presumably a black hole will emit precisely what it would accrete from a heat bath at the same temperature. In terms of our simple arguments before the probability of emission is now no longer simply proportional to the available phase space. Carter, Lin and Perry and myself [38] have recently made some rough calculations on the basis. We represented a heat bath of strongly interacting particles as a perfect fluid with an equation of state. The accretion problem is straight forward and the corresponding emission is described by a sort of stellar wind. The interesting thing about our calculation is that provided the high density gas has a reasonably hard equation of state the wind is essentially transparent. This indicates that interactions might not be very important. It is perhaps worth reemphasizing here that for higher spin ($S > \frac{3}{2}$) the external field theory approach breaks down.

In this thermodynamic vein it is amusing to note that in this light the old theory of black body radiation requires revision. Consider a cavity of fixed volume V . They fill it with more and more energy.

At first the cavity will contain an ordinary black body gas and its temperature will rise. As it does, however, a qualitatively different behavior sets in. We have to maximize the entropy, S , of the configuration subject to the energy, E , being held constant.

$$S = 4\pi M^2 + \frac{4}{3}aVT^3 \quad (15)$$

$$E = M + aVT^4 \quad (16)$$

where a is Stefan's constant.

If $x = M/E$ and $y = (aV/E^5)/3\pi$ this amounts to maximizing $F = x^2 - (1-x)^{3/4}$ on $[0, 1]$. For $y > 2^5 3^{-1} 5^{-5/4} = 1.4266$ has no turning points and its greatest value is attained at $x = 0$ (pure radiation). For $1.4266 > y > 1.01440$ there is a local minimum at $x < \frac{4}{5}$ and a local maximum at $x < \frac{4}{5}$ but that

$x = 0$ is still a global maximum. For $y < 1.01440$ the local maximum at $x > 0.97702$ is also a global maximum. Thus for a box with sufficient energy a black hole of mass $M = \frac{1}{8\pi T}$ will condense out. As the energy of the enclosure is further increased the temperature will drop. Thus for any volume V there is a maximum temperature $T_m = \frac{1}{8\pi} x_c^{3/4} a V^{-1/5} (3\pi y_c)^{4/5} = x_c^{-1/4} T_c$; $x_c = 0.97702$ and $y_c = 1.01440$.

The point of the example is to indicate how basic Hawking's discovery is in relation to our views of fundamental physics. What part the sort of speculations contained in [38] on the deeper role that black holes have to play in the scheme of things remains to be seen. With that comment I shall close this review — in doing so is appropriate for me to thank my many colleagues in this field but most especially S. W. Hawking for many discussions.

ADDENDUM

After completing the review I received the following preprints. The of which is most pertinent to the final section:

S. W. Hawking, "Black Holes and Thermodynamics"

T. Damour "On the Correspondence between Classical and Quantum Energy States in Stationary Geometries"

N. Deruelle, R. Ruffini "Klein Paradox in a Kerr Geometry"

W. Unruh "Notes on Black Hole Evaporation"

REFERENCES

- [1] R. Penrose. Riv. Nuovo Cimento **1** 252 (1969)
- [2] G. Denardo & A. Treves. Lett. al. Nuovo Cimento **8** 295 (1973)
- [3] G. Denardo & R. Ruffini. Phys. Lett. 45B 259 (1973); G. Denardo, L. Hively & R. Ruffini. Phys. Lett. 50B 270 (1974); cf. also Ya. B. Zeldovich & V. S. Popov. Uspekhi **14** 673 (1972); L. I. Schiff, H. Snyder & J. Weinberg, Phys. Rev. **57** 315 (1940)
- [4] C. Misner. Bull. Amer. Phys. Soc. **17** 472 (1972)
- [5] Ya. Zeldovich J.E.T.P. **35** (1972)
- [6] O. Klein, Zeit. fur Phys. **53** 157 (1929)
- [7] W. Unruh. Phys. Rev. Lett. **31** 1265 (1973)
- [8] P. A. M. Dirac. Quantum Mechanics O.U.P.
- [9] R. D. Feynman. Lectures on Physics, Vol. III
- [10] A. Einstein. Phys. Zeit **18** 121 (1917)
- [11] A. Starobinsky. J.E.T.P. **37** 28 (1973)
- [12] S. W. Hawking. M.N.R.A.S. **152** 75 (1971); B. J. Carr & S. W. Hawking, M.N.R.A.S. **168** 399 (1974)
- [13] Ya. Zeldovich. J.E.T.P. **446** (1962)
- [14] G. W. Gibbons & S. W. Hawking. Work reported at Warsaw Conference 1973 see "Gravitational Radiations and Gravitational Collapse" C. deWitt (ed.) Reidel (1974)
- [15] J. Schwinger. Phys. Rev. **82** 664 (1951)
- [16] J. Wheeler. "Geometrodynamics" Academic Press (1972)
- [17] W. T. Zaumen. Nature **247** 530 (1974)
- [19] N. Deruelle & R. Ruffini. Phys. Lett. **52B** 437 (1975)
- [20] T. Damour & R. Ruffini. "Quantum electrodynamic effects in Kerr-Newman Geometries" Princeton Preprint. Dec. 1974
- [21] W. Unruh. Phys. Rev. D **10** 3194 (1974) .

- [22] Cf. M. Castagnino, A. Verbeuri & R. A. Weder, *Nuovo Cimento* **26B** 396 (1975), *Phys. Lett.* **48A** 99 (1974); R. Wald. *Commun. Math. Phys.* **46** (1975)
- [23] L. Ford “Quantization of a Scalar Field in the Kerr Spacetime” Milwaukee Preprint UWM-4867-74-17 (1974)
- [24] S. W. Hawking. *Nature* **248** **30** (1974); S. W. Hawking. *Commun. Math. Phys.* **43** 199 (1975)
- [25] G. W. Gibbons. *Commun. M. Phys.* **45** 191 (1975)
- [26] E. Wigner. *Ann. of Math.* **40** (1939)
- [27] D. Boulware. *Phys. Rev. D* **11** 1406 (1975); D. Boulware. “Spin 1/2 Quantum Field Theory in Schwarzschild Space” Seattle Preprint RL-1388-689 (1975).
- [28] J. D. Beckenstein. *Phys. Rev.* **D7** 2333 (1972)
- [29] J. Bardeen, B. Carter & S. W. Hawking, *Commun. Math. Phys.* **31** 162 (1973)
- [30] G. W. Gibbons, *Commun. Math. Phys.* **44** 245 (1975)
- [31] U. Gerlach. “Mechanism of Black Body Radiation from an incipient Black Hole” Ohio Preprint.
- [32] J. G. Taylor & P. Davies. *Nature* **250** 37 (1974) .
- [33] P. W. Davies. *J. Phys. A* **8** 609 (1975).
- [34] B. deWitt. *Physics Reports* **29C** 295 (1975).
- [35] E.g. S. Deser and P. van Nieuwenhuizen. *Phys. Rev. D* **10** 411 (1974) .
- [36] L. Parker *Phys. Rev. D* **12** (1975).
- [37] B. Carter. *Phys. Rev. Lett.* **33** 558 (1974).
- [38] B. Carter, G. W. Gibbons, D. Lin & M. Perry “The Black Hole emission process in the high energy limit” in preparation.
- [39] J. Sarfatti. *Nature* **240** 101 (1972) .
- [40] B. J. Carr. “The Primordial Mass Spectrum” Caltech Preprint (1975).
- [41] G. W. Gibbons. *J. Phys. A* **9** 145 (1976).