

# From $\lambda$ -Calculus to Cartesian Closed Categories

J. Lambek

*Mathematics Department, McGill University*

*Montreal, P.Q. H3A 2K6 Canada.*

*Dedicated to Professor H. B. Curry on the occasion of his 80th Birthday \**

Haskell Curry may be surprised to hear that he has spent a lifetime doing fundamental work in category theory. The purpose of this account is to convince categorists that Cartesian closed categories (Eilenberg and Kelly, 1966) have been anticipated by logicians (Curry, 1930) by many years and, conversely, to persuade logicians that combinatory logic may benefit from being phrased in categorical language.

I have attempted to tell this story twice before (1972, 1974), but am not entirely satisfied with these earlier accounts. The present exposition is essentially my unscheduled talk at the 1977 Durham Symposium on applications of sheaf theory to logic, algebra and analysis.

I regret that limitations of space do not permit a discussion of illative combinatory logic (Curry and Feys, 1958) or combinatory type theory (Church, 1940) and applications thereof to the construction of free toposes.

Let me confess at once that I am not a historical scholar and that I have taken some liberties with the original material. Thus, I have taken the opportunity to present the early discoveries of combinatory logic in the language of universal algebra.

Our story begins in 1924, when Schönfinkel studied what would now be called an algebra  $\mathcal{A} = (|\mathcal{A}|, \cdot, I, S, K)$  consisting of a set  $|\mathcal{A}|$  equipped with a binary operation  $\cdot$  and constants  $I, S$  and  $K$ . These were to satisfy the following identities:

$$(1) \quad I \cdot a = a,$$

---

\*This is a remake of the paper *From  $\lambda$ -Calculus to Cartesian Closed Categories* originally published in R. Hindley and J. Seldin, editors, *To H.B. Curry: Essays in Combinatory Logic, Lambda Calculus and Formalisms*. Academic Press, 1980. This file was created in March 2023.

$$(2) \quad (K \circ a) \circ b = a,$$

$$(3) \quad ((S \circ f) \circ g) \circ c = (f \circ c) \circ (g \circ c),^\dagger$$

for all elements  $a, b, c, f$  and  $g$  of  $|\mathcal{A}|$ . Actually, Schönfinkel did not employ the language of universal algebra, and he defined  $I$  in terms of  $K$  and  $S$ , but of this we shall speak later. His main result would now be stated as follows.

**Proposition 1.** *Every polynomial  $\varphi(x)$  over a Schönfinkel algebra  $\mathcal{A}$  can be written in the form  $f \circ x$ , where  $f \in |\mathcal{A}|$ .*

Polynomials are of course formed as words in an indeterminate  $x$  and are subject to the same three identities. More precisely, equality  $\equiv_x$  between polynomials is the smallest equivalence relation  $\equiv$  between words in  $x$  which has the substitution property

$$(0_x) \quad \frac{\varphi(x) \equiv \psi(x) \quad \alpha(x) \equiv \beta(x)}{\varphi(x) \circ \alpha(x) \equiv \psi(x) \circ \beta(x)}$$

and which satisfies

$$(1_x) \quad I \circ \alpha(x) \equiv \alpha(x),$$

$$(2_x) \quad (K \circ \alpha(x)) \circ \beta(x) \equiv \alpha(x)$$

$$(3_x) \quad ((S \circ \varphi(x)) \circ \psi(x)) \circ \gamma(x) \equiv (\varphi(x) \circ \gamma(x)) \circ (\psi(x) \circ \gamma(x))$$

Alternatively, one may regard the polynomial  $\varphi(x)$  as an element of the Schönfinkel algebra  $\mathcal{A}[x]$ , which comes equipped with an element  $x$  and with a homomorphism  $h_x : \mathcal{A} \rightarrow \mathcal{A}[x]$  with the usual universal property: for every algebra  $\mathcal{B}$  every homomorphism  $f : \mathcal{A} \rightarrow \mathcal{B}$  and every element  $b \in |\mathcal{B}|$ , there exists a unique homomorphism  $f' : \mathcal{A}[x] \rightarrow \mathcal{B}$  such that  $f' \circ h_x = f$  and  $f'(x) = b$ . In the special case when  $\mathcal{B} = \mathcal{A}$  and  $f$  is the identity homomorphism  $\mathcal{A} \rightarrow \mathcal{A}$ ,  $f'$  is the *substitution* homomorphism which allows us to replace  $x$  in any polynomial by  $b$ . Similarly, when  $\mathcal{B} = \mathcal{A}[x]$  and  $f = h_x$ ,  $f'$  allows us to replace  $x$  by a polynomial  $\beta(x)$ .

The proof of Proposition 1 is remarkably simple. It proceeds by induction on the length of the word  $\varphi(x)$ , which must be either  $x$ , or a constant or

---

<sup>†</sup>Editor's note: Lambek uses this squiggle  $\circ$  to denote a binary operation for function application. It appears here and in his later book about this same subject. I made a best guess about how to translate this into  $\mathbb{E}\mathbb{T}\mathbb{E}\mathbb{X}$ .

of the form  $\psi(x) \cdot \chi(x)$  not constant, in which last case we may assume by inductual assumption that  $\psi(x) \equiv g \cdot x$  and  $\chi(x) \equiv h \cdot x$ . In the three cases we have

$$\begin{aligned}\varphi(x) &\equiv I \cdot x, \\ \varphi(x) &\equiv (K \cdot a) \cdot x, \\ \varphi(x) &\equiv (g \cdot x) \cdot (h \cdot x) \equiv ((S \cdot g) \cdot h) \cdot x,\end{aligned}$$

respectively.

This proof also yields an algorithm for converting every polynomial into the form  $f \cdot x$ . For example, one easily calculates

$$x \cdot x \equiv ((S \cdot I) \cdot I) \cdot x.$$

In 1930, Schönfinkel's result was rediscovered by Curry. However, Curry was interested in imposing an additional requirement:

$$(4) \quad \text{If } f \cdot x \equiv g \cdot x \text{ in } \mathcal{A}[x] \text{ then } f = g \text{ in } \mathcal{A}.$$

For example, from

$$((S \cdot K) \cdot I) \cdot x \equiv (K \cdot x) \cdot (I \cdot x) \equiv x \equiv I \cdot x$$

one could use (4) to obtain  $(S \cdot K) \cdot I = I$ , which equation cannot be derived from (1) to (3) alone. In the same way, one could deduce that  $(S \cdot K) \cdot K = I$ . In fact, Schönfinkel originally defined  $I$  by this equation. For reasons that will become clear later, we shall not follow him in this application of Occam's razor.

While (4) does not have the form of an identity, by which I mean an equation prefixed by universal quantifiers, Curry discovered that it could be replaced by a finite number of equations. These were later simplified, and Rosenbloom lists four, one of which reads:

$$(S \cdot ((S \cdot (K \cdot S)) \cdot K)) \cdot (K \cdot I) = I.$$

The reader will forgive us for not copying out the other three!

By a *Curry algebra* we shall mean a Schönfinkel algebra subject to certain additional equations or identities whose conjunction is equivalent to (4). Curry's result may then be formulated thus:

**Proposition 2.** *Over a Curry algebra  $\mathcal{A}$  every polynomial  $\varphi(x)$  may be uniquely written in the form  $f \cdot x$  with  $f \in |\mathcal{A}|$ .*

This property of Curry algebras is called *functional completeness*. It is an immediate consequence of the equivalence of (4) with the conjunction of the above mentioned four equations. For a proof we could refer the reader to the book by Rosenbloom. However, we prefer to give another proof, in the course of which we shall discover five equations whose conjunction is equivalent to (4).

*Proof.* Let  $\lambda_x \varphi(x)$  be defined by induction on the length of the word  $\varphi(x)$  thus:

- (i)  $\lambda_x x = I,$
- (ii)  $\lambda_x a = K^1 a,$  when  $a$  is a constant;
- (iii)  $\lambda_x (\psi(x)^1 \chi(x)) = (S^1 \lambda_x \psi(x))^1 \lambda_x \chi(x)$  when  $\psi(x)$  and  $\chi(x)$  are not both constant.

We shall prove below that the restriction on (iii) is not necessary. In view of the above proof of Proposition 1, we have

$$\varphi(x) = \lambda_x \varphi(x)^1 x$$

so the existence of  $f$  with  $\varphi(x) \equiv_x f^1 x$  is assured. It remains to prove its uniqueness. First, we claim that

$$(*) \quad \varphi(x) \equiv_x \psi(x) \quad \text{implies} \quad \lambda_x \varphi(x) = \lambda_x \psi(x),$$

so that  $\lambda_x \varphi(x)$  depends not just on the word  $\varphi(x)$  but on the polynomial  $\varphi(x)$ , that is, the word modulo the equivalence relation  $\equiv_x$ .

To prove (\*), we write  $\varphi(x) \equiv \psi(x)$  for  $\lambda_x \varphi(x) = \lambda_x \psi(x)$ . It is easily seen that  $\equiv$  is an equivalence relation between words which satisfies  $(0_x)$ , that is, a congruence relation. If we make sure that  $\equiv$  also satisfies  $(1_x)$  to  $(3_x)$  it will follow that  $\equiv$  contains  $\equiv_x$ , and this is what (\*) asserts.

To say that  $\equiv$  satisfies  $(1_x)$  means that

$$\lambda_x (I^1 \alpha(x)) = \lambda_x \alpha(x).$$

Writing  $a$  for  $\lambda_x \alpha(x)$  we may rewrite this, in view of the unrestricted (iii), as

$$(S^1 (K^1 I))^1 a = a,$$

an easy consequence of

$$(4.1) \quad S^1 (K^1 I) = I,$$

which itself has already been derived from (4).

In the same manner we may obtain consequences (4.2) and (4.3) of (4) which imply  $(2_x)$  and  $(3_x)$  respectively. We shall not bother to spell them out.

We now turn to the uniqueness of  $f$  in Proposition 2. Suppose also  $g^1x \equiv \varphi(x)$ , we claim that  $g = \lambda_x \varphi(x)$ . Now, by (\*)  $\lambda_x(g^1x) = \lambda_x \varphi(x)$ , so it suffices to prove that  $\lambda_x(g^1x) = g$ .

Now a small calculation shows that  $\lambda_x(g^1x) = (S^1(K^1g))^1I$ , so we require the identity

$$(S^1(K^1g))^1I = g$$

for all  $g$ , which is an easy consequence of (4).

This identity must remain valid if we adjoin an indeterminate  $y$  to the algebra, so we have

$$(S^1(K^1y))^1I \equiv y.$$

We may therefore replace the required identity by the equation

$$(4.4) \quad \lambda_y(S^1(K^1y))^1I = y.$$

Of course the  $A$  may be eliminated from this using (i) to (iii).

It remains to show the validity of (iii) when  $\psi(x)^1\chi(x)$  is constant, say  $b^1c$ . So we want to show that

$$\lambda_x(b^1c) = (S^1(K^1b))^1(K^1c).$$

But, by (ii),  $\lambda_x(b^1c) = K^1(b^1c)$ , so we are led to stipulate the identity

$$(S^1(K^1b))^1(K^1c) = K^1(b^1c)$$

for all  $b$  and  $c$ . gain, this is an easy consequence of (4). By the same argument as above, we may replace the stipulated identity by the equation

$$(4.5) \quad \lambda_x \lambda_y((S^1(K^1x))^1(K^1y)) = \lambda_x \lambda_y(K^1(x^1y)).$$

The proof of Proposition 2 is now complete, provided we adopt (4.1) to (4.5) as the five equations which a Curry algebra must satisfy in addition to the identities (1) to (3).  $\square$

From now on we shall write  $\lambda_x \varphi(x)$  for the unique  $f$  corresponding to  $\varphi(x)$  by Proposition 2, as we did in the proof. The properties of the new symbol  $\lambda_x$  are embodied in the  $\lambda$ -calculus of Church (1932). The equivalence of the systems of Curry and Church ( $\lambda K$ -calculus, 1941) are summed up as follows.

**Proposition 3.** *The identities and equations of Curry algebras imply and are implied by the following.<sup>1</sup>*

- (1)  $I = \lambda_x x,$
- (2)  $K = \lambda_x \lambda_y x,$
- (3)  $S = \lambda_u \lambda_v \lambda_z ((u^{\cdot} z)^{\cdot} (v^{\cdot} z)),$
- (4)  $\lambda_x (f^{\cdot} x) = f,$
- (5)  $(\lambda_x \varphi(x))^{\cdot} a = \varphi(a).$

The proof is almost straightforward. We shall only explain why (5) holds for Curry algebras. By Proposition 2, we have  $f^{\cdot} x \stackrel{\cdot}{=} \varphi(x)$  and we want to deduce from this that  $f^{\cdot} a = \varphi(a)$ . By the universal property of  $\mathcal{A}[x]$ , there exists a unique homomorphism  $h' : \mathcal{A}[x] \rightarrow \mathcal{A}$  such that  $h' = h_x$  and  $h'(x) = a$ . This is of course the substitution homomorphism which replaces  $x$  by  $a$ , hence yields the required equation from  $f^{\cdot} x \stackrel{\cdot}{=} \varphi(x)$ .

To recapture the traditional terminology, let us mention that the theory of Curry algebras is called *combinatory logic*. Proposition 2 may then be compressed into the slogan:

$$\text{combinatory logic} = \lambda\text{-calculus}.$$

Incidentally, *combinators* are the canonical elements of Curry (or Schönfinkel) algebras, that is, the elements of the free Curry algebra generated by the empty set.

Both Schönfinkel and Curry had intended to use combinatory logic for the foundations of mathematics. An obstacle arose in the following result.

**Proposition 4.** *In a Curry algebra every element  $f$  has a "fixpoint"  $a$  such that  $f^{\cdot} a = a$ .*

*Proof.* Let  $b = \lambda_x (f^{\cdot} (x^{\cdot} x))$  and put  $a = b^{\cdot} b$ . Then

$$f^{\cdot} a = f^{\cdot} (b^{\cdot} b) = b^{\cdot} b = a.$$

□

---

<sup>1</sup>Here  $=$  denotes equality in  $\mathcal{A}$ ,  $\mathcal{A}[x]$ ,  $\mathcal{A}[x, y]$ , etc. simultaneously, subscripts having been suppressed. It is an equivalence relation subject to the following rules:

$$\frac{f = g \quad a = b}{f^{\cdot} a = g^{\cdot} b}, \quad \frac{\varphi(x) = \psi(x)}{\lambda_x \varphi(x) = \lambda_x \psi(x)}$$

If  $f$  is negation, usually denoted by  $\neg$ , we have  $\neg a = a$ , so  $a$  cannot be a proposition. We must therefore distinguish between propositions and other entities; but even this distinction does not prevent Russell's paradox from raising its head (e.g. Curry and Feys, 1958).

It is of course well-known that Russell's paradox may be avoided by introducing types. In the following exposition of typed combinatory logic I shall follow Curry and Feys in principle, even though I shall reject their permissiveness in allowing symbols with ambiguous types.

First of all we replace a Curry algebra by a many-sorted algebra, whose elements, which we call entities, may belong to different sorts, which we call types. If  $A$  and  $B$  are types, so is  $B^A$ , the type of all "functions" from  $A$  to  $B$ . For typographical reasons, we write  $B \Leftarrow A$  in place of  $B^A$ . The binary operation symbol  $\cdot$  may not be placed between entities indiscriminately, but is subject to the following rule:

$$(0) \quad \frac{a \in A \quad f \in B \Leftarrow A}{f \cdot a \in B},$$

meaning that, if  $a$  is of type  $A$  and  $f$  of type  $B \Leftarrow A$  then  $f \cdot a$  is of type  $B$ .<sup>2</sup>

In place of the three constants  $I$ ,  $K$  and  $S$  we adopt three families of constants:

- (1)  $I_A \in A \Leftarrow A$  such that  $I_A \cdot a = a$
- (2)  $K_{A,B} \in (B \Leftarrow A) \Leftarrow A$  such that  $(K_{A,B} \cdot a) \cdot b = a$
- (3)  $S_{A,B,C} \in ((C \Leftarrow A) \Leftarrow (C \Leftarrow B)) \Leftarrow ((A \Leftarrow B) \Leftarrow C)$  such that  $((S_{A,B,C} \cdot f) \cdot g) \cdot c = (f \cdot c) \cdot (g \cdot c)$ .

It is assumed here that  $a \in A$ ,  $b \in B$ ,  $c \in C$ ,  $f \in (A \Leftarrow B) \Leftarrow C$  and  $g \in B \Leftarrow C$ .

A many-sorted algebra of the kind constructed above may be called a *typed Schönfinkel algebra*, or a *typed Curry algebra* if the appropriate equations are postulated.

Curry and Feys have pointed out that, if  $\Leftarrow$  is read as "if", the types of  $I_A$ ,  $K_{A,B}$ , and  $S_{A,B,C}$  are precisely the axioms of the intuitionist implicational calculus, while (0) bears an obvious relation to the usual rule of modus ponens.

Two remarks are in order. The same axioms also appear in the classical propositional calculus accompanied by an additional axiom involving

---

<sup>2</sup>Actually, the application symbol  $\cdot$  should carry subscripts  $A$  and  $B$ ; but we omit these whenever they are clear from the context, as they are throughout this paper.

negation. Nonetheless, the negationless formula  $A \Leftarrow (A \Leftarrow (B \Leftarrow A))$  is a theorem classically but not in the system without negation. This is why we call the present system intuitionistic.

Secondly, it should be pointed out that  $A \Leftarrow A$  is usually not taken as an axiom, but is deduced from the other two axioms. Nevertheless, we prefer to regard it as an axiom here.

Implicit in the the definition of a typed Schönfinkel algebra is a way of regarding each entity of type  $A$  as a proof of the formula  $A$ .

For example,  $S_{A,B,A}$  is by definition a proof of the axiom  $((A \Leftarrow A) \Leftarrow (B \Leftarrow A)) \Leftarrow ((A \Leftarrow B) \Leftarrow A)$  and  $K_{A,B}$  is by definition a proof of the axiom  $(A \Leftarrow B) \Leftarrow A$ . Hence  $S_{A,B,A} \circ K_{A,B}$  is a proof of the theorem  $(A \Leftarrow A) \Leftarrow (B \Leftarrow A)$  by modus ponens. Again,  $K_{A,C}$  is a proof of the axiom  $(A \Leftarrow C) \Leftarrow A$ . Take  $B = (A \Leftarrow C)$  then  $(S_{A,B,A} \circ K_{A,B}) \circ K_{A,C}$  is a proof of the theorem  $A \Leftarrow A$ . Since we regard  $A \Leftarrow A$  as axiom, another proof is  $I_A$ . Incidentally, we see here that the derivation of  $A \Leftarrow A$  the other axioms of the implicational calculus is nothing else then Schönfinkel's definition of  $I$  as  $(S \circ K) \circ K$ .

The association of entities with proofs becomes even more striking when we compare the free typed Schönfinkel algebra (generated by a set of letters) with pure intuitionistic implicational logic. Then

$$\text{combinators} = \text{proofs.}$$

The reader should note that throughout we distinguish between an axiom such as  $(A \Leftarrow B) \Leftarrow A$  and its proof  $K_{A,B}$ , a pedantic but necessary distinction.

Let us now look at Proposition 1 for typed Schönfinkel algebras. The reader will easily convince himself that the proposition and its proof remain valid, provided  $x$  is an indeterminate of type  $A$ , where  $A$  is any type of  $\mathcal{A}$ . It asserts that, if  $\varphi(x)$  is a polynomial of type  $B$  in the indeterminate  $x$  of type  $A$ , then there is a constant  $f$  of type  $B \Leftarrow A$  such that  $\varphi(x) \equiv_{\overline{x}} f \circ x$ .

In the proof-theoretic interpretation we should regard  $x$  as an assumption that  $A$  holds. Proposition 1 then becomes the usual deduction theorem with an extra punch at the end: if  $\varphi(x)$  is a proof of  $B$  from the assumption  $x$  that  $A$  holds, then there is a proof of  $B \Leftarrow A$  without any assumption such that  $\varphi(x) \equiv_{\overline{x}} f \circ x$ .

The extra bit at the end asserts the “equality” of two proofs. Perhaps it would have been better to speak only of “equivalence” of proofs.

As far as I know, this association between combinators and proofs is due to Curry and Feys (1958). It was developed further by Howard (see Stenlund,



1972) and the author (1969, 1972).<sup>3</sup>

Proposition 2 will also remain valid for typed Curry algebras. It asserts that the proof  $f$  of  $B \Leftarrow A$ , whose existence has been established in Proposition 1, is unique up to equivalence of proofs (which we called equality here).

Proposition 3 will remain valid for typed Curry algebras provided we use the typed  $\lambda$ -calculus and write  $f = \lambda_{x \in A} \varphi(x)$ .

Proposition 4 will not remain valid for typed Curry algebras without very strong restrictions. After all, the whole purpose of introducing types was to render expressions such as  $b \cdot b$  meaningless in general. If  $b \in B$ , it does have a meaning only if  $B = B \Leftarrow B$ . Proposition 4 remains valid for typed algebras in the following sense: if  $A$  is a type such that  $A \Leftarrow A = A$ , then every entity  $f$  of type  $A$  has a fixpoint  $a$  such that  $f \cdot a = a$ .

The fragment of propositional logic investigated up to now studiously avoids conjunction, which classically was usually defined in terms of implication and negation. Let us now turn to a fragment of logic, the positive intuitionist propositional calculus, which deals with implication  $\Leftarrow$ , conjunction  $\wedge$ , and truth  $\top$ . We shall present this as a deductive system, making use of the symbol  $\longrightarrow$  for entailment, in the spirit of Gentzen. Here are our axioms and rules of inference, suitably labelled:

- $$\begin{aligned}
 (1) \quad & A \xrightarrow{1_A} A, \quad \frac{A \xrightarrow{f} B \quad B \xrightarrow{g} C}{A \xrightarrow{gf} C} \\
 (2) \quad & A \xrightarrow{0_A} \top; \\
 (3) \quad & A \wedge B \xrightarrow{\pi_{A,B}} A, \quad A \wedge B \xrightarrow{\pi'_{A,B}} B, \quad \frac{C \xrightarrow{f} A \quad C \xrightarrow{g} B}{C \xrightarrow{\langle f, g \rangle} A \wedge B} \\
 (4) \quad & (A \Leftarrow B) \wedge B \xrightarrow{\varepsilon_{A,B}} A, \quad \frac{C \wedge B \xrightarrow{h} A}{C \xrightarrow{h^*} (A \Leftarrow B)}
 \end{aligned}$$

The labels are useful in naming proofs. For example, the commutative law is proved thus:

$$\frac{A \wedge B \xrightarrow{\pi'_{A,B}} B \quad A \wedge B \xrightarrow{\pi_{A,B}} A}{A \wedge B \xrightarrow{\langle \pi'_{A,B}, \pi_{A,B} \rangle} B \wedge A}$$

---

<sup>3</sup>For related ideas and further references see also the thought provoking paper by Scott (1970).

The label  $\langle \pi'_{A,B}, \pi_{A,B} \rangle$  appearing in the last line may be used to denote the whole proof.

We are also interested in an equivalence relation between proofs, which we may as well denote by the equality symbol  $=$ . We do not bother to write down the usual reflexive, symmetric, transitive and substitution laws for equality. However, we list the following equations, where in  $f = g$  it is assumed that  $f$  and  $g$  have the same source and target.

$$(1') \quad f1_A = f, 1_B f = f, (hg)f = g(hf), \text{ for all } f : A \longrightarrow B, g : B \longrightarrow C, \text{ and } h : C \longrightarrow D;$$

$$(2') \quad f = 0_A, \text{ for all } f : A \longrightarrow 1;$$

$$(3') \quad \pi_{A,B} \langle f, g \rangle = f, \pi'_{A,B} \langle f, g \rangle = g, \langle \pi_{A,B} h, \pi'_{A,B} h \rangle = h, \text{ for all } f : C \longrightarrow A, g : C \longrightarrow B, \text{ and } h : C \longrightarrow A \wedge B;$$

$$(4') \quad \varepsilon_{A,B} \langle h^* \pi_{C,B}, \pi'_{C,B} \rangle = h, (\varepsilon_{A,B} \langle k^* \pi_{C,B}, \pi'_{C,B} \rangle)^* = k, \text{ for all } h : C \wedge A \longrightarrow A, \text{ and } k : C \longrightarrow A \Leftarrow B.$$

The reader will recognize that (1) and (1') define a category. Moreover, (2) and (2') assert that  $\top$  is a terminal object, usually denoted by 1. Furthermore, (3) and (3') assert that  $A \wedge B$  is the Cartesian product of  $A$  and  $B$ , usually written  $A \times B$ . Finally, (4) and (4') express the fact that  $(-) \Leftarrow B$  is right adjoint to  $(-) \wedge B : \mathcal{A} \longrightarrow \mathcal{A}$ , which makes  $\Leftarrow$  an internal hom-functor, the usual notation for  $A \Leftarrow B$  being  $A^B$ . (1) to (3) and (1') to (3') define a Cartesian category, (1) to (4) and (1') to (4') define a Cartesian closed category.

Cartesian closed categories were introduced under this name by Eilenberg and Kelly (1966). Lawvere (1969) also pointed out that  $\times$  is right adjoint to the diagonal  $\mathcal{A} \longrightarrow \mathcal{A} \times \mathcal{A}$  and emphasized the analogy with propositional logic. Our definition of Cartesian closed categories is slightly different, in as much as 1,  $\times$ , and exponentiation are not only said to exist, but are regarded as part of the structure. In fact, our notion of Cartesian closed category is algebraic over the category of categories, or the category of graphs for that matter, in much the same way that the notion of group is algebraic over the category of sets.

For future reference, we also note the following consequence of conditions (1') to (4'):

$$(5') \quad \langle f, g \rangle h = \langle fh, gh \rangle.$$

This is proved thus: let  $k = \langle f, g \rangle h$ , then, omitting subscripts, we have

$$\pi k = fh, \pi' k = gh$$

by (3) and hence

$$\langle fh, gh \rangle = \langle \pi k, \pi' k \rangle = k,$$

by (3') again.

In the author's opinion, it is a tour de force to present propositional logic without conjunction. Curiously, the same tour de force is found in the paper by Eilenberg and Kelly, who went to some trouble to eliminate the Cartesian product from Cartesian closed categories. One could argue that closed categories without products are essentially the same as typed Curry algebras. It therefore seems reasonable to proceed from the study of typed Curry algebras to the study of Cartesian closed categories.

I had proved in 1972 that functional completeness holds for Cartesian closed categories, provided that these satisfied a finite set of equations, like those due to Curry, and conjectured that these equations are already a consequence of (1') to (4') above, that is, they hold in any Cartesian closed category. I proved this later (1974) and shall give another exposition of the proof here. The new versions of Propositions 1 and 2 will be called Theorems 1 and 2.

First, we must explain what it means to adjoin an indeterminate morphism  $x : 1 \rightarrow A$  to a Cartesian closed category  $\mathcal{A}$  with  $A$  being an object of  $\mathcal{A}$ , which we also regard as a formula. In the same spirit, we may regard  $x$  as an assumption that  $\top$  entails  $A$ . The objects of  $\mathcal{A}[x]$  are the same as those of  $\mathcal{A}$ , but the morphisms  $\varphi(x) : B \rightarrow C$  in  $\mathcal{A}[x]$  may be regarded as proofs that  $B$  entails  $C$  on the assumption  $x$ . Equality between proofs or polynomials is determined by (1') to (4') and is denoted by  $\overline{\equiv}_x$ . One must check of course that  $\mathcal{A}[x]$  thus constructed has the expected universal property.

**Theorem 5** (Deduction theorem). *With every proof  $\varphi(x) : B \rightarrow C$  from the assumption  $x : 1 \rightarrow A$  there is associated a proof  $f : A \wedge B \rightarrow C$  not depending on this assumption. Moreover  $f \langle x 0_B, 1_B \rangle \overline{\equiv}_x \varphi(x)$ .*

It should be pointed out that we have presented the positive intuitionist propositional calculus as a deductive system, so that the usual deduction theorem becomes absorbed in the rules governing the deduction symbol  $\rightarrow$ , thus:

$$\frac{A \wedge B \rightarrow C}{A \rightarrow C \Leftarrow B}$$

However, at a higher level the horizontal bar functions as a deduction symbol and Theorem 1 plays the role of a new deduction theorem.

*Proof.* Clearly, every polynomial  $\varphi(x)$  must have one of the five forms:  $k$ ,  $x$ ,  $\langle \psi(x), \chi(x) \rangle$ ,  $\chi(x)\psi(x)$ , or  $\psi(x)^*$ , where  $k$  is constant and where  $\psi(x)$  and

$\chi(x)$  are shorter polynomials. By inductional assumption we have, omitting subscripts:

$$\psi(x) \equiv_{\bar{x}} g\langle x0, 1 \rangle, \chi(x) \equiv_{\bar{x}} h\langle x0, 1 \rangle.$$

The result now follows by verifying the following equations:

$$\begin{aligned} k\pi'_{A,B}\langle x0_B, 1_B \rangle &\equiv_{\bar{x}} k, \\ \pi_{A,1}\langle x0_1, 1_1 \rangle &\equiv_{\bar{x}} x, \\ \langle g, h \rangle \langle x0_B, 1_B \rangle &\equiv_{\bar{x}} \langle \psi(x), \chi(x) \rangle, \\ h\langle \pi_{A,B}, g \rangle \langle x0_B, 1_B \rangle &\equiv_{\bar{x}} \chi(x)\psi(x), \\ (g\alpha_{A,B,D})^* \langle x0_B, 1_B \rangle &\equiv_{\bar{x}} \psi(x)^*, \end{aligned}$$

where  $\alpha_{A,B,D} : (A \wedge B) \wedge D \longrightarrow A \wedge (B \wedge D)$  is given by

$$\alpha_{A,B,D} = \langle \pi_{A,B}, \pi_{A \wedge B, D}, \langle \pi'_{A,B} \pi_{A \wedge B, D}, \pi'_{A \wedge B, D} \rangle \rangle$$

The last equation is proved by showing (omitting subscripts) that:

$$\varepsilon \langle (g\alpha)^* \langle x0, 1 \rangle \pi, \pi' \rangle \equiv_{\bar{x}} g\langle x0, 1 \rangle$$

which follows from a routine calculation, as in (Lambek 1974, p. 272). □

## References