

The Cepheid Distance to the Seyfert 1 Galaxy NGC 4151

W. YUAN,¹ M. M. FAUSNAUGH,^{2,3} S. L. HOFFMANN,⁴ L. M. MACRI,⁵ B. M. PETERSON,^{2,6,4} A. G. RIESS,^{1,4}
M. C. BENTZ,⁷ J. S. BROWN,⁸ E. DALLA BONTÀ,^{9,10} R. I. DAVIES,¹¹ G. DE ROSA,⁴ L. FERRARESE,¹² C. J. GRIER,^{2,13}
E. K. S. HICKS,¹⁴ C. A. ONKEN,^{15,16} R. W. POGGE,^{2,6} T. STORCHI-BERGMANN,¹⁷ AND M. VESTERGAARD^{18,13}

¹Department of Physics & Astronomy, Johns Hopkins University, Baltimore, MD 21218, USA

²Department of Astronomy, The Ohio State University, 140 W 18th Ave, Columbus, OH 43210, USA

³Kavli Institute for Space and Astrophysics Research, Massachusetts Institute of Technology,
77 Massachusetts Avenue, Cambridge, MA 02139, USA

⁴Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁵George P. and Cynthia W. Mitchell Institute for Fundamental Physics and Astronomy, Department of Physics and Astronomy,
Texas A&M University, College Station, TX 77843, USA

⁶Center for Cosmology and AstroParticle Physics, The Ohio State University, 191 West Woodruff Ave, Columbus, OH 43210, USA

⁷Department of Physics and Astronomy, Georgia State University, 25 Park Place, Suite 605, Atlanta, GA 30303, USA

⁸UCO/Lick Observatory, University of California, Santa Cruz, CA 95064, USA

⁹Dipartimento di Fisica e Astronomia “G. Galilei,” Università di Padova, Vicolo dell’Osservatorio 3, I-35122 Padova, Italy

¹⁰INAF-Osservatorio Astronomico di Padova, Vicolo dell’Osservatorio 5 I-35122, Padova, Italy

¹¹Max Planck Institut für extraterrestrische Physik, Postfach 1312, D-85741, Garching, Germany

¹²NRC Herzberg Astronomy and Astrophysics, National Research Council, 5071 West Saanich Road, Victoria, BC V9E 2E7, Canada

¹³Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ 85721, USA

¹⁴Department of Physics and Astronomy, University of Alaska Anchorage, AK 99508, USA

¹⁵Research School of Astronomy and Astrophysics, Australian National University, Canberra, ACT 2611, Australia

¹⁶Australian Research Council (ARC) Centre of Excellence in All-sky Astrophysics (CAASTRO)

¹⁷Departamento de Astronomia, Universidade Federal do Rio Grande do Sul, Av. Bento Goncalves 9500, 91501 Porto Alegre, RS, Brazil

¹⁸Dark Cosmology Centre, Niels Bohr Institute, University of Copenhagen, Lyngbyvej 2, 4. sal., 2100 Copenhagen Ø, Denmark

ABSTRACT

We derive a distance of 16.1 ± 0.5 Mpc to the archetypical Seyfert 1 galaxy NGC 4151 based on the near-infrared Cepheid Period–Luminosity relation and new *Hubble Space Telescope* multiband imaging. This distance determination, based on measurements of 35 long-period ($P > 25$ d) Cepheids, will support the absolute calibration of the supermassive black hole mass in this system, as well as studies of the dynamics of the feedback or feeding of its active galactic nucleus.

1. INTRODUCTION

Local active galactic nuclei (AGNs) provide unique laboratories to study the co-evolution of galaxies and their supermassive black holes (SMBHs). At redshifts $z < 0.005$ ($D \lesssim 20$ Mpc), the host galaxies are well resolved, enabling detailed studies of the structure and dynamics of the their interstellar gas and any connection to the AGN through feedback or feeding. Additionally, since the sphere of influence of the central supermassive black holes is spatially resolved in these local systems, the black hole mass can be measured using resolved dynamical methods. Local AGNs, therefore, are not only central to our understanding of SMBH scaling relations in AGNs, but are also crucial in calibrating SMBH mass measurements based on methods (e.g., reverberation mapping) applicable at higher redshifts (see Ferrarese & Ford 2005; McConnell & Ma 2013; Sahu et al. 2019; Zubovas & King 2019 for reviews).

However, these local AGNs are subject to a single systematic uncertainty that makes all physical parameters highly uncertain: their distances. With accurate knowledge of the local Hubble Constant H_0 (currently better than a few percent, though tension with results from the cosmic microwave background and baryon acoustic oscillations remains; Riess et al. 2019), the uncertainty of extragalactic distances is principally limited by peculiar velocities relative to the Hubble flow. At very low redshifts, the peculiar velocities can be large compared to the cosmological expansion rate. Even for galaxies as far away as 20 Mpc, peculiar velocities can introduce errors of 20% or more in their redshift-based distances and, of course, the problem is more severe for nearer galaxies.

Redshift-independent distances for the late-type hosts of local AGNs are available only through the Tully–Fisher relation (Tully & Fisher 1977), but nevertheless are typically uncertain by an irreducible 20%, and more

for galaxies at lower inclination. Without an accurate distance, measurements of many other parameters are of limited utility, hampering our ability to understand the impact of the AGN on the evolution of the host galaxy. Thus, we find ourselves in a paradoxical situation for these local AGNs: even while they are the best systems for detailed studies of the structure, dynamics, and energetics (including luminosity measures) of the AGN and host galaxy, these are precisely the objects for which a lack of accurate distances makes all physical parameters highly uncertain.

Understanding the masses of SMBH in nearby AGNs also depends on accurate distance measurements. AGN black hole masses are generally best determined by reverberation mapping (RM; Peterson 2014). As it is usually applied, RM yields black hole mass uncertainties at $\sim 0.3\text{--}0.4$ dex, but careful modeling of high-quality data can reduce the uncertainties to 0.1–0.2 dex (Pancoast et al. 2014b; Grier et al. 2017) and are distance-independent. However, verification of RM-based masses requires independent measurements using other techniques such as modeling stellar or gas dynamics and, more recently, interferometry (Gravity Collaboration et al. 2018), that are distance dependent and, for the most part, applicable only to the nearest AGNs. Accurate distances to nearby AGNs are thus crucial for verification of RM-based masses by other techniques, particularly as the mass measurements themselves become increasingly precise.

For galaxies out to ~ 50 Mpc, Cepheid variables provide an excellent means of distance determination, enabled by the Period–Luminosity Relation (PLR) or Leavitt Law (Leavitt & Pickering 1912). Recently, Bentz et al. (2019) applied this method at optical wavelengths to the Seyfert 1 galaxy NGC 6814, in order to compare the SMBH mass inferred from dynamical modeling and reverberation mapping. In this contribution, we employ Cepheid variables to determine the distance to the well-known nearby AGN NGC 4151.

NGC 4151 is one of the best-studied AGNs at all wavelengths (e.g., Ulrich 2000). Yet, despite the intensive scrutiny that it has received on account of its proximity, we do not know its distance to any acceptable degree of accuracy; indeed, the most extreme distance estimates differ by more than a factor of seven. The Extragalactic Distance Database (Tully et al. 2009) quotes a distance for NGC 4151 of 11.2 Mpc based on the average distances of its fellow group members. Only four galaxies contribute to this group-averaged distance, however, with individual distances ranging from 3.9 to 34 Mpc based on the Tully–Fisher line width–luminosity correlation. The object with the smallest estimated distance

is NGC 4151 itself at 3.9 Mpc; however, the total galaxy luminosity has not been corrected for the large contribution from the AGN itself, which would cause the galaxy to appear brighter, and therefore nearer, than it actually is. A distance of 29.2 ± 0.4 Mpc is based on dust reverberation (Yoshii et al. 2014), and one other measurement of 19 ± 2 Mpc is based on interband continuum lags and a thermal reprocessing model (Cackett et al. 2007). By combining an IR interferometric angular size measurement with a dust reverberation result, Höning et al. (2014) infer a distance of $19.0^{+2.4}_{-2.6}$ Mpc. Recently, Tsvetkov et al. (2019) analyzed the type II-P supernova 2018aoq, obtaining a distance of 20.0 ± 1.6 Mpc using the expanding photosphere method and 16.6 ± 1.1 Mpc using the standard candle method. To further illustrate the confused situation, distances to NGC 4151 adopted in recent works include 13.2 Mpc (Hicks & Malkan 2008), 13.3 Mpc (Storchi-Bergmann et al. 2010; Iserlohe et al. 2013), 13.9 Mpc (Onken et al. 2014; De Rosa et al. 2018), 14.9 Mpc (Esquej et al. 2014), 16.6 Mpc (Bentz et al. 2013), 16.9 Mpc (Burtscher et al. 2013), and 17.6 Mpc (Kishimoto et al. 2011).

An uncertain distance to NGC 4151 limits our understanding of the feeding and feedback processes that we can otherwise study in great detail. For example, the geometry, kinematics, and excitation (e.g., Winge et al. 1999; Das et al. 2005; Shimono et al. 2010; Storchi-Bergmann et al. 2009) of the spatially extended narrow-line region have been well-characterized. AGN-driven outflows have been mapped by Storchi-Bergmann et al. (2010), using near-infrared integral field spectroscopy, down to 8 pc from the nucleus. The UV absorption spectrum (e.g., Crenshaw & Kraemer 2007) demonstrates that there is considerable kinetic energy in outflows. The calculated mass outflow rate, as well as the power of the outflow, depend on the luminosity of the emitting gas as well as on the geometry of the outflow, both strongly dependent on the distance to the galaxy. It also appears that energetic feedback is occurring on large, resolvable scales. For example, Wang et al. (2010) study the X-ray emission that fills the “H I cavity” in the central ~ 2 kpc of NGC 4151 and consider four separate hypotheses on the origin of the X-rays and the energetics of the interaction between the AGN and the host galaxy. These estimates of the energetics depend linearly on distance, but they still constitute some of the best measurements of AGN feedback in any galaxy in the observable universe.

Finally, NGC 4151 is one of the few AGNs where the mass of the supermassive black hole has been measured by more than one technique: from stellar dynamics, $M_{\text{BH}} = 3.76 \pm 1.15 \times 10^7 M_{\odot}$ (Onken et al.

2014); from gas dynamics, $M_{\text{BH}} = 3.0^{+0.75}_{-2.2} \times 10^7 M_{\odot}$ (Hicks & Malkan 2008); and from reverberation mapping, $M_{\text{BH}} = 2.14^{+0.75}_{-0.55} \times 10^7 M_{\odot}$, with an additional systematic uncertainty of ~ 0.15 dex (De Rosa et al. 2018). The precision of the reverberation measurement is likely to improve with detailed modeling.

In this work, we derive the distance to NGC 4151 based on one of the most accurate and widely-used methods, the Cepheid PLR, using data obtained with the *Hubble Space Telescope* (*HST*). The rest of this article is organized as follows: in §2, we detail our observations, data reduction, and photometry; in §3, we describe the Cepheid search; we present our results in §4 and discuss our findings in §5.

2. OBSERVATIONS, DATA REDUCTION, AND PHOTOMETRY

We acquired multiband time-series images of NGC 4151 using the Wide Field Camera 3 (WFC3) onboard *HST*. The galaxy's optically bright disk fits well within the field of view of both the UVIS and IR detectors of this instrument. Thanks to the exceptional resolving power of *HST*, hundreds of thousands of bright stars, including long-period Cepheids, can be photometrically measured in the images. A color composite of NGC 4151 from these *HST* observations is shown in Fig. 1. We reduced these *HST* data and performed photometry using DAOPHOT (Stetson 1987) and related programs. We emphasize that the analysis methods used in this work are independent from (and should not be confused with) the methods used by the SH0ES project (Hoffmann et al. 2016; Riess et al. 2016).

2.1. Observations

In order to search for Cepheids and measure their periods, we made use of the long-pass optical filter *F350LP* to phase the light curves with 12 epochs across a baseline of 74 days, in the middle of which the telescope's orientation angle changed once. The *F350LP* filter transmission curve is significantly broader in wavelength than the combination of *F555W* and *F814W* filters, boosting the Cepheid discovery efficiency by a factor of ~ 2.5 compared to the filters used traditionally (Hoffmann et al. 2016). We also observed NGC 4151 in *F555W* (3 epochs), *F814W* (3 epochs), and *F160W* (6 epochs), to obtain well-calibrated Cepheid PLRs based on combinations of these filters (Riess et al. 2016). All these observations were dithered to enable hot-pixel and/or cosmic-ray rejection. The per-epoch exposure times were 1050, 1100, 1100, and 1106 s for *F350LP*, *F555W*, *F814W*, and *F160W*, respectively. A detailed observation log is presented in Table 1.

Table 1. Observation Log

Epoch	MJD	Dither×Exposure Time (seconds)			
		<i>F350LP</i>	<i>F555W</i>	<i>F814W</i>	<i>F160W</i>
1	57363.1	3×350	2×553
2	57370.7	3×350	2×550
3	57377.9	3×350	...	2×550	...
4	57384.7	3×350	2×553
5	57390.0	3×350	2×550
6	57392.8	3×350	2×553
7	57399.1	3×350	...	2×550	...
8	57405.9	3×350	2×553
9	57414.7	3×350	2×550
10	57423.6	3×350	2×553
11	57429.5	3×350	...	2×550	...
12	57436.9	3×350	2×553

2.2. Data Reduction

We retrieved calibrated images (flat-fielded, dark-subtracted, charge-transfer efficiency corrected, etc.) from the Mikulski Archive for Space Telescopes (MAST), then registered and drizzled them using the AstroDrizzle v2.2.6 package. All the optical images (*F350LP*, *F555W*, and *F814W*) were iteratively registered to the first *F350LP* image, while all the NIR images were registered to the first *F160W* image due to the different optics and pixel scales of the IR and UVIS detectors. We drizzle-combined the aligned images of each band into deep master images, with cosmic rays rejected for the optical bands. We also produced a single drizzled image for each epoch, which serves as the basis for the time-series photometry. We set the drizzle pixel scales to the native $0.04''$ for the optical images, and a 1/3-finer-than-native (i.e., $0.08''/\text{pixel}$) for the *F160W* ones.

We modeled the surface brightness gradients of all the drizzled images and subtracted these models prior to carrying out the photometry. NGC 4151 exhibits a high surface brightness in a large central region of the disk and a steep gradient in the outskirts, which hampers the ability of DAOPHOT to detect sources and accurately fit PSFs. We masked all point sources found in each image at $> 7\sigma$ significance, performed thin-plate spline fits to the background pixel counts across the image using the R implementation of the `Tps` function with default settings, and obtained a smooth surface brightness model. Fig. 2 shows the results for *F350LP* as an example. These models were also used for local crowding estimation for Cepheids, as detailed below.




Figure 1. Color image of NGC 4151 using our master frames in $F350LP$, $F555W$, $F814W$, and $F160W$ as the luminance layer and the blue, green, and red channels, respectively. A square-root scaling was applied to all layers to increase visibility of the outer disk. Color is only shown where all layers overlap, while grayscale is used in areas not covered in $F160W$.




Figure 2. The $F350LP$ surface brightness model shows large gradient and complex variations of the background level across NGC 4151, in units of count rate per square arcsecond. Note the z -axis is plotted using a log scale.

2.3. Photometry

Given their superior depth and resolution, the $F350LP$ data were used to derive the positions of point sources. This information was propagated to other bands for fixed-position photometry. This strategy not only allowed us to track stars in different bands with the same set of identifiers (IDs), but also provided more accurate flux measurements for other bands, especially in $F160W$ where the resolution is much worse. We summarize below the photometry procedures for each band.

$F350LP$ —We started by creating a source list from a two-pass 3σ source detection on the $F350LP$ master image using DAOPHOT and ALLSTAR (Stetson 1994), then performed simultaneous PSF photometry on the 12 epochs of $F350LP$ images using ALLFRAME (Stetson 1994). In this step, we allowed both the stellar positions and flux to vary until a global PSF fit of all the sources converged. We excluded from the photometry results spurious sources associated with the stray light of foreground bright stars or background extended sources, as well as stars with exceptionally large (top 0.38%) measurement errors for their magnitudes. We selected 57 bright, isolated, non-variable stars as secondary standards to calibrate the epoch-to-epoch magnitude zero-

Table 2. Secondary Standards

ID	R.A.	Decl.	$F350LP$	$F555W$	$F814W$
			[J2000]	[mag(mmag)]	
79502	182.61050	39.41246	26.237(17)	27.647(23)	27.210(22)
73817	182.61060	39.41312	25.443(18)	27.023(10)	27.046(28)
41386	182.61072	39.41676	24.514(25)	26.120(14)	25.980(19)
76745	182.61148	39.41319	24.180(18)	25.788(12)	25.006(10)
59463	182.61307	39.41580	25.214(12)	27.507(20)	25.217(14)
37771	182.61457	39.41887	24.647(26)	26.168(17)	25.918(29)
158737	182.61525	39.40575	23.987(21)	25.580(13)	25.294(17)
149856	182.61543	39.40682	25.981(18)	27.548(15)	26.883(35)
35238	182.61730	39.42037	25.395(16)	26.940(19)	26.669(27)
118806	182.61890	39.41185	25.562(12)	27.048(16)	26.451(26)
60500	182.61891	39.41829	24.209(23)	25.890(11)	25.572(16)
202324	182.61935	39.40209	26.180(19)	27.798(18)	27.990(50)

NOTE—Coordinates are based on the WCS solution of the first $F350LP$ image. Uncertainties are given in parentheses and are expressed in units of mmag. This table is available in its entirety in machine-readable form.

point offsets. Their positions and calibrated Vega magnitudes are listed in Table 2. We extracted the $F350LP$ light curves using the TRIAL program kindly provided by Peter Stetson.

$F555W$ & $F814W$ —We performed $F555W$ and $F814W$ photometry using a modified version of ALLFRAME which fixes the stellar positions during the PSF fitting. We ran ALLFRAME on the $F350LP$, $F555W$, and $F814W$ master images, with positions fixed to those derived from the aforementioned $F350LP$ time-series photometry. This method kept the stellar centroids from drifting during the PSF fit and provided high precision $F555W$ and $F814W$ flux measurements. We based the optical magnitude calibration on the photometric measurements from this step.

$F160W$ —While the $F350LP$ stellar catalog contained a complete set of detectable sources in $F555W$ and $F814W$, it did not include a large number of redder objects that appear only in the $F160W$ image. We simultaneously fit all the point sources in the $F160W$ image by combining the list of $F350LP$ objects that were detected in $F160W$ with those that were found only in the NIR. We first transformed the positions of the $F350LP$ catalog to the $F160W$ frame and performed a preliminary PSF fit. We then subtracted any detected stars from the $F160W$ master image and obtained a residual image. After that, we ran a two-pass 3σ source detection on the residual image to find additional sources that only appeared in $F160W$. Finally, we combined the lists to form a complete $F160W$ source input catalog and performed iterative PSF fitting using ALLSTAR in fix-position mode to obtain the final $F160W$ photometry.

3. CEPHEID SEARCH

We searched for Cepheids in NGC 4151 using a combination of template fitting and visual inspection.

3.1. Variable Candidate Selection

We selected candidate variables using the widely-used Stetson (1996) L index, which measures the significance of variability in a light curve taking measurement error into account. In our case, L is reduced to

$$L = \frac{\sum_i \text{sgn}(\delta_i^2 - 1)|\delta_i^2 - 1|^{1/2}}{0.798 \cdot 12 \cdot \sqrt{n}} \cdot \frac{\sum_i |\delta_i|}{\sqrt{\sum_i \delta_i^2}}$$

$$\delta_i = \sqrt{\frac{n}{n-1}} \frac{v_i - \bar{v}}{\sigma_i},$$

where sgn is a function that returns -1 for negative values and 1 otherwise, n , v_i , σ_i , and \bar{v} are the total number of measurements, the i th magnitude, the i th photometric uncertainty, and the weighted mean magnitude, respectively, for $F350LP$. Constant stars should have $L \sim 0$, while Cepheids should exhibit positive L (Stetson 1996).

In practice, the photometric uncertainties reported by ALLFRAME are known to exhibit nonlinear and magnitude-dependent biases (as illustrated in Fig. 3) which often lead to a nonuniform shape for the variability–magnitude ($L-m$) distribution (Kalogun et al. 1998). Instead of rescaling measurement errors, we subtracted the median curve of the $L-m$ relation. The original and adjusted relations are shown in Fig. 4. We conservatively chose $L > 0.5$ and selected 6,956 variable candidates out of 403,738 objects. For reference, Stetson (1996) adopted $L > 0.9$ while Hoffmann et al. (2016) used $L > 0.6 - 0.75$.

3.2. Cepheid Template Fit

Thanks to their periodicity and predictable light curve shapes, Cepheids with sparse sampling can be accurately characterized with template-fitting methods (e.g. Stetson 1996; Tanvir et al. 2005). We made use of the Yoachim et al. (2009) V-band Cepheid templates, which are well applicable to the $F350LP$ data given the similarity in effective wavelength between these two filters.

Similar to the practice of Shappee & Stanek (2011) and Hoffmann et al. (2016), we modified the Yoachim et al. (2009) fitting method by letting the Cepheid amplitude be a free parameter. Their templates were derived from eighth-order Fourier series, whose coefficients




Figure 3. Illustration of the magnitude-dependent errors in measurement errors for nonvariable stars ($-0.1 < L < 0.1$). The vertical axis represents the ratio of reported errors and true errors, obtained by rescaling the reduced χ^2 to unity. The red points and curve indicate the median values in each magnitude bin and their spline fit, respectively.




Figure 4. The original (top) and adjusted (bottom) L index against $F350LP$ magnitudes. The red dashed lines indicate the median curves of the $L-m$ relations. The red solid line in the lower panel indicates our threshold for variable selection. Points with $L > 2.5$ are indicated by arrows. Magenta symbols indicate the Cepheids described in §3.3.

were firstly transformed into 4 leading parameters obtained from principal component analysis (PCA) and then modeled as polynomial functions of period. This yields fixed template amplitudes for a given period. Since Cepheids with the same period span a range of amplitudes (Stetson 1996) we solved for the amplitude in our template-fitting procedure, expressed as




Figure 5. Color (left) and period (right) cuts (shown as red dashed lines) imposed on the visually-identified Cepheid candidates. The variable identified by the red circle in the right panel was excluded based on its PLR residual.

$$\begin{aligned}
 m &= \bar{m} + \frac{A}{\bar{A}(P)} \times \left(\sum_{i=1}^8 \alpha_i \sin(i\Phi) + \beta_i \cos(i\Phi) \right) \\
 \Phi &= 2\pi\left(\frac{t}{P} + \phi\right) \\
 \begin{pmatrix} \alpha \\ \beta \end{pmatrix} &= \bar{\mathbf{V}} + \sum_{j=1}^4 a_j \cdot \mathbf{V}_j \\
 a_j &= \text{Polynomial}_j(\log P),
 \end{aligned}$$

where \bar{m} , A , P , and ϕ are free parameters for mean magnitude, amplitude, period, and initial phase, respectively. $\bar{A}(P)$ is a period-dependent factor that scales the amplitudes of those original templates to unity. $\bar{\mathbf{V}}$ and \mathbf{V}_j are the mean and j th PCA vector, respectively. Polynomial_j represents the j th polynomial function as shown in Fig. 8 of Yoachim et al. (2009). The allowed ranges of the free parameters were $0.01 < A < 2$ mag, $10 < P < 120$ days, and $0 < \phi < 1$.

We fit the templates to all the variable candidates and derived the best-fit parameters, despite the fact that most of them are not Cepheids. We adopted the SLSQP method (Kraft 1988) to search for the least square fit. To find the global minimum of the χ^2 , in each fit we used 1500 initial guesses for the nonlinear spaces P and ϕ , forming a meshgrid of 150 logarithmically spaced trial periods and 10 evenly spaced trial phases. We determined a template error of 0.05 mag using the OGLE Cepheid light curves (Soszynski et al. 2008), and added it to the covariance matrix in quadrature. We obtained

the best-fit “periods” and “amplitudes” for all the candidate variables.

We rejected objects exhibiting unrealistic amplitudes or periods that are longer than the observation baseline by 20%, then visually inspected the remaining light curve fits to identify Cepheid candidates. We computed the $F350LP$ amplitudes (A) of template fits, and rejected objects with $A > 1.7$ mag or $A < 0.4$ mag. Given the limited time span of the observations, we also restricted the visual selection sample to objects with “periods” shorter than 90 days. We visually inspected the light curves of 4,544 objects that survived these cuts and subjectively identified 136 Cepheid candidates that fit well to the saw-tooth shaped templates, with a philosophy that sample cleanliness is more important than completeness.

3.3. Color & Period Cuts

We inspected the $F555W - F814W$ colors and the optical Wesenheit PLR of the Cepheid candidates and applied additional cuts on color and period shown in Fig. 5. We rejected 4 candidates with $F555W - F814W > 2$ mag, which are possibly due to extreme reddening or measurement errors. We found that the scatter of the optical Wesenheit PLR was significantly larger for $P < 25$ d. Since the optical color is used in the NIR Wesenheit distance determination, we excluded 96 candidates with $P < 25$ d. We further rejected one object that is $\sim 8\sigma$ off the PLR, leaving 35 long period ($P > 25$ d) Cepheids that survived all cuts. We present their light curves in Fig. 6.




Figure 6. $F350LP$ light curves of the 35 optically identified Cepheids ordered by period. The blue curves indicate the best-fit models. We plot two cycles of pulsation for visualization purposes.

4. RESULTS

As described in §2.3, our photometry strategy allowed us to track the *F160W* Cepheid measurements by ID. We measured the *F160W* magnitudes for 32 out of the 35 optically identified Cepheids, with one of them off the *F160W* frame and two that were not detected in the *F160W* frame. We computed their crowding bias in photometry for both *F160W* and optical data using artificial star tests, and calibrated their magnitudes onto the Vega system. Finally, we obtained the optical and NIR Wesenheit PLRs of these Cepheids and derived the distance to NGC 4151.

4.1. Crowding Correction

We corrected for photometric bias including crowding through artificial star tests. “Crowding” refers to the measurement bias in crowded-field PSF photometry. This effect is location- and image-specific and highly dependent on the exact method used to carry out the photometry. It is usually characterized via artificial star tests (e.g., Stetson & Harris 1988; Gallart et al. 1996; Holtzman et al. 2006). We estimated the bias of our photometric procedure by injecting and measuring one artificial star at a time in the vicinity of each Cepheid. We repeated this process 100 times (200 times for *F160W*) in order to obtain statistically solid estimates. We also measured the dispersion of the recovered magnitudes of the artificial stars to obtain an estimate of the measurement uncertainty, which was added in quadrature to the Cepheid photometric errors. The magnitudes of artificial stars were based on the periods of their counterpart Cepheids and best-fit Cepheid PLRs, and as a result multiple realizations of the artificial star tests were performed to update the Cepheid PLRs with iteratively-determined corrections. We found mean corrections of 0.011, 0.006, 0.010, and 0.047 mag for *F350LP*, *F555W*, *F814W*, and *F160W*, respectively. We once again note that these values are highly dependent on the adopted photometry strategy and thus cannot not be compared with crowding corrections obtained using different methods and/or applied to other data sets. The only meaningful comparison should be based on fully-corrected and calibrated magnitudes for the same stars in the same images.

4.2. Magnitude Calibration

We calibrated the Cepheid magnitudes onto the Vega system by adding the aforementioned crowding correction, as well as phase sampling correction, aperture correction, and Vega zeropoints, to be discussed below.

By convention, the mean magnitude of a Cepheid refers to its intensity mean over a full cycle. However,




Figure 7. Aperture corrections in *F350LP*, *F555W*, *F814W* and *F160W* (clockwise from upper left). Circles indicate magnitude offsets between PSF and aperture photometry of those stars selected from SH0ES images (see text). Red lines are the curves of growth obtained using PySynphot. Blue lines indicates the correction to infinite aperture. Vertical dashed lines indicate the maximum radii used in the curve-of-growth fits.

the directly measured magnitudes from the drizzle combined images are sparsely sampled and do not exactly cover one cycle. We computed the offset between the directly measured magnitude and full-cycle intensity mean using the best-fit *F350LP* template for each Cepheid. For the *F555W* and *F814W* band, we scaled the amplitude of the *F350LP* light curve by the relation in Table 2 of Hoffmann et al. (2016). By simulating *F160W* light curves (sinusoidal shape assumed) of a typical 0.3 mag amplitude, we found that the *F160W* phase corrections are marginal due to the small amplitude, and thus we did not include them in the calibration.

We derived the aperture correction and Vega zero-points using the PySynphot v0.9.12 package (STScI Development Team 2013). Due to the lack of isolated bright stars in the NGC 4151 field, we derived PSF models and their aperture corrections based on *HST* images taken with a similar configuration. We selected hundreds of suitable stars from the images acquired by the SH0ES team (Riess et al. 2016), who observed Cepheids in dozens of nearby galaxies. For *F160W*, we made use of archival observations of the standard star P330E to construct the PSF model. We measured the aperture magnitudes of a series of aperture sizes using DAOPHOT, and computed their offsets against the PSF magnitudes, as shown in Fig. 7. These offsets were then fit to the curve of growth obtained from PySynphot by adjusting

Table 3. Cepheid Properties

ID	R.A.	Decl.	<i>P</i>	Ampl.	<i>F350LP</i>	<i>F555W</i>	<i>F814W</i>	<i>F160W</i>
	[J2000]		[day]	[mag]		[mag(mmag)]		
317030	182.655415	39.402869	25.098	1.012	26.274(83)	26.489(122)	25.293(102)	...
395256	182.652095	39.392541	25.412	0.914	25.535(63)	25.819(81)	24.766(93)	23.877(483)
350220	182.650011	39.396713	26.405	0.494	26.133(95)	26.450(126)	25.352(107)	24.570(558)
142091	182.649362	39.422838	26.578	0.997	25.511(48)	25.811(65)	24.809(76)	24.267(177)
285952	182.628861	39.394500	27.044	1.130	27.265(93)	27.389(160)	26.188(157)	24.896(375)

NOTE—Coordinates are based on the WCS solution of the first *F350LP* image. This table is available in its entirety in machine-readable form.




Figure 8. Cepheid locations on the *F350LP* (left) and *F160W* (right) master images. The red circles indicate Cepheids with *F160W* local surface brightness beyond 300 counts/s/sq arcsec. Magenta crosses indicate Cepheids without valid *F160W* photometry.

only the *Y*-intercept of those red curves in Fig. 7. Since the offset between the magnitude measured in a certain aperture and an infinite aperture is fixed, and the offset between instrumental and Vega magnitudes is well established for these instruments and filters, we were able to convert the instrumental PSF magnitudes to infinite-aperture Vega magnitudes. We adopted a conservative 0.02 mag systematic uncertainty since we used the average of many calibrating stars in the aforementioned SH0ES fields to calibrate a different field, NGC 4151.

We list the properties and fully calibrated magnitudes of the final Cepheid sample in Table 3. Their locations in the *F350LP* and *F160W* images are shown in Fig. 8.

4.3. Cepheid PLRs

We adopted the Wesenheit indices and Cepheid PLR calibrations of Riess et al. (2019), who used the same combination of filters. Since the crowding level varies significantly across our field, we inspected the PLR residuals against crowding using local surface brightness (*SB*) as an indicator, as shown in Fig. 9. We observed an increased scatter at high SB levels, and thus restricted our sample to objects with $SB(F160W) < 300$ counts/s/sq arcsec for the W^H PLR. This surface brightness cut reduced the scatter in the W^H PLR by 23%. We present the final optical (W^I) and NIR (W^H) PLRs in Fig. 10 and in Table 4. We note that the *F160W* count-rate non-linearity correction is included in the LMC calibration of Riess et al. (2019).




Figure 9. W^H PLR for all Cepheids with NIR measurements (left) and their residuals against local surface brightness (SB, right). The black solid and dashed lines indicate the PLR fit and $\pm 1\sigma$ scatter, respectively. The red points indicate Cepheids rejected by the SB cut, indicated by the vertical red line.




Figure 10. W^I -band (left) and W^H -band (right) PLRs of the final Cepheid sample. The black solid and dashed lines indicate the PLR fit and $\pm 1\sigma$ scatter, respectively. The distance moduli are based on the LMC PLRs of Riess et al. (2019) and the LMC distance determined by Pietrzyński et al. (2019).

Table 4. Cepheid PLRs

Index	Expression	Slope	Intercept		Scatter	N	
			LMC	NGC 4151			
W^I	$F814W - 1.3$	($F555W - F814W$)	-3.31	15.935	28.426	0.348	35
W^H	$F160W - 0.386$	($F555W - F814W$)	-3.26	15.898	28.449	0.295	24

4.4. Distance to NGC 4151

We adopted the LMC as the single anchor for this work, as both its Cepheid PLRs and absolute distance have the highest precision to date (Riess et al. 2019; Pietrzynski et al. 2019). We fixed the slopes of the PLRs to those derived by Riess et al. (2019) and computed the PLR intercept offsets between the LMC and NGC 4151. Using the W^H PLR, we derived a relative distance modulus of $\Delta\mu = 12.55 \pm 0.07$ mag. Adding the geometrically determined LMC distance modulus of $\mu = 18.477$ mag (Pietrzynski et al. 2019), we obtained a distance modulus for NGC 4151 of $\mu = 31.03 \pm 0.07$ mag ($D = 16.07 \pm 0.53$ Mpc). We also calculated the distance modulus based on the optical measurements only (W^I PLR), which yielded $\mu = 30.97 \pm 0.08$ mag and is consistent with the NIR-based distance.

The leading error sources in our distance determination are: the random error due to PLR scatter, the uncertainty in the absolute distance to the LMC, the aperture corrections (see §4.2) and the metallicity dependence of Cepheid PLRs. To estimate the latter, we multiplied the scatter of [O/H] measurements in 21 galaxies presented by Riess et al. (2016) by the metallicity term coefficient in Equation 2 of Riess et al. (2016). We note that uncertainties due to the choice of reddening law are marginal. We present the full error budget in Table 5.

5. DISCUSSION

5.1. Black Hole Masses

One of the primary reasons for undertaking this investigation has been to facilitate comparisons among various methods of determining the masses of the central black holes in these galaxies. NGC 4151 and NGC 3227 are among a handful of galaxies for which the black hole radius of influence $r_{\text{BH}} = GM_{\text{BH}}/\sigma_*^2$, where σ_* is the stellar velocity dispersion, is resolved so masses based on stellar or gas dynamical modeling have high credibility. An accurate distance is required since masses based on stellar or gas dynamics are distance dependent. Masses based on reverberation mapping are, by contrast, distance-independent. Reverberation-based masses are computed from

$$M_{\text{BH}} = f \left(\frac{\Delta V^2 c \tau}{G} \right), \quad (1)$$

where τ is the reverberation time lag between continuum and emission-line flux variations, ΔV is the emission line width, and f is a dimensionless scaling factor of order unity that depends on the inclination, geometry,

Table 5. Error Budget

Source	W^I (mag)	W^H (mag)
PLR scatter	0.06	0.06
LMC distance	0.026	0.026
Magnitude calibration	0.02	0.02
Metallicity	0.03	0.02
Total	0.075	0.071

and kinematics of the broad emission-line region. The scale factor f can be determined by detailed modeling of reverberation data (e.g., Pancoast et al. 2014a; Grier et al. 2017; Horne et al. 2020) and seems to be most sensitive to inclination (Grier et al. 2017). In the absence of detailed modeling, it is customary to use an average scaling factor (Onken et al. 2004) that is determined by assuming that both quiescent and active galaxies follow the same relationship between black hole mass M_{BH} and bulge velocity dispersion σ_* (Ferrarese & Merritt 2000; Gebhardt et al. 2000; Gultekin et al. 2009). The most recent calibration gives $\langle \log f \rangle = 0.683$ (Batiste et al. 2017) with an error on the mean of 0.030 dex and a standard deviation of 0.150 dex. Since detailed reverberation models of NGC 3227 and NGC 4151 are not yet available, we use this value here, and note that this introduces a systematic uncertainty of ~ 0.150 dex.

In Table 6, we show various mass determinations for both NGC 3227 and NGC 4151 along with assumed distances for each investigation. Our preferred distance to NGC 3227 is based on a surface brightness fluctuation measurement of its interacting elliptical companion, NGC 3226, as determined by Tonry et al. (2001) with a correction by Blakeslee et al. (2010). Table 6 also shows mass measurements for the central black hole in both galaxies adjusted to our preferred distances of 23.7 Mpc for NGC 3227 and 16.1 Mpc for NGC 4151.

The mass measurements for the SMBH in NGC 4151 include a recent determination based on gas dynamics using integral field spectroscopy (Hicks et al., in preparation) and the various techniques applied to this system are in excellent agreement. On the other hand, even with the currently large systematic uncertainty in the reverberation-based masses, the corresponding estimate for NGC 3227 is much lower than the values based on stellar or gas dynamics. However, these differences can be understood as an inclination effect; indeed, unification models (Antonucci 1993) predict that Type 1

Table 6. Black Hole Mass Comparison

Source	Ref.	Method	D (Mpc)	$\log M/M_\odot$
(1)	(2)	(3)	(4)	(5)
NGC 3227	1	stars	17.0	$7.18^{+0.12}_{-0.33}$
NGC 3227	2	gas	15.6	$7.30^{+0.18}_{-0.10}$
NGC 4151	3	stars	13.9	$7.58^{+0.12}_{-0.16}$
NGC 4151	2	gas	13.2	$7.48^{+0.10}_{-0.58}$
NGC 3227	1,4,5	stars	23.7	$7.32^{+0.12}_{-0.33}$
NGC 3227	2,4,5	gas	23.7	$7.48^{+0.18}_{-0.10}$
NGC 3227	6	RM	...	$6.65 \pm 0.20 (\pm 0.15)$
NGC 4151	1,7	stars	16.1	$7.64^{+0.12}_{-0.16}$
NGC 4151	2,7	gas	16.1	$7.56^{+0.10}_{-0.58}$
NGC 4151	7,8	gas	16.1	$7.68^{+0.09}_{-0.11}$
NGC 4151	6	RM	...	$7.61 \pm 0.05 (\pm 0.15)$

NOTE—Columns are: 1: Source; 2: References; 3: “stars” refers to stellar dynamical modeling, “gas” refers to gas dynamical modeling, “RM” refers to reverberation mapping; 4: Distance assumed; 5: Black hole mass.

The top part of this table contains the original mass measurements assuming the distances listed, while the two lower parts show the adjusted masses.

References— 1: Davies et al. (2006); 2: Hicks & Malkan (2008); 3: Onken et al. (2014); 4: Tonry et al. (2001) (distance); 5: Blakeslee et al. (2010) (distance correction); 6: De Rosa et al. (2018); 7: this work (distance); 8: Hicks et al., in preparation (mass).

AGNs, like NGC 3227 and NGC 4151, are preferentially at lower rather than higher inclination. If $f \propto 1/\sin^2 i$, which is appropriate for a thin disk, this would imply $i \approx 11^\circ$, which is consistent with the 15° inclination of the narrow-line region (Fischer et al. 2013, though note in their paper the inclination given is the complement of the standard definition), but not with the less-reliable inclination of $33^\circ \pm 2^\circ$ inferred from the Fe K α profile (Patrick et al. 2012). However, the inclination of the narrow-line region in NGC 4151 is only 21° (Winge et al. 1999), so a similar inclination correction introduces some tension in the mass comparison. The conclusion we draw from this is that velocity-resolved reverberation mapping is required for a meaningful mass comparison in individual sources because the unknown geometry and inclination of the BLR makes reverberation-based mass measurement using only lag and line width too uncertain.

5.2. Other Implications

We noted in the Introduction that a wide range of distances have been reported for NGC 4151. The distance of $19.0^{+2.4}_{-2.6}$ Mpc derived by Hönig et al. (2014) by combining near-IR interferometry with dust reverberation is $\sim 18\%$ larger than our Cepheid-based distance, but statistically consistent. On the other hand, the dust-

reverberation distance of 29.2 ± 0.4 Mpc (Yoshii et al. 2014) is nearly a factor of two larger than our measurement, probably because at least one assumption in the theoretical prediction of the dust torus radius is incorrect; indeed, of the 17 objects studied, only two of 16 yielded distances consistent within the errors with the standard luminosity distance (we will discuss a Cepheid-based distance for the remaining source, NGC 4051, in a subsequent paper).

The improved distance allowed us to improve both mass and energetics in one of the nearest and best targets to study the physics of active nuclei on small scales. For example, Storchi-Bergmann et al. (2009) derived masses of ionized and hot molecular gas. These masses scale as D^2 , and our revised distance thus increases these masses by 46%. Similarly, the mass of ionized gas in the outflow that was calculated as $3.1 \times 10^6 M_\odot$ becomes $4.6 \times 10^6 M_\odot$. Storchi-Bergmann et al. (2010) derived the mass outflow rate and outflow power. These both scale with D , because the rates are calculated at a certain distance from the nuclear source; thus these quantities are proportional to D^2/D , and therefore should be increased by 21%.

6. SUMMARY

We obtained multiband time-series observations of the archetypical Seyfert 1 galaxy NGC 4151 with the *Hubble Space Telescope* and identified over 130 Cepheid candidates in this system. We used a subset of 35 long-period ($P > 25$ d) Cepheids to derive a distance of 16.1 ± 0.5 Mpc using the Near-Infrared Cepheid Period–Luminosity Relation.

The precise Cepheid-based distance enabled us to update previous dynamical mass determinations for the SMBH in this AGN, which yield consistent values of $\log M/M_\odot \sim 7.6$ dex. These, in turn, are in good agreement with mass estimates based on reverberation mapping.

ACKNOWLEDGMENTS

Support for *HST* program GO-13765 was provided by NASA through a grant from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. MV gratefully acknowledges support from the Independent Research Fund Denmark via grant number DFF 8021-00130. We thank K.Z. Stanek for helpful advice early in this project.

REFERENCES

- Antonucci, R. 1993, ARA&A, 31, 473,
doi: [10.1146/annurev.aa.31.090193.002353](https://doi.org/10.1146/annurev.aa.31.090193.002353)
- Batiste, M., Bentz, M. C., Raimundo, S. I., Vestergaard, M., & Onken, C. A. 2017, ApJL, 838, L10,
doi: [10.3847/2041-8213/aa6571](https://doi.org/10.3847/2041-8213/aa6571)
- Bentz, M. C., Ferrarese, L., Onken, C. A., Peterson, B. M., & Valluri, M. 2019, ApJ, 885, 161,
doi: [10.3847/1538-4357/ab48fb](https://doi.org/10.3847/1538-4357/ab48fb)
- Bentz, M. C., Denney, K. D., Grier, C. J., et al. 2013, ApJ, 767, 149, doi: [10.1088/0004-637X/767/2/149](https://doi.org/10.1088/0004-637X/767/2/149)
- Blakeslee, J. P., Cantiello, M., Mei, S., et al. 2010, ApJ, 724, 657, doi: [10.1088/0004-637X/724/1/657](https://doi.org/10.1088/0004-637X/724/1/657)
- Burtscher, L., Meisenheimer, K., Tristram, K. R. W., et al. 2013, A&A, 558, A149,
doi: [10.1051/0004-6361/201321890](https://doi.org/10.1051/0004-6361/201321890)
- Cackett, E. M., Horne, K., & Winkler, H. 2007, MNRAS, 380, 669, doi: [10.1111/j.1365-2966.2007.12098.x](https://doi.org/10.1111/j.1365-2966.2007.12098.x)
- Crenshaw, D. M., & Kraemer, S. B. 2007, ApJ, 659, 250,
doi: [10.1086/511970](https://doi.org/10.1086/511970)
- Das, V., Crenshaw, D. M., Hutchings, J. B., et al. 2005, AJ, 130, 945, doi: [10.1086/432255](https://doi.org/10.1086/432255)
- Davies, R. I., Thomas, J., Genzel, R., et al. 2006, ApJ, 646, 754, doi: [10.1086/504963](https://doi.org/10.1086/504963)
- De Rosa, G., Fausnaugh, M. M., Grier, C. J., et al. 2018, ApJ, 866, 133, doi: [10.3847/1538-4357/aadd11](https://doi.org/10.3847/1538-4357/aadd11)
- Esquej, P., Alonso-Herrero, A., González-Martín, O., et al. 2014, ApJ, 780, 86, doi: [10.1088/0004-637X/780/1/86](https://doi.org/10.1088/0004-637X/780/1/86)
- Ferrarese, L., & Ford, H. 2005, SSRv, 116, 523,
doi: [10.1007/s11214-005-3947-6](https://doi.org/10.1007/s11214-005-3947-6)
- Ferrarese, L., & Merritt, D. 2000, ApJL, 539, L9,
doi: [10.1086/312838](https://doi.org/10.1086/312838)
- Fischer, T. C., Crenshaw, D. M., Kraemer, S. B., & Schmitt, H. R. 2013, The Astrophysical Journal Supplement Series, 209, 1,
doi: [10.1088/0067-0049/209/1/1](https://doi.org/10.1088/0067-0049/209/1/1)
- Gallart, C., Aparicio, A., & Vilchez, J. M. 1996, AJ, 112, 1928, doi: [10.1086/118153](https://doi.org/10.1086/118153)
- Gebhardt, K., Bender, R., Bower, G., et al. 2000, ApJL, 539, L13, doi: [10.1086/312840](https://doi.org/10.1086/312840)
- Gravity Collaboration, Sturm, E., Dexter, J., et al. 2018, Nature, 563, 657, doi: [10.1038/s41586-018-0731-9](https://doi.org/10.1038/s41586-018-0731-9)
- Grier, C. J., Pancoast, A., Barth, A. J., et al. 2017, ApJ, 849, 146, doi: [10.3847/1538-4357/aa901b](https://doi.org/10.3847/1538-4357/aa901b)
- Gültekin, K., Richstone, D. O., Gebhardt, K., et al. 2009, ApJ, 698, 198, doi: [10.1088/0004-637X/698/1/198](https://doi.org/10.1088/0004-637X/698/1/198)
- Hicks, E. K. S., & Malkan, M. A. 2008, The Astrophysical Journal Supplement Series, 174, 31, doi: [10.1086/521650](https://doi.org/10.1086/521650)
- Hoffmann, S. L., Macri, L. M., Riess, A. G., et al. 2016, ApJ, 830, 10, doi: [10.3847/0004-637X/830/1/10](https://doi.org/10.3847/0004-637X/830/1/10)
- Holtzman, J. A., Afonso, C., & Dolphin, A. 2006, ApJS, 166, 534, doi: [10.1086/507074](https://doi.org/10.1086/507074)
- Hönig, S. F., Watson, D., Kishimoto, M., & Hjorth, J. 2014, Nature, 515, 528, doi: [10.1038/nature13914](https://doi.org/10.1038/nature13914)
- Horne, K., De Rosa, G., Peterson, B. M., et al. 2020, arXiv e-prints, arXiv:2003.01448.
<https://arxiv.org/abs/2003.01448>
- Iserlohe, C., Krabbe, A., Larkin, J. E., et al. 2013, A&A, 556, A136, doi: [10.1051/0004-6361/201219998](https://doi.org/10.1051/0004-6361/201219998)
- Kaluzny, J., Stanek, K. Z., Krockenberger, M., et al. 1998, AJ, 115, 1016, doi: [10.1086/300235](https://doi.org/10.1086/300235)
- Kishimoto, M., Hönig, S. F., Antonucci, R., et al. 2011, A&A, 527, A121, doi: [10.1051/0004-6361/201016054](https://doi.org/10.1051/0004-6361/201016054)
- Kraft, D. 1988, Forschungsbericht- Deutsche Forschungs- und Versuchsanstalt fur Luft- und Raumfahrt
- Leavitt, H. S., & Pickering, E. C. 1912, Harvard College Observatory Circular, 173, 1
- McConnell, N. J., & Ma, C.-P. 2013, ApJ, 764, 184,
doi: [10.1088/0004-637X/764/2/184](https://doi.org/10.1088/0004-637X/764/2/184)
- Onken, C. A., Ferrarese, L., Merritt, D., et al. 2004, ApJ, 615, 645, doi: [10.1086/424655](https://doi.org/10.1086/424655)
- Onken, C. A., Valluri, M., Brown, J. S., et al. 2014, ApJ, 791, 37, doi: [10.1088/0004-637X/791/1/37](https://doi.org/10.1088/0004-637X/791/1/37)
- Pancoast, A., Brewer, B. J., & Treu, T. 2014a, MNRAS, 445, 3055, doi: [10.1093/mnras/stu1809](https://doi.org/10.1093/mnras/stu1809)
- Pancoast, A., Brewer, B. J., Treu, T., et al. 2014b,
MNRAS, 445, 3073, doi: [10.1093/mnras/stu1419](https://doi.org/10.1093/mnras/stu1419)
- Patrick, A. R., Reeves, J. N., Porquet, D., et al. 2012,
MNRAS, 426, 2522,
doi: [10.1111/j.1365-2966.2012.21868.x](https://doi.org/10.1111/j.1365-2966.2012.21868.x)
- Peterson, B. M. 2014, SSRv, 183, 253,
doi: [10.1007/s11214-013-9987-4](https://doi.org/10.1007/s11214-013-9987-4)
- Pietrzyński, G., Graczyk, D., Gallenne, A., et al. 2019, Nature, 567, 200, doi: [10.1038/s41586-019-0999-4](https://doi.org/10.1038/s41586-019-0999-4)
- Riess, A. G., Casertano, S., Yuan, W., Macri, L. M., & Scolnic, D. 2019, ApJ, 876, 85,
doi: [10.3847/1538-4357/ab1422](https://doi.org/10.3847/1538-4357/ab1422)
- Riess, A. G., Macri, L. M., Hoffmann, S. L., et al. 2016,
ApJ, 826, 56, doi: [10.3847/0004-637X/826/1/56](https://doi.org/10.3847/0004-637X/826/1/56)
- Sahu, N., Graham, A. W., & Davis, B. L. 2019, ApJ, 887, 10, doi: [10.3847/1538-4357/ab50b7](https://doi.org/10.3847/1538-4357/ab50b7)
- Shappee, B. J., & Stanek, K. Z. 2011, ApJ, 733, 124,
doi: [10.1088/0004-637X/733/2/124](https://doi.org/10.1088/0004-637X/733/2/124)
- Shimono, A., Sugai, H., Kawai, A., et al. 2010, Publications of the Astronomical Society of Japan, 62, 225,
doi: [10.1093/pasj/62.2.225](https://doi.org/10.1093/pasj/62.2.225)
- Soszynski, I., Poleski, R., Udalski, A., et al. 2008, AcA, 58, 163. <https://arxiv.org/abs/0808.2210>

- Stetson, P. B. 1987, Publications of the Astronomical Society of the Pacific, 99, 191, doi: [10.1086/131977](https://doi.org/10.1086/131977)
- . 1994, Publications of the Astronomical Society of the Pacific, 106, 250, doi: [10.1086/133378](https://doi.org/10.1086/133378)
- . 1996, Publications of the Astronomical Society of the Pacific, 108, 851, doi: [10.1086/133808](https://doi.org/10.1086/133808)
- Stetson, P. B., & Harris, W. E. 1988, AJ, 96, 909, doi: [10.1086/114856](https://doi.org/10.1086/114856)
- Storchi-Bergmann, T., Lopes, R. D. S., McGregor, P. J., et al. 2010, MNRAS, 402, 819, doi: [10.1111/j.1365-2966.2009.15962.x](https://doi.org/10.1111/j.1365-2966.2009.15962.x)
- Storchi-Bergmann, T., McGregor, P. J., Riffel, R. A., et al. 2009, MNRAS, 394, 1148, doi: [10.1111/j.1365-2966.2009.14388.x](https://doi.org/10.1111/j.1365-2966.2009.14388.x)
- STScI Development Team. 2013, pysynphot: Synthetic photometry software package. <http://ascl.net/1303.023>
- Tanvir, N. R., Hendry, M. A., Watkins, A., et al. 2005, MNRAS, 363, 749, doi: [10.1111/j.1365-2966.2005.09466.x](https://doi.org/10.1111/j.1365-2966.2005.09466.x)
- Tonry, J. L., Dressler, A., Blakeslee, J. P., et al. 2001, ApJ, 546, 681, doi: [10.1086/318301](https://doi.org/10.1086/318301)
- Tsvetkov, D. Y., Baklanov, P. V., Potashov, M. S., et al. 2019, MNRAS, 487, 3001, doi: [10.1093/mnras/stz1474](https://doi.org/10.1093/mnras/stz1474)
- Tully, R. B., & Fisher, J. R. 1977, A&A, 500, 105
- Tully, R. B., Rizzi, L., Shaya, E. J., et al. 2009, AJ, 138, 323, doi: [10.1088/0004-6256/138/2/323](https://doi.org/10.1088/0004-6256/138/2/323)
- Ulrich, M. H. 2000, Astronomy and Astrophysics Review, 10, 135, doi: [10.1007/s001590000007](https://doi.org/10.1007/s001590000007)
- Wang, J., Fabbiano, G., Risaliti, G., et al. 2010, ApJ, 719, L208, doi: [10.1088/2041-8205/719/2/L208](https://doi.org/10.1088/2041-8205/719/2/L208)
- Winge, C., Axon, D. J., Macchetto, F. D., Capetti, A., & Marconi, A. 1999, ApJ, 519, 134, doi: [10.1086/307337](https://doi.org/10.1086/307337)
- Yoachim, P., McCommas, L. P., Dalcanton, J. J., & Williams, B. F. 2009, AJ, 137, 4697, doi: [10.1088/0004-6256/137/6/4697](https://doi.org/10.1088/0004-6256/137/6/4697)
- Yoshii, Y., Kobayashi, Y., Minezaki, T., Koshida, S., & Peterson, B. A. 2014, ApJ, 784, L11, doi: [10.1088/2041-8205/784/1/L11](https://doi.org/10.1088/2041-8205/784/1/L11)
- Zubovas, K., & King, A. R. 2019, General Relativity and Gravitation, 51, 65, doi: [10.1007/s10714-019-2549-7](https://doi.org/10.1007/s10714-019-2549-7)