

Holographic Derivation of Entanglement Entropy from AdS/CFT

Shinsei Ryu and Tadashi Takayanagi

Kavli Institute for Theoretical Physics,

University of California, Santa Barbara, CA 93106, USA

(Dated: 7 Mar 2006)

A holographic derivation of the entanglement entropy in quantum (conformal) field theories is proposed from AdS/CFT correspondence. We argue that the entanglement entropy in $d + 1$ dimensional conformal field theories can be obtained from the area of d dimensional minimal surfaces in AdS_{d+2} , analogous to the Bekenstein-Hawking formula for black hole entropy. We show that our proposal perfectly reproduces the correct entanglement entropy in 2D CFT when applied to AdS_3 . We also compare the entropy computed in $\text{AdS}_5 \times S^5$ with that of the free $\mathcal{N} = 4$ super Yang-Mills.

I. INTRODUCTION

One of the most remarkable successes in gravitational aspects of string theory is the microscopic derivation of the Bekenstein-Hawking entropy S_{BH}

$$S_{BH} = \frac{\text{Area of horizon}}{4G_N}, \quad (1.1)$$

for BPS black holes [1]. This idea relates the gravitational entropy with the degeneracy of quantum field theory as its microscopic description. Taking near horizon limit, we can regard this as a special example of AdS/CFT correspondence [2–4]. It claims that the $d + 1$ dimensional conformal field theories (CFT_{d+1}) are equivalent to the (super)gravity on $d + 2$ dimensional anti-deSitter space AdS_{d+2} . We expect that each CFT is sitting at the boundary of AdS space.

On the other hand, there is a different kind of entropy called entanglement entropy (von-Neumann entropy) in quantum mechanical systems. The entanglement entropy

$$S_A = -\text{tr}_A \rho_A \log \rho_A, \quad \rho_A = \text{tr}_B |\Psi\rangle\langle\Psi|, \quad (1.2)$$

provides us with a convenient way to measure how closely entangled (or how “quantum”) a given wave function $|\Psi\rangle$ is. Here, the total system is divided into two subsystems A and B and ρ_A is the reduced density matrix for the subsystem A obtained by taking a partial trace over the subsystem B of the total density matrix $\rho = |\Psi\rangle\langle\Psi|$. Intuitively, we can think S_A as the entropy for an observer who is only accessible to the subsystem A and cannot receive any signals from B . In this sense, the subsystem B is analogous to the inside of a black hole horizon for an observer sitting in A , i.e., outside of the horizon. Indeed, an original motivation of the entanglement entropy was its similarity to the Bekenstein-Hawking entropy [5, 6].

The entanglement entropy is of growing importance in many fields of physics in our exploration for better understanding of quantum systems. For example, in a modern trend of condensed matter physics it has been becoming clear that quantum phases of matter need to be characterized by their pattern of entanglement encoded in many-body wave functions of ground states, rather than conventional order parameters [7–9]. Recently, the entanglement entropy has been extensively studied in low-dimensional quantum many-body systems as a new tool to investigate the nature of quantum criticality (refer to [10] and references therein for example).

For one-dimensional (1D) quantum many-body systems at criticality (i.e. 2D CFT), it is known that the entanglement entropy is given by [10, 11]

$$S_A = \frac{c}{3} \cdot \log \left(\frac{L}{\pi a} \sin \left(\frac{\pi l}{L} \right) \right), \quad (1.3)$$

where l and L are the length of the subsystem A and the total system $A \cup B$ (both ends of $A \cup B$ are periodically identified), respectively; a is a ultra violet (UV) cutoff (lattice spacing); c is the central charge of the CFT. When we are away from criticality, Eq. (1.3) is replaced by [7, 10]

$$S_A = \frac{c}{6} \cdot \mathcal{A} \cdot \log \frac{\xi}{a}, \quad (1.4)$$

where ξ is the correlation length and \mathcal{A} is the number of boundary points of A (e.g. $\mathcal{A} = 2$ in the setup of (1.3)).

In spite of these recent developments, and its similarity to the black hole entropy, a comprehensive gravitational interpretation of the entanglement entropy has been lacking so far. Here, we present a simple proposal on this issue in the light of AdS/CFT duality. Earlier discussions from different viewpoints can be found in e.g. papers [12, 13]. Define the entanglement entropy S_A in a CFT on $\mathbb{R}^{1,d}$ (or $\mathbb{R} \times S^d$) for a subsystem A that has an arbitrary $d - 1$ dimensional boundary $\partial A \in \mathbb{R}^d$ (or S^d). In this setup we propose the following ‘area law’

$$S_A = \frac{\text{Area of } \gamma_A}{4G_N^{(d+2)}}, \quad (1.5)$$

where γ_A is the d dimensional static minimal surface in AdS_{d+2} whose boundary is given by ∂A , and $G_N^{(d+2)}$ is the $d + 2$ dimensional Newton constant. Intuitively, this suggests that the minimal surface γ_A plays the role of a holographic screen for an observer who is only accessible to the subsystem A . We show explicitly the relation (1.5) in the lowest dimensional case $d = 1$, where γ_A is given by a geodesic line in AdS_3 . We also compute S_A from the gravity side for general d and compare it with field theory results, which is successful at least qualitatively. From (1.5), it is readily seen that the basic properties of the entanglement entropy (i) $S_A = S_B$ (B is the complement of A) and (ii) $S_{A_1} + S_{A_2} \geq S_{A_1 \cup A_2}$ (subadditivity) are satisfied.

We can also define the entanglement entropy at finite temperature $T = \beta^{-1}$. E.g. in a 2D CFT on a infinitely long line, it is given by replacing L in Eq. (1.3) with $i\beta$ [10]. We argue that Eq. (1.5) still holds in $T > 0$ cases. Note that $S_A = S_B$ is no




FIG. 1: (a) AdS_3 space and CFT_2 living on its boundary and (b) a geodesics γ_A as a holographic screen.

longer true if $T > 0$ since ρ is in a mixed state generically. At high temperature, we will see that this occurs due to the presence of black hole horizon in the dual gravity description.

II. ENTANGLEMENT ENTROPY IN $\text{AdS}_3/\text{CFT}_2$

Let us start with the entanglement entropy in 2D CFTs. According to AdS/CFT correspondence, gravitational theories on AdS_3 space of radius R are dual to (1+1)D CFTs with the central charge [14]

$$c = \frac{3R}{2G_N^{(3)}}. \quad (2.1)$$

The metric of AdS_3 in the global coordinate (t, ρ, θ) is

$$ds^2 = R^2 \left(-\cosh \rho^2 dt^2 + d\rho^2 + \sinh \rho^2 d\theta^2 \right). \quad (2.2)$$

At the boundary $\rho = \infty$ of AdS_3 the metric is divergent. To regulate physical quantities we put a cutoff ρ_0 and restrict the space to the bounded region $\rho \leq \rho_0$. This procedure corresponds to the UV cutoff in the dual CFTs [15]. If L is the total length of the system with both ends identified, and a is the lattice spacing (or UV cutoff) in the CFTs, we have the relation (up to a numerical factor)

$$e^{\rho_0} \sim L/a. \quad (2.3)$$

The (1+1)D spacetime for the CFT_2 is identified with the cylinder (t, θ) at the boundary $\rho = \rho_0$. The subsystem A is the region $0 \leq \theta \leq 2\pi l/L$. Then γ_A in

Eq. (1.5) is identified with the static geodesic that connects the boundary points $\theta = 0$ and $2\pi l/L$ with t fixed, traveling inside AdS_3 (Fig. 1 (a)). With the cutoff ρ_0 introduced above, the geodesic distance L_{γ_A} is given by

$$\cosh\left(\frac{L_{\gamma_A}}{R}\right) = 1 + 2 \sinh^2 \rho_0 \sin^2 \frac{\pi l}{L}. \quad (2.4)$$

Assuming the large UV cutoff $e^{\rho_0} \gg 1$, the entropy (1.5) is expressed as follows, using Eq. (2.1)

$$S_A \simeq \frac{R}{4G_N^{(3)}} \log\left(e^{2\rho_0} \sin^2 \frac{\pi l}{L}\right) = \frac{c}{3} \log\left(e^{\rho_0} \sin \frac{\pi l}{L}\right). \quad (2.5)$$

This entropy precisely coincides with the known CFT result (1.3) after we remember the relation Eq. (2.3).

This proposed relation (1.5) suggests that the geodesic (or minimal surface in the higher dimensional case) γ_A is analogous to an event horizon as if B were a black hole, though the division into A and B is just artificial. In other words, the observer, who is not accessible to B , will probe γ_A as a holographic screen [16], instead of B itself (Fig. 1 (b)). The minimal surface provides the severest entropy bound when we fix its boundary condition. In our case it saturates the bound.

More generally, we can consider a subsystem A which consists of multiple disjoint intervals as follows

$$A = \{x | x \in [r_1, s_1] \cup [r_2, s_2] \cup \dots \cup [r_N, s_N]\}, \quad (2.6)$$

where $0 \leq r_1 < s_1 < r_2 < s_2 < \dots < r_N < s_N \leq L$. In the dual AdS_3 description, the region (2.6) corresponds to $\theta \in \cup_{i=1}^N [\frac{2\pi r_i}{L}, \frac{2\pi s_i}{L}]$ at the boundary. In this case it is not straightforward to determine minimal (geodesic) lines responsible for the entropy. However, we can find the answer from the entanglement entropy computed in the CFT side. The general prescription of calculating the entropy for such systems is given in [10] using conformal mapping. For our system (2.6), we find, when rewritten in the AdS_3 language, the following expression of S_A

$$S_A = \frac{\sum_{i,j} L_{r_j, s_i} - \sum_{i < j} L_{r_j, r_i} - \sum_{i < j} L_{s_j, s_i}}{4G_N^{(3)}}, \quad (2.7)$$

where $L_{a,b}$ is the geodesic distance between two boundary points a and b . We can think that the correct definition of minimal surface is given by the numerator in Eq. (2.7).

Next we turn to the entanglement entropy at finite temperature. We assume the spacial length of the total system L is infinitely long s.t. $\beta/L \ll 1$. At high temperature, the gravity dual of the CFT is the Euclidean BTZ black hole [17] with the metric given by

$$ds^2 = (r^2 - r_+^2)d\tau^2 + \frac{R^2}{r^2 - r_+^2}dr^2 + r^2d\varphi^2. \quad (2.8)$$

The Euclidean time is compactified as $\tau \sim \tau + \frac{2\pi R}{r_+}$ to obtain a smooth geometry in addition to the periodicity $\varphi \sim \varphi + 2\pi$. Looking at its boundary, we find the relation $\frac{\beta}{L} = \frac{R}{r_+} \ll 1$ between the CFT and the BTZ black hole.

The subsystem A is defined by $0 \leq \varphi \leq 2\pi l/L$ at the boundary. Then we expect that the entropy can be computed from the geodesic distance between the boundary points $\varphi = 0, 2\pi l/L$ at a fixed time. To find the geodesic line, it is useful to remember the familiar fact that the Euclidean BTZ black hole at temperature T_{BTZ} is equivalent to the thermal AdS_3 at temperature $1/T_{BTZ}$. This equivalence can be interpreted as a modular transformation in the CFT side [18]. If we define the new coordinates $r = r_+ \cosh \rho$, $r_+ \tau = R\theta$, $r_+ \varphi = Rt$, then the metric (2.8) indeed becomes the Euclidean version of AdS_3 metric (2.2). Thus the geodesic distance can be found in the same way as in Eq. (2.4): $\cosh(L_{\gamma_A}/R) = 1 + 2 \cosh^2 \rho_0 \sinh^2 \left(\frac{\pi l}{\beta} \right)$, where the UV cutoff is interpreted as $e^{\rho_0} \sim \beta/a$. Then the area law (1.5) reproduces the known CFT result [10]

$$S_A(\beta) = \frac{c}{3} \cdot \log \left(\frac{\beta}{\pi a} \sinh \left(\frac{\pi l}{\beta} \right) \right). \quad (2.9)$$

We can extend these arguments to the multi interval cases and find the same formula (2.7) as before.

It is instructive to repeat the same analysis in the Poincare metric $ds^2 = R^2 z^{-2}(dz^2 - dx_0^2 + dr^2)$. We define the subsystem A by the region $-l/2 \leq r \leq l/2$ at the boundary $z = 0$. The geodesic line γ_A is given by

$$(r, z) = \frac{l}{2}(\cos s, \sin s), \quad (\epsilon \leq s \leq \pi - \epsilon), \quad (2.10)$$

where the infinitesimal ϵ is the UV cutoff $\epsilon \sim 2a/l$ (or equally $z_{UV} \sim a$). We obtain the entropy S_A as follows

$$S_A = \frac{L_{\gamma_A}}{4G_N^{(3)}} = \frac{R}{2G_N^{(3)}} \int_{\epsilon}^{\pi/2} \frac{ds}{\sin s} = \frac{c}{3} \log \frac{l}{a}. \quad (2.11)$$

This reproduces the small l limit of Eq. (1.3) [11].

When we perturb a CFT by a relevant perturbation, the RG flow generically drives the theory to a trivial IR fixed point. We denote the correlation length ξ in the latter theory. In the AdS dual, this massive deformation corresponds to capping off the IR region, restricting the allowed values of z to $z \leq \xi$. In the large l limit, we find

$$S_A = \frac{1}{4G_N^{(3)}} \int_{\epsilon}^{2\xi/l} \frac{ds}{\sin s} = \frac{c}{6} \log \frac{\xi}{a}. \quad (2.12)$$

This agrees with the CFT result with $\mathcal{A} = 1$ (1.4) [7, 10].

III. HIGHER DIMENSIONAL CASES

Motivated by the success in our gravitational derivation of the entanglement entropy for $d = 1$, it is interesting to extend the idea to higher dimensional cases ($d \geq 2$). A natural thing to do is to replace geodesic lines with minimal surfaces. The computations are analogous to the evaluation of Wilson loops [19], though the dimension of relevant minimal surfaces is different.

We will work in the Poincare metric for AdS_{d+2}

$$ds^2 = R^2 z^{-2} (dz^2 - dx_0^2 + \sum_{i=1}^d dx_i^2). \quad (3.1)$$

We consider two examples for the shape of A . The first one is a straight belt $A_S = \{x_i | x_1 \in [-l/2, l/2], x_{2,3,\dots,d} \in [-\infty, \infty]\}$ at the boundary $z = 0$ (Fig. 2 (a)). In this case the minimal surface is defined by $dz/dx_1 = \sqrt{z_*^{2d} - z^{2d}}/z^d$, where z_* is determined by $l/2 = \int_0^{z_*} dz z^d (z_*^{2d} - z^{2d})^{-1/2} = z_* \sqrt{\pi} \Gamma(\frac{d+1}{2d}) / \Gamma(\frac{1}{2d})$. The area of this minimal surface is

$$\text{Area}_{A_S} = \frac{2R^d}{d-1} \left(\frac{L}{a}\right)^{d-1} - \frac{2^d \pi^{d/2} R^d}{d-1} \left(\frac{\Gamma(\frac{d+1}{2d})}{\Gamma(\frac{1}{2d})}\right)^d \left(\frac{L}{l}\right)^{d-1}, \quad (3.2)$$

where L is the length of A_S in the $x_{2,3,\dots,d}$ -direction.

The second example is the disk A_D defined by $A_D = \{x_i | r \leq l\}$ (Fig. 2(b)) in the polar coordinate $\sum_i dx_i^2 = dr^2 + r^2 d\Omega_{d-1}^2$. The minimal surface is a d dimen-




FIG. 2: Minimal surfaces in AdS_{d+2} : (a) A_S and (b) A_D .

sional ball B^d defined by (2.10). Its area is

$$\begin{aligned}
 \text{Area}_{A_D} &= C \int_{a/l}^1 dy \frac{(1-y^2)^{(d-2)/2}}{y^d} \\
 &= p_1 (l/a)^{d-1} + p_3 (l/a)^{d-3} + \dots \\
 &\dots + \begin{cases} p_{d-1} (l/a) + p_d + O(a/l), & d: \text{even}, \\ p_{d-2} (l/a)^2 + q \log(l/a) + O(1), & d: \text{odd}, \end{cases}
 \end{aligned} \tag{3.3}$$

where $C = 2\pi^{d/2} R^d / \Gamma(d/2)$ and $p_1/C = (d-1)^{-1}$ etc. For d even, $p_d/C = (2\sqrt{\pi})^{-1} \Gamma(\frac{d}{2}) \Gamma(\frac{1-d}{2})$ and for d odd, $q/C = (-)^{(d-1)/2} (d-2)!! / (d-1)!!$.

From these results, the entanglement entropy can be calculated by Eq. (1.5). Each of (3.2) and (3.3) has a UV divergent term $\sim a^{-d+1}$ that is proportional to the area of the boundary ∂A . This agrees with the known ‘area law’ of the entanglement entropy in quantum field theories [5, 6]. Note that this ‘area law’ is related to ours Eq. (1.5) via the basic property of holography.

We may prefer a physical quantity that is independent of the cutoff (i.e. universal). The second term in Eq. (3.2) has this property. In general, when A is a finite size, there is a universal and conformal invariant constant contribution to S_A if d is even (see [20] for properties of minimal surfaces in AdS). In (2+1)D topological field theories the constant contribution to S_A encodes the quantum dimension and is called the topological entanglement entropy [8, 9]. If d is odd, the coefficient of the logarithmic term $\sim \log(l/a)$ is universal as in Eq. (1.3).

Let us apply the previous results to a specific string theory setup. Type IIB string on $\text{AdS}_5 \times S^5$ is dual to 4D $\mathcal{N} = 4$ $SU(N)$ super Yang-Mills theory [2]. The radius of AdS_5 and S^5 are given by the same value $R = (4\pi g_s \alpha'^2 N)^{1/4}$. The 5D Newton constant is related to the 10D one via $G_N^{(10)} = \pi^3 R^5 G_N^{(5)}$. Then we obtain

from Eqs. (3.2) and (3.3)

$$S_{A_S} = \frac{N^2 L^2}{2\pi a^2} - 2\sqrt{\pi} \left(\frac{\Gamma(\frac{2}{3})}{\Gamma(\frac{1}{6})} \right)^3 \frac{N^2 L^2}{l^2}, \quad (3.4)$$

$$S_{A_D} = N^2 \left[\frac{l^2}{a^2} - \log \left(\frac{l}{a} \right) + O(1) \right]. \quad (3.5)$$

It is interesting to compare the finite universal term in Eq. (3.4) with the field theory one. For free real scalars and fermions in general dimensions, one way to compute S_{A_S} is presented in [21] (see also [22]). Indeed, this leads to the same behavior in a and l as in Eq. (3.4). Following this approach, we can estimate finite contributions from 6 scalars and 4 Majorana fermions in the $\mathcal{N} = 4$ Yang-Mills multiplet. In the end, we obtain $S_{finite}^{freeCFT} \sim -(0.068 + g) \cdot N^2 L^2 / l^2$, where g is the contribution from the gauge field ($g = 0.010$ if we treat the gauge field as 2 scalars). On the other hand, our AdS_5 result (3.4) leads to $S_{finite}^{AdS} \sim -0.051 \cdot N^2 L^2 / l^2$. We may think this is a good agreement if we remember that the gravity description corresponds to the strongly coupled gauge theory instead of the free theory as in [23].

We can also compute the entanglement entropy for the near horizon limit $AdS_4 \times S^7$ ($AdS_7 \times S^4$) of N M2-branes (M5-branes)

$$S_{A_S}^{M2} = \frac{\sqrt{2}}{3} N^{3/2} \left[\frac{L}{a} - \frac{4\pi^3}{\Gamma(1/4)^4} \frac{L}{l} \right], \quad (3.6)$$

$$S_{A_D}^{M2} = \frac{\sqrt{2}\pi}{3} N^{3/2} \left[\frac{l}{a} - 1 \right], \quad (3.7)$$

$$S_{A_S}^{M5} = \frac{2}{3\pi^2} N^3 \left[\frac{L^4}{a^4} - 16\pi^{5/2} \frac{\Gamma(3/5)^5}{\Gamma(1/10)^5} \frac{L^4}{l^4} \right], \quad (3.8)$$

$$S_{A_D}^{M5} = \frac{32}{9} N^3 \left[\frac{1}{4} \cdot \frac{l^4}{a^4} - \frac{3}{4} \cdot \frac{l^2}{a^2} + \frac{3}{8} \log \frac{l}{a} \right]. \quad (3.9)$$

Note that the constant terms in Eqs. (3.6), (3.7) and (3.8) do not depend on the choice of the cutoff a .

IV. YANG-MILLS AT FINITE TEMPERATURE

As the final example, we discuss the $\mathcal{N} = 4$ super Yang-Mills theory on \mathbb{R}^3 at a finite temperature T , which is dual to the AdS black hole defined by the metric




FIG. 3: (a) Minimal surfaces γ_A for various sizes of A . (b) γ_A and γ_B wrap the different parts of the horizon.

[24]

$$ds^2 = R^2 \left[\frac{du^2}{hu^2} + u^2 \left(-hdt^2 + dx_1^2 + dx_2^2 + dx_3^2 \right) + d\Omega_5^2 \right], \quad (4.1)$$

where $h = 1 - u_0^4/u^4$, $u_0 = \pi T$. For the straight belt A_S , the area is given by (putting the cut off $u \sim z^{-1} \sim a$)

$$\text{Area}_{A_S} = 2R^3 L^2 \int_{u_*}^{a^{-1}} \frac{du u^6}{\sqrt{(u^4 - u_0^4)(u^6 - u_*^6)}}, \quad (4.2)$$

where u_* satisfies $l/2 = \int_{u_*}^{\infty} du [(u^4 - u_0^4)(u^6/u_*^6 - 1)]^{-1/2}$. Eq. (4.2) contains the UV divergence $\sim a^{-2}$ as before. As in the analogous computation of Wilson loops [25], we also expect a term which is proportional to the area of A . Indeed, when l is large ($u_* \sim u_0$) we find the constant term $\sim \pi^3 R^3 L^2 l T^3$. This leads to the finite part of S_A

$$S_{finite} \simeq \frac{\pi^2}{2} N^2 T^3 L^2 l = \frac{\pi^2 N^2 T^3}{2} \times (\text{Area of } A_S). \quad (4.3)$$

We can regard this entropy as a part of the Bekenstein-Hawking entropy of black 5-branes [23], which is proportional to the area of horizon situated at $u = u_0$. Thus we can interpret the part (4.3) as a thermal entropy contribution to the total entanglement entropy at finite temperature. In our gravitational description, this part arises because the minimal surface wraps a part of the black hole horizon (Fig. 3 (a)). If we expand the size of A until it coincides with the total system (in the global coordinate), γ_A wraps the horizon completely and S_A becomes equal to the Bekenstein-Hawking entropy as expected. In a sense, the overall normalization in Eq. (1.5) is fixed from Eq. (1.1) once we consider the entanglement entropy at finite temperature. Note that at finite temperature, $S_A = S_B$ does not hold generically. In the present situation, this occurs because the surfaces γ_A and γ_B wrap two different parts of the horizon (Fig. 3 (b)).

As argued in [13, 26], the AdS black hole can be dual to an entanglement of two different CFTs at the two boundaries. It is interesting to look at the minimal surfaces that connect them. As a specific limit, we may think the black hole entropy is the same as the entanglement entropy of the CFTs as the minimal surface wrap the horizon.

V. CONCLUSION

In this paper we proposed a holographic description of the entanglement entropy in quantum (conformal) field theories via $\text{AdS}_{d+2}/\text{CFT}_{d+1}$ correspondence. This is summarized as the area law relation (1.5). Based on our proposal we computed the entanglement entropy for various systems, e.g. 2D CFTs and the 4D large- N $\mathcal{N} = 4$ super Yang-Mills theory. We checked that in the lowest dimensional case ($d = 1$), our formula exactly coincides with the entropy computed directly from CFT.

In higher dimensions ($d \geq 2$), a quantitative comparison is not easy because the gravity description corresponds to the strongly coupled gauge theory whose entanglement entropy is not known at present. Nevertheless, we found that our computation for $\text{AdS}_5 \times S^5$ reproduces the same functional form of entanglement entropy as the one in the 4D free $\mathcal{N} = 4$ super Yang-Mills. Their numerical coefficients only differ by a factor $\sim \frac{3}{2}$ in a rough estimation (assuming that the gauge field contribution is the same as that of two real scalar fields). This is similar to the well-known result of the thermal entropy [23].

Finally we computed the entanglement entropy at finite temperature employing AdS black hole geometry. In this case the minimal surface wraps a fraction of the black hole horizon and this part is responsible for the thermal contribution.

We would like to thank M. Einhorn, T. Harmark, G. Horowitz, Y. Hyakutake, D. Marolf, T. Okuda, S.-J. Rey, M. Shigemori and Y. Sugawara for useful discussions. This research was supported in part by the National Science Foundation under Grant No. PHY99-07949.

-
- [1] A. Strominger and C. Vafa, "Microscopic Origin of the Bekenstein-Hawking Entropy," *Phys. Lett. B* **379**, 99 (1996) [arXiv:hep-th/9601029].
 - [2] J. M. Maldacena, "The large N limit of superconformal field theories and supergravity," *Adv. Theor. Math. Phys.* **2**, 231 (1998) [*Int. J. Theor. Phys.* **38**, 1113 (1999)]

- [arXiv:hep-th/9711200].
- [3] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, "Gauge theory correlators from non-critical string theory," *Phys. Lett. B* **428**, 105 (1998) [arXiv:hep-th/9802109].
 - [4] E. Witten, "Anti-de Sitter space and holography," *Adv. Theor. Math. Phys.* **2**, 253 (1998) [arXiv:hep-th/9802150].
 - [5] L. Bombelli, R. K. Koul, J. H. Lee and R. D. Sorkin, "A Quantum Source Of Entropy For Black Holes," *Phys. Rev. D* **34**, 373 (1986).
 - [6] M. Srednicki, "Entropy and area," *Phys. Rev. Lett.* **71**, 666 (1993) [arXiv:hep-th/9303048].
 - [7] G. Vidal, J. I. Latorre, E. Rico and A. Kitaev, "Entanglement in quantum critical phenomena," *Phys. Rev. Lett.* **90**, 227902 (2003) [arXiv:quant-ph/0211074].
 - [8] A. Kitaev and J. Preskill, "Topological entanglement entropy," arXiv:hep-th/0510092.
 - [9] M. Levin and X. G. Wen, "Detecting topological order in a ground state wave function," arXiv:cond-mat/0510613.
 - [10] P. Calabrese and J. Cardy, "Entanglement entropy and quantum field theory," *J. Stat. Mech.* **0406**, P002 (2004) [arXiv:hep-th/0405152].
 - [11] C. Holzhey, F. Larsen and F. Wilczek, "Geometric and renormalized entropy in conformal field theory," *Nucl. Phys. B* **424**, 443 (1994) [arXiv:hep-th/9403108].
 - [12] S. Hawking, J. M. Maldacena and A. Strominger, "DeSitter entropy, quantum entanglement and AdS/CFT," *JHEP* **0105**, 001 (2001) [arXiv:hep-th/0002145].
 - [13] J. M. Maldacena, "Eternal black holes in Anti-de-Sitter," *JHEP* **0304**, 021 (2003) [arXiv:hep-th/0106112].
 - [14] J. D. Brown and M. Henneaux, "Central Charges In The Canonical Realization Of Asymptotic Symmetries," *Commun. Math. Phys.* **104**, 207 (1986).
 - [15] L. Susskind and E. Witten, "The holographic bound in anti-de Sitter space," arXiv:hep-th/9805114.
 - [16] G. 't Hooft, "Dimensional reduction in quantum gravity," arXiv:gr-qc/9310026; L. Susskind, "The World as a hologram," *J. Math. Phys.* **36**, 6377 (1995) [arXiv:hep-th/9409089]; R. Bousso, "A Covariant Entropy Conjecture," *JHEP* **9907** (1999) 004 [arXiv:hep-th/9905177].
 - [17] M. Banados, C. Teitelboim and J. Zanelli, "The Black hole in three-dimensional space-time," *Phys. Rev. Lett.* **69**, 1849 (1992) [arXiv:hep-th/9204099].
 - [18] J. M. Maldacena and A. Strominger, "AdS(3) black holes and a stringy exclusion principle," *JHEP* **9812**, 005 (1998) [arXiv:hep-th/9804085].
 - [19] S. J. Rey and J. T. Yee, "Macroscopic strings as heavy quarks in large N gauge theory and anti-de Sitter supergravity," *Eur. Phys. J. C* **22**, 379 (2001) [arXiv:hep-

- th/9803001]; J. M. Maldacena, "Wilson loops in large N field theories," *Phys. Rev. Lett.* **80**, 4859 (1998) [arXiv:hep-th/9803002]; D. Berenstein, R. Corrado, W. Fischler and J. M. Maldacena, "The operator product expansion for Wilson loops and surfaces in the large N limit," *Phys. Rev. D* **59**, 105023 (1999) [arXiv:hep-th/9809188].
- [20] C. R. Graham and E. Witten, "Conformal anomaly of submanifold observables in AdS/CFT correspondence," *Nucl. Phys. B* **546**, 52 (1999) [arXiv:hep-th/9901021].
- [21] H. Casini and M. Huerta, "Entanglement and alpha entropies for a massive scalar field in two dimensions," arXiv:cond-mat/0511014.
- [22] D. V. Fursaev, "Entanglement entropy in critical phenomena and analogue models of quantum gravity," arXiv:hep-th/0602134.
- [23] S. S. Gubser, I. R. Klebanov and A. W. Peet, "Entropy and Temperature of Black 3-Branes," *Phys. Rev. D* **54**, 3915 (1996) [arXiv:hep-th/9602135].
- [24] E. Witten, "Anti-de Sitter space, thermal phase transition, and confinement in gauge theories," *Adv. Theor. Math. Phys.* **2**, 505 (1998) [arXiv:hep-th/9803131].
- [25] S. J. Rey, S. Theisen and J. T. Yee, "Wilson-Polyakov loop at finite temperature in large N gauge theory and anti-de Sitter supergravity," *Nucl. Phys. B* **527**, 171 (1998) [arXiv:hep-th/9803135]; A. Brandhuber, N. Itzhaki, J. Sonnenschein and S. Yankielowicz, "Wilson loops in the large N limit at finite temperature," *Phys. Lett. B* **434**, 36 (1998) [arXiv:hep-th/9803137]; "Wilson loops, confinement, and phase transitions in large N gauge theories from supergravity," *JHEP* **9806**, 001 (1998) [arXiv:hep-th/9803263].
- [26] R. Brustein, M. B. Einhorn and A. Yarom, "Entanglement interpretation of black hole entropy in string theory," *JHEP* **0601**, 098 (2006) [arXiv:hep-th/0508217].