

THE SIMPLICIAL MODEL OF UNIVALENT FOUNDATIONS (AFTER VOEVODSKY)

KRZYSZTOF KAPULKIN AND PETER LEFANU LUMSDAINE

ABSTRACT. We present Voevodsky’s construction of a model of univalent type theory in the category of simplicial sets.

To this end, we first give a general technique for constructing categorical models of dependent type theory, using universes to obtain coherence. We then construct a (weakly) universal Kan fibration, and use it to exhibit a model in simplicial sets. Lastly, we introduce the Univalence Axiom, in several equivalent formulations, and show that it holds in our model.

As a corollary, we conclude that Martin-Löf type theory with one univalent universe (formulated in terms of contextual categories) is at least as consistent as ZFC with two inaccessible cardinals.

To the memory of Vladimir Voevodsky.

CONTENTS

Introduction	3
1. Models from Universes	6
1.1. The type theory under consideration	6
1.2. Contextual categories	6
1.3. Contextual categories from universes	12
1.4. Logical structure on universes	13
2. The Simplicial Model	20
2.1. A universe of Kan complexes	21
2.2. Kan fibrancy of the universe	24
2.3. Modelling type theory in simplicial sets	26
3. Univalence	30
3.1. Type-theoretic univalence	30
3.2. Simplicial univalence	31
3.3. Equivalence of type-theoretic and simplicial univalence	35
3.4. Univalence of the simplicial universes	42
3.5. Univalence and pullback representations	44
Appendix A. Syntax of Martin-Löf Type Theory	46
Appendix B. Logical structure on contextual categories	52

Date: November 2012; last revised September 2018.

2010 *Mathematics Subject Classification.* Primary 03B15, 55U10; Secondary 18C50, 55U35.

References

INTRODUCTION

The Univalent Foundations programme is a new proposed approach to foundations of mathematics, originally suggested by Vladimir Voevodsky in [Voe06], building on the systems of dependent type theory developed by Martin-Löf and others.

A major motivation for earlier work with such logical systems has been their well-suitedness to computer implementation. One notable example is the Coq proof assistant, based on the Calculus of Inductive Constructions (a closely related dependent type theory), which has shown itself feasible for large-scale formal verification of mathematics, with developments including formal proofs of the Four-Colour Theorem [Gon08] and the Feit-Thompson (Odd Order) Theorem [GAA⁺13].

One feature of dependent type theory which has previously remained comparatively unexploited, however, is its richer treatment of equality. In traditional foundations, equality carries no information beyond its truth-value: if two things are equal, they are equal in at most one way. This is fine for equality between elements of discrete sets; but it is unnatural for objects of categories (or higher-dimensional categories), or points of spaces. In particular, it is at odds with the informal mathematical practice of treating isomorphic (and sometimes more weakly equivalent) objects as equal; which is why this usage must be so often disclaimed as an abuse of language, and kept rigorously away from formal statements, even though it is so appealing.

In dependent type theory, equalities can carry information: two things may be equal in multiple ways. So the basic objects—the *types*—may behave not just like discrete sets, but more generally like higher groupoids (with equalities being morphisms in the groupoid), or spaces (with equalities being paths in the space). And, crucially, this is the *only* equality one can talk about within the logical system: one cannot ask whether elements of a type are “equal on the nose”, in the classical sense.¹ The logical language only allows one to talk about properties and constructions which respect its equality.

The *Univalence Axiom*, introduced by Voevodsky, strengthens this characteristic. In classical foundations one has sets of sets, or classes of sets, and uses these to quantify over classes of structures. Similarly, in type theory, types of types—*universes*—are a key feature of the language. The Univalence Axiom states that equality between types, as elements of a universe, is the same as equivalence between them, as types. It formalises the practice of treating equivalent structures as completely interchangeable; it ensures that one can only talk about properties of types, or more general structures, that respect such equivalence. In sum, it helps solidify the idea of types as some kind of spaces, in the homotopy-theoretic sense; and more practically—its original motivation—it provides for free many theorems (transfer along equivalences, naturality with respect to these, and so on) which must otherwise be re-proved by hand for each new construction.

The main goal of this paper is to justify the intuition outlined above, of types as spaces. To this end, we focus on the Quillen model category \mathbf{sSet} of simplicial sets, a well-studied model for topological spaces in homotopy theory; we construct a model of type theory in \mathbf{sSet} , and show that it satisfies the Univalence Axiom. The fibrations of this model structure, called

¹There is a strict equality, called *judgemental* or *definitional*, but there is no *type/proposition* in the system expressing it, just as with e.g. literal syntactic equality in traditional foundations.

Kan fibrations, will serve as an interpretation of type dependency. In particular the closed types will be interpreted as Kan complexes, which also serve as a model for ∞ -groupoids, for instance in Joyal and Lurie’s approach to higher category theory.

It follows from this model that Martin-Löf type theory plus the Univalence Axiom (presented in terms of contextual categories) is consistent, provided that the classical foundations we use are—precisely, ZFC together with the existence of two strongly inaccessible cardinals, or equivalently two Grothendieck universes.

As hinted above, there is one important technical caveat regarding our treatment of type theory: we state the model and consistency results in terms of contextual categories, not syntax, so as to avoid reliance on initiality results.

This paper therefore includes a mixture of logical and homotopy-theoretic ingredients; however, we have aimed to separate the two wherever possible. Good background references for the logical parts include [NPS90], a general introduction to the type theory; [Hof97], for the categorical semantics; and [ML84], the *locus classicus* for the logical rules. For the homotopy-theoretic aspects, [GJ09] and [Hov99] are both excellent and sufficient references. Finally, for the category-theoretic language used throughout, [ML98] is canonical.

Organisation. In Section 1 we consider general techniques for constructing models of type theory. After setting out (in Section 1.1) the specific type theory that we will consider, we review (Section 1.2) some fundamental facts about its intended semantics in contextual categories, following [Str91]. In Section 1.3, we use universes to construct contextual categories, representing the structural core of type theory; and in Section 1.4, we use categorical constructions on the universe to model the logical constructions of type theory. Together, these present a new solution to the *coherence problem* for modelling type theory (cf. [Hof95b]).

In Section 2, we turn towards constructing a model in the category of simplicial sets. Sections 2.1 and 2.2 are dedicated to the construction and investigation of a (weakly) universal Kan fibration (a “universe of Kan complexes”); in Section 2.3 we use this universe to apply the techniques of Section 1, giving a model of the full type theory in simplicial sets.

Section 3 is devoted to the Univalence Axiom. We formulate univalence first in type theory (Section 3.1), then directly in homotopy-theoretic terms (Section 3.2), and show that these definitions correspond under the simplicial model (Section 3.3). In Section 3.4, we show that the universal Kan fibration is univalent, and hence that the Univalence Axiom holds in the simplicial model. Finally, in Section 3.5 we discuss an alternative formulation of univalence, shedding further light on the universal property of the universe.

We include also two appendices, setting out in full the type theory under consideration: first a conventional syntactic presentation in Appendix A, and then in Appendix B its translation into algebraic structure on contextual categories.

History of the paper. This paper started life as notes by the current authors based on Vladimir Voevodsky’s lectures at the 2011 Oberwolfach workshop [AGMLV11] along with his associated manuscript [Voe12]. It was subsequently expanded with Voevodsky’s collaboration into the present full exposition of the simplicial model, and appeared as a preprint in 2012 with Voevodsky included as co-author.

In 2016, due to his dissatisfaction with the existing literature on type theory, which this paper took as background, Voevodsky asked us to remove him as co-author and delay publication until he had finished developing his own treatment of semantics of type theory (cf. [Voe14] and sequels) and presentation of the simplicial model in that framework.

Tragically, Voevodsky passed away in September 2017, before completing that project. This paper therefore remains the only account of Voevodsky’s construction of the simplicial model, so with the support of Daniel Grayson, Voevodsky’s academic executor, we have prepared it again for publication. We have made several changes to accommodate Voevodsky’s reservations regarding the treatment of semantics; most importantly, we present the initiality of syntax as a conjecture rather than a theorem (Conjecture 1.2.9), and give all main results in terms of contextual categories rather than syntax. Otherwise, the paper remains substantially unchanged from the original 2012 version.

The main results of the paper are due to Voevodsky, including Theorems 1.4.15, 2.3.4, 3.4.2 and 3.5.3. Mathematical contributions of Kapulkin and Lumsdaine include all of Section 3.3, along with streamlining various parts of the main constructions and completing portions omitted in [Voe12].

Out of respect for Voevodsky’s stated wishes, and following discussion with his executor, he remains absent as an author of the final version of this paper. However, we wish to leave no doubt regarding the share of the credit that is his due. We mourn the loss of an exceptional mathematician and mentor, and dedicate this paper to his memory.

Related work. While the present paper discusses just models of type theory with the univalence axiom, the major motivation for this is the actual development of mathematics within these foundations. Introductions to various aspects of this are given in [Gra18], [Voe15b], [PW14], and [Uni13], while large computer-formalised developments include the UniMath² and HoTT³ libraries, presented in [VAG⁺] and [BGL⁺17].

Earlier work on homotopy-theoretic models of type theory can be found in [HS98], [AW09], [War11]. Other current and recent work on such models includes [GvdB11], [AK11], and [Shu15]. Other general coherence theorems, for comparison with the results of Section 1, can be found in [Hof95b] and [LW15]. Univalence in homotopy-theoretic settings is also considered in [GK17]. (These references are, of course, far from exhaustive.)

Acknowledgements. First and foremost we would like to thank Vladimir Voevodsky: the creator of Univalent Foundations, a mentor to both the authors, and whose insight and ingenuity underlie not only the present paper but much subsequent work in the field. We are particularly indebted also to Michael Warren, whose illuminating seminars and discussions heavily influenced our understanding and presentation of the simplicial model. We also thank Daniel Grayson, Ieke Moerdijk, Mike Shulman, Raffael Stenzel, and Karol Szumilo, for helpful correspondence, conversations, and corrections to drafts and earlier versions, and Steve Awodey, for support and encouragement.

The first-named author was financially supported during this work by the NSF, Grant DMS-1001191 (P.I. Steve Awodey), and by a grant from the Benter Foundation (P.I. Thomas C. Hales); the second-named author, by an AARMS postdoctoral fellowship at Dalhousie

²<https://www.github.com/UniMath/UniMath>

³<https://www.github.com/HoTT/HoTT>

University, and grants from NSERC (P.I.'s Peter Selinger, Robert Dawson, and Dorette Pronk).

1. MODELS FROM UNIVERSES

In this section, we set up the machinery which we will use, in later sections, to model type theory in simplicial sets. The type theory we consider, and some of the technical machinery we use, are standard; the main original contribution is a new technique for solving the so-called coherence problem, using universes.

1.1. The type theory under consideration. Formally, the type theory we will consider is a slight variant of Martin-Löf's Intensional Type Theory, as presented in e.g. [ML84]. The rules of this theory are given in full in Appendix A; briefly, it is a dependent type theory, taking as basic constructors Π -, Σ -, Id -, and W -types, 0 , 1 , $+$, and one universe à la Tarski closed under these constructors.

A related theory of particular interest is the Calculus of Inductive Constructions, on which the Coq proof assistant is based ([Wer94]). CIC differs from Martin-Löf type theory most notably in its very general scheme for inductive definitions, and in its treatment of universes. We do not pursue the question of how our model might be adapted to CIC, but for some discussion and comparison of the two systems, see [PM96], [Bar12], and [Voe12, 6.2].

One abuse of notation that we should mention: we will sometimes write e.g. $A(x)$ or $t(x, y)$ to indicate free variables on which a term or type may depend, so that we can later write $A(g(z))$ to denote the substitution $[g(z)/x]A$ more readably. Note however that the variables explicitly shown need not actually appear; and there may also always be other free variables in the term, not explicitly displayed.

1.2. Contextual categories. Rather than working formally with the syntax of this type theory, we work instead in terms of *contextual categories*, a class of algebraic objects abstracting the key structure given by the syntax.⁴ The plain definition of a contextual category corresponds to the structural core of the syntax; further syntactic rules (logical constructors, etc.) correspond to extra algebraic structure that contextual categories may carry. Essentially, contextual categories are intended to provide a completely equivalent alternative to the syntactic presentation of type theory.

Why do we make this bait-and-switch? The trouble with the syntax is that it is very tricky to handle rigorously. Any full presentation must account for (among other complications) variable binding, capture-free substitution, and the possibility of multiple derivations of a judgement; and so any careful construction of an interpretation must deal with all of these, at the same time as tackling the details of the particular model in question. Contextual categories, by contrast, are a purely algebraic notion, with no such subtleties. The idea is therefore that given any contextual category \mathcal{C} with structure corresponding to the logical rules of some syntactic type theory \mathbf{T} , one should obtain an interpretation of the syntax of

⁴Contextual categories are not the only option; the closely related notions of *categories with attributes* [Car78, Mog91, Pit00], *categories with families* [Dyb96, Hof97], and *comprehension categories* [Jac93] would all also serve our purposes.

\mathbf{T} in \mathcal{C} ; and in proving this, one deals with the subtleties and bureaucracy of \mathbf{T} once and for all, giving a clear framework for subsequently constructing models of \mathbf{T} .

Such an “initiality theorem” (cf. Conjecture 1.2.9 below) has been proven for some specific rather small type theories, e.g. by Streicher in the Correctness Theorem of [Str91, Ch. III, p. 181]. For larger type theories such as the present one, however, its status is debatable, and at best unsatisfactory. The traditional view is that for suitable type theories, a straightforward extension of Streicher’s and other standard methods suffices, and therefore the theorem can be regarded as established. However, Voevodsky has argued persuasively that this is an unacceptably unrigorous attitude. No precise definition has been given of what “suitable type theories” the methods apply to; nor (to our knowledge) has the proof been even sketched in detail for any type theory beyond small toy examples; and while a “straightforward” extension of standard methods may indeed suffice for a type theory such as the present one, that sufficiency is far from obvious.

For the present paper, therefore, we work formally entirely in terms of contextual categories, and avoid relying on initiality theorems in any form.

Conversely, then, why bring up syntax at all, other than as motivation? The trouble on this side is that working with higher-order logical structure in contextual categories quickly becomes unreadable: compare, for instance, the statements of functional extensionality in Sections A.4 and B.3.

We therefore make free use of the syntax of type theory, as a *notation* for working in contextual categories. The situation is rather comparable to that of string diagrams, as used in monoidal and more elaborately structured categories [Sel11], or indeed of the traditional notations for differentiation and integration. In each case, one has a powerful, flexible, and intuitive notation, whose rigorous definition and validity requires quite non-trivial work to establish; but in lieu of such a general justification, one may still fruitfully make use of the notation, trusting the reader to translate it into the unproblematic algebraic form as required.

Definition 1.2.1 (Cartmell [Car78, Sec. 2.2], Streicher [Str91, Def. 1.2]). A *contextual category* \mathcal{C} consists of the following data:

- (1) a category \mathcal{C} ;
- (2) a grading of objects as $\text{Ob } \mathcal{C} = \coprod_{n \in \mathbb{N}} \text{Ob}_n \mathcal{C}$;
- (3) an object $1 \in \text{Ob}_0 \mathcal{C}$;
- (4) maps $\text{ft}_n : \text{Ob}_{n+1} \mathcal{C} \longrightarrow \text{Ob}_n \mathcal{C}$ (whose subscripts we usually suppress);
- (5) for each $X \in \text{Ob}_{n+1} \mathcal{C}$, a map $p_X : X \longrightarrow \text{ft } X$ (the *canonical projection* from X);
- (6) for each $X \in \text{Ob}_{n+1} \mathcal{C}$ and $f : Y \longrightarrow \text{ft}(X)$, an object $f^*(X)$ together with a map $q(f, X) : f^*(X) \longrightarrow X$;

such that:

- (7) 1 is the unique object in $\text{Ob}_0(\mathcal{C})$;
- (8) 1 is a terminal object in \mathcal{C} ;

- (9) for each $n > 0$, $X \in \text{Ob}_n \mathcal{C}$, and $f: Y \longrightarrow \text{ft}(X)$, we have $\text{ft}(f^*X) = Y$, and the square

$$\begin{array}{ccc} f^*X & \xrightarrow{q(f,X)} & X \\ p_{f^*X} \downarrow & \lrcorner & \downarrow p_x \\ Y & \xrightarrow{f} & \text{ft}(X) \end{array}$$

is a pullback (the *canonical pullback* of X along f); and

- (10) these canonical pullbacks are strictly functorial: that is, for $X \in \text{Ob}_{n+1} \mathcal{C}$, $1_{\text{ft } X}^* X = X$ and $q(1_{\text{ft } X}, X) = 1_X$; and for $X \in \text{Ob}_{n+1} \mathcal{C}$, $f: Y \longrightarrow \text{ft } X$ and $g: Z \longrightarrow Y$, we have $(fg)^*(X) = g^*(f^*(X))$ and $q(fg, X) = q(f, X)q(g, f^*X)$.

Contextual categories have also been studied under the name *C-systems* ([Voe14] et seq.)

Remark 1.2.2. Note that these may be seen as models of a multi-sorted essentially algebraic theory ([AR94, 3.34]), with sorts indexed by $\mathbb{N} + \mathbb{N} \times \mathbb{N}$.

This definition is best understood in terms of its prototypical example:

Example 1.2.3 (Cartmell [Car78, p2.6]; cf. also [Voe14], [Voe16b]). Let \mathbf{T} be the dependent type theory given by the structural rules of Section A.1, plus any selection of the subsequent logical rules.⁵ Then there is a contextual category $\mathcal{C}(\mathbf{T})$, described as follows:

- $\text{Ob}_n \mathcal{C}(\mathbf{T})$ consists of the contexts $[x_1:A_1, \dots, x_n:A_n]$ of length n , up to definitional equality and renaming of free variables;
- maps of $\mathcal{C}(\mathbf{T})$ are *context morphisms*, or *substitutions*, considered up to definitional equality and renaming of free variables. That is, a map

$$f: [x_1:A_1, \dots, x_n:A_n] \longrightarrow [y_1:B_1, \dots, y_m:B_m(y_1, \dots, y_{m-1})]$$

is an equivalence class of sequences of terms f_1, \dots, f_m such that

$$x_1:A_1, \dots, x_n:A_n \vdash f_1 : B_1$$

$$\vdots$$

$$x_1:A_1, \dots, x_n:A_n \vdash f_m : B_m(f_1, \dots, f_{m-1}),$$

and two such maps $[f_i]$, $[g_i]$ are equal exactly if for each i ,

$$x_1:A_1, \dots, x_n:A_n \vdash f_i = g_i : B_i(f_1, \dots, f_{i-1});$$

- composition is given by substitution, and the identity $\Gamma \longrightarrow \Gamma$ by the variables of Γ , considered as terms;
- 1 is the empty context $[\]$;
- $\text{ft}[x_1:A_1, \dots, x_{n+1}:A_{n+1}] = [x_1:A_1, \dots, x_n:A_n]$;
- for $\Gamma = [x_1:A_1, \dots, x_{n+1}:A_{n+1}]$, the map $p_\Gamma: \Gamma \longrightarrow \text{ft } \Gamma$ is the *dependent projection* context morphism

$$(x_1, \dots, x_n): [x_1:A_1, \dots, x_{n+1}:A_{n+1}] \longrightarrow [x_1:A_1, \dots, x_n:A_n],$$

simply forgetting the last variable of Γ ;

⁵Heuristically, \mathbf{T} may be “any type theory” here; but there is no established definition of what this means!

- for contexts

$$\Gamma = [x_1:A_1, \dots, x_{n+1}:A_{n+1}(x_1, \dots, x_n)],$$

$$\Gamma' = [y_1:B_1, \dots, y_m:B_m(y_1, \dots, y_{m-1})],$$

and a map $f = [f_i(\vec{y})]_{i \leq n}: \Gamma' \longrightarrow \text{ft } \Gamma$, the pullback $f^*\Gamma$ is the context

$$[y_1:B_1, \dots, y_m:B_m(y_1, \dots, y_{m-1}), y_{m+1}:A_{n+1}(f_1(\vec{y}), \dots, f_n(\vec{y}))],$$

(for some fresh y_{m+1}) and $q(\Gamma, f): f^*\Gamma \longrightarrow \Gamma$ is the map

$$[f_1, \dots, f_n, y_{m+1}].$$

Note that typed terms $\Gamma \vdash t : A$ of \mathbf{T} may be recovered from $\mathcal{C}(\mathbf{T})$, up to definitional equality, as sections of the projection $p_{[\Gamma, x:A]}: [\Gamma, x:A] \longrightarrow \Gamma$. For this reason, when working with contextual categories, we will often write just “sections” to refer to sections of dependent projections.

We will also use several other notations deserving of particular comment. For a fixed contextual category \mathcal{C} and an object $\Gamma \in \text{Ob}_n \mathcal{C}$, we write (Γ, A) for any object in $\text{Ob}_{n+1} \mathcal{C}$ with $\text{ft}(\Gamma, A) = \Gamma$, shall such object exist, and p_A for the dependent projection $p_{(\Gamma, A)}$. Similarly, we write (Γ, A, B) for an arbitrary object in $\text{Ob}_{n+2} \mathcal{C}$ with $\text{ft}(\Gamma, A, B) = (\Gamma, A)$, and so on.

Given a morphism $f: \Delta \longrightarrow \Gamma$ and an object (Γ, A) , we write (Δ, f^*A) for the canonical pullback $f^*(\Gamma, A)$ and similarly (Δ, f^*A, f^*B) for $f^*(\Gamma, A, B)$. We also extend the notation f^* to apply not only to the canonical pullbacks of appropriate objects, but also the pullbacks of maps between them.

As mentioned above, Definition 1.2.1 alone corresponds precisely to the basic judgements and structural rules of dependent type theory. Similarly, each logical rule or type- or term-constructor should correspond to certain extra structure on a contextual category. We state this intended correspondence precisely in Conjecture 1.2.9 below, once we have set up the appropriate definitions.

Definition 1.2.4 (cf. [Voe16a, Sec. 4]). A Π -type structure on a contextual category \mathcal{C} consists of:

- (1) for each $(\Gamma, A, B) \in \text{Ob}_{n+2} \mathcal{C}$, an object $(\Gamma, \Pi(A, B)) \in \text{Ob}_{n+1} \mathcal{C}$;
- (2) for each such (Γ, A, B) and section $b: (\Gamma, A) \longrightarrow (\Gamma, A, B)$ (of the dependent projection p_B), a section $\lambda(b): \Gamma \longrightarrow (\Gamma, \Pi(A, B))$ (of $p_{\Pi(A, B)}$);
- (3) for each (Γ, A, B) and pair of sections $k: \Gamma \longrightarrow (\Gamma, \Pi(A, B))$ and $a: \Gamma \longrightarrow (\Gamma, A)$, a section $\text{app}(k, a): \Gamma \longrightarrow (\Gamma, A, B)$ such that the following diagram commutes:

$$\begin{array}{ccc}
 & (\Gamma, A, B) & \\
 & \uparrow \text{app}(k, a) & \downarrow p_B \\
 & (\Gamma, A) & \\
 \Gamma & \nearrow a & \downarrow p_A \\
 \Gamma & \xlongequal{\quad} & \Gamma
 \end{array}$$

- (4) such that for all such (Γ, A, B) , $a: \Gamma \longrightarrow (\Gamma, A)$, and $b: (\Gamma, A) \longrightarrow (\Gamma, A, B)$, we have $\mathbf{app}(\lambda(b), a) = b \cdot a$;
- (5) and moreover such that all the above operations are stable under substitution: for any morphism $f: \Delta \longrightarrow \Gamma$, and suitable (Γ, A, B) , a, b, k , we have

$$\begin{aligned} (\Delta, f^* \Pi(A, B)) &= (\Delta, \Pi(f^* A, f^* B)), \\ \lambda(f^* b) &= f^* \lambda(b), \quad \mathbf{app}(f^* k, f^* a) = f^*(\mathbf{app}(k, a)). \end{aligned}$$

These are direct translations of the rules for Π -types given in Section A.2.⁶ Similarly, all the other logical rules of Appendix A may be routinely translated into structure on a contextual category; see Appendix B and [Hof97, 3.3] for more details and discussion.

Example 1.2.5. If \mathbf{T} is a type theory with Π -types, then $\mathcal{C}(\mathbf{T})$ carries an evident Π -type structure; similarly for Σ -types and the other constructors of Sections A.2 and A.3.

Remark 1.2.6. Note that all of these structures, like the definition of contextual categories themselves, are essentially algebraic in nature.

Definition 1.2.7. A map $F: \mathcal{C} \longrightarrow \mathcal{D}$ of contextual categories, or *contextual functor*, consists of a functor $\mathcal{C} \longrightarrow \mathcal{D}$ between underlying categories, respecting the gradings, and preserving (on the nose) all the structure of a contextual category.

Similarly, a map of contextual categories with Π -type structure, Σ -type structure, etc., is a contextual functor preserving the additional structure.

Remark 1.2.8. These are exactly the maps given by considering contextual categories as essentially algebraic structures.

We are now equipped to state precisely the sense in which the structures defined above are expected to correspond to the appropriate syntactic rules:

Conjecture 1.2.9. *Let \mathbf{T} be the type theory given by the structural rules of Section A.1, plus any combination of the logical rules of Sections A.2, A.3. Then $\mathcal{C}(\mathbf{T})$ is initial among contextual categories with the correspondingly-named extra structure.*

In other words, if \mathcal{C} is a contextual category with structure corresponding to the logical rules of a type theory \mathbf{T} , then there should be a unique homomorphism $\mathcal{C}(\mathbf{T}) \longrightarrow \mathcal{C}$, interpreting the syntax of \mathbf{T} in \mathcal{C} . As discussed at the beginning of this section, the Correctness Theorem of [Str91, Ch. III, p. 181] gives an analogous fact for a rather smaller type theory, while the status of the present conjecture is debated, accepted by some but not all in the field as a straightforward extension of that theorem.

Bearing this intended correspondence in mind, therefore, but avoiding relying on it, we take for the present paper the following definitions:

Definition 1.2.10. By *Martin-Löf Type Theory plus the Univalence Axiom* (MLTT + UA for short), we mean dependent type theory with Π -, Σ -, Id -, W -, unit , zero , and sum types, along with one universe closed under all these type formers and satisfying the Univalence Axiom, as set out in Appendix A.

⁶A partial exception is the stability axiom, which corresponds not to any explicitly given rule of the syntax, but rather to clauses for Π , λ , and \mathbf{app} in the inductive *definition* of substitution.

By a *model* of $\text{MLTT} + \text{UA}$, or more generally of dependent type theory with any selection of the logical rules of Appendix A, we mean a contextual category equipped with the corresponding structure from Appendix B. By the *contextual-category presentation* of such a type theory, we mean the essentially algebraic theory of such structures.

Note that by definition as an essentially algebraic theory, it is immediate that any such type theory has an initial model.

Definition 1.2.11. A dependent type theory of the form considered in Definition 1.2.10 and including the empty type 0 is *inconsistent* just if in the initial model, the map $p_{0_1} : (1, 0_1) \rightarrow 1$ admits a section, and is *consistent* if it is not inconsistent.

Assuming initiality, this corresponds to the usual type-theoretic sense of inconsistency: a closed term inhabiting the empty type. Readers who accept the initiality conjecture as true may therefore read Theorem 3.4.2 as providing an interpretation of the usual syntactic presentation of $\text{MLTT} + \text{UA}$, and Theorem 3.4.3 as its consistency in the usual type-theoretic sense.

1.3. Contextual categories from universes. The major difficulty in constructing models of type theories is the so-called *coherence problem*: the requirement for pullback to be strictly functorial, and for the logical structure to commute strictly with it. In most natural categorical situations, operations on objects commute with pullback only up to isomorphism, or even more weakly; and for constructors with weak universal properties, operations on maps (corresponding for example to the Id-ELIM rule) may also fail to commute with pullback. Hofmann [Hof95b] gives a construction which solves the issue for Π - and Σ -types, but Id -types in particular remain problematic with this method. Other methods exist for certain specific categories ([HS98], [War08]), but are not applicable to the present case.

In order to obtain coherence for our model, we thus use a construction based on *universes* (not necessarily the same as universes in the type-theoretic sense, though the two may sometimes coincide), studied in more detail in [Voe15a].

Definition 1.3.1 ([Voe15a, Def. 2.1]). Let \mathcal{C} be a category. A *universe* in \mathcal{C} is an object U together with a morphism $p: \tilde{U} \rightarrow U$, and for each map $f: X \rightarrow U$ a choice of pullback square

$$\begin{array}{ccc} (X; f) & \xrightarrow{Q(f)} & \tilde{U} \\ P_{(X,f)} \downarrow & \lrcorner & \downarrow p \\ X & \xrightarrow{f} & U. \end{array}$$

The intuition here is that the map p represents the generic family of types over the universe U .

By abuse of notation, we often refer to the universe simply as U , with p and the chosen pullbacks understood.

Given a map $q: Y \rightarrow X$, we will often write $\ulcorner q \urcorner$ (or $\ulcorner Y \urcorner$, if q is understood) for a map $X \rightarrow U$ such that $q \cong P_{(X, \ulcorner q \urcorner)}$ in \mathcal{C}/X . Also, for a sequence of maps $f_1: X \rightarrow U$, $f_2: (X; f_1) \rightarrow U$, etc., we write $(X; f_1, \dots, f_n)$ for $((\dots (X; f_1); \dots); f_n)$. (In particular, with $n = 0$, $(X;) = X$.)

Definition 1.3.2 ([Voe15a, Constr. 2.12]). Given a category \mathcal{C} , together with a universe U and a terminal object 1 , we define a contextual category \mathcal{C}_U as follows:

- $\text{Ob}_n \mathcal{C}_U := \{ (f_1, \dots, f_n) \in (\text{Mor } \mathcal{C})^n \mid f_i: (1; f_1, \dots, f_{i-1}) \longrightarrow U \ (1 \leq i \leq n) \}$;
- $\mathcal{C}_U((f_1, \dots, f_n), (g_1, \dots, g_m)) := \mathcal{C}((1; f_1, \dots, f_n), (1; g_1, \dots, g_m))$;
- $1_{\mathcal{C}_U} := ()$, the empty sequence;
- $\text{ft}(f_1, \dots, f_{n+1}) := (f_1, \dots, f_n)$;
- the projection $p_{(f_1, \dots, f_{n+1})}$ is the map $P_{(X, f_{n+1})}$ provided by the universe structure on U ;
- given (f_1, \dots, f_{n+1}) and a map $\alpha: (g_1, \dots, g_m) \longrightarrow (f_1, \dots, f_n)$ in \mathcal{C}_U , the canonical pullback $\alpha^*(f_1, \dots, f_{n+1})$ in \mathcal{C}_U is given by $(g_1, \dots, g_m, f_{n+1} \cdot \alpha)$, with projection induced by $Q(f_{n+1} \cdot \alpha)$:

$$\begin{array}{ccccc}
 & & \xrightarrow{Q(f_{n+1} \cdot \alpha)} & & \\
 (1; g_1, \dots, g_m, f_{n+1} \cdot \alpha) & \xrightarrow{\quad} & (1; f_1, \dots, f_{n+1}) & \xrightarrow{Q(f_{n+1})} & \tilde{U} \\
 \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow p \\
 (1; g_1, \dots, g_m) & \xrightarrow{\alpha} & (1; f_1, \dots, f_n) & \xrightarrow{f_{n+1}} & U
 \end{array}$$

Proposition 1.3.3 ([Voe15a, Constr. 2.12, Ex. 4.9]).

- (1) These data define a contextual category \mathcal{C}_U .
- (2) This contextual category is well-defined up to canonical isomorphism given just \mathcal{C} and $p: \tilde{U} \longrightarrow U$, independently of the choice of pullbacks and terminal object.

Proof. Routine computation. □

Justified by the second part of this proposition, we will not explicitly consider the choices of pullbacks and terminal object when we construct the universe in the category \mathbf{sSet} of simplicial sets.

As an aside, let us note that every small contextual category arises in this way:

Proposition 1.3.4 ([Voe15a, Constr. 5.2]). *Let \mathcal{C} be a small contextual category. Consider the universe U in the presheaf category $[\mathcal{C}^{\text{op}}, \mathbf{Set}]$ given by*

$$\begin{aligned}
 U(X) &= \{Y \mid \text{ft } Y = X\} \\
 \tilde{U}(X) &= \{(Y, s) \mid \text{ft } Y = X, \ s \text{ a section of } p_Y\},
 \end{aligned}$$

with the evident projection map, and any choice of pullbacks.

Then $[\mathcal{C}^{\text{op}}, \mathbf{Set}]_U$ is isomorphic, as a contextual category, to \mathcal{C} .

Proof. Straightforward, with liberal use of the Yoneda lemma. □

1.4. Logical structure on universes. Given a universe U in a category \mathcal{C} , we want to know how to equip \mathcal{C}_U with various logical structure— Π -types, Σ -types, and so on. For general \mathcal{C} , this is rather fiddly; but when \mathcal{C} is locally cartesian closed (as in our case of interest), it is more straightforward, since local cartesian closedness allows us to construct and manipulate “objects of U -contexts”, and hence to construct objects representing the premises of each rule.

In working with locally cartesian closed categories (LCCC's), we will follow topos-theoretic convention and write Σ_f and Π_f respectively for the left and right adjoints to the pullback functor f^* along a map $f: A \rightarrow B$:

$$\begin{array}{ccc} & \Sigma_f & \\ \mathcal{C}/A & \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{f^*} \end{array} & \mathcal{C}/B \\ & \Pi_f & \end{array}$$

Also, the intended map $A \rightarrow B$ is often clearly determined by the objects A and B , as some sort of associated projection; in such a case, we will write $\Sigma_{A \rightarrow B}$, $\Pi_{A \rightarrow B}$ for the functors arising from this map.

An alternative notation for locally cartesian closed categories is their internal logic, *extensional* dependent type theory [See84], [Hof95b]. While this language is convenient and powerful, we avoid it due to the difficulties of working clearly with two logical languages in parallel.

Returning to the question at hand, first consider Π -types.⁷ We know that dependent products exist in \mathcal{C} ; so informally, we need only to ensure that U (considered as a universe of types) is closed under such products. Specifically, given a type A in U over some base X (that is, a map $\ulcorner A \urcorner: X \rightarrow U$), and a dependent family of types B over A , again in U (i.e. a map $\ulcorner B \urcorner: A := (X; \ulcorner A \urcorner) \rightarrow U$), the product $\Pi_{A \rightarrow X} B$ of this family in the slice \mathcal{C}/X should again “live in U ”; that is, there should be a map $\ulcorner \Pi(A, B) \urcorner: X \rightarrow U$ such that $(X; \ulcorner \Pi(A, B) \urcorner) \cong \Pi_{A \rightarrow X} B$. Moreover, we need this construction to be strictly natural in X .

Due to the strict naturality requirement, we cannot simply provide this structure for each X and A, B individually. Instead, we construct an object U^Π representing such pairs (A, B) , and a generic such pair $(A_{\text{gen}}, B_{\text{gen}})$ based on U^Π . It is sufficient to define Π in this generic case $X = U^\Pi$; the construction then extends to other X by precomposition, and as such, is automatically strictly natural in X .

Precisely:

Definition 1.4.1. Given a universe U in an lccc \mathcal{C} , define

$$U^\Pi := \Sigma_{U \rightarrow 1} \Pi_{\tilde{U} \rightarrow U} (\pi_2: U \times \tilde{U} \rightarrow \tilde{U}).$$

(This definition can be expressed in several ways, according to one's preferred notation. In the internal language of \mathcal{C} as an LCCC, it can be written as $\llbracket A:U, B:[\tilde{U}_A, U] \rrbracket$, showing it more explicitly as an internalisation of the premises of the Π -FORM rule. Using a more traditional internal-hom notation, it could alternatively be written as $\mathbf{Hom}_U(\tilde{U}, U \times U)$.)

Pulling back \tilde{U} along the projection $U^\Pi \rightarrow U$ induces an object $A_{\text{gen}} = \tilde{U} \times_U U^\Pi$, along with a projection map $\alpha_{\text{gen}}: A_{\text{gen}} \rightarrow U^\Pi$. Similarly, pulling back \tilde{U} along the counit (the evaluation map of the internal hom)

$$A_{\text{gen}} = \tilde{U} \times_U \Pi_{\tilde{U} \rightarrow U} (U \times \tilde{U}) \rightarrow U \times \tilde{U} \rightarrow U$$

⁷The following construction is studied in considerable detail in [Voe17].

induces an object $(B_{\text{gen}}, \beta_{\text{gen}})$ over A_{gen} :

$$\begin{array}{ccc}
 B_{\text{gen}} & \longrightarrow & \tilde{U} \\
 \downarrow \lrcorner & & \downarrow \\
 A_{\text{gen}} & \longrightarrow & \tilde{U} \\
 \downarrow \lrcorner & & \downarrow \\
 U^\Pi & \longrightarrow & U
 \end{array}
 \quad
 \begin{array}{c}
 \xrightarrow{\quad} \\
 \text{curved arrow} \\
 \xrightarrow{\quad}
 \end{array}$$

Moreover, the universal properties of the LCCC structure ensure that for any sequence $B \rightarrow A \rightarrow \Gamma$ with maps $\Gamma \rightarrow U$, $A \rightarrow U$, $A \rightarrow \tilde{U}$, $B \rightarrow \tilde{U}$ exhibiting $A \rightarrow \Gamma$ and $B \rightarrow A$ as pullbacks of $\tilde{U} \rightarrow U$, there is a unique map $\ulcorner(A, B)\urcorner: \Gamma \rightarrow U^\Pi$ which induces the given sequence via precomposition and pullback:

$$\begin{array}{ccccc}
 B & \longrightarrow & B_{\text{gen}} & \longrightarrow & \tilde{U} \\
 \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow \\
 A & \longrightarrow & A_{\text{gen}} & \longrightarrow & \tilde{U} \\
 \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow \\
 \Gamma & \xrightarrow{\ulcorner(A, B)\urcorner} & U^\Pi & \longrightarrow & U
 \end{array}
 \quad
 \begin{array}{c}
 \xrightarrow{\quad} \\
 \text{curved arrow} \\
 \xrightarrow{\quad}
 \end{array}$$

So $B_{\text{gen}} \rightarrow A_{\text{gen}} \rightarrow U^\Pi$ is *generic* among such sequences, and U^Π *represents* the inputs for a Π operation (that is, the premises of the Π -FORM rule) on \mathcal{C}_U .

Definition 1.4.2 (cf. [Voe17, Def. 4.1]). A Π -structure on a universe U in a lccc \mathcal{C} consists of a map

$$\Pi: U^\Pi \rightarrow U.$$

whose realisation is a dependent product for the generic dependent family of types; that is, it is equipped with an isomorphism $\Pi^* \tilde{U} \cong \Pi_{\alpha_{\text{gen}}} B_{\text{gen}}$ over U^Π , or equivalently with a map $\tilde{\Pi}: \Sigma_{U^\Pi \rightarrow 1} \Pi_{\alpha_{\text{gen}}} B_{\text{gen}} \rightarrow \tilde{U}$ making the square

$$\begin{array}{ccc}
 \Sigma_{U^\Pi \rightarrow 1} \Pi_{\alpha_{\text{gen}}} B_{\text{gen}} & \xrightarrow{\tilde{\Pi}} & \tilde{U} \\
 \downarrow & & \downarrow \\
 U^\Pi & \xrightarrow{\Pi} & U
 \end{array}$$

a pullback.

The approach used here gives a template which we follow for all the other constructors, with extra subtleties entering the picture just in the cases of **Id**-types and (type-theoretic) universes, since these structures are not characterised by strict category-theoretic universal properties.

Definition 1.4.3. Take U^Σ to be the object representing the premises of the Σ -FORM rule:

$$U^\Sigma := \Sigma_{U \rightarrow 1} \Pi_{\tilde{U} \rightarrow U} (U \times \tilde{U})$$

Since these are the same as the premises of the Π -FORM rule, we have in this case that $U^\Sigma = U^\Pi$; and we have again the generic family of types $B_{\text{gen}} \longrightarrow A_{\text{gen}} \longrightarrow U^\Sigma$, as over U^Π .

Definition 1.4.4. A Σ -structure on a universe U in a lccc \mathcal{C} consists of a map

$$\Sigma: U^\Sigma \longrightarrow U$$

whose realisation is a dependent sum for the generic dependent family of types; that is, it is equipped with an isomorphism $\Sigma^* \tilde{U} \cong \Sigma_{\alpha_{\text{gen}}} B_{\text{gen}}$ over U^Σ (or again equivalently with a map $\tilde{\Sigma}: \Sigma_{U^\Sigma \rightarrow 1} \Sigma_{\alpha_{\text{gen}}} B_{\text{gen}} \longrightarrow \tilde{U}$ making the appropriate square a pullback).

Id-structure requires a few auxiliary definitions.⁸ Recall first the classical notion of weak orthogonality of maps:

Definition 1.4.5. For maps $i: A \longrightarrow B$, $f: Y \longrightarrow X$ in a category \mathcal{C} , say i is (*weakly*) *orthogonal* to f if any commutative square from i to f has some diagonal filler:

$$\begin{array}{ccc} A & \longrightarrow & Y \\ i \downarrow & \nearrow & \downarrow f \\ B & \longrightarrow & X \end{array}$$

or, in other words, if the function

$$\begin{aligned} \text{Hom}(B, Y) &\longrightarrow \text{Hom}(A, Y) \times_{\text{Hom}(A, X)} \text{Hom}(B, X) \\ g &\mapsto (g \cdot i, f \cdot g) \end{aligned}$$

has a section.

Say i is moreover *stably orthogonal* to f if for every object C of \mathcal{C} , $C \times i$ is orthogonal to f .

In a cartesian closed category, this notion has an internal analogue:

Definition 1.4.6. Given maps $i: A \longrightarrow B$, $f: Y \longrightarrow X$ in a cartesian closed category \mathcal{C} , an *internal lifting operation* for i against f is a section of the evident map $Y^B \longrightarrow Y^A \times_{X^A} X^B$.

The following proposition connects the classical and internal notions:

Proposition 1.4.7. *Given i, f as above, there exists an internal lifting operation for i against f if and only if i is stably orthogonal to f .*

Proof. If i is stably orthogonal to f , then an internal lifting operation may be obtained as (the exponential transpose of) a filler for the canonical square

$$\begin{array}{ccc} A \times (Y^A \times_{X^A} X^B) & \xrightarrow{\text{ev}_{A,Y} \cdot (A \times \pi_1)} & Y \\ i \times Y^A \times_{X^A} X^B \downarrow & & \downarrow f \\ B \times (Y^A \times_{X^A} X^B) & \xrightarrow{\text{ev}_{B,X} \cdot (B \times \pi_2)} & X. \end{array}$$

⁸We should thank here Michael Warren and Steve Awodey, who both strongly influenced the current presentation of the definition.

Conversely, any square from $C \times i$ to f induces a map $C \rightarrow Y^A \times_{X^A} X^B$; composing this with an internal lifting operation provides a map $C \rightarrow Y^B$, whose transpose is a filler for the square. \square

As shown in [AW09] and [GG08], the rules for Id-types can be understood roughly as follows. In a model where dependent types are interpreted as fibrations, the identity type over a type A (in any slice \mathcal{C}/Γ) is a factorisation of the diagonal $\Delta_A: A \rightarrow A \times_\Gamma A$ as a stable trivial cofibration, followed by a fibration. (Here, by a stable trivial cofibration, we mean a map which is stably orthogonal to fibrations, in \mathcal{C}/Γ .) Additionally, choices of all data (including liftings) must be given which commute with pullbacks in the base Γ .

In our case, the “fibrations” are just the pullbacks of p ; so it suffices to consider orthogonality between the first map of the factorisation and p itself. Moreover, as for Π - and Σ -structure above, we demand the structure just in the universal case where A is \tilde{U} , in the slice \mathcal{C}/U . Finally, an internal lifting operation turns out to be exactly the structure required to give chosen lifts commuting with pullbacks. We therefore define:

Definition 1.4.8. An *Id-structure* on a universe consists of maps

$$\text{Id}: U^{\text{Id}} := \tilde{U} \times_U \tilde{U} \rightarrow U, \quad r: \tilde{U} \rightarrow \text{Id}^* \tilde{U}$$

such that the triangle

$$\begin{array}{ccc} \tilde{U} & \xrightarrow{r} & \text{Id}^* \tilde{U} \\ \Delta_{\tilde{U}} \searrow & & \swarrow \text{Id}^* p \\ & \tilde{U} \times_U \tilde{U} & \end{array}$$

commutes, together with an internal lifting operation J for r against $p \times U$ in \mathcal{C}/U .

Remark 1.4.9. By virtue of Proposition 1.4.7, we could instead simply stipulate that r be stably orthogonal to $p \times U$. We choose the current version since it provides exactly the structure required for Theorem 1.4.15, without requiring any arbitrary choices.

Another alternative is described in [Voe15c, Sec. 2.3].

Definition 1.4.10. A *W-structure* on a universe consists of a map

$$W: U^W := \Sigma_{U \rightarrow 1} \Pi_{\tilde{U} \rightarrow U} (U \times \tilde{U}) \rightarrow U$$

such that $W^* \tilde{U}$ is an initial algebra for the polynomial endofunctor of \mathcal{C}/U^W specified by $\beta_{\text{gen}}: B_{\text{gen}} \rightarrow A_{\text{gen}}$, i.e. the endofunctor

$$\mathcal{C}/U^W \xrightarrow{\beta_{\text{gen}}^* \alpha_{\text{gen}}^*} \mathcal{C}/B_{\text{gen}} \xrightarrow{\Pi_{\beta_{\text{gen}}}} \mathcal{C}/A_{\text{gen}} \xrightarrow{\Sigma_{\alpha_{\text{gen}}}} \mathcal{C}/U^W.$$

(For details on polynomial endofunctors in logical settings, see [MP00], [GH04]. Intuitively, their initial algebras may be seen as types of well-founded trees, or of syntax over algebraic signatures.)

Definition 1.4.11. A *0-structure* on U consists of a map $\mathbf{0}: 1 \rightarrow U$ such that $\mathbf{0}^* \tilde{U} \cong 0$.

(By analogy with the preceding definitions, one might write $\mathbf{0}: U^0 \rightarrow U$ instead and similarly in the next two definitions. However, since U^0 is a terminal object, we choose not to do so simply for the sake of readability.)

Definition 1.4.12. A *1-structure* on U consists of a map $\mathbf{1}: 1 \rightarrow U$ such that $\mathbf{1}^*\tilde{U} \cong 1$.

Definition 1.4.13. A *+ -structure* on U consists of a map $+: U \times U \rightarrow U$, together with an isomorphism $+^*\tilde{U} \cong \pi_1^*\tilde{U} + \pi_2^*\tilde{U}$ in $\mathcal{C}/(U \times U)$.

Finally, we consider the structure on U needed to give a universe (in the type-theoretic sense) in \mathcal{C}_U . Here, for the first time, we need to consider a nested pair of universes, since the internal universe of \mathcal{C}_U must be some smaller universe U_0 in \mathcal{C} .

Definition 1.4.14. An *internal universe* (U_0, i) in U consists of arrows

$$u_0: 1 \rightarrow U \quad i: U_0 := u_0^*\tilde{U} \rightarrow U.$$

Given these, i induces by pullback a universe structure $(p_0, \tilde{U}_0, \dots)$ on U_0 . We say that U_0 is closed under Π -types in U if U_0 carries a Π -structure Π_0 , commuting with i in the sense that the square

$$\begin{array}{ccc} U_0^\Pi & \xrightarrow{i^\Pi} & U^\Pi \\ \Pi_0 \downarrow & & \downarrow \Pi \\ U_0 & \xrightarrow{i} & U \end{array}$$

commutes (where the top map is induced by the evident functoriality of U^Π in U).

Similarly, we say that U_0 is closed under Σ -types (resp. Id-types, etc.) if it carries a Σ -structure Σ_0 (resp. an Id-structure (Id_0, r_0) , etc.) commuting with i .

With these structures defined, we can now prove that they are fit for purpose:

Theorem 1.4.15 (cf. [Voe17, Constr. 4.3], [Voe15c, Sec. 2.4]). *A Π -structure (resp. Σ -structure, etc.) on a universe U induces Π -type structure (resp. Σ -type structure, etc.) on \mathcal{C}_U .*

Moreover, an internal universe (U_0, i) in U closed under any combination of Π -types, Σ -types, etc., induces a universe à la Tarski in \mathcal{C}_U closed under the corresponding constructors.

Proof. This proof is essentially a routine verification; we give the case of Π -types in full, and leave the rest mostly to the reader.

In a nutshell, the constructor Π is induced by the map Π ; and the constructors λ and app are induced by the corresponding lccc structure in \mathcal{C} .

Precisely, we treat the rules of Π -types (corresponding to the components of the desired Π -type structure) one at a time.

(Π -FORM): The premises

$$\Gamma \vdash A \text{ type} \quad \Gamma, x:A \vdash B \text{ type}$$

in \mathcal{C}_U correspond to data in \mathcal{C} of the form

$$\begin{array}{ccc} A & \longrightarrow & \tilde{U} \\ \downarrow & \lrcorner & \downarrow \\ \Gamma & \xrightarrow{\ulcorner A \urcorner} & U \end{array} \quad \begin{array}{ccc} B & \longrightarrow & \tilde{U} \\ \downarrow & \lrcorner & \downarrow \\ A & \xrightarrow{\ulcorner B \urcorner} & U \end{array}$$

and hence to a map

$$(\ulcorner A^\top, \ulcorner B^\top) : \Gamma \longrightarrow U^\Pi.$$

Then the composite $\Pi \cdot (\ulcorner A^\top, \ulcorner B^\top)$ gives a type $\Gamma \longrightarrow U$ which we take as $\Pi(A, B)$. By construction, this is stable under substitution along any map $f : \Delta \longrightarrow \Gamma$, since substitution in \mathcal{C}_U is again just composition in \mathcal{C} .

(Π -INTRO): Besides Γ, A, B as before, we have an additional premise

$$\Gamma, x:A \vdash t : B(x).$$

This is by definition a map $1_A \longrightarrow B$ in \mathcal{C}/A , corresponding by adjunction to a map $\hat{t} : 1_\Gamma \longrightarrow \Pi_{A \rightarrow \Gamma} B$ in \mathcal{C}/Γ . But

$$\begin{aligned} \Pi_{A \rightarrow \Gamma} B &\cong (\ulcorner A^\top, \ulcorner B^\top)^* \Pi_{A_{\text{gen}} \rightarrow U^\Pi} B_{\text{gen}} \\ &\cong (\ulcorner A^\top, \ulcorner B^\top)^* \Pi^* \tilde{U} \\ &\cong (\Pi \cdot (\ulcorner A^\top, \ulcorner B^\top))^* \tilde{U} \end{aligned}$$

so \hat{t} corresponds to a section of $\Pi(A, B)$ over Γ , which we take as $\lambda(t)$.

Stability under substitution follows by the uniqueness in the universal property of $\Pi_{A \rightarrow \Gamma} B$.

We could alternatively have defined λ more analogously to Π , by representing the premises as a single map $(\ulcorner A^\top, \ulcorner B^\top, t) : \Gamma \longrightarrow U^\lambda$ (where $U^\lambda := \Sigma_{U^\Pi \rightarrow 1} \Pi_{A_{\text{gen}} \rightarrow U^\Pi} B_{\text{gen}}$ represents the inputs of λ , i.e. the premises of Π -INTRO); then taking the transpose of the generic term t_{gen} over U^λ ; and then pulling this back along $(\ulcorner A^\top, \ulcorner B^\top, t)$. In fact, thanks to the uniqueness in the universal property of $\Pi_{A_{\text{gen}} \rightarrow U^\Pi} B_{\text{gen}}$, that would give the same result as the present, more straightforward, definition. However, the alternative definition has the advantage that its stability under substitution follows simply from properties of pullbacks; this becomes important for Id -types, whose universal property lacks a uniqueness condition.

(Π -APP): The premises now are

$$\begin{aligned} \Gamma \vdash A \text{ type} \quad \Gamma, x:A \vdash B \text{ type} \\ \Gamma \vdash f : \Pi(A, B) \quad \Gamma \vdash a : A \end{aligned}$$

corresponding to Γ, A, B as before, plus sections

$$\begin{array}{ccc} A & & \Pi(A, B) \cong \Pi_{A \rightarrow \Gamma} B \\ & \searrow a & \nearrow f \\ & \Gamma & \end{array}$$

Together, these give a section over Γ of $\Pi_{A \rightarrow \Gamma} B \times_\Gamma A$; so composing this with the evaluation map $\text{ev}_{A,B}$ of $\Pi_{A \rightarrow \Gamma} B$ gives a map $\Gamma \longrightarrow B$ lifting a , which we take to be $\text{app}(f, a)$.

(Π -COMP): here, we have premises Γ, A, B, t as in Π -INTRO, and a as in Π -APP; and we have formed $\text{app}(\lambda(t), a)$ as prescribed above. So, unwinding the isomorphism $\Pi(A, B) \cong \Pi_{A \rightarrow \Gamma} B$ used in each case,

$$\begin{aligned} \text{app}(\lambda(t), a) &= \text{ev}_{A,B} \cdot (\hat{t}, a) \\ &= t \cdot a \end{aligned}$$

as desired, by the usual rules of LCCCs.

This completes the proof for Π -structures.

As indicated above, the remaining constructors are for the most part entirely analogous; the only subtlety is in the case for the Id-ELIM rule. In this case, there are two ways that one could define the appropriate structure: one can either pull back to each specific context and then choose liftings, or choose a lifting in the universal context and then pull it back (as discussed following the Π -INTRO case above). The second of these is the correct choice: the first is not automatically stable under substitution. (For other constructors, this distinction does not arise, since their strict categorical universal properties canonically determine the maps involved.) And, in fact, the “universal lifting” required is precisely the internal lifting operation provided by the Id -structure on U .

□

2. THE SIMPLICIAL MODEL

In this section, we will apply the techniques of Section 1 to construct a model of type theory in the category \mathbf{sSet} . As mentioned in the Introduction, type dependency is interpreted using Kan fibrations and in particular the closed type will be Kan complexes. To this end, we construct (for any regular cardinal α) a Kan fibration $p_\alpha: \tilde{U}_\alpha \rightarrow U_\alpha$, weakly universal among Kan fibrations with α -small fibers, and investigate the key properties of U_α and p_α . We then show that U_α is a Kan complex, and (when α is inaccessible) carries the various logical structures defined in Section 1.4. Together, these yield our first main goal: a model of type theory in \mathbf{sSet} , with an internal universe.

2.1. A universe of Kan complexes. In constructing a universe U_α intended to represent α -small Kan fibrations, one might expect (by the Yoneda lemma) to simply define $(U_\alpha)_n$ as the set of α -small fibrations over $\Delta[n]$. This definition has two problems: firstly, it gives not sets, but proper classes; and secondly, it is not strictly functorial, since pullback is functorial only up to isomorphism.

Some extra technical device is therefore needed to resolve these issues. Several possible solutions exist⁹; we take the approach of passing to isomorphism classes, having first added well-orderings to the mix so that fibrations have no non-trivial automorphisms (without which the crucial Lemmas 2.1.4, 2.1.5 would fail). We emphasise, however, that this is the sole reason for introducing the well-orderings: they are of no intrinsic interest or significance.

Definition 2.1.1. A *well-ordered morphism* of simplicial sets consists of an ordinary map of simplicial sets $f: Y \rightarrow X$, together with a function assigning to each simplex $x \in X_n$ a well-ordering on the fiber $Y_x := f^{-1}(x) \subseteq Y_n$.

If $f: Y \rightarrow X$, $f': Y' \rightarrow X$ are well-ordered morphisms into a common base X , an *isomorphism* of well-ordered morphisms from f to f' is an isomorphism $Y \cong Y'$ over X preserving the well-orderings on the fibers.

Proposition 2.1.2. *Given two well-ordered sets, there is at most one isomorphism between them. Given two well-ordered morphisms over a common base, there is at most one isomorphism between them.*

⁹Other possible approaches include ones based on the general results of [Hof95b] and [LW15], or taking $U_{\alpha n}$ as $[(f\Delta[n])^{\text{op}}, \text{Set}_{<\alpha}]$ as in [HS99].

Proof. The first statement is classical (and immediate by induction); the second follows from the first, applied in each fiber. \square

Definition 2.1.3. Fix (for the remainder of this and the following section) a regular cardinal α . Say a map of simplicial sets $f: Y \rightarrow X$ is α -small if each of its fibers Y_x has cardinality $< \alpha$.

Given a simplicial set X , define $\mathbf{W}_\alpha(X)$ to be the set of isomorphism classes¹⁰ of α -small well-ordered morphisms $Y \rightarrow X$; together with the pullback action $\mathbf{W}_\alpha(f) := f^*: \mathbf{W}_\alpha(X) \rightarrow \mathbf{W}_\alpha(X')$, for $f: X' \rightarrow X$, this gives a contravariant functor $\mathbf{W}_\alpha: \mathbf{sSet}^{\text{op}} \rightarrow \mathbf{Set}$.

Lemma 2.1.4. \mathbf{W}_α preserves all limits: $\mathbf{W}_\alpha(\text{colim}_i X_i) \cong \lim_i \mathbf{W}_\alpha(X_i)$.

Proof. Suppose $F: J \rightarrow \mathbf{sSet}$ is some diagram, and $X = \text{colim}_J F$ is its colimit, with injections $\nu_i: F(i) \rightarrow X$. We need to show that the canonical map $\mathbf{W}_\alpha(X) \rightarrow \lim_J \mathbf{W}_\alpha(F(i))$ is an isomorphism.

To see that it is surjective, suppose we are given $[f_i: Y_i \rightarrow F(i)] \in \lim_J \mathbf{W}_\alpha(F(i))$. For each $x \in X_n$, choose some i and $\bar{x} \in F(i)$ with $\nu(\bar{x}) = x$, and set $Y_x := (Y_i)_{\bar{x}}$. By Proposition 2.1.2, this is well-defined up to canonical isomorphism, independent of the choices of representatives i, \bar{x}, Y_i, f_i . The total space of these fibers then defines a well-ordered morphism $f: Y \rightarrow X$, with fibers of size $< \alpha$, and with pullbacks isomorphic to f_i as required.

For injectivity, suppose f, f' are well-ordered morphisms over X , and $\nu_i^* f \cong \nu_i^* f'$ for each i . By Proposition 2.1.2, these isomorphisms must agree on each fiber, so together give an isomorphism $f \cong f'$. \square

Define the simplicial set W_α by

$$W_\alpha := \mathbf{W}_\alpha \cdot \mathbf{y}^{\text{op}}: \Delta^{\text{op}} \rightarrow \mathbf{Set},$$

where \mathbf{y} denotes the Yoneda embedding $\Delta \rightarrow \mathbf{sSet}$.

Lemma 2.1.5. The functor \mathbf{W}_α is representable, represented by W_α .

Proof. The functors \mathbf{W}_α and $\text{Hom}(-, W_\alpha)$ agree up to isomorphism on the standard simplices (by the Yoneda lemma), and send colimits in \mathbf{sSet} to limits; but every simplicial set is canonically a colimit of standard simplices. \square

Notation 2.1.6. Given an α -small well-ordered map $f: Y \rightarrow X$, the corresponding map $X \rightarrow W_\alpha$ will be denoted by $\ulcorner f \urcorner$.

Applying the natural isomorphism above to the identity map $W_\alpha \rightarrow W_\alpha$ yields a universal α -small well-ordered simplicial set $\widetilde{W}_\alpha \rightarrow W_\alpha$. Explicitly, n -simplices of \widetilde{W}_α are classes of pairs

$$(f: Y \rightarrow \Delta[n], s \in f^{-1}(1_{[n]}))$$

i.e. the fiber of \widetilde{W}_α over an n -simplex $\ulcorner f \urcorner \in W_\alpha$ is exactly (an isomorphic copy of) the main fiber of f . So, by construction:

¹⁰We use *isomorphism classes* in the sense of “Scott’s trick” [Sco55] for constructing proper class quotients. The class of *all* well-ordered morphisms isomorphic to a given one is a proper class, so one instead uses the subclass of such morphisms of *minimal rank*, which is a set.

Proposition 2.1.7. *The canonical projection $\widetilde{W}_\alpha \rightarrow W_\alpha$ is strictly universal for α -small well-ordered morphisms; that is, any such morphism can be expressed uniquely as a pullback of this projection.* \square

Corollary 2.1.8. *The canonical projection $\widetilde{W}_\alpha \rightarrow W_\alpha$ is weakly universal for α -small morphisms of simplicial sets: any such morphism can be given, not necessarily uniquely, as a pullback of this projection.*

Proof. By the well-ordering principle and the axiom of choice, one can well-order the fibers, and then use the universal property of W_α . \square

Definition 2.1.9. Let $U_\alpha \subseteq W_\alpha$ (respectively, $U_\alpha \subseteq W_\alpha$) be the subobject consisting of (isomorphism classes of) α -small well-ordered fibrations¹¹; and define $p_\alpha: \widetilde{U}_\alpha \rightarrow U_\alpha$ as the pullback:

$$\begin{array}{ccc} \widetilde{U}_\alpha & \longrightarrow & \widetilde{W}_\alpha \\ p_\alpha \downarrow & \lrcorner & \downarrow \\ U_\alpha & \hookrightarrow & W_\alpha \end{array}$$

Lemma 2.1.10. *The map $p_\alpha: \widetilde{U}_\alpha \rightarrow U_\alpha$ is a fibration.*

Proof. Consider a horn to be filled

$$\begin{array}{ccc} \Lambda^k[n] & \longrightarrow & \widetilde{U}_\alpha \\ \downarrow & & \downarrow p_\alpha \\ \Delta[n] & \xrightarrow{\lceil x \rceil} & U_\alpha \end{array}$$

for some $0 \leq k \leq n$. It factors through the pullback

$$\begin{array}{ccccc} \Lambda^k[n] & \longrightarrow & \bullet & \longrightarrow & \widetilde{U}_\alpha \\ \downarrow & & \downarrow x & \lrcorner & \downarrow p_\alpha \\ \Delta[n] & \xlongequal{\quad} & \Delta[n] & \xrightarrow{\lceil x \rceil} & U_\alpha \end{array}$$

where by the definition of U_α and \widetilde{U}_α , x is a fibration. Thus the left square admits a diagonal filler, and hence so does the outer rectangle. \square

Lemma 2.1.11. *An α -small well-ordered morphism $f: Y \rightarrow X \in W_\alpha(X)$ is a fibration if and only if $\lceil f \rceil: X \rightarrow W_\alpha$ factors through U_α .*

Proof. For ‘ \Rightarrow ’, assume that $f: Y \rightarrow X$ is a fibration. Then the pullback of f to any representable is certainly a fibration:

$$\begin{array}{ccc} \bullet & \longrightarrow & Y \\ x^*f \downarrow & \lrcorner & \downarrow f \\ \Delta[n] & \xrightarrow{x} & X. \end{array}$$

¹¹Here and throughout, by “fibration” we always mean “Kan fibration”.

so $\lceil f \rceil(x) = \lceil x^* f \rceil \in U_\alpha$, and hence $\lceil f \rceil$ factors through U_α .

Conversely, suppose $\lceil f \rceil$ factors through U_α . Then we obtain:

$$\begin{array}{ccccc} Y & \longrightarrow & \widetilde{U}_\alpha & \longrightarrow & \widetilde{W}_\alpha \\ \downarrow f & \lrcorner & \downarrow p_\alpha & \lrcorner & \downarrow \\ X & \longrightarrow & U_\alpha & \hookrightarrow & W_\alpha, \end{array}$$

where the lower composite is $\lceil f \rceil$, and the outer rectangle and the right square are by construction pullbacks. Hence so is the left square; so by Lemma 2.1.10 f is a fibration. \square

Corollary 2.1.12. *The functor U_α is representable, represented by U_α ; so $p_\alpha: \widetilde{U}_\alpha \rightarrow U_\alpha$ is strictly universal for α -small well-ordered fibrations, and weakly universal for α -small fibrations.* \square

In Section 3.5, we will strengthen this universal property, showing that while the representation of a fibration as a pullback of p_α may not be strictly unique, it is unique up to homotopy: precisely, the space of such representations is contractible.

2.2. Kan fibrancy of the universe. The previous section provides the main ingredients needed to use U_α as a universe in the sense of Section 1, and hence to give a model of the core type theory. However, to give additionally a type-theoretic universe within that model, we need to show that each U_α itself can be seen as a *type* of the model; in other words, that it is Kan. The main goal of this section is therefore to prove the following theorem:

Theorem 2.2.1. *The simplicial set U_α is a Kan complex.*

Before proceeding with the proof we will gather four useful lemmas. The first two concern *minimal fibrations*, which for the present purposes are a technical device whose details, beyond these two lemmas, are unimportant.

Lemma 2.2.2 (Quillen’s Lemma, [Qui68]). *Any fibration $f: Y \rightarrow X$ may be factored as $f = pg$, where p is a minimal fibration and g is a trivial fibration.*

Lemma 2.2.3 ([BGM59, III.5.6]; see also [May67, Cor. 11.7]). *Suppose X is contractible, with $x_0 \in X$, and $p: Y \rightarrow X$ is a minimal fibration with fiber $F := Y_{x_0}$. Then there is an isomorphism over X :*

$$\begin{array}{ccc} Y & \xrightarrow{g} & F \times X \\ & \searrow p & \swarrow \pi_2 \\ & X & \end{array}$$

For Lemma 2.2.5, the proof we give is due to André Joyal; we include details here since the original [Joy11] is not currently publicly available. For this, and again for Theorem 3.4.1 below, we make crucial use of exponentiation along cofibrations; so we pause first to establish some facts about this.

Lemma 2.2.4 (Cf. [Joy11, Lemma 0.2]). *For any map $i: A \rightarrow B$,*

1. $\Pi_i: \mathbf{sSet}/A \rightarrow \mathbf{sSet}/B$ *preserves trivial fibrations;*

and if moreover i is a cofibration, then:

2. the counit $i^* \Pi_i \rightarrow 1_{\mathbf{sSet}/A}$ is an isomorphism;
3. if $p: E \rightarrow A$ is α -small, then so is $\Pi_i p$.

Proof.

1. By adjunction, since i^* preserves cofibrations.
2. Since i is mono, $i^* \Sigma_i \cong 1_{\mathbf{sSet}/A}$; so by adjointness, $i^* \Pi_i \cong 1_{\mathbf{sSet}/A}$.
3. For any n -simplex $x: \Delta[n] \rightarrow B$, we have $(\Pi_i p)_x \cong \mathrm{Hom}_{\mathbf{sSet}/B}(x, \Pi_i p) \cong \mathrm{Hom}_{\mathbf{sSet}/B}(i^* x, p)$. As a subobject of $\Delta[n]$, $i^* x$ has only finitely many non-degenerate simplices, so $(\Pi_i p)_x$ injects into a finite product of fibers of p and is thus of size $< \alpha$. \square

Lemma 2.2.5 ([Joy11, Lemma 0.2]). *Trivial fibrations extend along cofibrations. That is, if $t: Y \rightarrow X$ is a trivial fibration and $j: X \rightarrow X'$ is a cofibration, then there exists a trivial fibration $t': Y' \rightarrow X'$ and a pullback square of the form:*

$$\begin{array}{ccc} Y & \xrightarrow{\quad} & Y' \\ t \downarrow & \lrcorner & \downarrow t' \\ X & \xrightarrow{j} & X'. \end{array}$$

Moreover, if t is α -small, then t' may be chosen to also be.

Proof. Take $t' := \Pi_j t$. By part 1 of Lemma 2.2.4, this is a trivial fibration; by part 2, $j^* Y' \cong Y$; and by part 3, it is α -small. \square

We are now ready to prove that U_α is a Kan complex.

Proof of Theorem 2.2.1. We need to show that we can extend any horn in U_α to a simplex:

$$\begin{array}{ccc} \Lambda^k[n] & \longrightarrow & U_\alpha \\ \downarrow & \nearrow & \\ \Delta[n] & & \end{array}$$

By Corollary 2.1.12, any such horn $\lceil q \rceil$ corresponds to an α -small well-ordered fibration $q: Y \rightarrow \Lambda^k[n]$. To extend $\lceil q \rceil$ to a simplex, we just need to construct an α -small fibration Y' over $\Delta[n]$ which restricts on the horn to Y :

$$\begin{array}{ccc} Y & \xrightarrow{\quad} & Y' \\ q \downarrow & \lrcorner & \downarrow q' \\ \Lambda^k[n] & \hookrightarrow & \Delta[n]. \end{array}$$

By the axiom of choice one can then extend the well-ordering of q to q' , so the map $\lceil q' \rceil: \Delta[n] \rightarrow U_\alpha$ gives the desired simplex.

By Quillen's Lemma, we can factor q as

$$Y \xrightarrow{q_t} Y_0 \xrightarrow{q_m} \Lambda^k[n],$$

where q_t is a trivial fibration and q_m is a minimal fibration. Both are still α -small: each fiber of q_t is a subset of a fiber of q , and since a trivial fibration is onto, each fiber of q_m is a quotient of a fiber of q .

By Lemma 2.2.3, we have an isomorphism $Y_0 \cong F \times \Delta^k[n]$, and hence a pullback diagram:

$$\begin{array}{ccc} Y_0 & \hookrightarrow & F \times \Delta[n] \\ \downarrow \lrcorner & & \downarrow \\ \Lambda^k[n] & \hookrightarrow & \Delta[n] \end{array}$$

By Lemma 2.2.5, we can then complete the upper square in the following diagram, with both right-hand vertical maps α -small fibrations:

$$\begin{array}{ccc} Y & \longrightarrow & Y' \\ \downarrow \lrcorner \scriptstyle q_t & & \downarrow \\ Y_0 & \hookrightarrow & F \times \Delta[n] \\ \downarrow \lrcorner \scriptstyle q_m & & \downarrow \\ \Lambda^k[n] & \hookrightarrow & \Delta[n] \end{array}$$

Since α is regular, the composite of the right-hand side is again α -small; so we are done. \square

2.3. Modelling type theory in simplicial sets. To prove that U_α carries the structure to model type theory, we will need a couple of further lemmas; firstly, that taking dependent products preserves fibrations:

Lemma 2.3.1. *Suppose $Z \xrightarrow{q} Y \xrightarrow{p} X$ are fibrations. Then the dependent product $\Pi_p q$ is a fibration over X .*

Proof. The pullback functor $p^* : \mathbf{sSet}/X \rightarrow \mathbf{sSet}/Y$ preserves trivial cofibrations (since \mathbf{sSet} is right proper and cofibrations are monomorphisms); so its right adjoint Π_p preserves fibrant objects. \square

Secondly, to model \mathbf{Id} -types, we will require well-behaved fibered path objects. The construction below may be found in [War08, Thm. 2.25]; we recall it in more elementary terms, which will be useful to us later.

Definition 2.3.2. Given a fibration $p : E \rightarrow B$, define the *fibered path object* $P_B(E)$ as the pullback

$$\begin{array}{ccc} P_B(E) & \longrightarrow & E^{\Delta[1]} \\ \downarrow \lrcorner & & \downarrow p^{\Delta[1]} \\ B & \xrightarrow{c} & B^{\Delta[1]}, \end{array}$$

the object of paths in E that are constant in B .

The “constant path” map $c: E \rightarrow E^{\Delta[1]}$ factors through $P_B(E)$; call the resulting map $r_p: E \rightarrow P_B(E)$. There are also evident source and target maps $s_p, t_p: P_B(E) \rightarrow E$. (On all of these maps, we will omit the subscripts when they are clear from context.)

Proposition 2.3.3. *For any fibration $p: E \rightarrow B$, the maps*

$$E \xrightarrow{r} P_B(E) \xrightarrow{(s,t)} E \times_B E$$

give a factorisation of the diagonal map $\Delta_p: E \rightarrow E \times_B E$ over B as a (trivial cofibration, fibration); and this is stable over B in that the pullback along any $B' \rightarrow B$ is again such a factorisation.

Proof. It is clear that these maps give a factorisation of Δ_p over B . To see that they are a trivial cofibration and a fibration respectively, consider the pullback construction of $P_B(E)$ via two intermediate stages:

$$\begin{array}{ccc} P_B(E) & \xrightarrow{\quad} & E^{\Delta[1]} \\ (s,t) \downarrow \lrcorner & & \downarrow (s, p^{\Delta[1]}, t) \\ E \times_B E & \xrightarrow{\quad} & E \times_B B^{\Delta[1]} \times_B E \\ \pi_1 \downarrow \lrcorner & & \downarrow (\pi_1, \pi_2) \\ E & \xrightarrow{\quad} & E \times_B B^{\Delta[1]} \\ \downarrow \lrcorner & & \downarrow \\ B & \xrightarrow{c} & B^{\Delta[1]} \end{array}$$

Now (s, t) is certainly a fibration, since it is a pullback of the map $E^{\Delta[1]} \rightarrow E \times_B B^{\Delta[1]} \times_B E \cong E^{1+1} \times_{B^{1+1}} B^{\Delta[1]}$, which is a fibration by the monoidal model category axioms [Hov99, Lemma 4.2.2(3)], applied to the cofibration $1 + 1 \rightarrow \Delta[1]$ and the fibration p .

Similarly, the source map $s: P_B(E) \rightarrow E$ is a trivial fibration, since it is a pullback of $E^{\Delta[1]} \rightarrow E^1 \times_{B^1} B^{\Delta[1]}$, which is one by the monoidal model category axioms. But s is a retraction of r , so r is a weak equivalence (by 2-out-of-3) and a monomorphism, so is a trivial cofibration as desired.

Finally, stability of these properties under pullback follows immediately from the stability (up to isomorphism) of the construction itself: for any $f: B' \rightarrow B$, there is a canonical isomorphism $P_{B'}(f^*E) \cong f^*P_B(E)$, commuting with the maps r, s, t . \square

We are now fully equipped for the main result of the present section:

Theorem 2.3.4. *Let α be an inaccessible cardinal¹². Then U_α carries Π -, Σ -, Id -, W -, 1 -, 0 -, and $+$ -structures.*

Moreover, if $\beta < \alpha$ is also inaccessible, then U_β gives an internal universe in U_α closed under all these constructors.

¹²I.e. infinite, regular, and strong limit; *strongly inaccessible* in some literature.

Proof. (Π -structure): Given a pair of α -small fibrations $Z \xrightarrow{q} Y \xrightarrow{p} X$, the dependent product $\Pi_p q$ in \mathbf{sSet}/X is again a fibration, by Lemma 2.3.1; it is also α -small, since α is inaccessible.

Hence by Corollary 2.1.12, the universal dependent product over $U_\alpha^{\Pi\text{-FORM}}$ is representable as the pullback of \tilde{U}_α along some map $\Pi: U_\alpha^{\Pi\text{-FORM}} \rightarrow U_\alpha$, giving the desired Π -structure.

(Σ -structure): Similarly, given α -small fibrations $Z \xrightarrow{q} Y \xrightarrow{p} X$, the composite $p \cdot q$ is again an α -small fibration. So the universal dependent sum over $U_\alpha^{\Sigma\text{-FORM}}$ is representable by some map $\Sigma: U_\alpha^{\Sigma\text{-FORM}} \rightarrow U_\alpha$.

(Id-structure): Given any α -small fibration $p: Y \rightarrow X$, consider the factorisation of the diagonal Δ_p as $Y \xrightarrow{r} P_X(Y) \xrightarrow{(s,t)} Y \times_X Y$. The fibration (s, t) is easily seen to be α -small; and by Proposition 2.3.3, r is stably orthogonal to (s, t) over X .

Applying this construction to $p_\alpha: \tilde{U}_\alpha \rightarrow U_\alpha$ itself yields, via Proposition 1.4.7, the desired Id-structure on U_α .

(W-structure): Given α -small fibrations $Z \xrightarrow{q} Y \xrightarrow{p} X$, the initial algebra $W_q \rightarrow X$ for the induced polynomial endofunctor on \mathbf{sSet}/X may be obtained as a transfinite colimit of iterations of the endofunctor; it can be shown from this description that it is again an α -small fibration (cf. [vdBM15, Thm. 3.4] and [vdBM18]).

(0-structure), (1-structure), (+-structure): straightforward.

(Internal universe.) Since $\beta < \alpha$, U_β is itself α -small; and by Theorem 2.2.1, it is Kan. So U_β is representable as the pullback of \tilde{U}_α along some $u_\beta: 1 \rightarrow U_\alpha$. Moreover, there is a natural inclusion $i: U_\beta \rightarrow U_\alpha$, with $\tilde{U}_\alpha[\beta] \cong i^* \tilde{U}_\alpha$ by construction. Together these give the desired internal universe (u_β, i) .

Finally, to see that (u_β, i) is closed under the appropriate constructors in i , note that for each of Π , Σ , and Id as constructed above, the image of the composite with i lies again in U_β , and hence factors through i ; for instance, in the case of Π ,

$$\begin{array}{ccc} U_\beta^{\Pi\text{-FORM}} & \xrightarrow{i^{\Pi\text{-FORM}}} & U_\alpha^{\Pi\text{-FORM}} \\ \downarrow \Pi & & \downarrow \Pi \\ U_\beta & \xrightarrow{i} & U_\alpha \end{array}$$

(Note that while we do already have a Π -structure (and so on) on U_β as constructed in the first parts of this theorem, those choices of the structure do not automatically commute with i .) \square

Corollary 2.3.5. *Let $\beta < \alpha$ be inaccessible cardinals. Then there is a model of dependent type theory in \mathbf{sSet}_{U_α} with all the logical constructors of Section A.2, and a universe (given by U_β) closed under these constructors.* \square

Assuming initiality (Conjecture 1.2.9), this implies the existence of a morphism $\mathcal{C}(\mathbf{T}) \rightarrow \mathbf{sSet}_{U_\alpha}$, interpreting the syntax of Martin-Löf type theory in simplicial sets. Even without assuming initiality, it gives us the operations with which to heuristically interpret individual judgements of the syntax by hand. We therefore freely make notational use of the interpretation, writing $\llbracket \mathcal{J} \rrbracket$ for the interpretation of a judgement \mathcal{J} .

In doing so, we will make several systematic abuses of notation. Firstly, referring in the syntax to fibrations, we will write E rather than $\lceil E \rceil$, and so on, whenever some choice of name $\lceil E \rceil: B \longrightarrow U_\alpha$ for the fibration is understood; and conversely, referring to the interpretation of a type $\Gamma \vdash T$ type, we use $\llbracket T \rrbracket$ to refer to the fibration over $\llbracket \Gamma \rrbracket$ given by pulling back \widetilde{U}_α along the literal interpretation $\llbracket \Gamma \vdash T \text{ type} \rrbracket: \llbracket \Gamma \rrbracket \longrightarrow U_\alpha$.

As a first characteristic of the model, we note that both of the extra principles on equality of functions hold.

Proposition 2.3.6. *The η -rule and functional extensionality rules of Section A.4 hold in the simplicial model.*

Proof. The η -rule follows immediately from our use of categorical exponentials to interpret Π -types, by the uniqueness in the categorical universal property.

For functional extensionality, Garner [Gar09, Sec. 5] shows that it holds just if each product of identity types,

$$f, g: \Pi_{x:A} B(x) \vdash \Pi_{x:A} \text{Id}_{B(x)}(\text{app}(f, x), \text{app}(g, x)) \text{ type}$$

admits the structure given by the rules for the identity type on the corresponding product types,

$$f, g: \Pi_{x:A} B(x) \vdash \text{Id}_{\Pi_{x:A} B(x)}(f, g) \text{ type}.$$

So it is enough to show that for any pair of (α -small, well-ordered) fibrations $Z \xrightarrow{q} Y \xrightarrow{p} X$, given by names

$$\lceil Y \rceil: X \longrightarrow U_\alpha, \quad \lceil Z \rceil: Y \longrightarrow U_\alpha,$$

the interpretation of the product of identity types

$$\llbracket \Pi_{x:Y} \text{Id}_{Z(x)}(\text{app}(f, x), \text{app}(g, x)) \rrbracket \cong \Pi_p(\text{P}_Y Z),$$

gives a suitably stable path object for the interpretation of the product types,

$$\llbracket \text{Id}_{\Pi_{x:Y} Z(x)}(f, g) \rrbracket \cong \Pi_p Z.$$

For this, it is clear that $\Pi_p(s, t): \Pi_p(\text{P}_Y Z) \longrightarrow \Pi_p(Z \times_Y Z) \cong \Pi_p Z \times_X \Pi_p Z$ is a fibration, since Π_p preserves fibrations (Lemma 2.3.1). Similarly, $\Pi_p r_q: \Pi_p Z \longrightarrow \Pi_p(\text{P}_Y Z)$ is a cofibration since Π_p preserves monomorphisms; and it is a weak equivalence, since Π_p preserves trivial fibrations (Lemma 2.2.5), and so the retraction $\Pi_p s_q: \Pi_p(\text{P}_Y Z) \longrightarrow \Pi_p Z$ is again a trivial fibration. Finally, by the Beck-Chevalley condition in an LCCC, the entire construction is stable under pullback in X , as required. \square

It now remains only to show that the Univalence Axiom holds in this model.

3. UNIVALENCE

In this section, we will introduce the Univalence Axiom, and show that it holds in the simplicial model.

The proof of this involves both simplicial and type-theoretic components; we keep these separate, as far as possible. First of all (Section 3.1), we define univalence type-theoretically and state the Univalence Axiom; next, we define an analogous simplicial concept of univalence (Section 3.2); we then show that via the simplicial model, the two notions coincide

(Section 3.3). Finally, in Section 3.4, we prove our main theorem: that U_α is univalent (using the simplicial sense), and hence that the Univalence Axiom holds in the simplicial model of type theory. Lastly, in Section 3.5, we discuss an alternative formulation of simplicial univalence, and so obtain an up-to-homotopy uniqueness statement for the weak universal property of U_α .

Once again, we freely use the syntax of type theory as a notation; to avoid formal dependence on Conjecture 1.2.9, the reader should translate each individual syntactic expression used into the language of contextual categories.

3.1. Type-theoretic univalence. To state the univalence axiom, we first need to define a few basic notions in the type theory.

Definition 3.1.1 (Joyal). Let $f: A \rightarrow B$ be a function in some context Γ , i.e. $\Gamma \vdash f: [A, B]$ (where the function type $[A, B]$ is defined using Π , as described in Section A.2).

- A *left homotopy inverse* for f is a function $g: B \rightarrow A$, together with a homotopy $g \cdot f \simeq 1_A$. Formally, we define the type $\text{LInv}(f)$ of left homotopy inverses to f :

$$\Gamma \vdash \text{LInv}(f) := \Sigma_{g:[B,A]} \Pi_{x:A} \text{Id}_A(g(f(x)), x) \text{ type}$$

- Analogously, we define the type $\text{RInv}(f)$ of *right homotopy inverses*:

$$\Gamma \vdash \text{RInv}(f) := \Sigma_{g:[B,A]} \Pi_{y:B} \text{Id}_B(f(g(y)), y) \text{ type}$$

- We say $f: A \rightarrow B$ is a *homotopy isomorphism* (or more briefly, an h-isomorphism) if it is equipped with both a left and a right inverse:

$$\Gamma \vdash \text{isHlso}(f) := \text{LInv}(f) \times \text{RInv}(f) \text{ type}$$

- For any types A and B , we thus have the type of h-isomorphisms from A to B :

$$\Gamma \vdash \text{Hlso}(A, B) := \Sigma_{f:[A,B]} \text{isHlso}(f)$$

It may perhaps be surprising that we use homotopy isomorphisms rather than the more familiar homotopy equivalences, with a single two-sided homotopy inverse. The reason is that while a map carries either structure if and only if it carries the other, the type, or object, of such structures on a map is different. In particular, the analogue of Lemma 3.3.4 for homotopy equivalences does not hold; for further discussion of these issues, see [Uni13, Ch. 4].

Example 3.1.2. For any type B , the identity function on B is canonically an h-isomorphism.

Suppose now that A is any type, and $x: A \vdash B(x)$ type a family of types over A . By the identity elimination rule, we can derive

$$x, y: A, u: \text{Id}_A(x, y) \vdash w_{x,y,u}: \text{Hlso}(B(x), B(y)).$$

This can equivalently be seen as a map

$$x, y: A \vdash w_{x,y}: [\text{Id}_A(x, y), \text{Hlso}(B(x), B(y))].$$

Definition 3.1.3. We say the family $B(x)$ is *univalent* if for each x, y , the map $w_{x,y}$ is itself a homotopy isomorphism:

$$\vdash \text{isUnivalent}(x: A. B(x)) := \Pi_{x,y:A} \text{isHlso}(w_{x,y}).$$

Axiom 3.1.4. *The Univalence Axiom, for a given type-theoretic universe U , is the statement that the canonical family \mathbf{El} of types over U is univalent.*

Informally, the Univalence Axiom says that just as elements of the universe correspond to types, so equalities in the universe correspond to equivalences between types. In particular, since every statement or construction must respect propositional equality, the Univalence Axiom stipulates that the language can never distinguish between equivalent types.

3.2. Simplicial univalence. To define a simplicial notion of univalence, we first need to construct the *object of weak equivalences* between fibrations $p_1: E_1 \rightarrow B$ and $p_2: E_2 \rightarrow B$ over a common base. In other words, we want an object representing the functor sending $(X, f) \in \mathbf{sSet}/B$ to the set $\mathbf{Eq}_X(f^*E_1, f^*E_2)$. As we did for \mathbf{U}_α , we proceed in two steps, first exhibiting it as a subfunctor of a functor more easily seen (or already known) to be representable.

For the remainder of the section, fix fibrations E_1, E_2 as above over a base B . Since \mathbf{sSet} is locally Cartesian closed, we can construct the exponential object between them:

Definition 3.2.1. Let $\mathbf{Hom}_B(E_1, E_2) \rightarrow B$ denote the internal hom from E_1 to E_2 in \mathbf{sSet}/B .

Then for any X , a map $X \rightarrow \mathbf{Hom}_B(E_1, E_2)$ corresponds to a map $f: X \rightarrow B$, together with a map $u: f^*E_1 \rightarrow f^*E_2$ over X .

Together with the Yoneda lemma, this implies the explicit description: an n -simplex of $\mathbf{Hom}_B(E_1, E_2)$ is a pair

$$(b: \Delta[n] \rightarrow B, u: b^*E_1 \rightarrow b^*E_2).$$

Lemma 3.2.2. $\mathbf{Hom}_B(E_1, E_2) \rightarrow B$ is a Kan fibration.

Proof. Follows immediately from Lemma 2.3.1, since the exponential is a special case of dependent products. \square

Within $\mathbf{Hom}_B(E_1, E_2)$, we now want to construct the subobject of weak equivalences.

Lemma 3.2.3. *Let $f: E_1 \rightarrow E_2$ be a weak equivalence over B , and suppose $g: B' \rightarrow B$. Then the induced map between pullbacks $g^*E_1 \rightarrow g^*E_2$ is a weak equivalence.*

Proof. The pullback functor $g^*: \mathbf{sSet}/B \rightarrow \mathbf{sSet}/B'$ preserves trivial fibrations; so by Ken Brown's Lemma [Hov99, Lemma 1.1.12], it preserves all weak equivalences between fibrant objects. \square

Thus, weak equivalences from E_1 to E_2 form a subfunctor of the functor of maps from E_1 to E_2 . To show that this is representable, we need just to show:

Lemma 3.2.4. *Let $f: E_1 \rightarrow E_2$ be a morphism over B . If for each simplex $b: \Delta[n] \rightarrow B$ the induced map $f_b: b^*E_1 \rightarrow b^*E_2$ is a weak equivalence, then f is a weak equivalence.*

Proof. Without loss of generality, B is connected; otherwise, apply the result over each connected component separately. Take some vertex $b: \Delta[0] \rightarrow B$, and set $F_i := b^*E_i$. Now

for any vertex $e: \Delta[0] \rightarrow F_1$, and any $n \geq 1$, we have by the long exact sequence for a fibration:

$$\begin{array}{ccccccccc} \pi_{n+1}(B, b) & \longrightarrow & \pi_n(F_1, e) & \longrightarrow & \pi_n(E_1, e) & \longrightarrow & \pi_n(B, b) & \longrightarrow & \pi_{n-1}(F_1, e) \\ \downarrow 1 & & \downarrow \pi_n(f_b) & & \downarrow \pi_n(f) & & \downarrow 1 & & \downarrow \pi_{n-1}(f_b) \\ \pi_{n+1}(B, b) & \longrightarrow & \pi_n(F_2, f(e)) & \longrightarrow & \pi_n(E_2, f(e)) & \longrightarrow & \pi_n(B, b) & \longrightarrow & \pi_{n-1}(F_2, f(e)) \end{array}$$

Each $\pi_n(f_b)$ is an isomorphism, so by the Five Lemma, so is $\pi_n(f)$, for $n \geq 1$.

The case $n = 0$ is the same in spirit, but requires a little more work since the Five Lemma is unavailable. We have a square

$$\begin{array}{ccc} \pi_0(F_1) & \xrightarrow{\pi_0(i_1)} & \pi_0(E_1) \\ \downarrow \pi_0(f_b) & & \downarrow \pi_0(f) \\ \pi_0(F_2) & \xrightarrow{\pi_0(i_2)} & \pi_0(E_2) \end{array}$$

with both horizontal arrows surjections, and $\pi_0(f_b)$ an isomorphism. To show that $\pi_0(f)$ is an isomorphism, it therefore suffices to show that for each $x \in \pi_0(E_1)$, the restriction of f_b to a map of fibers

$$\pi_0(i_1)^{-1}(x) \longrightarrow \pi_0(i_2)^{-1}(\pi_0(f)(x))$$

is again an isomorphism. But this follows from the continuation of the long exact sequence to an exact sequence of pointed sets:

$$\begin{array}{ccccc} \pi_1(B, b) & \longrightarrow & \pi_0(F_1, e) & \xrightarrow{\pi_0(i_1)} & \pi_0(E_1, x) \\ \downarrow 1 & & \downarrow \pi_0(f_b) & & \downarrow \pi_0(f) \\ \pi_1(B, b) & \longrightarrow & \pi_0(F_2, f(e)) & \xrightarrow{\pi_0(i_2)} & \pi_0(E_2, f(e)) \end{array}$$

where e is any point of F_1 such that $[e] = x$.

Thus $\pi_n(f)$ is an isomorphism for each $n \geq 0$ and basepoint $e \in E_1$; so f is a weak equivalence. \square

Definition 3.2.5. Take $\mathbf{Eq}_B(E_1, E_2)$ to be the subobject of $\mathbf{Hom}_B(E_1, E_2)$ consisting of all n -simplices

$$(b: \Delta[n] \rightarrow B, w: b^*E_1 \rightarrow b^*E_2)$$

such that w is a weak equivalence. (By Lemma 3.2.3, this indeed defines a simplicial subset.)

From Lemma 3.2.4, we immediately have:

Corollary 3.2.6. *Let $(f, u): X \rightarrow \mathbf{Hom}_B(E_1, E_2)$. Then u is a weak equivalence if and only if (f, u) factors through $\mathbf{Eq}_B(E_1, E_2)$.*

Thus, maps $X \rightarrow \mathbf{Eq}_B(E_1, E_2)$ correspond to pairs of maps

$$(f: X \rightarrow B, w: f^*E_1 \rightarrow f^*E_2),$$

where w is a weak equivalence. \square

While Lemma 3.2.4 was stated just as required by representability, its proof actually gives a slightly stronger statement:

Lemma 3.2.7. *Let $f: E_1 \rightarrow E_2$ be a morphism over B . If for some vertex $b: \Delta[0] \rightarrow B$ in each connected component the map of fibers $f_b: b^*E_1 \rightarrow b^*E_2$ is a weak equivalence, then f is a weak equivalence.* \square

Corollary 3.2.8. *The map $\mathbf{Eq}_B(E_1, E_2) \rightarrow B$ is a fibration.*

Proof. Suppose we wish to fill a square:

$$\begin{array}{ccc} \Lambda^k[n] & \longrightarrow & \mathbf{Eq}_B(E_1, E_2) \\ \downarrow i & \nearrow \text{dotted} & \downarrow \\ \Delta[n] & \xrightarrow{b} & B \end{array}$$

By the universal property of $\mathbf{Eq}_B(E_1, E_2)$ this corresponds to showing that we can extend a weak equivalence $w: i^*b^*E_1 \rightarrow i^*b^*E_2$ over $\Lambda^k[n]$ to a weak equivalence $\bar{w}: b^*E_1 \rightarrow b^*E_2$ over $\Delta[n]$.

By Lemma 3.2.2, we can certainly find some map \bar{w} extending w . But then since $\Delta[n]$ is connected, Lemma 3.2.7 implies that \bar{w} is a weak equivalence. \square

While on the subject, we collect a proposition which is not required for the definition of univalence, but which will be useful later:

Proposition 3.2.9. *Suppose that E_1, E'_1, E_2, E'_2 are fibrations over a common base B , and $w_1: E'_1 \rightarrow E_1$, $w_2: E_2 \rightarrow E'_2$ are weak equivalences over B . Then the induced map $\mathbf{Eq}_B(w_1, w_2): \mathbf{Eq}_B(E_1, E_2) \rightarrow \mathbf{Eq}_B(E'_1, E'_2)$ is a weak equivalence.*

$$\begin{array}{ccccc} E'_1 & \xrightarrow{w_1} & E_1 & & E_2 & \xrightarrow{w_2} & E'_2 \\ & \searrow p'_1 & \downarrow p_1 & & \downarrow p_2 & \swarrow p'_2 & \\ & & B & & & & \end{array}$$

Proof. As weak equivalences between fibrations, w_1 and w_2 are fibered homotopy equivalences over B . Choosing fibered homotopy inverses v_1, v_2 for w_1 and w_2 respectively gives a homotopy inverse $\mathbf{Hom}_B(v_1, v_2)$ for $\mathbf{Hom}_B(w_1, w_2): \mathbf{Hom}_B(E_1, E_2) \rightarrow \mathbf{Hom}_B(E'_1, E'_2)$. But by Lemma 3.2.7, the image of a homotopy in \mathbf{Hom} whose endpoints lie in \mathbf{Eq} must lie entirely in \mathbf{Eq} ; so the restriction $\mathbf{Eq}_B(v_1, v_2)$ gives a homotopy inverse for $\mathbf{Eq}_B(w_1, w_2)$, as desired. \square

We are now ready to define univalence.

Let $p: E \rightarrow B$ be a fibration. We then have two fibrations over $B \times B$, given by pulling back E along the projections. Call the object of weak equivalences between these $\mathbf{Eq}(E) := \mathbf{Eq}_{B \times B}(\pi_1^*E, \pi_2^*E)$. Concretely, simplices of $\mathbf{Eq}(E)$ are triples

$$(b_1, b_2 \in B_n, w: b_1^*E \rightarrow b_2^*E).$$

By Corollary 3.2.6, a map $f: X \rightarrow \mathbf{Eq}(E)$ corresponds to a pair of maps $f_1, f_2: X \rightarrow B$ together with a weak equivalence $f_1^*E \rightarrow f_2^*E$ over X . In particular, there is a “diagonal” map $\delta_E: B \rightarrow \mathbf{Eq}(E)$ corresponding to the triple $(1_B, 1_B, 1_E)$, sending a simplex $b \in B_n$ to the triple $(b, b, 1_{E_b})$.

There are also source and target maps $s, t: \mathbf{Eq}(E) \rightarrow B$, given by the composites $\mathbf{Eq}(E) \rightarrow B \times B \xrightarrow{\pi_i} B$, sending (b_1, b_2, w) to b_1 and b_2 respectively. These are both retractions of δ ; and by Corollary 3.2.8, if B is fibrant then they are moreover fibrations.

Definition 3.2.10. A fibration $p: E \rightarrow B$ is *univalent* if the diagonal map $\delta_E: B \rightarrow \mathbf{Eq}(E)$ is a weak equivalence.

Since δ_E is always a monomorphism (thanks to its retractions), this is equivalent to saying that $B \rightarrow \mathbf{Eq}(E) \rightarrow B \times B$ is a (trivial cofibration, fibration) factorisation of the diagonal $\Delta_B: B \rightarrow B \times B$, i.e. that $\mathbf{Eq}(E)$ is a *path object* for B .

We conclude this section with a few examples, and non-examples, of univalent fibrations.

Examples 3.2.11.

- (1) The canonical map $X \rightarrow 1$ is univalent if and only if the space of homotopy auto-equivalences of X is contractible.
- (2) The identity map $X \rightarrow X$ is univalent if and only if X is either empty or contractible. In particular, the identity map $1 + 1 \rightarrow 1 + 1$ is *not* univalent: it has two fibers which are equivalent, over points that are not connected by any path.
- (3) Any fibration weakly equivalent to a univalent fibration is itself univalent (essentially, by Proposition 3.2.9).

3.3. Equivalence of type-theoretic and simplicial univalence. Having defined the type-theoretic and simplicial notions of univalence, we now wish to show that they coincide. As ever, we make essential use of representability; in particular, we work with the interpretations of type-theoretic notions entirely via their universal properties. With this in view, we need to define what are represented by the interpretations of \mathbf{LInv} , \mathbf{Hlso} , etc.

Definition 3.3.1. Let $p_1: E_1 \rightarrow B$, $p_2: E_2 \rightarrow B$ be fibrations over a common base (as in Definition 3.2.1).

Define $\mathbf{HomLInv}_B(E_1, E_2)$ to be the set of *maps with a left homotopy inverse* from E_1 to E_2 , i.e. triples (f, g, H) , where $f: E_1 \rightarrow E_2$ and $g: E_2 \rightarrow E_1$ are maps over B , and H is a fibred homotopy from $g \cdot f$ to 1_{E_1} , defined using the fibred path space $P_B(E_1)$ (as used for the \mathbf{Id} -structure in the proof of Theorem 2.3.4).

Similarly, define $\mathbf{HomRInv}_B(E_1, E_2)$ to consist of triples (f, g, H) , where f, g are as before, and H is now a fibred homotopy from $f \cdot g$ to 1_{E_2} , defined using $P_B(E_2)$.

Finally, these both come with evident projections to $\mathbf{Hom}_B(E_1, E_2)$; define $\mathbf{Hlso}_B(E_1, E_2)$ as the pullback $\mathbf{HomLInv}_B(E_1, E_2) \times_{\mathbf{Hom}_B(E_1, E_2)} \mathbf{HomRInv}_B(E_1, E_2)$.

Lemma 3.3.2. *Let B, E_1, E_2 be as above; additionally, suppose they are given by names $\ulcorner B \urcorner: 1 \rightarrow U_\alpha$, $\ulcorner E_i \urcorner: B \rightarrow U_\alpha$. Then for any $f: X \rightarrow B$, there are horizontal isomorphisms as in the diagram below, making the diagram commute, and natural in (X, f) .*

$$\begin{array}{ccc}
\mathrm{Hom}_B(X, \llbracket \mathbf{H} \mathbf{I} \mathbf{s} \mathbf{o} (E_1, E_2) \rrbracket) & \cdots \cong \cdots & \mathbf{H} \mathbf{I} \mathbf{s} \mathbf{o}_X(f^* E_1, f^* E_2) \\
\swarrow \quad \searrow & & \swarrow \quad \searrow \\
\mathrm{Hom}_B(X, \llbracket \mathbf{H} \mathbf{o} \mathbf{m} \mathbf{L} \mathbf{I} \mathbf{n} \mathbf{v} (E_1, E_2) \rrbracket) & \cdots \cong \cdots & \mathbf{H} \mathbf{o} \mathbf{m} \mathbf{L} \mathbf{I} \mathbf{n} \mathbf{v}_X(f^* E_1, f^* E_2) \\
\searrow \quad \swarrow & & \searrow \quad \swarrow \\
\mathrm{Hom}_B(X, \llbracket \mathbf{H} \mathbf{o} \mathbf{m} \mathbf{R} \mathbf{I} \mathbf{n} \mathbf{v} (E_1, E_2) \rrbracket) & \cdots \cong \cdots & \mathbf{H} \mathbf{o} \mathbf{m} \mathbf{R} \mathbf{I} \mathbf{n} \mathbf{v}_X(f^* E_1, f^* E_2) \\
\swarrow \quad \searrow & & \swarrow \quad \searrow \\
\mathrm{Hom}_B(X, \llbracket [E_1, E_2] \rrbracket) & \cdots \cong \cdots & \mathrm{Hom}_X(f^* E_1, f^* E_2)
\end{array}$$

(Here $\llbracket - \rrbracket$ denotes the interpretation of type theory, as described following Corollary 2.3.5; and $[-, -]$ is the ordinary function type, taken as a special case of Π -types.)

Proof. This is essentially a routine verification; we prove just the first case, that of $\llbracket [E_1, E_2] \rrbracket$. For this, we need to give a natural isomorphism $\mathrm{Hom}_B(X, \llbracket [E_1, E_2] \rrbracket) \cong \mathrm{Hom}_X(f^* E_1, f^* E_2)$; in other words, to show that $\llbracket [E_1, E_2] \rrbracket$ is the exponential between E_1 and E_2 in \mathbf{sSet}/B .

Recall that by definition, $\llbracket [E_1, E_2] \rrbracket$ is constructed as the pullback of \tilde{U}_α along the map $\Pi \cdot \ulcorner (E_1, E_2) \urcorner: B \longrightarrow U_\alpha$:

$$\begin{array}{ccccc}
\llbracket [E_1, E_2] \rrbracket & \longrightarrow & \Pi_{A_{\mathrm{gen}} \rightarrow B_{\mathrm{gen}}} B_{\mathrm{gen}} & \longrightarrow & \tilde{U}_\alpha \\
\downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow \\
B & \xrightarrow{\ulcorner (E_1, E_2) \urcorner} & U_\alpha & \xrightarrow{\Pi\text{-FORM}} & U_\alpha
\end{array}$$

$\llbracket [E_1, E_2] \rrbracket$ is thus a pullback of the dependent product of the universal pair of fibrations over $U_\alpha^{\Pi\text{-FORM}}$, and so by the Beck-Chevalley condition is a dependent product for the pullbacks of these fibrations along $\ulcorner (E_1, E_2) \urcorner$. But these pullbacks are isomorphic to E_1 , $E_1 \times_B E_2$, by the two pullbacks lemma and the construction of A_{gen} , B_{gen} as pullbacks of $\tilde{U}_\alpha \rightarrow U_\alpha$.

$$\begin{array}{ccccc}
& E_1^* E_2 & \longrightarrow & B_{\mathrm{gen}} & \\
\Pi_{E_1}(E_1^* E_2) & \searrow \lrcorner & \longrightarrow & \Pi_{A_{\mathrm{gen}}} B_{\mathrm{gen}} & \searrow \lrcorner \\
& E_1 & \longrightarrow & A_{\mathrm{gen}} & \\
\downarrow \lrcorner & \swarrow \lrcorner & & \swarrow \lrcorner & \\
B & \longrightarrow & U_\alpha^{\Pi\text{-FORM}} & &
\end{array}$$

So $\llbracket [E_1, E_2] \rrbracket$ is the dependent product of $E_1 \times_B E_2 \rightarrow E_1$ along $E_1 \rightarrow B$; but this is exactly the usual construction of exponentials in slices from dependent products [Joh02, A1.5.2]. \square

We also note, from the proof of the preceding lemma:

Corollary 3.3.3. *There is a natural isomorphism over B :*

$$\llbracket [E_1, E_2] \rrbracket \cong \mathbf{Hom}_B(E_1, E_2).$$

\square

Following this, we define $\mathbf{HIso}_B(E_1, E_2) := \llbracket \mathbf{Hiso}_B(E_1, E_2) \rrbracket$, and $\mathbf{HomLInv}$, $\mathbf{HomRInv}$ similarly.

Lemma 3.3.4. *The map $\mathbf{HIso}_B(E_1, E_2) \rightarrow \mathbf{Hom}_B(E_1, E_2)$ factors through $\mathbf{Eq}_B(E_1, E_2)$; and the resulting map $\mathbf{HIso}_B(E_1, E_2) \rightarrow \mathbf{Eq}_B(E_1, E_2)$ is a trivial fibration.*

Proof. The given map $\mathbf{HIso}_B(E_1, E_2) \rightarrow [E_1, E_2] \cong \mathbf{Hom}_B(E_1, E_2)$ corresponds, under the isomorphisms of Lemma 3.3.2, to the maps on hom-sets

$$(1) \quad \begin{aligned} \mathrm{Hom}_B(X, \mathbf{HIso}_B(E_1, E_2)) &\cong \mathrm{Hiso}_X(f^*E_1, f^*E_2) \\ &\rightarrow \mathrm{Hom}_X(f^*E_1, f^*E_2) \\ &\cong \mathrm{Hom}_B(X, \mathbf{Hom}_B(E_1, E_2)) \end{aligned}$$

where the middle map just forgets the chosen homotopy inverses of an h-isomorphism. But since any map admitting both homotopy inverses is a weak equivalence, the natural map

$$\mathrm{Hiso}_X(f^*E_1, f^*E_2) \rightarrow \mathrm{Hom}_X(f^*E_1, f^*E_2)$$

factors through $\mathrm{Eq}_X(f^*E_1, f^*E_2)$; so by Yoneda, $\mathbf{HIso}_B(E_1, E_2) \rightarrow \mathbf{Hom}_B(E_1, E_2)$ factors through $\mathbf{Eq}_B(E_1, E_2)$.

Thus, we obtain the desired map $\mathbf{HIso}_B(E_1, E_2) \rightarrow \mathbf{Eq}_B(E_1, E_2)$, corresponding to the forgetful function $\mathrm{Hiso}_X(f^*E_1, f^*E_2) \rightarrow \mathrm{Eq}_X(f^*E_1, f^*E_2)$.

Combining this with the left-hand pullback square in Lemma 3.3.2, we can consider $\mathbf{HIso}_B(E_1, E_2)$ as the pullback:

$$\begin{array}{ccccc} & & \mathbf{HIso}_B(E_1, E_2) & & \\ & \swarrow & \downarrow & \searrow & \\ \mathbf{EqLInv}_B(E_1, E_2) & & & & \mathbf{EqRInv}_B(E_1, E_2) \\ \downarrow \swarrow & & \mathbf{Eq}_B(E_1, E_2) & & \downarrow \searrow \\ \mathbf{HomLInv}_B(E_1, E_2) & & & & \mathbf{HomRInv}_B(E_1, E_2) \\ & \swarrow \downarrow \searrow & \downarrow & \swarrow \downarrow \searrow & \\ & \mathbf{Hom}_B(E_1, E_2) & & & \end{array}$$

where \mathbf{EqLInv} , \mathbf{EqRInv} are defined by the pullbacks above, and represent weak equivalences equipped with a left (resp. right) homotopy inverse. To show that $\mathbf{HIso}_B(E_1, E_2) \rightarrow \mathbf{Eq}_B(E_1, E_2)$ is a trivial fibration, it thus suffices to show that the maps

$$\begin{aligned} \mathbf{EqLInv}_B(E_1, E_2) &\rightarrow \mathbf{Eq}_B(E_1, E_2) \\ \mathbf{EqRInv}_B(E_1, E_2) &\rightarrow \mathbf{Eq}_B(E_1, E_2) \end{aligned}$$

are each trivial fibrations. We show this in the following two lemmas.

Lemma 3.3.5. *For B , E_1 , E_2 as above, the map*

$$\mathbf{EqLInv}_B(E_1, E_2) \rightarrow \mathbf{Eq}_B(E_1, E_2)$$

So since $F_1 \longrightarrow X$ is a fibration, we just need to show that the left-hand map of pushouts, induced by

$$\begin{array}{ccccc}
 & i^*F_1 & \longrightarrow & F_1 & \\
 & \downarrow \iota_1 & & \downarrow \iota_1 & \\
 i^*F_1 & \xrightarrow{\iota_0} & i^*F_1 \times \Delta[1] & & \\
 \downarrow w & \searrow & \searrow & \searrow & \\
 i^*F_2 & & & F_1 \xrightarrow{\iota_0} F_1 \times \Delta[1] & \\
 & \searrow & & \downarrow \bar{w} & \\
 & & & F_2 &
 \end{array}$$

is a trivial cofibration. For convenience, call this map t .

To see that t is a weak equivalence, consider it in the square

$$\begin{array}{ccc}
 (i^*F_1 \times \Delta[1]) +_{i^*F_1} F_1 & \longrightarrow & F_1 \times \Delta[1] \\
 \downarrow & & \downarrow \\
 i^*F_2 +_{i^*F_1} ((i^*F_1 \times \Delta[1]) +_{i^*F_1} F_1) & \xrightarrow{t} & F_2 +_{F_1} (F_1 \times \Delta[1]).
 \end{array}$$

The top map is a trivial cofibration by the pushout-product property; the vertical maps are pushouts of w and \bar{w} along cofibrations, so are also weak equivalences; and so by 2-out-of-3, t is a weak equivalence.

On that other hand, to see that t is a cofibration, consider it as induced by maps t_0, t_1 as in:

$$\begin{array}{ccc}
 i^*F_1 & \longrightarrow & F_1 \\
 \downarrow & & \downarrow t_1 \\
 i^*F_2 +_{i^*F_1} (i^*F_1 \times \Delta[1]) & \xrightarrow{t_0} & F_2 +_{F_1} (F_1 \times \Delta[1]).
 \end{array}$$

Here t_0 is isomorphic to the inclusion

$$i^*(F_2 +_{F_1} (F_1 \times \Delta[1])) \hookrightarrow F_2 +_{F_1} (F_1 \times \Delta[1])$$

(since pulling back preserves products and pushouts), so is mono. Next, i_0 and i_1 have disjoint images, so t_1 is also mono. Finally, the intersection of the images of t_0 and t_1 is exactly the image of i^*F_1 ; so t , as the induced map from $(i^*F_2 +_{i^*F_1} (i^*F_1 \times \Delta[1])) +_{i^*F_1} F_1$, is mono as desired.

Thus t is a trivial cofibration, completing the proof of the lemma. \square

Lemma 3.3.6. *For B, E_1, E_2 as above, the map*

$$\mathbf{EqRInv}_B(E_1, E_2) \longrightarrow \mathbf{Eq}_B(E_1, E_2)$$

is a trivial fibration. Equivalently, right homotopy inverses to equivalences between fibrant objects extend along cofibrations.

Proof. We must provide lifts against any cofibration $i: Y \hookrightarrow X$:

$$\begin{array}{ccc} Y & \longrightarrow & \mathbf{EqRInv}_B(E_1, E_2) \\ i \downarrow & \nearrow & \downarrow \\ X & \longrightarrow & \mathbf{Eq}_B(E_1, E_2) \end{array}$$

Analogously to the previous lemma, and again writing $f: X \rightarrow B$, $F_i := f^*E_i$, the square corresponds to a weak equivalence $\bar{w}: F_1 \rightarrow F_2$ over X together with a fibered right homotopy inverse to $w := i^*\bar{w}$, i.e. $r: i^*F_2 \rightarrow i^*F_1$ and a homotopy $H: w \cdot r \simeq 1_{i^*F_2}$ over Y ;

$$\begin{array}{ccccc} i^*F_1 & \xrightarrow{\quad} & F_1 & \xrightarrow{\quad \bar{w} \quad} & F_2 \\ & \searrow w & & & \\ & & i^*F_2 & \xrightarrow{\quad} & F_2 \\ & \nearrow r & & & \\ & & Y & \xrightarrow{\quad} & X \end{array}$$

and a filler corresponds to a fibered right homotopy inverse (\bar{r}, \bar{H}) for \bar{w} , extending (r, H) .

Again, putting these conditions together, we see that they correspond to filling another square:

$$\begin{array}{ccc} i^*F_2 & \xrightarrow{(r, H)} i^*F_1 \times_{i^*F_2} P_Y(i^*F_2) & \longrightarrow F_1 \times_{F_2} P_X F_2 \\ \downarrow & \nearrow (\bar{r}, \bar{H}) & \downarrow \text{ev}_1 \cdot \pi_1 \\ F_2 & \xrightarrow{1} & F_2 \end{array}$$

where the pullbacks are just the fibered mapping path spaces.

$$\begin{array}{ccc} i^*F_1 \times_{i^*F_2} P_Y(i^*F_2) & \longrightarrow & P_Y(i^*F_2) \\ \downarrow \lrcorner & & \downarrow \text{ev}_0 \\ i^*F_1 & \xrightarrow{w} & i^*F_2 \end{array} \quad \begin{array}{ccc} F_1 \times_{F_2} P_X F_2 & \longrightarrow & P_X F_2 \\ \downarrow \lrcorner & & \downarrow \text{ev}_0 \\ F_1 & \xrightarrow{\bar{w}} & F_2 \end{array}$$

Now $i^*F_2 \hookrightarrow F_2$ is certainly a cofibration; so to provide the filler, it suffices to show that the right-hand map is a trivial fibration. As the target map from a mapping path space, it is certainly a fibration. To see that it is a weak equivalence, consider the triangle

$$\begin{array}{ccc} F_1 & \xrightarrow{x \mapsto (x, c_{\bar{w}}x)} & F_1 \times_{F_2} P_X F_2 \\ & \searrow \bar{w} & \downarrow \text{ev}_0 \\ & & F_2 \end{array}$$

The top map is the inclusion of a deformation retraction, so is a weak equivalence; so by 2-out-of-3, the source map ev_0 is a weak equivalence. But ev_1 is homotopic to ev_0 , so is also a weak equivalence, as required. \square

Putting these two lemmas together concludes the proof of Lemma 3.3.4: **HIso** is trivially fibrant over **Eq**. \square

Theorem 3.3.7. *Let B be a Kan complex, $p: E \rightarrow B$ a fibration; choose some names $\ulcorner B \urcorner: 1 \rightarrow U_\alpha$, $\ulcorner E \urcorner: B \rightarrow U_\alpha$ for these. Then E is simplicially univalent if and only if the type $\text{isUnivalent}(E)$ is inhabited in the model.*

Proof. By definition, $p: E \rightarrow B$ is type-theoretically univalent when there exists a section of the type $\llbracket x_1, x_2: B \vdash \text{isHlso}(w_{x_1, x_2}) \text{ type} \rrbracket$ over $B \times B$ (where w_{x_1, x_2} is as in Definition 3.1.3). By Lemma 3.3.2 this is equivalent to the map

$$w_E := \llbracket x_1, x_2: B, \text{ld}_B(x_1, x_2) \vdash w_{x_1, x_2}(p) : \text{Hlso}(E(x_1), E(x_2)) \rrbracket$$

admitting the structure of a homotopy isomorphism, or equivalently being a weak equivalence.

$$\begin{array}{ccc} \llbracket x_1, x_2: B, p: \text{ld}_B(x_1, x_2) \rrbracket & \xrightarrow{w_E} & \llbracket x_1, x_2: B, f: \text{Hlso}(E(x_1), E(x_2)) \rrbracket \\ & \searrow & \swarrow \\ & B \times B & \end{array}$$

By Lemma 3.3.2, we may fit w_E into the following diagram.

$$\begin{array}{ccccc} B & \xrightarrow{r_B} & \mathbf{P}(B) & \xrightarrow{w_E} & \mathbf{Hlso}_{B \times B}(\pi_1^* E, \pi_2^* E) \\ & & \searrow & & \downarrow \\ & & & & \mathbf{Eq}_{B \times B}(\pi_1^* E, \pi_2^* E) \\ & & & & \downarrow \\ & & & & \mathbf{Hom}_{B \times B}(\pi_1^* E, \pi_2^* E) \\ & & \searrow \Delta_B & & \downarrow \\ & & & & B \times B \end{array}$$

Then by the **ld-COMP** rule applied to the definition of w_{x_1, x_2} , the overall composite map $B \rightarrow \mathbf{Hom}_{B \times B}(\pi_1^* E, \pi_2^* E)$ is the interpretation of $\llbracket x: B \vdash \lambda y: E(x). y : [E(x), E(x)] \rrbracket$, which corresponds under the universal property of **Hom** to $(\Delta_B, 1_E)$. So the composite map $B \rightarrow \mathbf{Eq}_{B \times B}(\pi_1^* E, \pi_2^* E)$ is exactly δ_E of Definition 3.2.10. But by definition, E is univalent precisely if δ_E is a weak equivalence; and by 2-out-of-3 and Lemma 3.3.4, δ_E is a weak equivalence if and only if w_E is. So we are done. \square

3.4. Univalence of the simplicial universes.

Theorem 3.4.1. *The fibration $p_\alpha: \tilde{U}_\alpha \rightarrow U_\alpha$ is univalent.*

Proof. We will show that the target map $t: \mathbf{Eq}(\tilde{U}_\alpha) \rightarrow U_\alpha$ is a trivial fibration. Since t is a retraction of $\delta_{\tilde{U}_\alpha}$, this implies by 2-out-of-3 that $\delta_{\tilde{U}_\alpha}$ is a weak equivalence.

So, we need to fill a square

$$\begin{array}{ccc} A & \xrightarrow{\quad} & \mathbf{Eq}(\tilde{U}_\alpha) \\ i \downarrow & \nearrow & \downarrow t \\ B & \xrightarrow{\quad} & U_\alpha \end{array}$$

where $i: A \hookrightarrow B$ is a cofibration.

By the universal properties of U_α and $\mathbf{Eq}(\widetilde{U}_\alpha)$, these data correspond to a weak equivalence $w: E_1 \rightarrow E_2$ between α -small well-ordered fibrations over A , and an extension \overline{E}_2 of E_2 to an α -small, well-ordered fibration over B ; and a filler corresponds to an extension \overline{E}_1 of E_1 , together with a weak equivalence \overline{w} extending w :

$$\begin{array}{ccccc}
 E_1 & \xrightarrow{\quad w \quad} & \overline{E}_1 & & \\
 \searrow & \swarrow & \searrow & \swarrow & \\
 & E_2 & \xrightarrow{\quad \quad} & \overline{E}_2 & \\
 \swarrow & \nwarrow & \swarrow & \nwarrow & \\
 A & \xrightarrow{\quad \quad} & B & &
 \end{array}$$

As usual, it is sufficient to construct this first without well-orderings on \overline{E}_1 ; these can then always be chosen so as to extend those of E_1 .

Recalling Lemmas 2.2.4–2.2.5, we define \overline{E}_1 and \overline{w} as the pullback

$$\begin{array}{ccc}
 \overline{E}_1 & \longrightarrow & \Pi_i E_1 \\
 \overline{w} \downarrow & \lrcorner & \downarrow \Pi_i w \\
 \overline{E}_2 & \xrightarrow{\quad \eta \quad} & \Pi_i E_2
 \end{array}$$

in \mathbf{sSet}/B , where η is the unit of $i^* \dashv \Pi_i$ at \overline{E}_2 . To see that this construction works, it remains to show:

- (a) $i^* \overline{E}_1 \cong E_1$ in \mathbf{sSet}/A , and under this, $i^* \overline{w}$ corresponds to w ;
- (b) \overline{E}_1 is α -small over B ;
- (c) \overline{E}_1 is a fibration over B , and \overline{w} is a weak equivalence.

For (a), pull the defining diagram of \overline{E}_1 back to \mathbf{sSet}/A ; by Lemma 2.2.4 part 2, we get a pullback square

$$\begin{array}{ccc}
 i^* \overline{E}_1 & \longrightarrow & E_1 \\
 i^* \overline{w} \downarrow & \lrcorner & \downarrow w \\
 E_2 & \xrightarrow{\quad 1_{E_2} \quad} & E_2
 \end{array}$$

in \mathbf{sSet}/A , giving the desired isomorphism.

For (b), Lemma 2.2.4 part 3 gives that $\Pi_i E_1$ is α -small over B , so \overline{E}_1 is a subobject of a pullback of α -small maps.

For (c), note first that by factoring w , we may reduce to the cases where it is either a trivial fibration or a trivial cofibration.

In the former case, by Lemma 2.2.4 part 1 $\Pi_i w$ is also a trivial fibration, and hence so is \overline{w} ; so \overline{E}_1 is fibrant over \overline{E}_2 , hence over B .

In the latter case, E_1 is then a deformation retract of E_2 over A ; we will show that \overline{E}_1 is also a deformation retract of \overline{E}_2 over B . Let $H: E_2 \times \Delta[1] \rightarrow E_2$ be a deformation retraction of E_2 onto E_1 . We want some homotopy $\overline{H}: \overline{E}_2 \times \Delta[1] \rightarrow \overline{E}_2$ extending H on

$E_2 \times \Delta[1]$, $1_{\overline{E}_1} \times \Delta[1]$ on $\overline{E}_1 \times \Delta[1]$, and $1_{\overline{E}_2}$ on $\overline{E}_2 \times \{0\}$. Since these three maps agree on the intersections of their domains, this is exactly an instance of the homotopy lifting extension property, i.e. a square-filler

$$\begin{array}{ccc} (E_2 \times \Delta[1]) \cup (\overline{E}_1 \times \Delta[1]) \cup (\overline{E}_2 \times \{0\}) & \xrightarrow{H \cup 1 \cup 1} & \overline{E}_2 \\ \downarrow & \nearrow \overline{H} & \downarrow \\ \overline{E}_2 \times \Delta[1] & \xrightarrow{\quad\quad\quad} & B \end{array}$$

which exists since the left-hand map is a trivial cofibration.

For \overline{H} to be a deformation retraction, we need to see that $\overline{H}_{\{1\}}: \overline{E}_2 \rightarrow \overline{E}_2$ factors through \overline{E}_1 . By the definition of \overline{E}_1 , a map $f: X \rightarrow \overline{E}_2$ over $b: X \rightarrow B$ factors through \overline{E}_1 just if the pullback $i^*f: i^*X \rightarrow E_2$ factors through E_1 . In the case of $\overline{H}_{\{1\}}$, the pullback is by construction $i^*(\overline{H}_{\{1\}}) = (i^*\overline{H})_{\{1\}} = H_{\{1\}}: E_2 \rightarrow E_2$, which factors through E_1 since H was a deformation retraction onto E_1 .

So \overline{w} embeds \overline{E}_1 as a deformation retract of \overline{E}_2 over B ; thus \overline{E}_1 is a fibration over B and \overline{w} a weak equivalence, as desired. \square

Putting this together with Corollary 2.3.5, we obtain our main theorem:

Theorem 3.4.2. *Let $\beta < \alpha$ be inaccessible cardinals. Then there is a model of dependent type theory in \mathbf{sSet}_{U_α} with all the logical constructors of Section A.2, and a universe (given by U_β) closed under these constructors and satisfying the Univalence Axiom.* \square

From this, we can immediately deduce:

Theorem 3.4.3. *Assuming the existence of two inaccessible cardinals, the contextual category presentation of $\mathbf{MLTT} + \mathbf{UA}$ (as given in Definition 1.2.10) is consistent.* \square

In practice one often considers a type theory with a sequence of n or ω univalent universes. We expect that the techniques used in the proof of Theorem 3.4.3 can be adapted to yield a consistency proof for such a theory, relative to set theory with suitable many inaccessible cardinals; but we do not pursue that here.

Remark 3.4.4. One can prove, within the type theory, that the Univalence Axiom together with the Π - η rule implies functional extensionality; see [V⁺], [Uni13, Sec. 4.9]. So we could have omitted functional extensionality from Proposition 2.3.6, and instead deduced it here as a corollary of univalence.

3.5. Univalence and pullback representations. We are now ready to give a uniqueness statement for the representation of an α -small fibration as a pullback of $p_\alpha: \widetilde{U}_\alpha \rightarrow U_\alpha$: we define the space of such representations, and show that it is contractible.

In fact, we work a bit more generally. Given fibrations q, p , we define a space $P_{q,p}$ of representations of q as a pullback of p ; and we show that a fibration p over a Kan base is univalent exactly when for every q , $P_{q,p}$ is either empty or contractible.

Let $p: E \rightarrow B$ and $q: Y \rightarrow X$ be fibrations. We define a functor

$$\mathbf{P}_{q,p}: \mathbf{sSet}^{\mathrm{op}} \rightarrow \mathbf{Set},$$

setting $\mathbf{P}_{q,p}(S)$ to be the set of pairs of a map $f: S \times X \rightarrow B$, and a weak equivalence $w: S \times Y \rightarrow f^*E$ over $S \times X$; equivalently, the set of squares

$$\begin{array}{ccc} S \times Y & \xrightarrow{f'} & E \\ S \times q \downarrow & & \downarrow p \\ S \times X & \xrightarrow{f} & B \end{array}$$

such that the induced map $S \times Y \rightarrow f^*E$ is a weak equivalence. Lemma 3.2.3 ensures that this is functorial in S , by pullback.

Lemma 3.5.1. *The functor $\mathbf{P}_{q,p}$ is representable, represented by the object*

$$\mathbf{P}_{q,p} := \Pi_{X \rightarrow 1} \Sigma_{\pi_1} \mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E).$$

$$\begin{array}{ccccc} & \pi_1^*Y & & \pi_2^*E & \\ & \swarrow & & \searrow & \\ Y & & X \times B & & E \\ & \searrow \pi_1 & & \swarrow \pi_2 & \\ & X & & B & \\ & \downarrow q & & \downarrow p & \\ & 1 & & & \end{array}$$

Proof. For any S , we have:

$$\begin{aligned} & \text{Hom}(S, \Pi_{X \rightarrow 1} \Sigma_{\pi_1} \mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E)) \\ & \cong \text{Hom}_X(X \times S, \Sigma_{\pi_1} \mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E)) \\ & \cong \{(\hat{f}, \hat{w}) \mid \hat{f}: X \times S \rightarrow X \times B \text{ over } X, \\ & \quad \hat{w}: X \times S \rightarrow \mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E) \text{ over } X \times B\} \\ & \cong \{(f, w) \mid f: X \times S \rightarrow B, w: Y \times S \rightarrow f^*E \text{ w.e. over } X \times S\} \\ & \cong \mathbf{P}_{q,p}(S) \end{aligned}$$

□

Remark 3.5.2. By Yoneda, we see from this that $(\mathbf{P}_{q,p})_n \cong \mathbf{P}_{q,p}(\Delta[n])$.

Theorem 3.5.3. *Let $p: E \rightarrow B$ be a fibration, with B Kan. Then p is univalent if and only if for every fibration $q: Y \rightarrow X$, $\mathbf{P}_{q,p}$ is either empty or contractible.*¹³

¹³Constructively-minded readers might prefer to phrase this as: if $\mathbf{P}_{q,p}$ is inhabited, then it is contractible. In the language of [Uni13], it is a *mere proposition*.

Proof. First, suppose that p is univalent. Take any q such that $P_{q,p}$ is non-empty; then we have some map $1 \rightarrow P_{q,p}$, corresponding to a square

$$\begin{array}{ccccc} Y & \xrightarrow{w} & f^*E & \longrightarrow & E \\ & \searrow q & \downarrow f^*p & & \downarrow p \\ & & X & \xrightarrow{f} & B \end{array}$$

We claim that $P_{q,p} \rightarrow 1$ is a trivial fibration, and hence $P_{q,p}$ is contractible. Π -functors preserve trivial fibrations (since their left adjoints, pullback, preserve cofibrations), so it is enough to show that

$$\mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E) \rightarrow X \times B \xrightarrow{\pi_1} X$$

is a trivial fibration.

For this, first note that w , as a weak equivalence between fibrations, is a homotopy equivalence over X , so induces a homotopy equivalence

$$(w \cdot -): \mathbf{Eq}_{X \times B}((\pi_1^*(f^*E), \pi_2^*E) \rightarrow \mathbf{Eq}_{X \times B}(\pi_1^*Y, \pi_2^*E).$$

So it is enough to show that $\mathbf{Eq}_{X \times B}((\pi_1^*(f^*E), \pi_2^*E) \rightarrow X \times B \xrightarrow{\pi_1} X$ is a trivial fibration; but this follows since it is the pullback along f of the “source” map $\mathbf{Eq}(E) = \mathbf{Eq}_{B \times B}(\pi_1^*E, \pi_2^*E) \rightarrow B \times B \xrightarrow{\pi_1} B$, which is a trivial fibration since p is univalent and B is Kan.

Conversely, suppose that for every fibration q , $P_{q,p}$ is either empty or contractible; now, we wish to show p univalent. For this, it is enough to show that the source map $s: \mathbf{Eq}(E) \rightarrow B$ is a trivial fibration, which will hold if each of its fibers is contractible.

So, take some $f: 1 \rightarrow B$, and consider the fiber $f^*\mathbf{Eq}(E)$. By the universal property of $\mathbf{Eq}(E)$, this is isomorphic to $P_{f^*p,p}$; and it is certainly non-empty, containing the pair $(f, 1_{f^*E})$; so by assumption, it is contractible, as desired.

$$\begin{array}{ccc} f^*\mathbf{Eq}(E) & \longrightarrow & \mathbf{Eq}(E) \\ \downarrow \lrcorner & & \downarrow s \\ 1 & \xrightarrow{f} & B \end{array}$$

□

Corollary 3.5.4. *For any α -small fibration q , the simplicial set P_{q,p_α} of representations of q as a pullback of p_α is contractible.* □

APPENDIX A. SYNTAX OF MARTIN-LÖF TYPE THEORY

A full introduction to Martin-Löf type theory is beyond the scope of this paper. For the reader new to type theory, we recommend [ML84] or [NPS90] as a general introduction, and [Hof97] for a more detailed presentation of the syntax.

However, there are many variant presentations of the theory; in this appendix, we lay out the one we have in mind for the present paper, and which is intended to correspond to the contextual categories described in Appendix B.

We consider the syntax as constructed in two stages: first the *raw* or *untyped* syntax of the theory—the set of expressions that are at least parseable, but not necessarily meaningful—and then the *derivable judgements*, certain inductively-generated predicates picking out the genuinely meaningful contexts, types, and terms.

The raw syntax may be constructed as certain strings of symbols, or alternatively, certain labelled trees. On this, one then defines *alpha-equivalence* (i.e. syntactic identity modulo renaming of bound variables), and the operation of (*capture-free*) *substitution*. This step is well-standardised in the literature.

At the second stage, one defines on the raw syntax several multi-place relations, picking out the *derivable judgements* of the theory. For instance, “ $\Gamma \vdash a : A$ ” will be a relation on triples (Γ, a, A) of a raw context, term, and type expression respectively, to be read as “ a is a term of type A , in context Γ ”. These relations are defined by mutual induction, as the smallest family of relations closed under a bevy of specified closure conditions, the *inference rules* of the theory.

Details of the judgements and inference rules used vary somewhat; we therefore set our choice out here in full. For the structural rules, our presentation is based largely on [Hof97]; our selection of logical rules, and in particular our treatment of the universe, follows [ML84].

We take as basic four judgement forms:

$$\Gamma \vdash A \text{ type} \quad \Gamma \vdash A = A' \text{ type} \quad \Gamma \vdash a : A \quad \Gamma \vdash a = a' : A.$$

We take the context judgement as defined from these: that is, if Γ is a list $(x_i : A_i)_{i < n}$, with x_i distinct variables and A_i raw type expressions, then $\vdash \Gamma$ cxt is an abbreviation for the statement that for each $i < n$, $(x_j : A_j)_{j < i} \vdash A_i$ type.

Derivability, for these four basic judgements, is then defined as the smallest family of relations closed under the closure conditions specified by the inference rules below. These are given in “rule-notation”, so for instance

$$\frac{\Gamma \vdash a : A \quad \Gamma \vdash A = B \text{ type}}{\Gamma \vdash a : B}$$

expresses the closure condition “for all suitable raw expressions Γ, a, A, B , if the judgements $\Gamma \vdash a : A$ and $\Gamma \vdash A = B$ are derivable, then so is $\Gamma \vdash a : B$ ”.

The inference rules fall into two groups: the *structural rules*, which we assume are always included, and the *logical rules*, which different type theories may include different subsets of.

A.1. Structural Rules. The structural rules of the type theory are (where \mathcal{J} may be the conclusion of any of the judgement forms):

$$\begin{array}{c} \frac{\vdash \Gamma, x:A, \Delta \text{ cxt}}{\Gamma, x:A, \Delta \vdash x : A} \text{VBLE} \qquad \frac{\Gamma \vdash a : A \quad \Gamma, x:A, \Delta \vdash \mathcal{J}}{\Gamma, \Delta[a/x] \vdash \mathcal{J}[a/x]} \text{SUBST} \\[2ex] \frac{\Gamma \vdash A \text{ type} \quad \Gamma, \Delta \vdash \mathcal{J}}{\Gamma, x:A, \Delta \vdash \mathcal{J}} \text{WKG} \end{array}$$

Definitional equality (also known as syntactic or judgemental equality):

$$\begin{array}{c}
\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A = A \text{ type}} \quad \frac{\Gamma \vdash A = B \text{ type}}{\Gamma \vdash B = A \text{ type}} \quad \frac{\Gamma \vdash A = B \text{ type} \quad \Gamma \vdash B = C \text{ type}}{\Gamma \vdash A = C \text{ type}} \\
\\
\frac{\Gamma \vdash a : A}{\Gamma \vdash a = a : A} \quad \frac{\Gamma \vdash a = b : A}{\Gamma \vdash b = a : A} \quad \frac{\Gamma \vdash a = b : A \quad \Gamma \vdash b = c : A}{\Gamma \vdash a = c : A} \\
\\
\frac{\Gamma \vdash a : A \quad \Gamma \vdash A = B \text{ type}}{\Gamma \vdash a : B} \quad \frac{\Gamma \vdash a = b : A \quad \Gamma \vdash A = B \text{ type}}{\Gamma \vdash a = b : B}
\end{array}$$

A.2. Logical Constructors. In this and subsequent sections, we present rules introducing various type- and term-constructors. For each such constructor, we assume (besides the explicitly stated rules introducing and governing it) a *congruence* rule stating that it preserves definitional equality in each of its arguments; for instance, along with the Π -INTRO rule introducing the constructor λ , we assume the rule

$$\frac{\Gamma \vdash A = A' \text{ type} \quad \Gamma, x:A \vdash B(x) = B'(x) \text{ type} \quad \Gamma, x:A \vdash b(x) = b'(x) : B(x)}{\Gamma \vdash \lambda x:A. b(x) = \lambda x:A'. b'(x) : \Pi_{x:A} B(x)} \lambda\text{-EQ}$$

The rules fall naturally into groups according to the various logical constructors. Many of the constructors considered (Σ -, Id -, W -, 0 -, 1 -, and $+$ -types) follow a common pattern, as *inductive types/families* with constructors and an elimination principle; here, as in [ML84], this pattern is purely heuristic, but in approaches such as the Calculus of Inductive Constructions [Wer94] they are unified formally as instances of a single scheme.

Π -types. (Dependent products; dependent function types).

$$\begin{array}{c}
\frac{\Gamma, x:A \vdash B(x) \text{ type}}{\Gamma \vdash \Pi_{x:A} B(x) \text{ type}} \Pi\text{-FORM} \quad \frac{\Gamma, x:A \vdash B(x) \text{ type} \quad \Gamma, x:A \vdash b(x) : B(x)}{\Gamma \vdash \lambda x:A. b(x) : \Pi_{x:A} B(x)} \Pi\text{-INTRO} \\
\\
\frac{\Gamma \vdash f : \Pi_{x:A} B(x) \quad \Gamma \vdash a : A}{\Gamma \vdash \text{app}(f, a) : B(a)} \Pi\text{-APP} \\
\\
\frac{\Gamma, x:A \vdash B(x) \text{ type} \quad \Gamma, x:A \vdash b(x) : B(x) \quad \Gamma \vdash a : A}{\Gamma \vdash \text{app}(\lambda x:A. b(x), a) = b(a) : B(a)} \Pi\text{-COMP}
\end{array}$$

As a special case of this, when B does not depend on x , we obtain the ordinary function type $[A, B] := \Pi_{x:A} B$.

Σ -types. (Dependent sums; type-indexed disjoint sums.)

$$\begin{array}{c}
\frac{\Gamma \vdash A \text{ type} \quad \Gamma, x:A \vdash B(x) \text{ type}}{\Gamma \vdash \Sigma_{x:A} B(x) \text{ type}} \Sigma\text{-FORM} \\
\\
\frac{\Gamma \vdash A \text{ type} \quad \Gamma, x:A \vdash B(x) \text{ type}}{\Gamma, x:A, y:B(x) \vdash \text{pair}(x, y) : \Sigma_{x:A} B(x)} \Sigma\text{-INTRO}
\end{array}$$

$$\frac{\Gamma, z:\Sigma_{x:A}B(x) \vdash C(z) \text{ type} \quad \Gamma, x:A, y:B(x) \vdash d(x, y) : C(\text{pair}(x, y))}{\Gamma, z:\Sigma_{x:A}B(x) \vdash \text{split}_d(z) : C(z)} \Sigma\text{-ELIM}$$

$$\frac{\Gamma, z:\Sigma_{x:A}B(x) \vdash C(z) \text{ type} \quad \Gamma, x:A, y:B(x) \vdash d(x, y) : C(\text{pair}(x, y))}{\Gamma, x:A, y:B(x) \vdash \text{split}_d(\text{pair}(x, y)) = d(x, y) : C(\text{pair}(x, y))} \Sigma\text{-COMP}$$

Again, the special case where B does not depend on x is of particular interest: this gives the cartesian product $A \times B := \Sigma_{x:A}B$.

Id-types. (Identity types, equality types.)

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma, x, y:A \vdash \text{Id}_A(x, y) \text{ type}} \text{Id-FORM} \qquad \frac{\Gamma \vdash A \text{ type}}{\Gamma, x:A \vdash \text{refl}_A(x) : \text{Id}_A(x, x)} \text{Id-INTRO}$$

$$\frac{\Gamma, x, y:A, u:\text{Id}_A(x, y) \vdash C(x, y, u) \text{ type} \quad \Gamma, z:A \vdash d(z) : C(z, z, \text{refl}_A(z))}{\Gamma, x, y:A, u:\text{Id}_A(x, y) \vdash J_{z.d}(x, y, u) : C(x, y, u)} \text{Id-ELIM}$$

$$\frac{\Gamma, x, y:A, u:\text{Id}_A(x, y) \vdash C(x, y, u) \text{ type} \quad \Gamma, z:A \vdash d(z) : C(z, z, r(z))}{\Gamma, x:A \vdash J_{z.d}(x, x, \text{refl}_A(x)) = d(x) : C(x, x, \text{refl}_A(x))} \text{Id-COMP}$$

W-types. (Types of well-founded trees; free term algebras.)

$$\frac{\Gamma, x:A \vdash B(x) \text{ type}}{\Gamma \vdash W_{x:A}B(x) \text{ type}} \text{W-FORM}$$

$$\frac{\Gamma, x:A \vdash B(x) \text{ type}}{\Gamma, x:A, y:[B(x), W_{u:A}B(u)] \vdash \text{sup}(x, y) : W_{u:A}B(u)} \text{W-INTRO}$$

$$\frac{\Gamma, w:W_{x:A}B(x) \vdash C(w) \text{ type} \quad \Gamma, x:A, y:[B(x), W_{u:A}B(u)], z:\prod_{u:B(x)} C(\text{app}(y, u)) \vdash d(x, y, z) : C(\text{sup}(x, y))}{\Gamma, w:W_{x:A}B(x) \vdash \text{wrec}_d(w) : C(w)} \text{W-ELIM}$$

$$\frac{\Gamma, w:W_{x:A}B(x) \vdash C(w) \text{ type} \quad \Gamma, x:A, y:[B(x), W_{u:A}B(u)], z:\prod_{u:B(x)} C(\text{app}(y, u)) \vdash d(x, y, z) : C(\text{sup}(x, y))}{\Gamma, x:A, y:[B(x), W_{u:A}B(u)] \vdash \text{wrec}_d(\text{sup}(x, y)) = d(x, y, \lambda u:B(x). \text{wrec}_d(\text{app}(y, u))) : C(\text{sup}(x, y))} \text{W-COMP}$$

0. (Empty type.)

$$\frac{\vdash \Gamma \text{ cxt}}{\Gamma \vdash 0 \text{ type}} \text{0-FORM} \quad (\text{No introduction rules.})$$

$$\frac{\Gamma, x:0 \vdash C(x) \text{ type}}{\Gamma, x:0 \vdash \text{case}(x) : C(x)} \text{0-ELIM} \quad (\text{No computation rules.})$$

1. (Unit type, singleton type.)

$$\begin{array}{c} \frac{\vdash \Gamma \text{ cxt}}{\Gamma \vdash 1 \text{ type}} \text{1-FORM} \quad \frac{\vdash \Gamma \text{ cxt}}{\Gamma \vdash * : 1} \text{1-INTRO} \quad \frac{\Gamma, x:1 \vdash C(x) \text{ type} \quad \Gamma \vdash d : C(*)}{\Gamma, x:1 \vdash \text{rec}_d(x) : C(x)} \text{1-ELIM} \\ \\ \frac{\Gamma, x:1 \vdash C(x) \text{ type} \quad \Gamma \vdash d : C(*)}{\Gamma \vdash \text{rec}_d(*) = d : C(*)} \text{1-COMP} \end{array}$$

+types. (Binary disjoint sums.)

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A + B \text{ type}} \text{+-FORM} \quad \frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma, x:A \vdash \text{inl}(x) : A + B} \text{+-INTRO 1.}$$

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma, y:B \vdash \text{inr}(y) : A + B} \text{+-INTRO 2.}$$

$$\frac{\Gamma, z:A + B \vdash C(z) \text{ type} \quad \Gamma, x:A \vdash d_l(x) : C(\text{inl}(x)) \quad \Gamma, y:B \vdash d_r(y) : C(\text{inr}(y))}{\Gamma, z:A + B \vdash \text{case}_{d_l, d_r}(z) : C(z)} \text{+-ELIM}$$

$$\frac{\Gamma, z:A + B \vdash C(z) \text{ type} \quad \Gamma, x:A \vdash d_l(x) : C(\text{inl}(x)) \quad \Gamma, y:B \vdash d_r(y) : C(\text{inr}(y))}{\Gamma, x:A \vdash \text{case}_{d_l, d_r}(\text{inl}(x)) = d_l(x) : C(\text{inl}(x))} \text{+-COMP 1.}$$

$$\frac{\Gamma, z:A + B \vdash C(z) \text{ type} \quad \Gamma, x:A \vdash d_l(x) : C(\text{inl}(x)) \quad \Gamma, y:B \vdash d_r(y) : C(\text{inr}(y))}{\Gamma, y:B \vdash \text{case}_{d_l, d_r}(\text{inr}(y)) = d_r(y) : C(\text{inr}(y))} \text{+-COMP 2.}$$

A.3. Universes. A universe within the theory may be closed under some or all of the logical constructors of the theory; we include below the rules corresponding to closure under all of the constructors given above.

$$\overline{\vdash U \text{ type}}$$

$$\overline{x:U \vdash \text{El}(x) \text{ type}}$$

$$\frac{\Gamma \vdash a : U \quad \Gamma, x:\text{El}(a) \vdash b(x) : U}{\Gamma \vdash \pi(a, x.b(x)) : U}$$

$$\frac{\Gamma \vdash a : U \quad \Gamma, x:\text{El}(a) \vdash b(x) : U}{\Gamma \vdash \text{El}(\pi(a, x.b(x))) = \prod_{x:\text{El}(a)} \text{El}(b(x)) \text{ type}}$$

$$\begin{array}{c}
\frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma, x:\mathsf{El}(a) \vdash b(x) : \mathbb{U}}{\Gamma \vdash \sigma(a, x.b(x)) : \mathbb{U}} \quad \frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma, x:\mathsf{El}(a) \vdash b(x) : \mathbb{U}}{\Gamma \vdash \mathsf{El}(\sigma(a, x.b(x))) = \Sigma_{x:\mathsf{El}(a)} \mathsf{El}(b(x)) \text{ type}} \\
\\
\frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma \vdash b, c : \mathsf{El}(a)}{\Gamma \vdash \mathsf{id}_A(b, c) : \mathbb{U}} \quad \frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma \vdash b, c : \mathsf{El}(a)}{\Gamma \vdash \mathsf{El}(\mathsf{id}_a(b, c)) = \mathsf{Id}_{\mathsf{El}(a)}(b, c) \text{ type}} \\
\\
\overline{\vdash z : \mathbb{U}} \quad \overline{\vdash \mathsf{El}(z) = 0 \text{ type}} \quad \overline{\vdash o : \mathbb{U}} \quad \overline{\vdash \mathsf{El}(o) = 1 \text{ type}} \\
\\
\frac{\Gamma \vdash a, b : \mathbb{U}}{\Gamma \vdash a + b : \mathbb{U}} \quad \frac{\Gamma \vdash a, b : \mathbb{U}}{\Gamma \vdash \mathsf{El}(a + b) = \mathsf{El}(a) + \mathsf{El}(b) \text{ type}} \\
\\
\frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma, x:\mathsf{El}(a) \vdash b(x) : \mathbb{U}}{\Gamma \vdash w(a, x.b(x)) : \mathbb{U}} \quad \frac{\Gamma \vdash a : \mathbb{U} \quad \Gamma, x:\mathsf{El}(a) \vdash b(x) : \mathbb{U}}{\Gamma \vdash \mathsf{El}(w(a, x.b(x))) = W_{x:\mathsf{El}(a)} \mathsf{El}(b(x)) \text{ type}}
\end{array}$$

A.4. Further rules. The rules above are somewhat weak in their implications for equality of functions. To this end, some further rules are often adopted: the η -rule for Π -types, and the *functional extensionality* rule(s). Our formulation of the latter is taken from [Gar09]; see also [Hof95a].

$$\begin{array}{c}
\frac{\Gamma \vdash f : \Pi_{x:A} B(x)}{\Gamma \vdash f = \lambda x:A. \mathsf{app}(f, x) : \Pi_{x:A} B(x)} \Pi\text{-}\eta \\
\\
\frac{\Gamma \vdash f, g : \Pi_{x:A} B(x) \quad \Gamma \vdash h : \Pi_{x:A} \mathsf{Id}_{B(x)}(\mathsf{app}(f, x), \mathsf{app}(g, x))}{\Gamma \vdash \mathsf{ext}(f, g, h) : \mathsf{Id}_{\Pi_{x:A} B(x)}(f, g)} \Pi\text{-EXT} \\
\\
\frac{\Gamma, x:A \vdash b : B(x)}{\Gamma \vdash \mathsf{ext-comp}(x.b) : \mathsf{Id}_{\Pi_{x:A} B(x)}(\mathsf{ext}(\lambda x:A. b, \lambda x:A. b, \lambda x:A. \mathsf{refl} b), \mathsf{refl}(\lambda x:A. b))} \Pi\text{-EXT-COMP-PROP}
\end{array}$$

APPENDIX B. LOGICAL STRUCTURE ON CONTEXTUAL CATEGORIES

We give here full translations of the various type-theoretic rules and axioms into the language of contextual categories: the logical rules of Section A.2, the universe rules of Section A.3, the extensionality and η -rules of Section A.4, and Axiom 3.1.4, the Univalence Axiom.

B.1. Logical structure.

Definition B.1.1. A Π -type structure on a contextual category \mathcal{C} consists of:

- (1) for each $(\Gamma, A, B) \in \mathsf{Ob}_{n+2} \mathcal{C}$, an object $(\Gamma, \Pi(A, B)) \in \mathsf{Ob}_{n+1} \mathcal{C}$;
- (2) for each such (Γ, A, B) and section $b : (\Gamma, A) \longrightarrow (\Gamma, B)$, a section $\lambda(b) : (\Gamma, A) \longrightarrow (\Gamma, \Pi(A, B))$;
- (3) for each such (Γ, A, B) and pair of sections $k : \Gamma \longrightarrow (\Gamma, \Pi(A, B))$, $a : \Gamma \longrightarrow (\Gamma, A)$, a section $\mathsf{app}(k, a) : \Gamma \longrightarrow (\Gamma, B)$ such that $p_B \cdot \mathsf{app}(k, a) = a$,

- (4) such that for each (Γ, A, B) , $a: \Gamma \longrightarrow (\Gamma, A)$ and $b: (\Gamma, A) \longrightarrow (\Gamma, A, B)$, we have $\mathbf{app}(\lambda(b), a) = b \cdot a$;
 (5) and for any $f: \Gamma' \longrightarrow \Gamma$, and all appropriate arguments as above,

$$\begin{aligned} f^*(\Gamma, \Pi(A, B)) &= (\Gamma', \Pi(f^*A, f^*B)), \\ f^*\lambda(b) &= \lambda(f^*b), \quad f^*(\mathbf{app}(k, a)) = \mathbf{app}(f^*k, f^*a). \end{aligned}$$

Given a Π -structure on \mathcal{C} , and (Γ, A, B) as above, write $\mathbf{app}_{A,B}$ for the morphism

$$\begin{aligned} q(q(p_{\Pi(A,B)} \cdot p_{p_{\Pi(A,B)}^*A}, A), B) \cdot \\ \mathbf{app}_{(p_{\Pi(A,B)} \cdot p_{p_{\Pi(A,B)}^*A})^*A, (p_{\Pi(A,B)} \cdot p_{p_{\Pi(A,B)}^*A})^*B} ((1, p_{p_{\Pi(A,B)}^*A}), (1, q(p_{\Pi(A,B)}, A))) \\ : (\Gamma, \Pi(A, B), p_{\Pi(A,B)}^*A) \longrightarrow (\Gamma, A, B); \end{aligned}$$

the general form $\mathbf{app}_{A,B}(k, a)$ can be re-derived from these instances. Also, for objects (Γ, A) , (Γ, B) in \mathcal{C} , write $(\Gamma, [A, B])$ for $(\Gamma, \Pi(A, p_A^*B))$.

Definition B.1.2. A Σ -type structure on a contextual category \mathcal{C} consists of:

- (1) for each $(\Gamma, A, B) \in \text{Ob}_{n+2} \mathcal{C}$, an object $(\Gamma, \Sigma(A, B)) \in \text{Ob}_{n+1} \mathcal{C}$;
- (2) for each such (Γ, A, B) , a morphism $\mathbf{pair}_{A,B}: (\Gamma, A, B) \longrightarrow (\Gamma, \Sigma(A, B))$ over Γ ;
- (3) for each (Γ, A, B) , object $(\Gamma, \Sigma(A, B), C)$, and map $d: (\Gamma, A, B) \longrightarrow (\Gamma, \Sigma(A, B), C)$ with $p_C \cdot d = \mathbf{pair}_{A,B}$, a section $\mathbf{split}_d: (\Gamma, \Sigma(A, B)) \longrightarrow (\Gamma, \Sigma(A, B), C)$, with $\mathbf{split}_d \cdot \mathbf{pair}_{A,B} = d$;
- (4) such that for $f: \Gamma' \longrightarrow \Gamma$, and all appropriate arguments as above,

$$\begin{aligned} f^*(\Gamma, \Sigma(A, B)) &= (\Gamma', \Sigma(f^*A, f^*B)), \\ f^*\mathbf{pair}_{A,B} &= \mathbf{pair}_{f^*A, f^*B}, \quad f^*\mathbf{split}_d = \mathbf{split}_{f^*d}. \end{aligned}$$

Definition B.1.3. An Id -type structure on a contextual category \mathcal{C} consists of:

- (1) for each (Γ, A) , an object $(\Gamma, A, p_A^*A, \text{Id}_A)$;
- (2) for each (Γ, A) , a morphism $\mathbf{refl}_A: (\Gamma, A) \longrightarrow (\Gamma, A, p_A^*A, \text{Id}_A)$, such that $p_{\text{Id}_A} \cdot \mathbf{refl}_A = (1_A, 1_A): (\Gamma, A) \longrightarrow (\Gamma, A, p_A^*A)$;
- (3) for each $(\Gamma, A, p_A^*A, \text{Id}_A, C)$ and $d: (\Gamma, A) \longrightarrow (\Gamma, A, p_A^*A, \text{Id}_A, C)$ with $p_C \cdot d = \mathbf{refl}_A$, a section $\mathbf{J}_{C,d}: (\Gamma, A, p_A^*A, \text{Id}_A) \longrightarrow (\Gamma, A, p_A^*A, \text{Id}_A, C)$, such that $\mathbf{J}_{C,d} \cdot \mathbf{refl}_A = d$;
- (4) such that for $f: \Gamma' \longrightarrow \Gamma$, and all appropriate arguments as above,

$$\begin{aligned} f^*(\Gamma, A, p_A^*A, \text{Id}_A) &= (\Gamma', f^*A, (p_{f^*A})^*(f^*A), \text{Id}_{f^*A}), \\ f^*\mathbf{refl}_A &= \mathbf{refl}_{f^*A}, \quad f^*\mathbf{J}_{C,d} = \mathbf{J}_{f^*C, f^*d}. \end{aligned}$$

Definition B.1.4. Given a contextual category \mathcal{C} equipped with a Π -type structure, a W -type structure on \mathcal{C} consists of:

- (1) for each (Γ, A, B) , an object $(\Gamma, W(A, B))$;
- (2) for each (Γ, A, B) , a map over Γ

$$\mathbf{sup}_{A,B}: (\Gamma, A, \Pi(B, p_B^*p_A^*W(A, B))) \longrightarrow (\Gamma, W(A, B));$$

(3) for each $(\Gamma, W(A, B), C)$ and map

$$d: (\Gamma, A, [B, p_A^* W(A, B)], \Pi(p_{[B, p_A^* W(A, B)]}^* B, \text{app}(p_{p_{[B, p_A^* W(A, B)]}^* B}, q(p_{[B, p_A^* W(A, B)]}, B))^* C)) \longrightarrow (\Gamma, W(A, B), C)$$

such that

$$p_C \cdot d = \sup_{A, B} \cdot p \Pi(p_{[B, p_A^* W(A, B)]}^* B, \text{app}(p_{p_{[B, p_A^* W(A, B)]}^* B}, q(p_{[B, p_A^* W(A, B)]}, B))^* C),$$

a section $\text{wrec}_{C, d}: (\Gamma, W(A, B)) \longrightarrow (\Gamma, W(A, B), C)$, such that

$$\text{wrec}_{C, d} \cdot \sup_{A, B} =$$

$$d \cdot \lambda(\text{wrec}_{C, d} \cdot \text{app}(p_{p_{[B, p_A^* W(A, B)]}^* B}, q(p_{[B, p_A^* W(A, B)]}, B)));$$

(4) and such that for $f: \Gamma' \longrightarrow \Gamma$, and all appropriate arguments as above,

$$f^*(\Gamma, W(A, B)) = (\Gamma', W(f^* A, f^* B)),$$

$$f^* \sup_{A, B} = \sup_{f^* A, f^* B}, \quad f^* \text{wrec}_{C, d} = \text{wrec}_{f^* C, f^* d}.$$

Definition B.1.5. A *zero-type structure* on a contextual category \mathcal{C} consists of:

- (1) for each Γ , an object $(\Gamma, 0_\Gamma)$;
- (2) for any object $(\Gamma, 0_\Gamma, C)$, a section $\text{case}_C: (\Gamma, 0_\Gamma) \longrightarrow (\Gamma, 0_\Gamma, C)$;
- (3) such that for each $f: \Gamma' \longrightarrow \Gamma$, $f^*(\Gamma, 0_\Gamma) = (\Gamma', 0_{\Gamma'})$; and for each such f and $(\Gamma, 0_\Gamma, C)$ as above, $f^*(\text{case}_C) = \text{case}_{f^* C}$.

Definition B.1.6. A *unit-type structure* on a contextual category \mathcal{C} consists of:

- (1) for each Γ , an object $(\Gamma, 1_\Gamma)$;
- (2) a section $*_\Gamma: (\Gamma) \longrightarrow (\Gamma, 1_\Gamma)$;
- (3) for each object $(\Gamma, 1_\Gamma, C)$, and map $d: \Gamma \longrightarrow (\Gamma, 1_\Gamma, C)$ with $p_C \cdot d = *_\Gamma$, a section $\text{rec}_{C, d}: (\Gamma, 1_\Gamma) \longrightarrow (\Gamma, 1_\Gamma, C)$, such that $\text{rec}_{C, d} \cdot *_\Gamma = d$;
- (4) such that for $f: \Gamma' \longrightarrow \Gamma$, and appropriate arguments as above,

$$f^*(\Gamma, 1_\Gamma) = (\Gamma', 1_{\Gamma'}),$$

$$f^* *_\Gamma = *_{\Gamma'}, \quad f^* \text{rec}_{C, d} = \text{rec}_{f^* C, f^* d}.$$

Definition B.1.7. A *sum-type structure* on a contextual category \mathcal{C} consists of:

- (1) for any objects (Γ, A) and (Γ, B) , an object $(\Gamma, A + B)$;
- (2) for each such (Γ, A) , (Γ, B) , maps $\text{inl}_{A, B}: (\Gamma, A) \longrightarrow (\Gamma, A + B)$ and $\text{inr}_{A, B}: (\Gamma, B) \longrightarrow (\Gamma, A + B)$, over Γ ;
- (3) for each object $(\Gamma, A + B, C)$, and maps $d_l: (\Gamma, A) \longrightarrow (\Gamma, A + B, C)$, $d_r: (\Gamma, B) \longrightarrow (\Gamma, A + B, C)$ with $p_C \cdot d_l = \text{inl}_{A, B}$ and $p_C \cdot d_r = \text{inr}_{A, B}$, a section $\text{case}_{C, d_l, d_r}: (\Gamma, A + B) \longrightarrow (\Gamma, A + B, C)$, such that $\text{case}_{C, d_l, d_r} \cdot \text{inl}_{A, B} = d_l$ and $\text{case}_{C, d_l, d_r} \cdot \text{inr}_{A, B} = d_r$;
- (4) such that for $f: \Gamma' \longrightarrow \Gamma$, and appropriate arguments as above,

$$f^*(\Gamma, A + B) = (\Gamma', f^* A + f^* B), \quad f^*(\text{inl}_{A, B}) = \text{inl}_{f^* A, f^* B},$$

$$f^*(\text{inr}_{A, B}) = \text{inr}_{f^* A, f^* B}, \quad f^*(\text{case}_{C, d_l, d_r}) = \text{case}_{f^* C, f^* d_l, f^* d_r}.$$

B.2. Universes. As in Section A.3, one may consider a universe closed under some or all of the logical structure of the ambient contextual category; that is, carrying operations reflecting the global operations on types.

Definition B.2.1. A *universe* in a contextual category \mathcal{C} is a distinguished object $(U, \text{El}) \in \text{Ob}_2 \mathcal{C}$.

Assuming a Π -type structure on \mathcal{C} , we say (U, El) is *closed under Π -types* if for all maps $a: \Gamma \rightarrow U$ and $b: (\Gamma, a^* \text{El}) \rightarrow U$, we are given a map $\pi(a, b): \Gamma \rightarrow U$, such that $(\Gamma, \pi(a, b)^* \text{El}) = (\Gamma, \Pi(a^* \text{El}, b^* \text{El}))$, and moreover stably in Γ , i.e. such that for $f: \Gamma' \rightarrow \Gamma$ and a, b as above, $f^*(\pi(a, b)) = \pi(f^*a, f^*b)$.

Similarly, given a Σ -type structure on \mathcal{C} , say (U, El) is *closed under Σ -types* if for each $a: \Gamma \rightarrow U$ and $b: (\Gamma, a^* \text{El}) \rightarrow U$, we are given $\sigma(a, b): \Gamma \rightarrow U$, such that $(\Gamma, \sigma(a, b)^* \text{El}) = (\Gamma, \Sigma(a^* \text{El}, b^* \text{El}))$, and such that for $f: \Gamma' \rightarrow \Gamma$ and a, b as above, $f^*(\sigma(a, b)) = \sigma(f^*a, f^*b)$.

Given an Id -type structure on \mathcal{C} , say (U, El) is *closed under Id -types* if for each $a: \Gamma \rightarrow U$, we are given a map $\text{id}_a: (\Gamma, a^* \text{El}, p_{a^* \text{El}}^* a^* \text{El}) \rightarrow U$, such that $(\Gamma, a^* \text{El}, p_{a^* \text{El}}^* a^* \text{El}, \text{id}_a^* \text{El}) = (\Gamma, a^* \text{El}, p_{a^* \text{El}}^* a^* \text{El}, \text{Id}_{a^* \text{El}})$, and such that for $f: \Gamma' \rightarrow \Gamma$ and a as above, $f^*(\text{id}_a) = \text{id}_{f^*a}$.

Given a zero-type (resp. unit-type) structure on \mathcal{C} , say (U, El) *contains 0* (resp. *1*) if we are given a map $z: 1 \rightarrow U$ (resp. \circ) such that $z^* \text{El} = 0$ ($\circ^* \text{El} = 1$).

Given a sum-type structure on \mathcal{C} , say (U, El) is *closed under sum types* if for each pair of maps $a, b: \Gamma \rightarrow U$, we are given $a + b: \Gamma \rightarrow U$, such that $(a + b)^* \text{El} = a^* \text{El} + b^* \text{El}$, and moreover such that $f: \Gamma' \rightarrow \Gamma$ and a, b as above, $f^*(a + b) = f^*a + f^*b$.

Given W -type structure on \mathcal{C} , say (U, El) is *closed under W -types* if for each $a: \Gamma \rightarrow U$ and $b: (\Gamma, a^* \text{El}) \rightarrow U$, we are given $w(a, b): \Gamma \rightarrow U$, such that $(\Gamma, w(a, b)^* \text{El}) = (\Gamma, W(a^* \text{El}, b^* \text{El}))$, and such that for $f: \Gamma' \rightarrow \Gamma$ and a, b as above, $f^*(w(a, b)) = w(f^*a, f^*b)$.

B.3. Extensionality and Univalence. For the following group of rules, let \mathcal{C} be a contextual category equipped with chosen Π - and Id -type structures.

Definition B.3.1. Say that \mathcal{C} satisfies the Π - η rule if for any (Γ, A, B) , the “ η -expansion” map

$$q(p_{\Pi(A, B)}, \Pi(A, B)) \cdot \lambda(1_{p_{\Pi(A, B)}^* A}, \text{app}_{A, B}): (\Gamma, \Pi(A, B)) \rightarrow (\Gamma, \Pi(A, B))$$

is equal to $1_{(\Gamma, \Pi(A, B))}$.

A Π -EXT structure on \mathcal{C} is an operation giving for each (Γ, A, B) a map

$$\text{ext}_{A, B}: (\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B), \text{Htp}_{A, B}) \rightarrow (\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B), \text{Id}_{A, B})$$

over $(\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B))$, stably in Γ , where $\text{Htp}_{A, B}$ is the object

$$\left(\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B), \Pi \left((p_{\Pi(A, B)} \cdot p_{p_{\Pi(A, B)}^* \Pi(A, B)}^* A, (\text{app}_{A, B} \cdot q(p_{\Pi(A, B)}, A), \text{app}_{A, B} \cdot q(q(p_{\Pi(A, B)}, \Pi(A, B)), A))^* \text{Id}_B \right) \right).$$

Given a Π -EXT structure on \mathcal{C} , a Π -EXT-COMP-PROP *structure* for it is an operation giving, for each (Γ, A, B) and section $f: \Gamma \longrightarrow (\Gamma, \Pi(A, B))$, a map

$$\text{ext-comp}(f): \Gamma \longrightarrow$$

$$(\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B), \text{Id}_{\Pi(A, B)}, p_{\text{Id}_{\Pi(A, B)}}^* \text{Id}_{\Pi(A, B)}, \text{Id}_{\text{Id}_{\Pi(A, B)}})$$

over the pair of maps

$$\begin{aligned} \text{ext}_{A, B}(f, g) \cdot \lambda(1_A, \text{refl}_B \cdot p_{p_A^* B}^* \cdot \text{app}((1_A, f), (1_A, 1_A))) \quad , \quad \text{refl}_{\Pi(A, B)} \cdot f \\ : \Gamma \longrightarrow (\Gamma, \Pi(A, B), p_{\Pi(A, B)}^* \Pi(A, B), \text{Id}_{\Pi(A, B)}), \end{aligned}$$

stably as ever in Γ .

Before defining the Univalence Axiom, we first (as in Section 3.1) set up several auxiliary definitions. We assume, from here on, that \mathcal{C} is equipped also with Σ - and Id -type structures.

Definition B.3.2. For objects (Γ, X) , (Γ, Y) of \mathcal{C} , let

$$\text{exch}_{X, Y}: (\Gamma, X, p_X^* Y) \longrightarrow (\Gamma, Y, p_Y^* X)$$

be the evident “exchange” map, $(q(p_X, Y), p_{p_X^* Y})$.

Definition B.3.3. Let (Γ, A) , (Γ, B) be objects of \mathcal{C} .

Take $(\Gamma, [A, B], \text{LInv}_{A, B})$ to be the object

$$\begin{aligned} (\Gamma, [A, B], \Sigma(p_{[A, B]}^* [B, A], \Pi((p_{[A, B]} \cdot p_{p_{[A, B]}^* [B, A]}^*)^* A, \\ (q(p_B, A) \cdot \text{app}_{B, A} \cdot q(p_{p_{[B, A]}^* A, p_{[B, A]}^* B) \cdot \text{app}_{p_{[B, A]}^* A, p_{[B, A]}^* B} \\ \cdot q(\text{exch}_{[A, B], [B, A]}, A), q(p_{[A, B]} \cdot p_{p_{[A, B]}^* [B, A]}^*)^* \text{Id}_A))) \end{aligned}$$

Similarly, take $(\Gamma, [A, B], \text{RInv}_{A, B})$ to be the object

$$\begin{aligned} (\Gamma, [A, B], \Sigma(p_{[A, B]}^* [B, A], \Pi((p_{[A, B]} \cdot p_{p_{[A, B]}^* [B, A]}^*)^* B, \\ (q(p_A, B) \cdot \text{app}_{A, B} \cdot q(p_{p_{[A, B]}^* B, p_{[A, B]}^* A) \cdot \text{app}_{p_{[A, B]}^* B, p_{[A, B]}^* A} \\ q(p_{[A, B]} \cdot p_{p_{[A, B]}^* [B, A]}^*)^* \text{Id}_B))) \end{aligned}$$

Now, set

$$\begin{aligned} (\Gamma, [A, B], \text{isHlso}_{A, B}) &:= (\Gamma, [A, B], \text{LInv}_{A, B} \times \text{RInv}_{A, B}), \\ (\Gamma, \text{Hlso}(A, B)) &:= (\Gamma, \Sigma([A, B], \text{isHlso}_{A, B})). \end{aligned}$$

Definition B.3.4. For any (Γ, A) , there is a canonical “identity” section

$$\text{id}_A := \lambda((1_A, 1_A)): \Gamma \longrightarrow (\Gamma, [A, A]);$$

and this moreover lifts to a section $\text{id}_A^{\text{isHlso}}: \Gamma \longrightarrow (\Gamma, [A, A], \text{isHlso}_{A, A})$, given in full by

$$\begin{aligned} \text{pair}_{\text{LInv}_{A, A}, \text{RInv}_{A, A}} \cdot \left(\text{pair}_{p_{[A, A]}^* [A, A], \text{H}_L} \cdot q((\text{id}_A, \text{id}_A), \text{H}_L) \cdot \lambda((1_A, \text{refl}_A)), \right. \\ \left. \text{pair}_{p_{[A, A]}^* [A, A], \text{H}_R} \cdot q((\text{id}_A, \text{id}_A), \text{H}_R) \cdot \lambda((1_A, \text{refl}_A)) \right), \end{aligned}$$

where H_L is the object

$$\begin{aligned} & \left(\Gamma, [A, A], p_{[A,A]}^*[A, A], \Pi \left((p_{[A,A]} \cdot p_{p_{[A,A]}^*[A,A]}^*)^* A, \right. \right. \\ & \quad \left. \left(q(p_A, A) \cdot \text{app}_{A,A} \cdot q(p_{p_{[A,A]}^*[A,A]}^* A, p_{[A,A]}^* A) \cdot \text{app}_{p_{[A,A]}^* A, p_{[A,A]}^* A} \right. \right. \\ & \quad \left. \left. \cdot q(\text{exch}_{[A,A],[A,A]}, A), q(p_{[A,A]} \cdot p_{p_{[A,A]}^*[A,A]}^*[A,A], A) \right)^* \text{Id}_A \right) \end{aligned}$$

and H_R is the same but with $q(\text{exch}_{[A,A],[A,A]}, A)$ omitted.

Lastly, set $\text{id}^{\text{Hlso}} := \text{pair}_{[A,A], \text{isHlso}_{A,A}} \cdot \text{id}_A^{\text{isHlso}} : \Gamma \longrightarrow (\Gamma, \text{Hlso}(A, A))$.

Definition B.3.5. For any object (Γ, A, B) , we can now define a map

$$w_{A,B} : (\Gamma, A, p_A^* A, \text{Id}_A) \longrightarrow (\Gamma, A, p_A^* A, \text{Hlso}(p_{p_A^* A}^* B, q(p_A, A)^* B))$$

by

$$\begin{aligned} w_{A,B} &:= q(p_{\text{Id}_A}, \text{Hlso}(p_{p_A^* A}^* B, q(p_A, A)^* B)) \cdot \\ & \quad J(p_{\text{Id}_A}^* \text{Hlso}(p_{p_A^* A}^* B, q(p_A, A)^* B), \\ & \quad q(\text{refl}_A, p_{\text{Id}_A}^* \text{Hlso}(p_{p_A^* A}^* B, q(p_A, A)^* B)) \cdot \text{id}_B^{\text{Hlso}}) \end{aligned}$$

Given this, define the predicate “ B is a univalent family over A ” as:

$$\begin{aligned} (\Gamma, \text{isUvt}(A, B)) &:= \\ & (\Gamma, \Pi(A, \Pi(p_A^* A, \lambda((1_{\text{Id}_A}, w_{A,B}))^* \text{isHlso}_{\text{Id}_A, \text{Hlso}(p_{p_A^* A}^* B, q(p_A, A)^* B)}))) \end{aligned}$$

Definition B.3.6. Given a universe (U, El) in \mathcal{C} , say (U, El) *satisfies the Univalence Axiom* if \mathcal{C} is equipped with a map $\text{uvt}_{U, \text{El}} : 1 \longrightarrow \text{isUvt}(U, \text{El})$.

REFERENCES

- [AGMLV11] Steve Awodey, Richard Garner, Per Martin-Löf, and Vladimir Voevodsky (eds.), *Mini-workshop: The homotopy interpretation of constructive type theory*, vol. 8, Oberwolfach Reports, no. 1, European Mathematical Society, 2011, [doi:10.4171/OWR/2011/11](#).
- [AK11] Peter Arndt and Krzysztof Kapulkin, *Homotopy-theoretic models of type theory*, Typed lambda calculi and applications, Lecture Notes in Comput. Sci., vol. 6690, Springer, Heidelberg, 2011, pp. 45–60, [doi:10.1007/978-3-642-21691-6_7](#).
- [AR94] Jiří Adámek and Jiří Rosický, *Locally presentable and accessible categories*, London Mathematical Society Lecture Note Series, vol. 189, Cambridge University Press, Cambridge, 1994, [doi:10.1017/CB09780511600579](#).
- [AW09] Steve Awodey and Michael A. Warren, *Homotopy theoretic models of identity types*, Math. Proc. Cambridge Philos. Soc. **146** (2009), no. 1, 45–55, [arXiv:0709.0248](#), [doi:10.1017/S0305004108001783](#).
- [Bar12] Bruno Barras, *Semantical investigations in intuitionistic set theory and type theories with inductive families*, Habilitation, Université Paris 7 (Denis Diderot), 2012.
- [BGL⁺17] Andrej Bauer, Jason Gross, Peter LeFanu Lumsdaine, Mike Shulman, Matthieu Sozeau, and Bas Spitters, *The HoTT library: A formalization of homotopy type theory in Coq*, Proc. 6th ACM SIGPLAN Conference on Certified Programs and Proofs (CPP) (Yves Bertot and Viktor Vafeiadis, eds.), ACM, 2017, pp. 164–172, [doi:10.1145/3018610.3018615](#).
- [BGM59] Michael G. Barratt, Victor K. A. M. Gugenheim, and John C. Moore, *On semisimplicial fibre-bundles*, Amer. J. Math. **81** (1959), 639–657.

- [Car78] John Cartmell, *Generalised algebraic theories and contextual categories*, Ph.D. thesis, Oxford, 1978.
- [Dyb96] Peter Dybjer, *Internal type theory*, Types for proofs and programs (Torino, 1995), Lecture Notes in Comput. Sci., vol. 1158, Springer, Berlin, 1996, pp. 120–134.
- [GAA⁺13] Georges Gonthier, Andrea Asperti, Jeremy Avigad, Yves Bertot, Cyril Cohen, François Garillot, Stéphane Le Roux, Assia Mahboubi, Russell O’Connor, Sidi Ould Biha, Ioana Pasca, Laurence Rideau, Alexey Solovyev, Enrico Tassi, and Laurent Théry, *A machine-checked proof of the odd order theorem*, Lecture Notes in Computer Science, vol. 7998, Springer, 2013, pp. 163–179, [doi:10.1007/978-3-642-39634-2_14](https://doi.org/10.1007/978-3-642-39634-2_14).
- [Gar09] Richard Garner, *On the strength of dependent products in the type theory of Martin-Löf*, Ann. Pure Appl. Logic **160** (2009), no. 1, 1–12, [doi:10.1016/j.apal.2008.12.003](https://doi.org/10.1016/j.apal.2008.12.003).
- [GG08] Nicola Gambino and Richard Garner, *The identity type weak factorisation system*, Theoretical Computer Science **409** (2008), no. 1, 94–109, [arXiv:0803.4349](https://arxiv.org/abs/0803.4349), [doi:10.1016/j.tcs.2008.08.030](https://doi.org/10.1016/j.tcs.2008.08.030).
- [GH04] Nicola Gambino and Martin Hyland, *Wellfounded trees and dependent polynomial functors*, Types for proofs and programs, Lecture Notes in Comput. Sci., vol. 3085, Springer, Berlin, 2004, pp. 210–225, [doi:10.1007/978-3-540-24849-1_14](https://doi.org/10.1007/978-3-540-24849-1_14).
- [GJ09] Paul G. Goerss and John F. Jardine, *Simplicial homotopy theory*, Modern Birkhäuser Classics, Birkhäuser, 2009.
- [GK17] David Gepner and Joachim Kock, *Univalence in locally cartesian closed ∞ -categories*, Forum Math. **29** (2017), no. 3, 617–652, <https://doi.org/10.1515/forum-2015-0228>.
- [Gon08] Georges Gonthier, *Formal proof—the four-color theorem*, Notices Amer. Math. Soc. **55** (2008), no. 11, 1382–1393.
- [Gra18] Daniel R. Grayson, *An introduction to univalent foundations for mathematicians*, Bull. Amer. Math. Soc. (2018), online first.
- [GvdB11] Richard Garner and Benno van den Berg, *Types are weak ω -groupoids*, Proc. Lond. Math. Soc. (3) **102** (2011), no. 2, 370–394, [doi:10.1112/plms/pdq026](https://doi.org/10.1112/plms/pdq026).
- [Hof95a] Martin Hofmann, *Extensional concepts in intensional type theory*, Ph.D. thesis, University of Edinburgh, 1995.
- [Hof95b] ———, *On the interpretation of type theory in locally Cartesian closed categories*, Computer science logic (Kazimierz, 1994), Lecture Notes in Comput. Sci., vol. 933, Springer, Berlin, 1995, pp. 427–441, [doi:10.1007/BFb0022273](https://doi.org/10.1007/BFb0022273).
- [Hof97] ———, *Syntax and semantics of dependent types*, Semantics and logics of computation (Cambridge, 1995), Publ. Newton Inst., vol. 14, Cambridge Univ. Press, Cambridge, 1997, pp. 79–130, [doi:10.1017/CB09780511526619.004](https://doi.org/10.1017/CB09780511526619.004).
- [Hov99] Mark Hovey, *Model categories*, Mathematical Surveys and Monographs, vol. 63, American Mathematical Society, Providence, Rhode Island, 1999.
- [HS99] Martin Hofmann and Thomas Streicher, *Lifting Grothendieck universes*, Unpublished note, 199?, <http://www.mathematik.tu-darmstadt.de/~streicher/>.
- [HS98] ———, *The groupoid interpretation of type theory*, Twenty-Five Years of Constructive Type Theory (Venice, 1995), Oxford Logic Guides, vol. 36, Oxford Univ. Press, New York, 1998, pp. 83–111.
- [Jac93] Bart Jacobs, *Comprehension categories and the semantics of type dependency*, Theoret. Comput. Sci. **107** (1993), no. 2, 169–207, [doi:10.1016/0304-3975\(93\)90169-T](https://doi.org/10.1016/0304-3975(93)90169-T).
- [Joh02] Peter T. Johnstone, *Sketches of an elephant: a topos theory compendium. Vol. 1*, Oxford Logic Guides, vol. 43, The Clarendon Press Oxford University Press, New York, 2002.
- [Joy11] André Joyal, *A result on Kan fibrations*, unpublished note, 2011.
- [LW15] Peter LeFanu Lumsdaine and Michael A. Warren, *The local universes model: an overlooked coherence construction for dependent type theories*, ACM Trans. Comput. Log. **16** (2015), no. 3, Art. 23, 31, [arXiv:1411.1736](https://arxiv.org/abs/1411.1736), [doi:10.1145/2754931](https://doi.org/10.1145/2754931).
- [May67] J. Peter May, *Simplicial objects in algebraic topology*, Mathematical Studies, vol. 11, Van Nostrand, 1967.

- [ML84] Per Martin-Löf, *Intuitionistic type theory*, Studies in Proof Theory. Lecture Notes, vol. 1, Bibliopolis, Naples, 1984.
- [ML98] Saunders Mac Lane, *Categories for the working mathematician*, 2 ed., Graduate Texts in Mathematics, vol. 5, Springer-Verlag, New York, 1998.
- [Mog91] Eugenio Moggi, *A category-theoretic account of program modules*, Math. Structures Comput. Sci. **1** (1991), no. 1, 103–139.
- [MP00] Ieke Moerdijk and Erik Palmgren, *Wellfounded trees in categories*, Proceedings of the Workshop on Proof Theory and Complexity, PTAC’98 (Aarhus), vol. 104, Ann. Pure Appl. Logic, no. 1-3, 2000, pp. 189–218, [doi:10.1016/S0168-0072\(00\)00012-9](https://doi.org/10.1016/S0168-0072(00)00012-9).
- [NPS90] Bengt Nordström, Kent Petersson, and Jan M. Smith, *Programming in Martin-Löf’s type theory*, International Series of Monographs on Computer Science, vol. 7, The Clarendon Press Oxford University Press, New York, 1990.
- [Pit00] Andrew M. Pitts, *Categorical logic*, Handbook of logic in computer science, Vol. 5, Handb. Log. Comput. Sci., vol. 5, Oxford Univ. Press, New York, 2000, pp. 39–128.
- [PM96] Christine Paulin-Mohring, *Inductive definitions in higher-order type theory*, Habilitation, Université Claude Bernard (Lyon I), 1996.
- [PW14] Álvaro Pelayo and Michael A. Warren, *Homotopy type theory and Voevodsky’s univalent foundations*, Bull. Amer. Math. Soc. (N.S.) **51** (2014), no. 4, 597–648, [arXiv:1210.5658](https://arxiv.org/abs/1210.5658), [doi:10.1090/S0273-0979-2014-01456-9](https://doi.org/10.1090/S0273-0979-2014-01456-9).
- [Qui68] Daniel G. Quillen, *The geometric realization of a Kan fibration is a Serre fibration*, Proc. Amer. Math. Soc. **19** (1968), 1499–1500.
- [Sco55] Dana Scott, *Definitions by abstraction in axiomatic set theory*, Bulletin Amer. Math. Soc. **61** (1955), 442, abstract only.
- [See84] Robert A. G. Seely, *Locally Cartesian closed categories and type theory*, Math. Proc. Cambridge Philos. Soc. **95** (1984), no. 1, 33–48, [doi:10.1017/S0305004100061284](https://doi.org/10.1017/S0305004100061284).
- [Sel11] Peter Selinger, *A survey of graphical languages for monoidal categories*, New structures for physics, Lecture Notes in Phys., vol. 813, Springer, Heidelberg, 2011, pp. 289–355, [doi:10.1007/978-3-642-12821-9_4](https://doi.org/10.1007/978-3-642-12821-9_4).
- [Shu15] Michael Shulman, *Univalence for inverse diagrams and homotopy canonicity*, Math. Structures Comput. Sci. **25** (2015), no. 5, 1203–1277, [arXiv:1203.3253](https://arxiv.org/abs/1203.3253), [doi:10.1017/S0960129514000565](https://doi.org/10.1017/S0960129514000565).
- [Str91] Thomas Streicher, *Semantics of type theory*, Progress in Theoretical Computer Science, Birkhäuser Boston Inc., Boston, MA, 1991, [doi:10.1007/978-1-4612-0433-6](https://doi.org/10.1007/978-1-4612-0433-6).
- [Uni13] The Univalent Foundations Program, *Homotopy type theory: Univalent foundations of mathematics*, <http://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.
- [V⁺] Vladimir Voevodsky et al., *Univalent foundations repository*, ongoing Coq development, <https://github.com/UnivalentMathematics/Foundations>.
- [VAG⁺] Vladimir Voevodsky, Benedikt Ahrens, Daniel Grayson, et al., *UniMath — a computer-checked library of univalent mathematics*, available at <https://github.com/UniMath/UniMath>, <https://github.com/UniMath/UniMath>.
- [vdBM15] Benno van den Berg and Ieke Moerdijk, *W-types in homotopy type theory*, Math. Structures Comput. Sci. **25** (2015), no. 5, 1100–1115, <https://doi.org/10.1017/S0960129514000516>.
- [vdBM18] ———, *W-types in homotopy-type theory—corrigendum [MR3340536]*, Math. Structures Comput. Sci. **28** (2018), no. 1, 140, <https://doi.org/10.1017/S0960129516000025>.
- [Voe06] Vladimir Voevodsky, *A very short note on homotopy λ -calculus*, notes from seminars given at Stanford University, 2006, http://math.ucr.edu/home/baez/Voevodsky_note.ps.
- [Voe12] ———, *Notes on type systems*, unpublished notes, 2012, http://www.math.ias.edu/~vladimir/Site3/Univalent_Foundations_files/expressions_current.pdf.
- [Voe14] ———, *C-system of a module over a monad on sets*, submitted, 2014, [arXiv:1407.3394](https://arxiv.org/abs/1407.3394).
- [Voe15a] ———, *A C-system defined by a universe category*, Theory Appl. Categ. **30** (2015), No. 37, 1181–1215, [arXiv:1409.7925](https://arxiv.org/abs/1409.7925).

- [Voe15b] ———, *An experimental library of formalized mathematics based on the univalent foundations*, Math. Structures Comput. Sci. **25** (2015), no. 5, 1278–1294, <https://doi.org/10.1017/S0960129514000577>.
- [Voe15c] ———, *Martin-Löf identity types in the C-systems defined by a universe category*, submitted, 2015, [arXiv:1505.06446](https://arxiv.org/abs/1505.06446).
- [Voe16a] ———, *Products of families of types and (Π, λ) -structures on C-systems*, Theory Appl. Categ. **31** (2016), Paper No. 36, 1044–1094.
- [Voe16b] ———, *Subsystems and regular quotients of C-systems*, Conference on Mathematics and its Applications, (Kuwait City, 2014), no. 658, 2016, [arXiv:1406.5389](https://arxiv.org/abs/1406.5389).
- [Voe17] ———, *The (Π, λ) -structures on the C-systems defined by universe categories*, Theory Appl. Categ. **32** (2017), Paper No. 4, 113–121.
- [War08] Michael A. Warren, *Homotopy theoretic aspects of constructive type theory*, Ph.D. thesis, Carnegie Mellon University, 2008.
- [War11] ———, *The strict ω -groupoid interpretation of type theory*, Models, logics, and higher-dimensional categories, CRM Proc. Lecture Notes, vol. 53, Amer. Math. Soc., Providence, RI, 2011, pp. 291–340.
- [Wer94] Benjamin Werner, *Une théorie des constructions inductives*, Ph.D. thesis, Université Paris 7 (Denis Diderot), May 1994.

(Krzysztof Kapulkin) UNIVERSITY OF PITTSBURGH

Email address: krk56@pitt.edu

(Peter LeFanu Lumsdaine) INSTITUTE FOR ADVANCED STUDY, PRINCETON

Email address: plumsdaine@ias.edu