

Heisenberg's uncertainty principle for simultaneous measurement of positive-operator-valued measures

Takayuki Miyadera^{*} and Hideki Imai[†]

Research Center for Information Security (RCIS),

National Institute of Advanced Industrial Science and Technology (AIST).

Daibiru building 1102, Sotokanda, Chiyoda-ku, Tokyo, 101-0021, Japan.

(Dated: 22 Nov 2008)

Abstract

A limitation on simultaneous measurement of two arbitrary positive operator valued measures is discussed. In general, simultaneous measurement of two noncommutative observables is only approximately possible. Following Werner's formulation, we introduce a distance between observables to quantify an accuracy of measurement. We derive an inequality that relates the achievable accuracy with noncommutativity between two observables. As a byproduct a necessary condition for two positive operator valued measures to be simultaneously measurable is obtained.

PACS numbers: 03.65.Ta, 03.67.-a

I. INTRODUCTION

Heisenberg's uncertainty principle is often considered as one of the most important features of quantum theory. In every text book on quantum theory one can find its explanation proposed by Heisenberg himself [1] and its "derivation" by Robertson [2]. However, as recently claimed by several researchers, the above explanation and the derivation have a certain gap between them. On the one hand Heisenberg is concerned with a simultaneous measurement of position and momentum, on the other hand the Robertson's formulation treats two distinct measurements each on position and momentum. The later formulation has been actively investigated since then, and there are a several different inequalities for general observables depending on quantities that characterize the uncertainty of probability distributions [3–7]. In contrast to it, it seems quite recent that the former formulation began being reflected extensively. In this formulation one must deal with a simultaneous (or joint) measurement of two or more observables

^{*} miyadera-takayuki@aist.go.jp

[†] Also at Graduate School of Science and Engineering, Chuo University. 1-13-27 Kasuga, Bunkyo-ku, Tokyo 112-8551, Japan .

and somehow estimate its accuracy compared with individual measurements of the observables. Appleby, in his pioneering work [8], investigating simultaneous measurement of position and momentum of quantum mechanical particles, introduced error operators and disturbance operators and derived simple inequalities between them. Ozawa [9] treated a pair of general self-adjoint observables and considered a tradeoff relation between his error operator and disturbance operator that have an interpretation in the context of his extended notion of simultaneous measurement. Werner [10] formulated the problem from an operational viewpoint and derived a beautiful inequality between position and momentum operator of quantum mechanical particles. Janssens' work [11] is related with it and showed a nice inequality on added variances between arbitrary self-adjoint observables in a simple manner. Busch and Pearson [12] introduced a notion of error bar that represents a resolution of the measurements, and discuss a tradeoff relation related with position and momentum. Busch and Heinosaari [13] estimated a tradeoff between observables in a single qubit in detail. Busch, Heinonen and Lahti gave a review on this topic in [14].

In this paper, we deal with simultaneous measurements of two arbitrary observables (positive operator valued measures). In general, a pair of noncommutative observables is not simultaneously measurable. We derive an inequality that relates the limitation on the simultaneous measurements with the noncommutativity of observables. The paper is organized as follow: In section II, we explain our formulation of the problem. For that purpose we introduce notions of distance between observables and simultaneous measurability of observables. In section III, we derive our main result. Tradeoff relations related with two types of distances are explained. As a byproduct we give a necessary condition for two positive operator valued measures to be simultaneously measurable. A simple example is also discussed.

II. FORMULATION

In this paper, we follow a formulation introduced by Werner [10] which has a clear operational meaning. Suppose that we have a quantum system described by a Hilbert space \mathcal{H} and a pair of observables; A and B on it. A and B are represented as positive operator valued measures (POVMs) $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. That is, for each a and b , $0 \leq A_a, B_b \leq \mathbf{1}$ holds and $\sum_a A_a = \sum_b B_b = \mathbf{1}$ is satisfied. (For simplicity, we hereafter assume the sets of outcomes (say Ω_A) finite.) Given a state ω , we can compute a probability distribution $\{\omega(A_a)\}_a$, where $P_A^\omega(a) := \omega(A_a)$ is interpreted as a probability to obtain an outcome a when we make a measurement on A with respect to the state ω . If A_a and B_b commute with each other for all a and b , $A \circ B := \{A_a B_b\}_{(a,b) \in \Omega_A \times \Omega_B}$ again defines a POVM. This newly defined POVM gives a probability distribution

$P_{A \circ B}^\omega(a, b) = \omega(A_a B_b)$ whose marginal distributions coincide exactly with P_A^ω and P_B^ω for any state ω . That is, in this case one can achieve a simultaneous measurement of A and B perfectly by using $A \circ B$. This is not always the case in general for noncommutative pairs. What we are interested in is that impossibility. What is the quantitative limitation on simultaneous measurements of general pairs of observables?

To formulate this problem quantitatively, one has to introduce a proper distance between two probability distributions. Suppose that we have a space $\mathcal{P}(\Omega)$ of probability distributions over a (finite) sample space Ω . (i.e., $\mathcal{P}(\Omega) := \{p : \Omega \rightarrow [0, 1] \mid \sum_x p(x) = 1\}$)

A function $d : \mathcal{P}(\Omega) \times \mathcal{P}(\Omega) \rightarrow \mathbf{R}_+$ is called a distance if the following conditions are satisfied:

- (i) $d(p, q) = d(q, p)$,
- (ii) $d(p, p) = 0 \Leftrightarrow p \equiv 0$,
- (iii) $d(q, p) + d(p, r) \geq d(q, r)$ (triangle inequality).

For instance, l_∞ distance (uniform distance) is defined as

$$d_\infty(p, q) := \max_{x \in \Omega} |p(x) - q(x)|,$$

and l_1 distance is defined as

$$d_1(p, q) := \frac{1}{2} \sum_{x \in \Omega} |p(x) - q(x)|.$$

(For later convenience we put a coefficient $\frac{1}{2}$.) Once we fix a distance d over the probability spaces, we can define a distance between two observables. Suppose that POVMs $A = \{A_a\}_{a \in \Omega}$ and $B = \{B_b\}_{b \in \Omega}$ have an identical set of outcomes. We define a distance between A and B with respect to d as

$$D_d(A, B) := \sup_{\omega} d(P_A^\omega, P_B^\omega),$$

where the supremum is taken over all the states. One can easily show that $D_d(A, B) = 0$ if and only if $A = B$. All other conditions for distance also follow easily. Thus D_d becomes a distance between a pair of POVMs which have an identical set of outcomes.

Let us recall the previous example in which all the elements of $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$ commute with each other. In such a case one can define $A \circ B$ that has $\Omega_A \times \Omega_B$ as the set of outcomes. If we define $f_A : \Omega_A \times \Omega_B \rightarrow \Omega_A$ as $f_A(a, b) = a$ and $f_B : \Omega_A \times \Omega_B \rightarrow \Omega_B$ as $f_B(a, b) = b$ for all $a \in \Omega_A$ and $b \in \Omega_B$, $A_a = \sum_{(a', b')^{f(a', b')=a}} A_{a'} B_{b'}$ and $B_b = \sum_{(a', b')^{f(a', b')=b}} A_{a'} B_{b'}$ hold. That is, truncating

outcomes of $A \circ B$ with some functions reproduces both A and B . This can be generalized to the following procedure. Let us consider a POVM $F := \{F_x\}_{x \in \Omega}$. $f_A : \Omega \rightarrow \Omega_A$ defines a new POVM $f_A(F) := \{f_A(F)_a\}_{a \in \Omega_A}$ as

$$f_A(F)_a := \sum_{x: f_A(x)=a} F_x.$$

Also, an other function $f_B : \Omega \rightarrow \Omega_B$ defines a POVM $f_B(F)$. Here it is appropriate to say that POVMs $f_A(F)$ and $f_B(F)$ are simultaneously measurable (through a POVM F). Actually, the following definition of the simultaneously measurable observables is a standard one (See e.g. [14, 15]).

Definition 1 *A pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$ is called simultaneously measurable if and only if there exists a POVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$ such that $A = f_A(F)$ and $B = f_B(F)$ hold.*

Having introduced the notions of distance and simultaneous measurement, we can formulate our problem as follows. For given arbitrary pair of POVMs $A = \{A_a\}$ and $B = \{B_b\}$, we choose a proper POVM $F = \{F_x\}$ and functions f_A and f_B that approximately reconstruct them. How close can we make $f_A(F)$ and $f_B(F)$ to A and B ? Once we fix F , f_A and f_B , we can define the closeness of them to A and B as $D_d(f_A(F), A)$ and $D_d(f_B(F), B)$. One may expect that if one chooses F to make $D_d(f_A(F), A)$ small, $D_d(f_B(F), B)$ becomes large. This tradeoff relation is what we are interested in.

III. RESULTS

A. Uncertainty principle in l_∞ distance

To proceed with the analysis, we have to fix a distance d first. The simplest l_∞ distance d_∞ has a clear meaning and in addition makes our problem tractable. Indeed, in this case the corresponding distance between two observables $A = \{A_a\}_{a \in \Omega}$ and $A' = \{A'_a\}_{a \in \Omega}$ can be written as [13]

$$D_{d_\infty}(A, A') = \sup_{\omega} \max_a |P_A^\omega(a) - P_{A'}^\omega(a)| = \max_a \|A_a - A'_a\|,$$

where $\|\cdot\|$ represents an operator norm. Suppose that we fix a POVM F and f_A and f_B . $D_{d_\infty}(A, f_A(F)) = \max_a \|A_a - f_A(F)_a\|$ and $D_{d_\infty}(B, f_B(F)) = \max_b \|B_b - f_B(F)_b\|$ naturally follow.

The second observation that makes our analysis easier is a representation of POVMs by completely positive maps (CP maps) [16]. When we have an Abelian von Neumann algebra \mathcal{M} and a unital (completely [17]) positive linear map $T : \mathcal{M} \rightarrow \mathbf{B}(\mathcal{H})$, a decomposition of unity (POVM) in \mathcal{M} , $\mathbf{1} = \sum_{x \in \Omega} m_x$ with

$m_x \geq 0$, defines a POVM $\{T(m_x)\}_{x \in \Omega}$ in $\mathbf{B}(\mathcal{H})$. The inverse is also true. When we have a POVM $F = \{F_x\}_{x \in \Omega}$, we define \mathcal{M}_Ω a set of all the diagonal matrices acting on $\mathbf{C}^{|\Omega|}$. For each $x \in \Omega$, a projection onto the canonical basis vector $e_x := |x\rangle\langle x|$ is an element of \mathcal{M}_Ω and $e := \{e_x\}_{x \in \Omega}$ is a decomposition of unity in \mathcal{M}_Ω . One can define $T : \mathcal{M}_\Omega \rightarrow \mathbf{B}(\mathcal{H})$ as

$$T\left(\sum_{x \in \Omega} c_x e_x\right) = \sum_{x \in \Omega} c_x F_x.$$

It is easy to see that thus defined T is indeed a CP map.

Let us begin the analysis. Suppose we have a pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. Take an arbitrary POVM $F = \{F_x\}_{x \in \Omega}$ and functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$. We represent the POVM F in terms of a CP map $T : \mathcal{M}_\Omega \rightarrow \mathbf{B}(\mathcal{H})$. That is, $F_x = T(e_x)$ holds for each $x \in \Omega$. Since a CP map is linear, if we define $E_a^A := \sum_x^{f_A(x)=a} e_x = f_A(e)$ and $E_b^B := \sum_x^{f_B(x)=b} e_x = f_B(e)$ for each $a \in \Omega_A$ and $b \in \Omega_B$, we have

$$\begin{aligned} f_A(F)_a &= T(E_a^A) \\ f_B(F)_b &= T(E_b^B). \end{aligned}$$

Since what we are interested in is $D_{d_\infty}(f_A(F), A)$ and $D_{d_\infty}(f_B(F), B)$, we shall estimate $\|T(E_a^A) - A_a\|$ and $\|T(E_b^B) - B_b\|$. They are represented in simple forms if one defines *error operators*,

$$\begin{aligned} \epsilon_a^A &:= T(E_a^A) - A_a \\ \epsilon_b^B &:= T(E_b^B) - B_b. \end{aligned}$$

That is, $D_{d_\infty}(f_A(F), A) = \max_a \|\epsilon_a^A\|$ and $D_{d_\infty}(f_B(F), B) = \max_b \|\epsilon_b^B\|$ hold. We consider a commutator $[T(E_a^A), T(E_b^B)] = [\epsilon_a^A + A_a, \epsilon_b^B + B_b] = [\epsilon_a^A, \epsilon_b^B] + [\epsilon_a^A, B_b] + [A_a, \epsilon_b^B] + [A_a, B_b]$. Taking the norm on both sides of the equation, $\|B_b, A_a\| = \|\epsilon_a^A, \epsilon_b^B\| + \|\epsilon_a^A, B_b\| + \|A_a, \epsilon_b^B\| + \|T(E_b^B), T(E_a^A)\|$, we have

$$\|[B_b, A_a]\| \leq \|[\epsilon_a^A, \epsilon_b^B]\| + \|[\epsilon_a^A, B_b]\| + \|[\epsilon_b^B, A_a]\| + \|[T(E_b^B), T(E_a^A)]\|, \quad (1)$$

where we have used the triangle inequality for norm. Each term in the right hand side is bounded as follows. The first term is $\|[\epsilon_a^A, \epsilon_b^B]\| \leq 2\|\epsilon_a^A\|\|\epsilon_b^B\|$. The second term is, by use of a relation $\|X\| = \sup_{\psi: \|\psi\|=1} |\langle \psi | X | \psi \rangle|$ for self-adjoint operator X ,

$$\begin{aligned} \|[\epsilon_a^A, B_b]\| &= \|i[\epsilon_a^A, B_b]\| \\ &= \sup_{\psi} |\langle \psi | [\epsilon_a^A, B_b] | \psi \rangle| \\ &\leq 2 \sup_{\psi} \langle \psi | (\epsilon_a^A)^2 | \psi \rangle^{1/2} \langle \psi | (\Delta B_b)^2 | \psi \rangle^{1/2} \\ &\leq 2\|\epsilon_a^A\| \sup_{\psi} \langle \psi | (\Delta B_b)^2 | \psi \rangle^{1/2}, \end{aligned}$$

where $\Delta X := X - \langle \psi | X | \psi \rangle$ and we used the Robertson uncertainty relation. Due to $0 \leq B_b \leq \mathbf{1}$, $\langle \psi | (\Delta B_b)^2 | \psi \rangle^{1/2} \leq 1/2$ holds and we have

$$\|[\epsilon_a^A, B_b]\| \leq \|\epsilon_a^A\|.$$

For the third term, we have in the similar manner,

$$\|[\epsilon_b^B, A_a]\| \leq \|\epsilon_b^B\|.$$

Thus (1) is bounded as

$$\|[A_a, B_b]\| \leq 2\|\epsilon_a^A\|\|\epsilon_b^B\| + \|\epsilon_a^A\| + \|\epsilon_b^B\| + \|[T(E_a^A), T(E_b^B)]\|. \quad (2)$$

To estimate the last term in the right hand side of the above inequality we use the following lemma proved by Janssens[11].

Lemma 1 *Let \mathcal{A} and \mathcal{B} be von Neumann algebras and $T : \mathcal{B} \rightarrow \mathcal{A}$ a CP map. Let B, \tilde{B} be commuting Hermitian operators in \mathcal{B} , then,*

$$\|T(B^2) - T(B)^2\|^{1/2} \|T(\tilde{B}^2) - T(\tilde{B})^2\|^{1/2} \geq \frac{1}{2} \|[T(B), T(\tilde{B})]\|$$

holds.

The proof is done by a direct application of the Cauchy-Schwarz inequality for operators. Since our E_a^A and E_b^B commute with each other, we can apply the above lemma to our inequality to obtain

$$\|[T(E_a^A), T(E_b^B)]\| \leq 2\|T(E_a^A) - T(E_a^A)^2\|^{1/2} \|T(E_b^B) - T(E_b^B)^2\|^{1/2}, \quad (3)$$

where we utilized $(E_a^A)^2 = E_a^A$ and $(E_b^B)^2 = E_b^B$. To delete $T(E_a^A)$ and $T(E_b^B)$ from the above inequality, we use $T(E_a^A) = \epsilon_a^A + A_a$ and $T(E_b^B) = \epsilon_b^B + B_b$. It derives

$$\begin{aligned} \|T(E_a^A) - T(E_a^A)^2\| &= \|\epsilon_a^A + A_a - (\epsilon_a^A)^2 - A_a^2 - \epsilon_a^A A_a - A_a \epsilon_a^A\| \\ &\leq \|\epsilon_a^A - (\epsilon_a^A)^2 - \epsilon_a^A A_a - A_a \epsilon_a^A\| + \|A_a - A_a^2\|, \end{aligned}$$

whose first term in the right hand side is further bounded as follows:

$$\begin{aligned} \|\epsilon_a^A - (\epsilon_a^A)^2 - \epsilon_a^A A_a - A_a \epsilon_a^A\| &= \|(1 - A_a)\epsilon_a^A - \epsilon_a^A A_a - (\epsilon_a^A)^2\| \\ &= \|[A_a, \epsilon_a^A] - A_a \epsilon_a^A + (1 - A_a)\epsilon_a^A - (\epsilon_a^A)^2\| \\ &\leq \|[A_a, \epsilon_a^A]\| + \|(1 - 2A_a - \epsilon_a^A)\epsilon_a^A\|. \end{aligned}$$

Here we use the relation $\|[A_a, \epsilon_a^A]\| \leq \|\epsilon_a^A\|$ and $1 - 2A_a - \epsilon_a^A = 1 - (A_a + T(E_a^A))$ to derive

$$\|[\epsilon_a^A, A_a]\| + \|(1 - 2A_a - \epsilon_a^A)\epsilon_a^A\| \leq \|\epsilon_a^A\| + \|1 - (A_a + T(E_a^A))\| \|\epsilon_a^A\| \leq 2\|\epsilon_a^A\|,$$

where we used a relation $\|1 - (A_a + T(E_a^A))\| \leq 1$ that is derived from $0 \leq A_a + T(E_a^A) \leq \mathbf{21}$. Finally we obtain

$$\|T(E_a^A) - T(E_a^A)^2\| \leq 2\|\epsilon_a^A\| + \|A_a - A_a^2\|. \quad (4)$$

We are ready to state the following theorem.

Theorem 1 Suppose that we have a pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. For any choice of a POVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$,

$$2D_{d_\infty}(A, f_A(F))D_{d_\infty}(B, f_B(F)) + D_{d_\infty}(A, f_A(F)) + D_{d_\infty}(B, f_B(F)) \\ + 2(2D_{d_\infty}(A, f_A(F)) + V(A))^{1/2}(2D_{d_\infty}(B, f_B(F)) + V(B))^{1/2} \geq \max_{a,b} \|[A_a, B_b]\|$$

holds, where $V(A) := \max_a \|A_a - A_a^2\|$ represents an intrinsic uncertainty of a POVM A (and similarly for $V(B) := \max_b \|B_b - B_b^2\|$).

Proof: Combining (2), (3) and (4), we obtain

$$\|[A_a, B_b]\| \leq 2\|\epsilon_a^A\| \|\epsilon_b^B\| + \|\epsilon_a^A\| + \|\epsilon_b^B\| \\ + 2(2\|\epsilon_a^A\| + \|A_a - A_a^2\|)^{1/2}(2\|\epsilon_b^B\| + \|B_b - B_b^2\|)^{1/2}. \quad (5)$$

We take its maximum over a and b to obtain the theorem. Q.E.D.

The intrinsic uncertainty of a POVM A satisfies $0 \leq V(A) \leq \frac{1}{4}$. As corollaries, we obtain some observations.

Corollary 1 Suppose we have a pair of projection valued measures (PVMs) $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. For any choice of a POVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$,

$$2D_{d_\infty}(A, f_A(F))D_{d_\infty}(B, f_B(F)) + D_{d_\infty}(A, f_A(F)) + D_{d_\infty}(B, f_B(F)) \\ + 4D_{d_\infty}(A, f_A(F))^{1/2}D_{d_\infty}(B, f_B(F))^{1/2} \geq \max_{a,b} \|[A_a, B_b]\|$$

holds.

Proof: For projection P , $P = P^2$ and thus $V(A) = V(B) = 0$ hold. Q.E.D.

From this corollary one can see that a pair of PVMs that have noncommutative elements is not simultaneously measurable. This tradeoff relation is, what we may call, Heisenberg's uncertainty principle.

On the other hand, it is possible for a pair of POVMs to be simultaneously measurable even if they are noncommutative with each other.

Corollary 2 Suppose that we have a pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. If they are simultaneously measurable, their intrinsic uncertainties $V(A)$ and $V(B)$ satisfy

$$V(A)^{1/2}V(B)^{1/2} \geq \frac{1}{2} \max_{a,b} \|[A_a, B_b]\|.$$

Proof: Put $D_{d_\infty}(f_A(F), A) = D_{d_\infty}(f_B(F), B) = 0$ in Theorem 1. Q.E.D.

The simultaneous measurability of noncommutative POVMs is not surprising at all. In fact, suppose that we have a doubly indexed POVM $F = \{F_{(a,b)}\}_{(a,b) \in \Omega_A \times \Omega_B}$, we can construct a pair of simultaneously measurable POVMs $f_A(F)$ and $f_B(F)$ by functions f_A and f_B with $f_A(a, b) = a$ and $f_B(a, b) = b$. The above corollary says that any such a pair must have sufficiently large intrinsic uncertainties.

The following result is also easy to obtain. We restrict the observable employed for the simultaneous measurement to PVM.

Corollary 3 *Suppose that we have a pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. For any choice of a PVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$,*

$$2D_{d_\infty}(A, f_A(F))D_{d_\infty}(B, f_B(F)) + D_{d_\infty}(A, f_A(F)) + D_{d_\infty}(B, f_B(F)) \geq \max_{a,b} \|[A_a, B_b]\|$$

holds.

Proof: In (2), we can put $[T(E_a^A), T(E_b^B)] = 0$. Q.E.D.

B. Uncertainty principle in l_1 distance

Next we consider another distance D_{d_1} induced by the l_1 distance. The following observation [10] plays a crucial role for the analysis. For a pair of POVMs $A = \{A_a\}_{a \in \Omega}$ and $A' = \{A'_a\}_{a \in \Omega}$,

$$D_{d_1}(A, A') = \sup_{\omega} \frac{1}{2} \sum_{a \in \Omega} |P_A^\omega(a) - P_{A'}^\omega(a)| = \max_{\Delta \subset \Omega} \left\| \sum_{a \in \Delta} A_a - \sum_{a \in \Delta} A'_a \right\|$$

holds. Thus if we define for each $\Delta_A \subset \Omega_A$ and $\Delta_B \subset \Omega_B$, $A_{\Delta_A} := \sum_{a \in \Delta_A} A_a$, $E_{\Delta_A}^A := \sum_{a \in \Delta_A} E_a^A$ and $B_{\Delta_B} := \sum_{b \in \Delta_B} B_b$, $E_{\Delta_B}^B := \sum_{b \in \Delta_B} E_b^B$, error operators should be introduced as $\epsilon_{\Delta_A}^A := T(E_{\Delta_A}^A) - A_{\Delta_A}$ and $\epsilon_{\Delta_B}^B := T(E_{\Delta_B}^B) - B_{\Delta_B}$, and the analysis to obtain equation (5) works just by replacing as follows:

$$\begin{aligned} A_a &\rightarrow A_{\Delta_A} \\ B_b &\rightarrow B_{\Delta_B} \\ E_a^A &\rightarrow E_{\Delta_A}^A \\ E_b^B &\rightarrow E_{\Delta_B}^B \\ \epsilon_a^A &\rightarrow \epsilon_{\Delta_A}^A \\ \epsilon_b^B &\rightarrow \epsilon_{\Delta_B}^B. \end{aligned}$$

That is, it holds

$$\begin{aligned} \|[A_{\Delta_A}, B_{\Delta_B}]\| &\leq 2\|\epsilon_{\Delta_A}^A\| \|\epsilon_{\Delta_B}^B\| + \|\epsilon_{\Delta_A}^A\| + \|\epsilon_{\Delta_B}^B\| \\ &\quad + 2(2\|\epsilon_{\Delta_A}^A\| + \|A_{\Delta_A} - A_{\Delta_A}^2\|)^{1/2} (2\|\epsilon_{\Delta_B}^B\| + \|B_{\Delta_B} - B_{\Delta_B}^2\|)^{1/2}. \end{aligned}$$

Taking the maximum over the subsets Δ_A and Δ_B , we obtain the following theorem.

Theorem 2 Suppose that we have a pair of POVMs $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. For any choice of a POVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$,

$$\begin{aligned} &2D_{d_1}(A, f_A(F))D_{d_1}(B, f_B(F)) + D_{d_1}(A, f_A(F)) + D_{d_1}(B, f_B(F)) \\ &+ 2(2D_{d_1}(A, f_A(F)) + V_1(A))^{1/2} (2D_{d_1}(B, f_B(F)) + V_1(B))^{1/2} \\ &\geq \max_{\Delta_A \subset \Omega_A, \Delta_B \subset \Omega_B} \left\| \sum_{a \in \Delta_A} \sum_{b \in \Delta_B} [A_a, B_b] \right\| \end{aligned}$$

holds, where $V_1(A) := \max_{\Delta_A \subset \Omega_A} \|\sum_{a \in \Delta_A} A_a(\mathbf{1} - \sum_{a \in \Delta_A} A_a)\|$ represents an intrinsic uncertainty of a POVM A (and similarly for $V_1(B)$).

The corresponding corollaries can be derived easily. For instance, the following statement hold.

Corollary 4 Suppose we have a pair of projection valued measures (PVMs) $A = \{A_a\}_{a \in \Omega_A}$ and $B = \{B_b\}_{b \in \Omega_B}$. For any choice of a POVM $F = \{F_x\}_{x \in \Omega}$ and a pair of functions $f_A : \Omega \rightarrow \Omega_A$ and $f_B : \Omega \rightarrow \Omega_B$,

$$\begin{aligned} &2D_{d_1}(A, f_A(F))D_{d_1}(B, f_B(F)) + D_{d_1}(A, f_A(F)) + D_{d_1}(B, f_B(F)) \\ &+ 4D_{d_1}(A, f_A(F))^{1/2} D_{d_1}(B, f_B(F))^{1/2} \geq \max_{\Delta_A \subset \Omega_A, \Delta_B \subset \Omega_B} \left\| \sum_{a \in \Delta_A} \sum_{b \in \Delta_B} [A_a, B_b] \right\| \end{aligned}$$

holds.

C. Example: A qubit

As the most simple example, we study a pair of PVMs for a single qubit. Each projection operator is parameterized by a Bloch sphere as, $E(\mathbf{n}) := \frac{1}{2}(\mathbf{1} + \mathbf{n} \cdot \boldsymbol{\sigma})$ for $\mathbf{n} \in \mathbf{R}^3$ with $|\mathbf{n}| = 1$. Let us consider two PVMs, $A_{\mathbf{n}} = \{E(\mathbf{n}), E(-\mathbf{n})\}$ and $A_{\mathbf{m}} = \{E(\mathbf{m}), E(-\mathbf{m})\}$ with $\angle(\mathbf{n}, \mathbf{m}) = \theta$ ($0 \leq \theta \leq \frac{\pi}{2}$). They satisfy the following inequality.

Corollary 5 For any POVM F and functions $f_{A_{\mathbf{n}}}$ and $f_{A_{\mathbf{m}}}$,

$$\begin{aligned} &2D_{d_\infty}(A_{\mathbf{n}}, f_{A_{\mathbf{n}}}(F))D_{d_\infty}(A_{\mathbf{m}}, f_{A_{\mathbf{m}}}(F)) + D_{d_\infty}(A_{\mathbf{n}}, f_{A_{\mathbf{n}}}(F)) + D_{d_\infty}(A_{\mathbf{m}}, f_{A_{\mathbf{m}}}(F)) \\ &+ 4D_{d_\infty}(A_{\mathbf{n}}, f_{A_{\mathbf{n}}}(F))^{1/2} D_{d_\infty}(A_{\mathbf{m}}, f_{A_{\mathbf{m}}}(F))^{1/2} \geq \frac{\sin \theta}{2} \end{aligned}$$

holds.

Proof: With $[E(\mathbf{n}), E(\mathbf{m})] = \frac{i}{2} \boldsymbol{\epsilon} \cdot (\mathbf{n} \times \mathbf{m})$, it is immediate from Corollary 1. Q.E.D.

This bound should be compared with a bound obtained in [13]. They derived an inequality,

$$D_{d_\infty}(A_{\mathbf{n}}, f_{A_{\mathbf{n}}}(F)) + D_{d_\infty}(A_{\mathbf{m}}, f_{A_{\mathbf{m}}}(F)) \geq \sqrt{\frac{1}{2}} \left(\cos \frac{\theta}{2} + \sin \frac{\theta}{2} - 1 \right). \quad (6)$$

In Figure 1 the contours of their admissible regions are shown. One can see that in some region ours is better and in other region worse. It may be interesting that our method gives a nonlinear estimate in contrast with [13].




FIG. 1. Admissible region of $X := D_{d_0}(f_A(F), A)$ (x -axis) and $Y := D_{d_0}(f_B(F), B)$ (y -axis) for $\theta = \frac{\pi}{2}$. A curved line is a contour of admissible region obtained by our method. A straight line is a contour of admissible region obtained in (6).

IV. SUMMARY

In this paper, a limitation on simultaneous measurement of two arbitrary (discrete) positive operator valued measures was investigated. Following Werner's work [10], we introduced the distance between observables by using distance between probability distributions. Introduction of the error operators simplified the analysis. We derived a novel inequality (Theorem 1), a possible representation of Heisenberg's uncertainty principle, that relates the limitation with noncommutativity. As a byproduct, we obtained a corollary indicating a necessary condition for a pair of POVMs to be simultaneously measurable. Compared with the previous works on this subject, the broad applicability of our result to an arbitrary discrete pair of POVMs is an advantage. Extension of our result to other distances will be a future problem.

Acknowledgments

The authors thank an anonymous referee for fruitful comments.

- [1] W. Heisenberg, Z. Phys. **43**, 172 (1927).
- [2] H. P. Robertson, Phys. Rev. **34**, 163 (1929).
- [3] H. J. Landau and H. O. Pollak, Bell Syst. Tech. J. **40**, 65 (1961).
- [4] D. Deutsch, Phys. Rev. Lett. **50**, 631 (1983).
- [5] H. Maassen and J. B. M. Uffink, Phys. Rev. Lett. **60**, 1103 (1988).
- [6] M. Krishna and K. R. Parthasarathy, Sankya, Ser. A **64**, 842 (2002).
- [7] T. Miyadera and H. Imai, Phys. Rev. A **76**, 062108 (2007).
- [8] D. M. Appleby, Int.J.Theor.Phys. **37**, 1491 (1998); *ibid.* **37** 2557 (1998).
- [9] M. Ozawa, Ann. Phys. **311**, 350 (2004).
- [10] R. F. Werner, Quantum Inf. Comp. **4**, 546 (2004).
- [11] B. Janssens, quant-ph/0606093.
- [12] P. Busch and D. B. Pearson, J. Math. Phys. **48**, 082103 (2007).
- [13] P. Busch and T. Heinosaari, Quantum Inf. Comp. **8**, 0797 (2008).
- [14] P. Busch, T. Heinonen and P. Lahti, Phys. Rep. **452**, 155 (2007).
- [15] H. Araki, *Mathematical Theory of Quantum Fields*, (Oxford Press, 2000).
- [16] B. Janssens, Master thesis, quant-ph/0503009.
- [17] A map Γ from a von Neumann algebra \mathcal{A} to a von Neumann algebra \mathcal{B} is called a unital completely positive map if (i) it is linear, (ii) $\Gamma(\mathbf{1}) = \mathbf{1}$ and (iii) its extension $\Gamma \otimes \text{id} : \mathcal{A} \otimes M(\mathbf{C}^N) \rightarrow \mathcal{B} \otimes M(\mathbf{C}^N)$ ($M(\mathbf{C}^N)$ is a set of all the $N \times N$ matrices) for any positive integer N , is positive. The condition (iii) is called complete positivity. If \mathcal{A} is Abelian, the complete positivity of the map Γ automatically follows from its positivity.