

On Spectral Triples in Quantum Gravity II

Johannes AASTRUP^{a1}, Jesper Møller GRIMSTRUP^{b2}
& Ryszard NEST^{c3}

^a SFB 478 “Geometrische Strukturen in der Mathematik”
Hittorfstr. 27, 48149 Münster, Germany

^b The Niels Bohr Institute
Blegdamsvej 17, DK-2100 Copenhagen, Denmark

^c Matematisk Institut
Universitetsparken 5, DK-2100 Copenhagen, Denmark

Abstract

A semifinite spectral triple for an algebra canonically associated to canonical quantum gravity is constructed. The algebra is generated by based loops in a triangulation and its barycentric subdivisions. The underlying space can be seen as a gauge fixing of the unconstrained state space of Loop Quantum Gravity. This paper is the second of two papers on the subject.

¹email: johannes.aastrup@uni-muenster.de

²email: grimstrup@nbi.dk

³email: rnest@math.ku.dk

Contents

1	Introduction	3
1.1	Content of the paper	3
2	Spaces of connections	6
2.1	Graphs	7
2.2	Completing spaces of connections	10
2.2.1	Smooth connections	11
2.2.2	Completing the group of gauge transformations	13
2.3	Group actions on the completions	14
2.4	Measure theory and Hilbert spaces	15
3	The Dirac operator on the completed space of connections	15
3.1	The case of subdividing an edge into two edges	16
3.2	A Riemannian metric	18
3.3	Some special covector fields	20
3.4	Construction of the Dirac operator	21
3.5	Gauge invariance	24
4	A semifinite spectral triple	26
4.1	The algebra	27
4.2	The trace and the von Neumann algebra	28
4.3	A coordinate change and selfadjointness of the Dirac operator	29
4.4	Affiliation of D	31
4.5	The main theorem	32
5	Appendix	36
5.1	The $U(1)$ -case	36
5.2	The $SU(2)$ -case	38

1 Introduction

One of the most striking application of Noncommutative Geometry [9] is Connes' derivation of the Standard Model of high energy physics [8,10]. In this derivation the Lagrangian of the full Standard Model coupled to gravity emerges from the spectral action principle [7] applied to a specific almost commutative geometry.

This formulation is, however, essentially classical and does, at a fundamental level, not involve quantization. This raises the question how the quantization procedure of Quantum Field Theory fits into the framework on Noncommutative Geometry. Since Connes' formulation of the Standard Model at its root is tied up to gravitation, a quantization scheme within the framework of Noncommutative Geometry must be expected to involve, at some level, quantum gravity.

In the papers [2] and [3] we started studying the question of formulating a quantization scheme within Noncommutative Geometry using canonical quantum gravity. Specifically, we were inspired by techniques applied in Loop Quantum Gravity. The concrete aim was to construct a spectral triple of a noncommutative algebra naturally associated to the unconstrained state space of Loop Quantum Gravity. Due to technical difficulties we only succeeded in constructing a spectral triple on a space closely related to the state space of Loop Quantum Gravity.

This paper is one of two papers presenting a more satisfactory solution to the problem of constructing a spectral triple on the state space of Loop Quantum Gravity as well as its physical interpretations. The paper [1] deals with the physical background and interpretations of the construction, and we therefore refer the reader to that paper for more thorough discussion. This paper deals with the concise mathematical construction.

1.1 Content of the paper

The unconstrained state space of Loop Quantum Gravity is the space of $SU(2)$ -connections in a trivial principal bundle on a three dimensional manifold. Since the construction we do works for arbitrary manifolds and arbitrary compact Lie-groups, and since we for physical applications might need it for later use, we will formulate it in this generality.

We let G be a compact Lie group and assume we have some principal G -fiber bundle P over a manifold M . The algebra associated to the space \mathcal{A} of connections in P we want to consider is the following:

Given a representation of G in M_N and given a loop L in M define a matrix

valued function on \mathcal{A} by

$$L(\nabla) = \text{Hol}(\nabla, L),$$

where $\text{Hol}(\nabla, L)$ denotes the holonomy of ∇ along L . Let \mathcal{B} be the algebra generated by all loops based in a fixed point in M . This is the type of algebra we want to consider.

The algebra \mathcal{B} is an algebra of matrix valued functions over \mathcal{A} . We will for technical reasons explained below consider a smaller algebra than \mathcal{B} . We will take a triangulation of M and consider the algebra \mathcal{B}_Δ which is constructed like \mathcal{B} but only includes loops lying in the edges of the triangulations, or one of its barycentric subdivisions. \mathcal{B}_Δ is also an algebra of matrix valued functions over \mathcal{A} . Since any loop can be approximated by a loop lying in the edges of the triangulations, or one of its barycentric subdivisions, \mathcal{B}_Δ can be considered as an approximation to \mathcal{B} . The crucial difference is that the groups of all diffeomorphisms preserving the base point acts on \mathcal{B} whereas only the group diffeomorphisms preserving the chosen triangulation and its barycentric subdivisions acts on \mathcal{B}_Δ .

In order to construct a spectral triple over \mathcal{B}_Δ we need a representation of \mathcal{B}_Δ . We therefore need to construct $L^2(\mathcal{A})$. A construction of $L^2(\mathcal{A})$ already exists within Loop Quantum Gravity, due to Ashtekar and Lewandowski. It turns out that this construction depends on a completion of \mathcal{A} and that this completion depends on a choice of a system \mathcal{S} of graphs. In the case of \mathcal{B}_Δ the relevant system of graphs is given by finite subgraphs of the triangulation and its barycentric subdivision. The system of graphs considered in Loop Quantum Gravity is that of graphs made up of piecewise real analytic edges (of course assuming M to be real analytic).

Seen from a graph Γ with edges $\{e_i\}_{i=1,\dots,n}$ the space of connections \mathcal{A} looks like G^n via

$$\mathcal{A} \ni \nabla \rightarrow (\text{Hol}(\nabla, e_1), \dots, \text{Hol}(\nabla, e_n)).$$

We will also denote $\mathcal{A}_\Gamma := G^n$. Of course \mathcal{A}_Γ tells little about \mathcal{A} . However, by letting the complexity of the graph grow we get a more and more refined picture of \mathcal{A} . This is implemented by noting that an embedding $\Gamma_1 \subset \Gamma_2$ naturally gives a map

$$P_{\Gamma_1\Gamma_2} : \mathcal{A}_{\Gamma_2} \rightarrow \mathcal{A}_{\Gamma_1},$$

and then simply define the completion of \mathcal{A} as the projective limit of \mathcal{A}_Γ , i.e.

$$\overline{\mathcal{A}}^{\mathcal{S}} := \varprojlim_{\Gamma \in \mathcal{S}} \mathcal{A}_\Gamma,$$

We will prove that under some condition on \mathcal{S} that \mathcal{A} is densely embedded in $\overline{\mathcal{A}}^{\mathcal{S}}$, and hence justifies the term completion.

The construction of $L^2(\mathcal{A})$ is straightforward from this completion. Since $\mathcal{A}_\Gamma = G^n$ we define $L^2(\mathcal{A}_\Gamma)$ as square integrable functions on G^n with respect to the Haar measure. The map $P_{\Gamma_1\Gamma_2}$ induces an embedding

$$P_{\Gamma_1\Gamma_2} : L^2(\mathcal{A}_{\Gamma_1}) \rightarrow L^2(\mathcal{A}_{\Gamma_2}),$$

and $L^2(\mathcal{A})$ is defined as an inductive limit

$$L^2(\mathcal{A}) := \varinjlim_{\Gamma \in \mathcal{S}} L^2(\mathcal{A}_\Gamma).$$

The idea for constructing the Dirac operator is that $\mathcal{A}_\Gamma = G^n$ is a classical geometry and therefore has a canonical Gauß-Bonnet-Dirac operator. In order to ensure that this defines an operator on the inductive limit we have to make sure that the operator is compatible with the structure maps of the projective limit. It is here the technical advantages of the triangulation compared to piecewise real analytic graphs shows up. The triangulation narrows down the types of structure maps appearing in the projective limit. With this we can define a Dirac operator compatible with the structure maps. It turns out that there is a lot freedom in the construction. Going from one level in the inductive limit to the next level we add new copies of G which corresponds to new degrees of freedom. Each of these degrees of freedom can be scaled. The entire construction therefore comes with a sequence of real non zero numbers $\{a^{j,k}\}_{k \leq j}$, where j, k is just a labelling of the degrees of freedoms convenient for the explicit construction.

However, the constructed Dirac operator together with the algebra \mathcal{B}_Δ will not fulfil the requirement of a spectral triple. The problem comes from the infinite dimensionality of the Clifford bundle of the space of connection. We therefore need to treat the identity operator on the Clifford bundle as a finite rank operator. The setting of semifinite spectral triples will allow exactly this. The von Neumann algebra \mathcal{N} appearing from this construction is a tensor product of the bounded operators on a separable Hilbert space and the weak closure of the CAR-algebra.

In order to obtain a semifinite spectral triple we still need to perturb the operator. At each level in the inductive limit we have to add a bounded perturbation roughly speaking of the form $b_j P_j$, where b_j is a real number and P_j is the projection onto a subspace related to the kernel of a part of the Dirac operator at the j 'th level.

The main result in the paper is [4.5.1](#) stating

Theorem *There exist sequences $\{a^{j,k}\}$ and $\{b_j\}$ such that \mathcal{B}_Δ together with the perturbed Dirac operator is a semifinite spectral triple with respect to the trace on \mathcal{N} .*

In the appendix we will demonstrate the case of $U(1)$ and show that for $SU(2)$ the perturbations are not needed.

Acknowledgements

We thank the following colleagues for fruitful discussions: Christian Fleischhack, Victor Gayral, Gerd Grubb, Troels Harmark, Thordur Jonsson, Mario Paschke, Adam Rennie, Carlo Rovelli, Thomas Schucker, Christian Voigh and Raimar Wulkenhaar. Furthermore, we are grateful to the following institutes for hospitality during visits: the Institute of Mathematics in Reykjavik, Iceland; The Max Planck Institute for Mathematics in the Sciences, Leipzig, Germany; the Isaac Newton Institute for Mathematical Sciences, Cambridge, UK; the Institute of Theoretical Physics in Marseilles, France.

Johannes Aastrup was funded by the German Research Foundation (DFG) within the research projects *Deformation Theory for Boundary Value Problems* and *Geometrische Strukturen in der Mathematik* (SFB 478).

2 Spaces of connections

In this section we will consider the space \mathcal{A} of connections in a principal fiber bundle. We will address the problem of topologizing this space together with the development of a measure theory. It turns out that the constructions, which we introduce in order to address these problems, depends on a completion of \mathcal{A} with respect to a system of graphs on the manifold. Different choices of graphs give different completions.

The material is standard from Loop Quantum Gravity, but we have chosen to write a more or less selfcontained exposition, not assuming prior knowledge of Loop Quantum Gravity, since we want to put emphasize on several different completions and their mutual interplay. The original techniques were developed by Ashtekar and Lewandowski in [4]. For a survey of Loop Quantum Gravity see [5] and for a detailed account see [14].

Let M be a manifold and P a G -principal bundle over M , where G is a compact connected Lie-group. We will for simplicity assume that P is the trivial bundle, i.e. P is isomorphic to $M \times G$.

2.1 Graphs

Definition 2.1.1 Let γ be a continuous piecewise smooth map $\gamma : [0, 1] \rightarrow M$ such that if $\gamma(t_1) = \gamma(t_2)$ then $t_1 = t_2$ or $t_1, t_2 \in \{0, 1\}$. We call γ a simple curve if $\dot{\gamma}(t)$ is non vanishing for all $t \in [0, 1]$. This requirement includes that the left and right derivative of γ is non vanishing in the non smooth points.

We will call $\gamma(0)$ the starting point and $\gamma(1)$ the endpoint.

If γ_1 and γ_2 are simple curves where the endpoint of γ_1 coincides with the starting point of γ_2 , then the composition is defined by

$$(\gamma_1 \circ \gamma_2)(t) = \begin{cases} \gamma_1(2t) & t \in [0, \frac{1}{2}] \\ \gamma_2(2t - 1) & t \in [\frac{1}{2}, 1] \end{cases}.$$

Note that the composition of two simple curves is not always a simple curve.

Definition 2.1.2 Two simple curves γ_1 and γ_2 are equivalent if there exist an increasing piecewise smooth bijection

$$\phi : [0, 1] \rightarrow [0, 1]$$

with

$$\gamma_1(\phi(t)) = \gamma_2(t), \quad t \in [0, 1],$$

and with nowhere vanishing derivative. This requirement includes that the left and right derivatives in the non smooth points are both non vanishing.

An equivalence class of simple curves is called a simple path.

Note that the relation defined above is in fact an equivalence relation.

The notion of a simple path chosen here implies that a simple path has an orientation. The inverse of a simple path represented by γ is the path represented by γ^{-1} , where

$$\gamma^{-1}(t) = \gamma(1 - t),$$

i.e. just the path with inverse orientation.

Definition 2.1.3 A graph Γ is the union of finite many simple paths $\{e_i\}$ where the only intersection points are start or end points of the simple paths.

The paths are called the edges of the graph. We denote the set of edges of Γ by E_Γ . The points $\{e_i(0), e_j(1)\}$ are called the vertices of Γ . We denote the set of vertices by V_Γ .

A graph is called connected if $\cup_{i,t} e_i(t)$ is connected.

We will consider paths on the graph. These are simply compositions of edges and inverses of edges. We will also think of each vertex as a path in the graph. Furthermore, we would like $e_i \circ (e_i)^{-1}$ to be equal to the path $e_i(0)$. We fix this with the following

Definition 2.1.4 Let Γ be a graph and let $\mathcal{P}(\Gamma)$ be the set of paths in Γ . We define an equivalence relation \sim on $\mathcal{P}(\Gamma)$ to be generated by

$$(e_i)^{-1} \circ e_i \sim e_i(1) \text{ and } e_i \circ (e_i)^{-1} \sim e_i(0).$$

The hoop groupoid $\mathcal{HG}(\Gamma)$ of path on Γ is defined by

$$\mathcal{HG}(\Gamma) = \mathcal{P}(\Gamma) / \sim.$$

This equivalence relation implies that if $p_1, p_2, p_3 \in \mathcal{P}(\Gamma)$ and $p_1 \sim p_2$ then $p_3 \circ p_1 \sim p_3 \circ p_2$ and $p_1 \circ p_3 \sim p_2 \circ p_3$ whenever the compositions are defined. The units for $\mathcal{HG}(\Gamma)$ are the vertices of Γ . It is easy to see that $\mathcal{HG}(\Gamma)$ is a groupoid.

Definition 2.1.5 Let Γ_1, Γ_2 be two graphs. We say that Γ_1 is a subgraph of Γ_2 , and we write $\Gamma_1 \subset \Gamma_2$, if the edges of Γ_1 are compositions of the edges or their inverses of Γ_2 .

Note that $\Gamma_1 \subset \Gamma_2$ implies that $\mathcal{HG}(\Gamma_1)$ is a subgroupoid of $\mathcal{HG}(\Gamma_2)$.

The relation \subset equips the set of graphs with a partial order. We are interested in subsystems of the set of all graphs.

Definition 2.1.6 A system \mathcal{S} of graphs is called directed if there for any $\Gamma_1, \Gamma_2 \in \mathcal{S}$ exists $\Gamma_3 \in \mathcal{S}$ with $\Gamma_1, \Gamma_2 \subset \Gamma_3$.

Definition 2.1.7 A system of graphs \mathcal{S} is called dense if there for every point $m \in M$ exists a coordinate chart $\mathbf{x} = (x_1, \dots, x_n)$ around m such that for all open subset U containing m in this coordinate chart there exists a collection of edges $e_1, \dots, e_n \subset U$ belonging to graphs in \mathcal{S} such that:

1. the e_i 's are straight lines with respect to the coordinate chart,
2. the tangent vectors of the e_i 's are linearly independent.

The definition of dense might appear awkward. The main purpose of the definition is to ensure proposition 2.2.4. Furthermore it is easy to check the definition in the concrete examples we have in mind. Certainly the requirement of straight lines can be eased.

We will now give three examples of dense systems of graphs, which will be important in the rest of the paper. These systems of graphs differ particularly in the size of their corresponding symmetry groups.

Example 1 Let \mathcal{S}_s be the system of all graphs. This system is clearly dense. The system is however not directed since we can have two simple paths e_1, e_2 with infinitely many isolated intersection points. Hence for graphs Γ_1, Γ_2 , where $e_1 \in \Gamma_1$ and $e_2 \in \Gamma_2$, there does not exist a graph Γ_3 containing Γ_1 and Γ_2 . The system \mathcal{S}_s admits a natural action of the diffeomorphism group $Diff(M)$. The system \mathcal{S}_s is the same as the piecewise immersed system defined in [11].

Example 2 Let M be a real analytic manifold. Let \mathcal{S}_a be the system of graph made up of real analytic simple curves. This system is dense and it is also directed since piecewise analytic curves have only finitely many isolated intersection points. The system carries a natural action of the group $Diff_a(M)$ of real analytic diffeomorphism, but no action of the full diffeomorphism group $Diff(M)$. This system was, in a base pointed version, first considered in [4].

Example 3 Let \mathcal{T} be a triangulation of M . We let Γ_0 be the graph consisting of all the edges in this triangulation. Strictly speaking this is not a graph if the manifold is not compact, but in this case we can consider Γ_0 as a system of graphs instead. Let \mathcal{T}_n be the triangulation obtained by barycentric subdividing each of the simplices in \mathcal{T} n times. The graph Γ_n is the graph consisting of the edges of \mathcal{T}_n . In this way we get a directed and dense system $\mathcal{S}_\Delta = \{\Gamma_n\}$ of graphs.

The important feature of \mathcal{S}_Δ which we are going to use in this paper is the following: **The step from Γ_n to Γ_{n+1} involves:**

1. **new edges are added**
2. **the edges of Γ_n get subdivided into two edges.**

The system \mathcal{S}_Δ only admits an action of the diffeomorphisms $Diff(\Delta)$ that maps edges in $\cup \mathcal{T}_n$ to edges in $\cup \mathcal{T}_n$. Hence this is a much more restrictive class of diffeomorphism than in the first two examples. Contrary to the first two examples, the system \mathcal{S}_Δ is countable.

Definition 2.1.8 Let \mathcal{S} be a directed system of graphs. We define the hoop groupoid $\mathcal{HG}(\mathcal{S})$ of \mathcal{S} to be the inductive limit

$$\mathcal{HG}(\mathcal{S}) = \varinjlim_{\Gamma \in \mathcal{S}} \mathcal{HG}(\Gamma).$$




Figure 1:

2.2 Completing spaces of connections

Given a graph Γ define the space

$$\mathcal{A}_\Gamma = \text{Hom}(\mathcal{HG}(\Gamma), G)$$

where G is the compact connected Lie-group. If $\Gamma_1 \subset \Gamma_2$ we have the embedding of groupoids $\mathcal{HG}(\Gamma_1) \rightarrow \mathcal{HG}(\Gamma_2)$, and we hence get a surjection

$$P_{\Gamma_1\Gamma_2} : \mathcal{A}_{\Gamma_2} \rightarrow \mathcal{A}_{\Gamma_1}.$$

Therefore, for a system of graphs \mathcal{S} we have a projective system

$$\{\mathcal{A}_\Gamma\}_{\Gamma \in \mathcal{S}}.$$

Definition 2.2.1 Let \mathcal{S} be a system of graphs. The space of generalized connections with respect to \mathcal{S} , denoted by $\overline{\mathcal{A}}^\mathcal{S}$, is defined by

$$\overline{\mathcal{A}}^\mathcal{S} = \varprojlim_{\Gamma \in \mathcal{S}} \mathcal{A}_\Gamma.$$

The projections from $\overline{\mathcal{A}}^\mathcal{S}$ to \mathcal{A}_Γ will be denoted by P_Γ .

For the systems $\mathcal{S}_s, \mathcal{S}_a, \mathcal{S}_\Delta$ we will denote the corresponding spaces of generalized connections by $\overline{\mathcal{A}}^s, \overline{\mathcal{A}}^a, \overline{\mathcal{A}}^\Delta$.

Note that when \mathcal{S} is directed we have the equality

$$\overline{\mathcal{A}}^\mathcal{S} = \text{Hom}(\mathcal{HG}(\mathcal{S}), G).$$

Given an element ∇ in \mathcal{A}_Γ we can associate to it $\Phi_\Gamma(\nabla) \in G^{n_\Gamma}$ where n_Γ is the numbers of edges in Γ . This is done by numbering the edges in Γ as e_1, \dots, e_{n_Γ} , and then defining Φ_Γ by

$$\Phi_\Gamma(\nabla) = (\nabla(e_1), \dots, \nabla(e_{n_\Gamma})).$$

Lemma 2.2.2 *The map $\Phi_\Gamma : \mathcal{A}_\Gamma \rightarrow G^{n_\Gamma}$ is a bijection.*

Proof. Follows since $\mathcal{HG}(\Gamma)$ is freely generated by the edges. \triangleleft

The bijection Φ_Γ gives a topology on \mathcal{A}_Γ by requiring Φ_Γ to be a homeomorphism. The topology is independent of the chosen numbering. The projection maps $P_{\Gamma_1\Gamma_2} : \mathcal{A}_{\Gamma_2} \rightarrow \mathcal{A}_{\Gamma_1}$ are continuous. In fact

$$\Phi_{\Gamma_1} \circ P_{\Gamma_1\Gamma_2} \circ (\Phi_{\Gamma_2})^{-1} : G^{n_{\Gamma_2}} \rightarrow G^{n_{\Gamma_1}}$$

is given by composition of one or more of the following operations:

- Multiplying g_{i_1} and g_{i_2} .
- Inverting g_i .
- Leaving out some g_i in $(g_1, \dots, g_{n_{\Gamma_2}}) \in G^{n_{\Gamma_2}}$.

Since $\overline{\mathcal{A}}^\mathcal{S}$ is a projective limit of $\{\mathcal{A}_\Gamma\}_{\Gamma \in \mathcal{S}}$, we define the topology on $\overline{\mathcal{A}}^\mathcal{S}$ as the projective limit topology. This topology is characterized by the following property:

Let X be a topological space and assume we have continuous maps $\phi_\Gamma : X \rightarrow \mathcal{A}_\Gamma$ for all $\Gamma \in \mathcal{S}$ such that $P_{\Gamma_1\Gamma_2} \circ \phi_{\Gamma_2} = \phi_{\Gamma_1}$ for all $\Gamma_1 \subset \Gamma_2$. Then there is a unique continuous map $\phi : X \rightarrow \overline{\mathcal{A}}^\mathcal{S}$ with $P_\Gamma \circ \phi = \phi_\Gamma$ for all $\Gamma \in \mathcal{S}$.

2.2.1 Smooth connections

Let \mathcal{A} denote the space of smooth connections in the principal bundle P . There is a natural map

$$\chi_\Gamma : \mathcal{A} \rightarrow \mathcal{A}_\Gamma = \text{Hom}(\mathcal{HG}(\Gamma), G)$$

defined by

$$\chi_\Gamma(\nabla)(p) = \text{Hol}(p, \nabla), \quad p \in \mathcal{HG}(\Gamma),$$

where $\text{Hol}(p, \nabla)$ denotes the holonomy of ∇ along p . Clearly $P_{\Gamma_1\Gamma_2} \circ \chi_{\Gamma_2} = \chi_{\Gamma_1}$ when $\Gamma_1 \subset \Gamma_2$, and hence by the property of the projective limit we get a unique map

$$\chi_\mathcal{S} : \mathcal{A} \rightarrow \overline{\mathcal{A}}^\mathcal{S}.$$

Proposition 2.2.3 *When \mathcal{S} is directed, $\chi_{\mathcal{S}}(\mathcal{A})$ is dense in $\overline{\mathcal{A}}^{\mathcal{S}}$.*

Proof. We first prove that each χ_{Γ} is surjective. The composition

$$\Phi_{\Gamma} \circ \chi_{\Gamma} : \mathcal{A} \rightarrow G^{n_{\Gamma}}$$

is given by

$$\Phi_{\Gamma}(\chi_{\Gamma}(\nabla)) = (\text{Hol}(e_1, \nabla), \dots, \text{Hol}(e_{n_{\Gamma}}, \nabla)).$$

Let $(g_1, \dots, g_{n_{\Gamma}})$ be given. Since G is connected we can for each i find a connection ∇_i with

$$\text{Hol}(e_i, \nabla_i) = g_i.$$

It is furthermore easy to see that since the e_i 's only intersect in the endpoints we can arrange that $\nabla_i = 0$ on all the edges in Γ apart from e_i . Hence

$$\Phi_{\Gamma} \left(\chi_{\Gamma} \left(\sum \nabla_i \right) \right) = (g_1, \dots, g_{n_{\Gamma}}).$$

Therefore χ_{Γ} is surjective.

From the directedness of \mathcal{S} and the surjectivity of all χ_{Γ} the density follows.

◁

The system \mathcal{S}_s is not directed. It is however possible with more elaborate methods to prove that \mathcal{A} is dense in $\overline{\mathcal{A}}^{\mathcal{S}_s}$ when G is semi-simple, see for example the discussion in [12].

We now turn to the question of injectivity of χ .

Proposition 2.2.4 *When \mathcal{S} is dense, $\chi_{\mathcal{S}}$ is injective.*

Proof. Let ∇_1, ∇_2 be two different smooth connections. Hence there is a point $m \in M$ where $\nabla_1(m) \neq \nabla_2(m)$. Let us choose a coordinate chart $\mathbf{x} = (x_1, \dots, x_n)$ around m according to the density of \mathcal{S} , where m corresponds to $\mathbf{x} = \mathbf{0}$. We can then write

$$\nabla_k = \sum_j \mathfrak{g}_j^k(\mathbf{x}) dx_j, \quad k \in \{1, 2\},$$

where $\mathfrak{g}_j^i(\mathbf{x})$ is a smooth function of \mathbf{x} with values in the Lie-algebra of G . Let U be a neighbourhood of $\mathbf{0}$ such that we can assume that the functions $\mathfrak{g}_j^i(\mathbf{x})$ are constant in U with sufficiently good approximation. Let e_1, \dots, e_n be the edges in \mathcal{S} which are straight lines with respect to the coordinate chart, which belongs to U and where the tangent vectors $\mathbf{t}^1, \dots, \mathbf{t}^n$ are linearly independent. We assume

that the edges are parametrized by arc lengths. Since $\nabla_1(m) \neq \nabla_2(m)$ there is a j such that

$$\sum_i \mathfrak{g}_i^1(\mathbf{0})t_i^j \neq \sum_i \mathfrak{g}_i^2(\mathbf{0})t_i^j.$$

With sufficiently good approximation we have

$$\text{Hol}(e_j, \nabla_k) = \exp\left(\sum_i \mathfrak{g}_i^1(\mathbf{0})t_i^j\right), \quad k \in \{1, 2\}.$$

Hence, if we had chosen U small enough, we conclude

$$\text{Hol}(e_j, \nabla_1) \neq \text{Hol}(e_j, \nabla_2).$$

Thus $\chi_S(\nabla_1) \neq \chi_S(\nabla_2)$. ◁

2.2.2 Completing the group of gauge transformations

Let U be an element in the gauge group \mathcal{G} of $M \times P$, i.e. $U : M \rightarrow G$ is a smooth function. Given a connection $\nabla \in \mathcal{A}$, U induces a gauge transformed connection $\tilde{\nabla}$. Given a path p on M with startpoint x_0 and endpoint x_1 the holonomy along e transforms according to

$$\text{Hol}(p, \tilde{\nabla}) = U(x_0)\text{Hol}(p, \nabla)U^{-1}(x_1).$$

This leads to the following

Definition 2.2.5 *Let Γ be a graphs and $U : V_\Gamma \rightarrow G$ be a map. Define $U_* : \mathcal{A}_\Gamma \rightarrow \mathcal{A}_\Gamma$ by*

$$U_*(\nabla)(e) = U(e(0))\nabla(e)U(e(1))^{-1} \text{ for all } e \in E_\Gamma.$$

Since $\mathcal{HG}(\Gamma)$ is freely generated by E_Γ , this is well defined.

We denote by \mathcal{G}_Γ the group of all maps $U : V_\Gamma \rightarrow G$.

Note that via U_* we get a left action of \mathcal{G}_Γ on \mathcal{A}_Γ .

Like with spaces of connections there are natural projections

$$P_{\Gamma_1\Gamma_2} : \mathcal{G}_{\Gamma_2} \rightarrow \mathcal{G}_{\Gamma_1},$$

when $\Gamma_1 \subset \Gamma_2$ and

$$P_{\Gamma_1\Gamma_2}(U_*(\nabla)) = P_{\Gamma_1\Gamma_2}(U)_*(P_{\Gamma_1\Gamma_2}(\nabla)), \quad \nabla \in \mathcal{A}_{\Gamma_2}, U \in \mathcal{G}_{\Gamma_2}. \quad (1)$$

Definition 2.2.6 Let \mathcal{S} be a system of graphs. Put

$$\overline{\mathcal{G}}^{\mathcal{S}} = \{U|U : \cup_{\Gamma \in \mathcal{S}} V_{\Gamma} \rightarrow G\}.$$

Define the left action of $\overline{\mathcal{G}}^{\mathcal{S}}$ on $\overline{\mathcal{A}}^{\mathcal{S}}$ by

$$U \cdot \nabla = (U|_{V_{\Gamma}})_*(\nabla), \quad \nabla \in \mathcal{A}_{\Gamma}.$$

Due to (1) this is well defined.

A gauge transformation $U \in \mathcal{G}$ naturally gives an element in $\overline{\mathcal{G}}^{\mathcal{S}}$. When \mathcal{S} is dense this induces an embedding $\mathcal{G} \rightarrow \overline{\mathcal{G}}^{\mathcal{S}}$. Via this embedding we get an action of \mathcal{G} on $\overline{\mathcal{A}}^{\mathcal{S}}$, which extends the action of \mathcal{G} on \mathcal{A} . We will therefore call $\overline{\mathcal{G}}^{\mathcal{S}}$ the completed gauge group, or simply the gauge group.

2.3 Group actions on the completions

If a system \mathcal{S}_1 is contained in \mathcal{S}_2 we get a surjective continuous map

$$P_{\mathcal{S}_1 \mathcal{S}_2} : \overline{\mathcal{A}}^{\mathcal{S}_2} \rightarrow \overline{\mathcal{A}}^{\mathcal{S}_1}.$$

In particular we have the commutative diagram

$$\begin{array}{ccccc} \overline{\mathcal{A}}^{\mathcal{S}} & \xrightarrow{P_{\mathcal{S}_a \mathcal{S}_s}} & \overline{\mathcal{A}}^a & \xrightarrow{P_{\mathcal{S}_{\Delta} \mathcal{S}_a}} & \overline{\mathcal{A}}^{\Delta} \\ & \searrow & & \searrow & \\ & & & & P_{\mathcal{S}_{\Delta} \mathcal{S}_s} \end{array}$$

Of course the map $P_{\mathcal{S}_{\Delta} \mathcal{S}_a}$ only exists if the triangulation is real analytic.

The group of all diffeomorphisms $Diff(M)$ acts on \mathcal{A} . The question is what kind of groups acts on the different completions $\overline{\mathcal{A}}^{\mathcal{S}}$. In general a diffeomorphism d preserving \mathcal{S} induces a homeomorphism $d^* : \overline{\mathcal{A}}^{\mathcal{S}} \rightarrow \overline{\mathcal{A}}^{\mathcal{S}}$. Thus in this way there is an action of $Diff(\mathcal{S})$, the diffeomorphisms preserving \mathcal{S} , on $\overline{\mathcal{A}}^{\mathcal{S}}$. We have the following diagram

$$\begin{array}{ccccc} & & \overline{\mathcal{A}}^{\mathcal{S}} & \xleftarrow{\quad} & Diff(M) \\ & \nearrow \chi_{\mathcal{S}_s} & \downarrow P_{\mathcal{S}_a \mathcal{S}_s} & \searrow P_{\mathcal{S}_{\Delta} \mathcal{S}_s} & \uparrow \\ \mathcal{A} & \xrightarrow{\chi_{\mathcal{S}_a}} & \overline{\mathcal{A}}^a & \xleftarrow{\quad} & Diff_a(M) \\ & \searrow \chi_{\mathcal{S}_{\Delta}} & \downarrow P_{\mathcal{S}_{\Delta} \mathcal{S}_a} & \searrow & \uparrow \\ & & \overline{\mathcal{A}}^{\Delta} & \xleftarrow{\quad} & Diff(\Delta) \end{array}$$

We see that the size of the completion of \mathcal{A} is strongly related to the size of the symmetry group. The spaces $\overline{\mathcal{A}}^s, \overline{\mathcal{A}}^a$ are non separable and the symmetry groups are large, whereas $\overline{\mathcal{A}}^\Delta$ is separable and the symmetry group $\text{Diff}(\Delta)$ is comparatively small. In this way we can think of $\overline{\mathcal{A}}^\Delta$ as being $\overline{\mathcal{A}}^s$ (or $\overline{\mathcal{A}}^a$) subjected to a kind of gauge fixing of $\text{Diff}(M)$ (or $\text{Diff}_a(M)$).

2.4 Measure theory and Hilbert spaces

We will here recall the construction of measures and Hilbert space structures on completions of spaces of connections. The construction first appeared in [4]. See also [13] for a different approach.

Because of lemma 2.2.2 we can identify \mathcal{A}_Γ with G^{n_Γ} via Φ_Γ . On G^{n_Γ} there is a canonical normalized measure, namely the Haar measure $\mu_{G^{n_\Gamma}}$. Denote by μ_Γ the image measure of $\mu_{G^{n_\Gamma}}$ under Φ_Γ^{-1} .

Lemma 2.4.1 *Let $\Gamma_1 \subset \Gamma_2$. The image measure of μ_{Γ_2} under $P_{\Gamma_2\Gamma_1}$ is μ_{Γ_1} .*

Proof. See for example lemma I.2.9 in [14]. ◁

Lemma 2.4.1 ensures that $\{\mathcal{A}_\Gamma, \mu_\Gamma\}_{\Gamma \in \mathcal{S}}$ is a projective system of measure spaces. Therefore, according to Theorem I.2.10 in [14], there is a unique measure μ on $\overline{\mathcal{A}}^S$ such that $P_\Gamma(\mu) = \mu_\Gamma$ for all $\Gamma \in \mathcal{S}$. Also lemma 2.4.1 ensures that

$$P_{\Gamma_2\Gamma_1}^* : L^2(\mathcal{A}_{\Gamma_1}) \rightarrow L^2(\mathcal{A}_{\Gamma_2})$$

is an embedding of Hilbert spaces when $\Gamma_1 \subset \Gamma_2$.

Proposition 2.4.2 *Let*

$$L^2(\mathcal{A}) = \varinjlim L^2(\mathcal{A}_\Gamma),$$

the inductive limit of the Hilbert spaces $\{L^2(\mathcal{A}_\Gamma)\}_{\Gamma \in \mathcal{S}}$. Then $L^2(\mathcal{A}) = L^2(\overline{\mathcal{A}}^S)$, where the latter is with respect to the measure μ .

Proof. This follows directly from the construction. See for example section I.2.4 in [14]. ◁

3 The Dirac operator on the completed space of connections

In this section we will only work with the completed space $\overline{\mathcal{A}}^\Delta$. We will construct a Dirac type operator acting on a Hilbert space \mathcal{H} which can naturally be seen as

square integrable functions on $\overline{\mathcal{A}}^\Delta$ with values in an infinite dimensional vector bundle.

The idea is to construct a Dirac operator on each \mathcal{A}_Γ . Since these look like classical geometries we more or less only have to put a Riemannian metric on each of these. Next we need to check that the construction made on each \mathcal{A}_Γ is consistent with maps between the different graphs.

In this section we have no restrictions on the manifold M . In particular we can have infinitely many simplices in \mathcal{T}_0 .

To simplify the notation objects indexed by the graph Γ_k will be indexed simply by k for the rest of the paper.

3.1 The case of subdividing an edge into two edges

The construction of the Dirac operator, which will be carried out in the following subsections, looks cumbersome, but is really forced upon us by the requirement of consistency with the maps between different graphs. Therefore, in order to present the construction without the full notational weight, we will first demonstrate the case where the triangulation consists of one edge and where we consider the step going to the first barycentric subdivision. We thus have $\mathcal{A}_0 = G$, $\mathcal{A}_1 = G^2$ and $P_{0,1}(g_1, g_2) = g_1 g_2$.

For Γ_0 we choose a left and right invariant metric $\langle \cdot, \cdot \rangle$ on G . Let \hat{e}_i be an orthonormal basis for $T_{id}G$ and let $\hat{E}_i(g) = L_g \hat{e}_i$ be the corresponding left translated vector fields. Define the Dirac operator

$$D_0(\xi) = \sum \hat{E}_i \nabla_{\hat{E}_i}(\xi),$$

where ∇ is some $SO(\dim(G))$ -valued connection and $\xi \in L^2(G, Cl(TG))$, where $Cl(TG)$ denotes the Clifford bundle.

We want to construct D_1 acting on $L^2(G^2, Cl(G^2))$ on the form

$$D_1(\xi) = \sum_j K_j \nabla_{K_j}(\xi),$$

where K_j is an orthonormal frame in TG^2 , such that

$$P_{0,1}^*(D_0(\xi)) = D_1(P_{0,1}^*(\xi)).$$

First we have to make sense of $P_{0,1}^*$. Since $P_{0,1}$ induces maps

$$(P_{0,1})_* : TG^2 \rightarrow TG, \quad \text{and} \quad P_{0,1}^* : T^*G \rightarrow T^*G^2$$

between tangent and cotangent spaces it is also natural to let the Dirac operators act on $Cl(T^*G)$, resp. $Cl(T^*G^2)$ instead of $Cl(TG)$, resp. $Cl(TG^2)$. This is easily done once we have chosen a metric on T^*G and resp. T^*G^2 . In order for

$$P_{0,1}^* : L^2(G, Cl(T^*G)) \rightarrow L^2(G^2, Cl(T^*G^2))$$

to be an embedding of Hilbert spaces, and in fact to be defined at the level of Clifford bundles, the map induced by $P_{0,1}$ at the level of cotangent bundles, also denoted $P_{0,1}^*$, must be metric.

Let $w \in T_g^*G$. It is easy to see that

$$P_{0,1}^*(w) = (R_{g_2^{-1}}w, L_{g_1^{-1}}w), \quad g_1g_2 = g,$$

where

$$R_{g_2^{-1}}(w)(v) = w(R_{g_2}v)$$

and

$$L_{g_1^{-1}}(w)(v) = w(L_{g_1}v)$$

. We ensure that $P_{0,1}^* : T^*G \rightarrow T^*G^2$ is metric by defining the inner product on T^*G^2

$$\langle (w_1, w_2), (w_3, w_4) \rangle_2 = \frac{1}{2}(\langle w_1, w_3 \rangle + \langle w_2, w_4 \rangle).$$

Denote by E_i the cotangent vector field which is dual to \hat{E}_i via $\langle \cdot, \cdot \rangle$. Using the inner product on G^2 we get from $P_{0,1}^*(E_i)$ a vector field on G^2 . A small computation shows that this vector field equals

$$(L_{g_1}L_{g_2}R_{g_2^{-1}}\hat{e}_i, L_{g_2}\hat{e}_i) =: (\hat{\mathcal{E}}_i^1, \hat{\mathcal{E}}_i^2). \quad (2)$$

Put $\hat{\mathcal{E}}_i^+ = (\hat{\mathcal{E}}_i^1, \hat{\mathcal{E}}_i^2)$ and $\hat{\mathcal{E}}_i^- = (\hat{\mathcal{E}}_i^1, -\hat{\mathcal{E}}_i^2)$. Since $\{\hat{\mathcal{E}}_i^\pm\}_i$ is an orthonormal frame for TG^2 it is natural to try to define

$$D_1(\xi) = \sum_i \mathcal{E}_i^+ \nabla_{\hat{\mathcal{E}}_i^+} \xi + \sum_i \mathcal{E}_i^- \nabla_{\hat{\mathcal{E}}_i^-} \xi.$$

where $\xi \in L^2(G^2, Cl(T^*G^2))$ and where $\{\hat{\mathcal{E}}_i^\pm\}$ is the corresponding orthonormal frame for T^*G^2 . A small computation shows that

$$\begin{aligned} \hat{\mathcal{E}}_i^+(P_{0,1}^*\xi) &= 2P_{0,1}^*(\hat{E}_i(\xi)), & \xi &\in L^2(G) \\ \hat{\mathcal{E}}_i^-(P_{0,1}^*\xi) &= 0, & \xi &\in L^2(G). \end{aligned}$$

We therefore define

$$D_1(\xi) = \frac{1}{2} \left(\sum_i \mathcal{E}_i^+ \nabla_{\mathcal{E}_i^+} \xi + \sum_i \mathcal{E}_i^- \nabla_{\mathcal{E}_i^-} \xi \right).$$

Next, write an element in $\xi \in L^2(G, T^*G)$ as

$$\xi = \sum_i \xi_i E_i,$$

where $\xi_i \in L^2(G)$. Hence

$$P_{0,1}^*(\xi) = \sum_i P_{0,1}^*(\xi_i) \mathcal{E}_i^+.$$

We calculate

$$\begin{aligned} & D_1(P_{0,1}^*(\xi)) \\ &= \sum_{i,j} \left(P_{0,1}^*(\hat{E}_i(\xi_j)) \mathcal{E}_i^+ \mathcal{E}_j^+ + \frac{1}{2} \left(P_{0,1}^*(\xi_j) \mathcal{E}_i^+ \nabla_{\mathcal{E}_i^+} \mathcal{E}_j^+ + P_{0,1}^*(\xi_j) \mathcal{E}_i^- \nabla_{\mathcal{E}_i^-} \mathcal{E}_j^+ \right) \right). \end{aligned}$$

If we therefore require ∇ to be a $SO(2\dim(G))$ -valued connection in T^*G^2 satisfying

$$\begin{aligned} \nabla_{\mathcal{E}_i^+}(\mathcal{E}_j^+) &= P_{0,1}^*(\nabla_{\hat{E}_i}(E_j)) \\ \nabla_{\mathcal{E}_i^-}(\mathcal{E}_j^+) &= 0 \end{aligned}$$

we see that $P_{0,1}^*(D_0(\xi)) = D_1(P_{0,1}^*(\xi))$ for all $\xi \in L^2(G, Cl(T^*G))$.

3.2 A Riemannian metric

Due to lemma 2.2.2 we have at each level the identification $\mathcal{A}_k = G^{n_k}$, where n_k is the number of edges in Γ_k . We want to construct a Riemannian metric on $T\mathcal{A}_k = TG^{n_k}$. However, we require the metric to be consistent with the embeddings of graphs. Embeddings of graphs $\Gamma_k \subset \Gamma_{k+1}$ induces a surjective smooth map

$$P_{k,k+1} : \mathcal{A}_{k+1} \rightarrow \mathcal{A}_k,$$

and therefore a map of tangent bundles

$$(P_{k,k+1})_* : T\mathcal{A}_{k+1} \rightarrow T\mathcal{A}_k.$$

Dualizing this map we get an embedding

$$P_{k,k+1}^* : T^* \mathcal{A}_k \rightarrow T^* \mathcal{A}_{k+1}$$

of cotangent bundles.

We want to construct the Hilbert space on which the Dirac operator acts as a inductive limit of Hilbert spaces. It is hence natural to construct the metric on the cotangent bundle of \mathcal{A}_k , since we have canonical maps

$$P_{k,k+1}^* : L^2(\mathcal{A}_k, T^* \mathcal{A}_k) \rightarrow L^2(\mathcal{A}_{k+1}, T^* \mathcal{A}_{k+1}).$$

Definition 3.2.1 Let $\langle \cdot, \cdot \rangle$ be a left and right invariant Riemannian metric on T^*G . Let Γ_k be the graph consisting of the edges in \mathcal{T}_k , the k 's barycentric subdivision of \mathcal{T}_0 . On $T^* \mathcal{A}_k = T^* G^{n_k}$ define

$$\langle \cdot, \cdot \rangle_k = \frac{1}{2^k} \langle \cdot, \cdot \rangle^{n_k}, \quad (3)$$

where $\langle \cdot, \cdot \rangle^{n_k}$ is the product metric on $T^* G^{n_k}$.

Proposition 3.2.2 The map

$$P_{k,k+1}^* : T^* \mathcal{A}_k \rightarrow T^* \mathcal{A}_{k+1}$$

preserves the metric (3).

Proof. We subdivide $G^{n_{k+1}}$ as $G^{2n_k} \times G^{n_2}$, where G^{2n_k} corresponds to subdivision of the edges in Γ_k and G^{n_2} corresponds to the new edges added when going from k to $k+1$. Write

$$T_{(g_1, \dots, g_{n_{k+1}})}^* \mathcal{A}_{k+1} = T_{g_1}^* G \oplus \dots \oplus T_{g_{n_{k+1}}}^* G.$$

We choose orientations of the edges in such a way that

$$P_{k,k+1}(g_1, \dots, g_{n_{k+1}}) = (g_1 g_2, g_3 g_4, \dots, g_{2n_k-1} g_{2n_k}).$$

For a tangent vector $\mathbf{v} = (v_1, \dots, v_{n_{k+1}})$ in $T^* \mathcal{A}_{k+1}$ we have

$$(P_{k,k+1})_*(\mathbf{v}) = (L_{g_1} v_2 + R_{g_2} v_1, L_{g_3} v_4 + R_{g_4} v_3, \dots, L_{g_{2n_k-1}} v_{2n_k} + R_{g_{2n_k}} v_{2n_k-1}),$$

where L_g means left translation of tangent vectors and R_g right translation. Hence for a cotangent vector $\mathbf{w} = (w_1, \dots, w_{n_k})$ in $T_{(g_1 g_2, \dots, g_{2n_k-1} g_{2n_k})}^* \mathcal{A}_k$ we have

$$P_{k,k+1}^*(\mathbf{w}) = (R_{g_2}^{-1} w_1, L_{g_1}^{-1} w_1, \dots, R_{g_{2n_k}}^{-1} w_{n_k}, L_{g_{2n_k-1}} w_{n_k}, 0, \dots, 0),$$

where by definition

$$L_g(w)(v) = w(L_{g^{-1}}(v)), \quad R_g(w)(v) = w(R_{g^{-1}}(v)), \quad v \in TG, \quad w \in T^*G.$$

From this and the left and right invariance of $\langle \cdot, \cdot \rangle$ the proposition follows. \triangleleft

By proposition 3.2.2 the map $P_{k,k+1}^*$ induces a map, also denoted $P_{k,k+1}^*$, from $Cl(T^*\mathcal{A}_k)$ to $Cl(T^*\mathcal{A}_{k+1})$, where $Cl(T^*\mathcal{A}_k)$ denotes the Clifford bundle of $T^*\mathcal{A}_k$ with respect to $\langle \cdot, \cdot \rangle_k$. Furthermore this map is isometric. We thus get embeddings of Hilbert spaces

$$P_{k,k+1}^* : L^2(\mathcal{A}_k, Cl(T^*\mathcal{A}_k)) \rightarrow L^2(\mathcal{A}_{k+1}, Cl(T^*\mathcal{A}_{k+1})).$$

Definition 3.2.3 Put $\mathcal{H}_k = L^2(\mathcal{A}_k, Cl(T^*\mathcal{A}_k))$. Define

$$\mathcal{H} = \varinjlim \mathcal{H}_k,$$

the inductive limit of the system $\{\mathcal{H}_k, P_{k,k+1}^*\}_k$.

Due to proposition 2.4.2 we can consider \mathcal{H} as $L^2(\overline{\mathcal{A}}^\Delta, Cl(\overline{\mathcal{A}}^\Delta))$ or more freely written $L^2(\mathcal{A}, Cl(\mathcal{A}))$.

3.3 Some special covector fields

For the general construction of the Dirac operator we will need some special covector fields generalizing duals of the vector fields (2). For notational simplicity we will only present the construction on a single edge which is then subdivided infinitely many times. Thus $\mathcal{A}_n = G^{2^n}$ and the structure maps are given by

$$P_{n-1,n}(g_1, g_2, \dots, g_{2^{n-1}}, g_{2^n}) = (g_1 g_2, \dots, g_{2^{n-1}} g_{2^n}).$$

Let $\{e_i\}$ be an orthonormal basis of T_{id}^*G . Define covector fields on $\mathcal{A}_0 = G$ by

$$\mathcal{E}_i^{0,1}(g) = L_g(e_i).$$

The construction of the covector fields on \mathcal{A}_n will be by induction. Assume that covector fields

$$\{\mathcal{E}_i^{j,k}\}_{j \leq n-1, k \leq 2^{j-1}}$$

on \mathcal{A}_{n-1} has been defined (For $j = 0$ the set $k \leq 2^{j-1}$ is $\{1\}$). We adopt the notation

$$\begin{aligned}\mathcal{E}_i^1 &= L_{g_1 g_2 \cdots g_{2^n}} R_{(g_2 g_3 \cdots g_{2^n})^{-1}} e_i \\ \mathcal{E}_i^2 &= L_{g_2 g_3 \cdots g_{2^n}} R_{(g_3 g_4 \cdots g_{2^n})^{-1}} e_i \\ &\vdots\end{aligned}$$

for covector fields on G depending on g_1, \dots, g_{2^n} . Define covector fields on \mathcal{A}_n by

$$\{P_{n-1,n}^*(\mathcal{E}_i^{j,k})\}_{j \leq n-1, k \leq 2^{j-1}}$$

and

$$\begin{aligned}\mathcal{E}_i^{n,1} &= 2^{n-1}(\mathcal{E}_i^1, -\mathcal{E}_i^2, 0, \dots, 0) \\ \mathcal{E}_i^{n,2} &= 2^{n-1}(0, 0, \mathcal{E}_i^3, -\mathcal{E}_i^4, 0, \dots, 0) \\ &\vdots\end{aligned}$$

With sloppy notation we will also write $\mathcal{E}_i^{j,k}$ instead of $P_{n-1,n}^*(\mathcal{E}_i^{j,k})$.

Lemma 3.3.1 $\{\mathcal{E}_i^{j,k}\}_{j \leq n, k \leq 2^{j-1}}$ is an orthonormal frame for $T^*\mathcal{A}_n$ with respect to the inner product $\langle \cdot, \cdot \rangle_n$.

Proof. Since $P_{n-1,n}^*$ preserves the inner product by proposition 3.2.2 it is enough to check that $\{\mathcal{E}_i^{n,k}\}_{k \leq 2^{n-1}}$ are orthonormal and are orthogonal to $\{\mathcal{E}_i^{j,k}\}_{j \leq n-1, k \leq 2^{j-1}}$.

It is clear that $\{\mathcal{E}_i^{n,k}\}_{k \leq 2^{n-1}}$ are orthonormal. By induction and the computation in the proof of proposition 3.2.2 we see that the vectors in $\{\mathcal{E}_i^{j,k}\}_{j \leq n-1, k \leq 2^{j-1}}$ are of the form

$$(c_1 \mathcal{E}_i^1, c_1 \mathcal{E}_i^2, c_2 \mathcal{E}_i^3, c_2 \mathcal{E}_i^4, \dots, c_{2^{n-1}-1} \mathcal{E}_i^{2^{n-1}-1}, c_{2^{n-1}-1} \mathcal{E}_i^{2^{n-1}}).$$

From this the lemma follows. \triangleleft

3.4 Construction of the Dirac operator

We want to construct a Dirac type operator acting on \mathcal{H} . For this we construct Dirac type operators acting on \mathcal{H}_n which are consistent with $P_{n,n+1}^*$. We first need to determine sufficient conditions on the connections which permit the existence of the Dirac operator.

Let $\hat{\mathcal{E}}_i^{j,k}$ be the vector field obtained from $\mathcal{E}_i^{j,k}$ by using $\langle \cdot, \cdot \rangle_n$ to identify $T^* \mathcal{A}_n$ with $T \mathcal{A}_n$. Also let \hat{e}_i denote the basis in $T_{id} G$ obtained from e_i by using $\langle \cdot, \cdot \rangle$ to identify $T_{id} G$ with $T_{id}^* G$. Note also under this identification the covectorfield

$$(0, \dots, 0, \mathcal{E}_i^j, 0, \dots)$$

gets mapped to

$$\frac{1}{2^n} (0, \dots, 0, L_{g_j \dots g_{2^n}} R_{(g_{j+1} \dots g_{2^n})^{-1}} \hat{e}_i, 0, \dots, 0).$$

Definition 3.4.1 A system of connections $\{\nabla^n\}_n$, where ∇^n is a connection in $T^* \mathcal{A}_n$ is called admissible if ∇^n is a $SO(2^n \cdot \dim(G))$ -valued connection and if

$$\begin{aligned} \nabla_{\hat{\mathcal{E}}_i^{j,k}}^n(\mathcal{E}_l^{m,p}) &= P_{n-1,n}^*(\nabla_{\hat{\mathcal{E}}_i^{j,k}}^{n-1}(\mathcal{E}_l^{m,p})) \quad j, m < n \\ \nabla_{\hat{\mathcal{E}}_i^{n,k}}^n(\mathcal{E}_l^{m,p}) &= 0 \quad m < n \end{aligned}$$

We want to define the Dirac operator in the usual fashion using an admissible family of connection, i.e. at each level it should be on the form

$$D_n = \sum \mathcal{E}_i^{j,k} \cdot \nabla_{\hat{\mathcal{E}}_i^{j,k}}^n.$$

Proposition 3.4.2 Let $\{\nabla^n\}$ be an admissible system of connections and let $\{a^{j,k}\}_{k \leq 2^{j-1}}$ be a sequence of complex numbers. For $\xi \in \mathcal{H}_n$ define

$$D_n(\xi) = \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\hat{\mathcal{E}}_i^{j,k}}^n \xi.$$

Then

$$P_{n,n+1}^*(D_n(\xi)) = D_{n+1}(P_{n,n+1}^*(\xi))$$

and hence the system of operators $\{D_n\}$ defines a densely defined operator D on \mathcal{H} .

Proof. We first check the identity on functions, i.e. $\xi \in L^2(\mathcal{A}_n)$:

$$\begin{aligned} & P_{n,n+1}^*(D_n(\xi))(g_1, g_2, \dots, g_{2^{n+1}}) \\ &= P_{n,n+1}^* \left(\sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot \hat{\mathcal{E}}_i^{j,k}(\xi) \right) (g_1, \dots, g_{2^{n+1}}) \\ &= \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot P_{n,n+1}^*(\hat{\mathcal{E}}_i^{j,k}(\xi))(g_1, \dots, g_{2^{n+1}}) \\ &= \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot (\hat{\mathcal{E}}_i^{j,k}(\xi))(g_1 g_2, \dots, g_{2^{n+1}-1} g_{2^{n+1}}). \end{aligned}$$

Let γ_i be a curve in G with $\dot{\gamma}_i(0) = \hat{e}_i$. Write

$$\mathcal{E}_i^{j,k} = (c_1 \mathcal{E}_i^1, c_2 \mathcal{E}_i^2, \dots, c_{2^n} \mathcal{E}_i^{2^n}).$$

Thus

$$\begin{aligned} & \hat{\mathcal{E}}_i^{j,k}(\xi)(g_1, \dots, g_{2^n}) \\ &= \frac{1}{2^n} \sum_l c_l \frac{d}{dt} \xi(g_1, \dots, g_{l-1}, g_l \cdots g_{2^n} \gamma_i(g_{l+1} \cdots g_{2^n})^{-1}, \\ & \quad g_{l+1}, \dots, g_{2^n}), \end{aligned}$$

and

$$\begin{aligned} & \hat{\mathcal{E}}_i^{j,k}(\xi)(g_1 g_2, \dots, g_{2^{n+1}-1} g_{2^{n+1}}) \\ &= \frac{1}{2^n} \frac{d}{dt} \sum_l c_l \xi(g_1 g_2, \dots, g_{2(l-1)-1} g_{2(l-1)}, g_{2l-1} \\ & \quad \cdots g_{2^{n+1}} \gamma_i(g_{2l+1} \cdots g_{2^{n+1}})^{-1}, g_{l+1}, \dots, g_{2^{n+1}}). \end{aligned}$$

On the other hand

$$\begin{aligned} & D_{n+1}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \\ &= \sum_{j \leq n+1, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot \hat{\mathcal{E}}_i^{j,k}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \\ &= \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot \hat{\mathcal{E}}_i^{j,k}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \\ & \quad + \sum_{j=n+1, k, i} a^{j,k} \mathcal{E}_i^{j,k} \cdot \hat{\mathcal{E}}_i^{j,k}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \end{aligned}$$

The terms in the last sum are

$$\begin{aligned} & \hat{\mathcal{E}}_i^{n+1,k}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \\ &= \frac{1}{2} \frac{d}{dt} (\xi(g_1 g_2, \dots, g_{2k-1} \cdots g_{2^{n+1}} \gamma_i(g_{2k} \cdots g_{2^{n+1}})^{-1} g_{2k}, \dots, g_{2^{n+1}-1} g_{2^{n+1}}) \\ & \quad - \xi(g_1 g_2, \dots, g_{2k-1} g_{2k} \cdots g_{2^{n+1}} \gamma_i(g_{2k+1} \cdots g_{2^{n+1}})^{-1}, \\ & \quad g_{2k+1}, \dots, g_{2^{n+1}-1} g_{2^{n+1}})) \\ &= 0 \end{aligned}$$

The terms in the first sum are

$$\begin{aligned}
& \hat{\mathcal{E}}_i^{j,k}(P_{n,n+1}^*(\xi))(g_1, \dots, g_{2^{n+1}}) \\
&= \frac{1}{2^{n+1}} \frac{d}{dt} \sum_l c_l(\xi(g_1 g_2, \dots, \\
&\quad g_{2l-1} \cdots g_{2^{n+1}} \gamma_i(g_{2l} \cdots g_{2^{n+1}})^{-1} g_{2l}, \dots, g_{2^{n+1}-1} g_{2^{n+1}}) \\
&\quad + \xi(g_1 g_2, \dots, g_{2l-1} g_{2l} \cdots g_{2^{n+1}} \gamma_i(g_{2l+1} \cdots g_{2^{n+1}})^{-1}, \dots, g_{2^{n+1}-1} g_{2^{n+1}})) \\
&= \frac{1}{2^n} \frac{d}{dt} \sum_l c_l \xi(g_1 g_2, \dots, g_{2(l-1)-1} g_{2(l-1)}, g_{2l-1} \\
&\quad \cdots g_{2^{n+1}} \gamma_i(g_{2l+1} \cdots g_{2^{n+1}})^{-1}, g_{l+1}, \dots, g_{2^{n+1}}).
\end{aligned}$$

This proves the compatibility for functions.

Because of the derivation property of the connection on the Clifford bundle it only remains to prove compatibility on vectors of the form $\mathcal{E}_i^{j,k}$, $j \leq n$. This follows from

$$\begin{aligned}
& P_{n,n+1}^*(D_n(\mathcal{E}_l^{m,p})) \\
&= P_{n,n+1}^*\left(\sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\hat{\mathcal{E}}_i^{j,k}}^n \mathcal{E}_l^{m,p}\right) \\
&= \sum_{j \leq n+1, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\hat{\mathcal{E}}_i^{j,k}}^{n+1}(\mathcal{E}_l^{m,p}) \\
&= D_{n+1}(\mathcal{E}_l^{m,p})
\end{aligned}$$

where we have used the admissibility condition for the connections. \triangleleft

3.5 Gauge invariance

In section 2.2.2 we have constructed a left action of $\bar{\mathcal{G}}^\Delta$ on $\bar{\mathcal{A}}^\Delta$. It follows from the construction that this action preserves the inner product on $L^2(\bar{\mathcal{A}}^\Delta)$, and hence is a unitary action of $\bar{\mathcal{G}}^\Delta$ on $L^2(\bar{\mathcal{A}}^\Delta)$. In order to talk about gauge invariance of the Dirac operator we first need to extend this action to a unitary action on $L^2(\bar{\mathcal{A}}^\Delta, Cl(\bar{\mathcal{A}}^\Delta))$.

Let U be a gauge transformation written as

$$U_n(g_1, \dots, g_{2^n}) = (u_0 g_1 u_1^{-1}, \dots, u_{2^n-1} g_{2^n} u_{2^n}^{-1}),$$

on \mathcal{A}_n , where $u_0, \dots, u_{2^n} \in G$. Since the metric is left and right invariant in each copy of G , we see that $(U_n)^* : T^* \mathcal{A}_n \rightarrow T^* \mathcal{A}_n$ preserves the metric and therefore

U extends to a unitary on $L^2(\overline{\mathcal{A}}^\Delta, Cl(\overline{\mathcal{A}}^\Delta))$. This unitary will also be denoted by U .

A vectorfield of the form

$$(g_1, \dots, g_{2^n}) \rightarrow (0, \dots, 0, L_{g_1 \dots g_{2^n}} R_{(g_{j+1} \dots g_{2^n})^{-1}} \hat{e}_i, 0, \dots, 0)$$

gets mapped to

$$(u_0 g_1 u_1^{-1}, \dots, u_{2^n-1} g_{2^n} u_{2^n}^{-1}) \rightarrow (0, \dots, 0, L_{u_{j-1} g_j \dots g_{2^n}} R_{(u_j^{-1} g_{j+1} \dots g_{2^n})^{-1}} \hat{e}_i, 0, \dots, 0),$$

under $(U_n)_*$. Written differently

$$(g_1, \dots, g_{2^n}) \rightarrow (0, \dots, 0, L_{g_j \dots g_{2^n}} R_{(g_{j+1} \dots g_{2^n})^{-1}} u_{2^n} \hat{e}_i u_{2^n}^{-1}, 0, \dots, 0).$$

Therefore $(U_n)^*$ is, on a covector field of the form

$$(g_1, \dots, g_{2^n}) \rightarrow (0, \dots, 0, L_{g_j \dots g_{2^n}} R_{(g_{j+1} \dots g_{2^n})^{-1}} e_i, 0, \dots, 0),$$

given by

$$(g_1, \dots, g_{2^n}) \rightarrow (0, \dots, 0, L_{g_j \dots g_{2^n}} R_{(g_{j+1} \dots g_{2^n})^{-1}} u_{2^n}^{-1} e_i u_{2^n}, 0, \dots, 0).$$

Put $f_i = u_{2^n} e_i u_{2^n}$. Since $\{f_i\}$ be another orthonormal basis for the dual of the Lie algebra of G , there exist a matrix $O_{ij} \in O(\dim(G))$ with

$$f_i = \sum_j O_{ij} e_j.$$

From the construction of the covector fields $\{\mathcal{E}_i^{j,k}\}$ it follows that

$$U(\mathcal{E}_i^{j,k}) = \sum_l O_{il} \mathcal{E}_l^{j,k}.$$

It turns out that the operator D is not always gauge invariant. We therefore need

Definition 3.5.1 *An admissible system of connections is gauge admissible if the conditions*

$$\sum_{r,i,q} O_{ql} O_{ir} \mathcal{E}_r^{j,k} U(\nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_q^{m,p}) = \sum_i \mathcal{E}_i^{j,k} \nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_l^{m,p}$$

hold for all n, j, k and all gauge transformations $U \in \overline{\mathcal{G}}^\Delta$.

Note that the system of trivial connections with respect to the trivializations given by $\{\mathcal{E}_i^{j,k}\}$ is a gauge admissible system of connections.

Proposition 3.5.2 *When D is constructed from a gauge admissible connection, D is gauge invariant, i.e. $D = UDU^*$ for all $U \in \overline{\mathcal{G}}^\Delta$.*

Proof. Let $\xi \in L^2(\mathcal{A}_n)$. We compute

$$\begin{aligned}
UDU^*(\xi) &= \sum_{j \leq n, k, i} a^{j,k} (U_n)^* (\mathcal{E}_i^{j,k}) d_{(U_n^{-1})^* (\mathcal{E}_i^{j,k})} \xi \\
&= \sum_{j \leq n, k} a^{j,k} \sum_{i, l, m} O_{il} \mathcal{E}_l^{j,k} d_{O_{im} \mathcal{E}_m^{j,k}} \xi \\
&= \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} d_{\mathcal{E}_i^{j,k}} \xi \\
&= D(\xi)
\end{aligned}$$

Due to the derivation and product structures we now only need to check gauge invariance on the covectorfields $\{\mathcal{E}_i^{j,k}\}$. Here we get

$$\begin{aligned}
UDU^*(\mathcal{E}_l^{m,p}) &= U \left(\sum_{j \leq n, k, i, q} a^{j,k} \mathcal{E}_i^{j,k} O_{ql} \nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_q^{m,p} \right) \\
&= \sum_{j \leq n, k, i, q} a^{j,k} U(\mathcal{E}_i^{j,k} O_{ql}) U(\nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_q^{m,p}) \\
&= \sum_{j \leq n, k} a^{j,k} \left(\sum_{r, i, q} O_{ql} O_{ir} \mathcal{E}_r^{j,k} U(\nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_q^{m,p}) \right) \\
&= \sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\mathcal{E}_i^{j,k}}^n \mathcal{E}_l^{m,p} \\
&= D(\mathcal{E}_l^{m,p}),
\end{aligned}$$

where we have used gauge admissibility of ∇^n . \triangleleft

4 A semifinite spectral triple

In this section we will construct a semifinite spectral associated to $\overline{\mathcal{A}}^\Delta$ or rather to the algebra of holonomy loops, see definition 4.1.1. The following definition

first appeared [6].

Definition 4.0.1 Let \mathcal{N} be a semifinite von Neumann algebra with a semifinite trace τ . Let \mathcal{K}_τ be the τ -compact operators. A semifinite spectral triple $(\mathcal{B}, \mathcal{H}, D)$ is a $*$ -subalgebra \mathcal{B} of \mathcal{N} , a representation of \mathcal{N} on the Hilbert space \mathcal{H} and an unbounded densely defined self adjoint operator D on \mathcal{H} affiliated with \mathcal{N} satisfying

1. $b(\lambda - D)^{-1} \in \mathcal{K}_\tau$ for all $b \in \mathcal{B}$ and $\lambda \notin \mathbb{R}$.
2. $[b, D]$ is densely defined and extends to a bounded operator.

4.1 The algebra

Let v be a vertex in \mathcal{S}_Δ . Denote by $\mathcal{HG}_v(S_\Delta)$ the subgroupoid of $\mathcal{HG}(S_\Delta)$ of loops based in v , i.e. paths starting and ending in v . Let $G \rightarrow M_N$ be a unitary matrix representation of G and let $\mathcal{H}' = \mathcal{H} \otimes M_N$. Hence

$$\mathcal{H}' = \varinjlim (\mathcal{H}_n \otimes M_N).$$

Definition 4.1.1 Let L be a loop in $\mathcal{HG}_v(S_\Delta)$. Let $\xi \in \mathcal{H}_n \otimes M_N$, where n is such that $L \in \mathcal{HG}(n)$. Define

$$L(\xi)(\nabla) = \nabla(L)\xi(\nabla), \quad \nabla \in \mathcal{A}_n = \text{Hom}(\mathcal{HG}(n), G).$$

Since

$$P_{n,m}^*(L(\xi)) = L(P_{n,m}^*(\xi))$$

we get a densely defined operator on \mathcal{H}' , also denoted L . Clearly L is bounded.

Denote by \mathcal{B}_v the $*$ -algebra generated by $\{L\}_{L \in \mathcal{HG}_v(S_\Delta)}$. We will call \mathcal{B}_v the algebra of holonomy loops.

Proposition 4.1.2 For $L \in \mathcal{HG}_v(S_\Delta)$ define the function $HL : \mathcal{A} \rightarrow M_N$ by

$$HL(\nabla) = \text{Hol}(L, \nabla).$$

The $*$ -algebra generated by $\{HL\}_{L \in \mathcal{HG}_v(S_\Delta)}$ equipped with the sup norm is isomorphic to \mathcal{B}_v as normed $*$ -algebra.

Proof. Follows from the dense embedding $\mathcal{A} \rightarrow \overline{\mathcal{A}}^\Delta$. \triangleleft

4.2 The trace and the von Neumann algebra

Let

$$A_n = \mathcal{K}(L^2(\mathcal{A}_n)) \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes End(M_N).$$

where $\mathcal{K}(L^2(\mathcal{A}_n))$ denotes the compact operators on $L^2(\mathcal{A}_n)$. We define maps, with an abuse of notation also denoted $P_{n,n+1}^*$, from A_n to A_{n+1} in the following way

- On $End(M_N)$, $P_{n,n+1}^*$ is the identity.
- On $\mathcal{K}(L^2(\mathcal{A}_n))$, $P_{n,n+1}^*$ is the map induced by the embedding

$$P_{n,n+1}^* : L^2(\mathcal{A}_n) \rightarrow L^2(\mathcal{A}_{n+1}),$$

i.e. if $P_{n,n+1}$ denotes the projection in $L^2(\mathcal{A}_{n+1})$ onto $P_{n,n+1}^*(L^2(\mathcal{A}_n))$ we have

$$\mathcal{K}(L^2(\mathcal{A}_n)) \ni a \rightarrow aP_{n,n+1} \in \mathcal{K}(L^2(\mathcal{A}_{n+1})).$$

- We use the map

$$P_{n,n+1} : Cl(T_{id}^* \mathcal{A}_n) \rightarrow Cl(T_{id}^* \mathcal{A}_{n+1}),$$

which has already been defined in subsection 3.2 and restrict it to the identity. If we write

$$T_{id}^* \mathcal{A}_{n+1} = T_{id}^* \mathcal{A}_n \oplus V_{n,n+1},$$

and hence

$$Cl(T_{id}^* \mathcal{A}_{n+1}) = Cl(T_{id}^* \mathcal{A}_n) \hat{\otimes} Cl(V_{n,n+1}),$$

then

$$P_{n,n+1}^* : Cl(T_{id}^* \mathcal{A}_n) \rightarrow Cl(T_{id}^* \mathcal{A}_{n+1})$$

is given by

$$P_{n,n+1}^*(\mathcal{E}) = \mathcal{E} \otimes \mathbf{1}_{Cl(V_{n,n+1})}.$$

Note that $P_{n,n+1}^*$ is a C^* -algebra homomorphism. Therefore $\{A_n\}$ is an inductive system of C^* -algebras. Let

$$A = \lim A_n$$

be the inductive limit. By

$$P_{n,\infty}^* : A_n \rightarrow A,$$

we denote the induced embeddings.

From the construction of A it follows that

$$A = \mathcal{K}(L^2(\mathcal{A})) \otimes B \otimes M_N, \quad (4)$$

where B is a UHF-algebra. Since the dimension of the Clifford algebra is a power of 2 when $n \geq 1$, B is the CAR-algebra.

There is a trace Tr on A_k defined by

$$Tr = Tr_o \otimes Tr_n \otimes Tr_o,$$

where Tr_o are the operator traces on $\mathcal{K}(L^2(\mathcal{A}_k))$, resp. $End(M_N)$ and Tr_n is the normalized trace on $Cl(T_{id}^* \mathcal{A}_k)$. By construction

$$Tr(a) = Tr(P_{k,k+1}^*(a)),$$

and hence defines a densely defined trace on A .

Using the trivialization $\{\mathcal{E}_i^{j,k}\}$ of $T_{id}^* \mathcal{A}_n$ we see that A acts on \mathcal{H}' . In fact \mathcal{H}' factors like (4) into

$$\mathcal{H}' = L^2(\mathcal{A}) \otimes Cl(T_{id}^* \mathcal{A}) \otimes M_N.$$

Note that the action of B on $Cl(T_{id}^* \mathcal{A})$ is just the GNS-representation of B with respect to the normalized trace on B .

Definition 4.2.1 *Let \mathcal{N} be the weak closure of A in $\mathcal{B}(\mathcal{H}')$. The trace $\tau : \mathcal{N} \rightarrow \mathbb{C}$ is defined as the extension of the trace Tr on A to \mathcal{N} .*

Note that τ is a semifinite trace, since it is the tensor product of the usual semifinite trace on $\mathcal{B}(\mathcal{H})$, \mathcal{H} separable, and the finite trace on the hyperfinite II_1 factor \overline{B}^w .

4.3 A coordinate change and selfadjointness of the Dirac operator

We define the coordinate transformation

$$\Theta_n : \mathcal{A}_n = G^{2^n} \rightarrow G^{2^n}$$

by

$$\Theta_n(g_1, \dots, g_{2^n}) = (g_1 g_2 \cdots g_{2^n}, g_2 g_3 \cdots g_{2^n}, \dots, g_{2^n-1} g_{2^n}, g_{2^n}).$$

It is easy to see that Θ_n preserves the Haar measure on G^{2^n} .

The inverse of θ_n is given by

$$\Theta_n^{-1}(g_1, \dots, g_{2^n}) = (g_1 g_2^{-1}, g_2 g_3^{-1}, \dots, g_{2^n-1} g_{2^n}^{-1}, g_{2^n}),$$

The main purpose of Θ_n is the following:

\mathcal{A}_n is a trivial $G^{2^{n-1}}$ -principal fiber bundle over \mathcal{A}_{n-1} , where the action of $G^{2^{n-1}}$ on \mathcal{A}_n is given by

$$(g_1, \dots, g_{2^{n-1}})(g'_1, \dots, g'_{2^n}) = (g'_1 g_1^{-1}, g_1 g'_2, \dots, g'_{2^{n-1}} g_{2^{n-1}}^{-1}, g_{2^{n-1}} g'_{2^n}).$$

Combining this with Θ_n we get the following commutative diagram

$$\begin{array}{ccccc} & & \vdots & & \vdots \\ G^{2^n} & \rightarrow & \mathcal{A}_n & \xrightarrow{\Theta_n} & G^{2^n} \\ & & \downarrow & & \downarrow pr_o \\ G^{2^{n-1}} & \rightarrow & \mathcal{A}_{n-1} & \xrightarrow{\Theta_{n-1}} & G^{2^{n-1}} \\ & & \vdots & & \vdots \\ G & \rightarrow & \mathcal{A}_1 & \xrightarrow{\Theta_1} & G^2 \\ & & \downarrow & & \downarrow pr_o \\ & & \mathcal{A}_0 & = & G \end{array}$$

where pr_o means projection onto the odd coordinates, i.e.

$$pr_o(g_1, g_2, \dots, g_{2^{n-1}}, g_{2^n}) = (g_1, g_3, \dots, g_{2^n-1}).$$

In other words $\{\Theta_n\}$ is just a consistent way of trivializing the principal bundles.

Lemma 4.3.1 *Let $\hat{E}_i(g) = L_g(\hat{e}_i)$, the left translate of \hat{e}_i . On \mathcal{A}_n we have*

$$((\Theta_n)_*(\hat{\mathcal{E}}_i^{j,k}))(g_1, \dots, g_{2^n}) = -\frac{1}{2}(0, \dots, 0, \hat{E}_i(g_{2^k}), 0, \dots, 0),$$

and if we write

$$\hat{\mathcal{E}}_i^{j,k} = (c_1 \hat{\mathcal{E}}_i^1, c_1 \hat{\mathcal{E}}_i^2, c_2 \hat{\mathcal{E}}_i^3, c_2 \hat{\mathcal{E}}_i^4, \dots, c_{2^{n-1}} \hat{\mathcal{E}}_i^{2^n-1}, c_{2^{n-1}} \hat{\mathcal{E}}_i^{2^n}), \quad j < n,$$

then

$$\begin{aligned} ((\Theta_n)_*(\hat{\mathcal{E}}_i^{j,k}))(g_1, \dots, g_{2^n}) = & \\ & \left(2 \sum_{j=1} c_j \hat{E}_i(g_1), (c_1 + 2 \sum_{j=2} c_j) \hat{E}_i(g_2), 2 \sum_{j=2} c_j \hat{E}_i(g_3), \dots, \right. \\ & \left. 2c_{2^{n-1}} \hat{E}_i(g_{2^n-1}), c_{2^{n-1}} \hat{E}_i(g_{2^n}) \right) \end{aligned}$$

Proof. Straightforward computation. \triangleleft

We will also write $\hat{E}_i^j(g_1, \dots, g_{2^n}) = (0, \dots, \hat{E}_i(g_j), \dots, 0)$.

Proposition 4.3.2 *Let $\{\nabla^n\}$ be an admissible system of connections. Assume that D_0 is self adjoint,*

$$\nabla_{\mathcal{E}_i^{j,k}}^n(\mathcal{E}_l^{n,m}) = 0 \text{ for all } i, j, k, m, l, \text{ and all } n \geq 1$$

and that $a^{j,k}$ are real and non zero for all j, k . Then D is self adjoint.

Proof. We will prove that D_n is formally self adjoint for each n . From this the statement follows because D_n is an elliptic pseudo differential operator on a compact manifold and therefore by elliptic regularity D_n is self adjoint. We can even find an orthonormal basis for \mathcal{H}'_n which diagonalizes D_n with real eigenvalues. We can therefore find an orthonormal basis for \mathcal{H}' which diagonalizes D with real eigenvalues. Hence D is self adjoint.

Write

$$D_n = \sum_i a^{0,1} \mathcal{E}_i^{0,1} \nabla_{\mathcal{E}_i^{0,1}}^n + \sum_{1 \leq j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\mathcal{E}_i^{j,k}}^n.$$

By the assumptions on the connection and lemma 4.3.1 the first summand on the right hand side is a linear combination of D_0 's acting on the different copies of G , and thus by assumption self adjoint. The second summand is, in the trivialization induced by $\{\mathcal{E}_i^{j,k}\}$ and transported by Θ_n on the form

$$\sum_{1 \leq j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} d_{(\Theta_n)_*(\mathcal{E}_i^{j,k})},$$

where d is the exterior derivative. Since $(\Theta_n)_*(\mathcal{E}_i^{j,k})$ is a left invariant vector field according to lemma 4.3.1, $d_{(\Theta_n)_*(\mathcal{E}_i^{j,k})}$ is formally skew self adjoint. The formal selfadjointness now follows since Clifford multiplication with $\mathcal{E}_i^{j,k}$ is also skew self adjoint and commutes with $d_{(\Theta_n)_*(\mathcal{E}_i^{j,k})}$. \triangleleft

4.4 Affiliation of D

The spectral projections of D_n will by construction belong to

$$\mathcal{K}(L^2(\mathcal{A}_n)) \otimes \text{End}(Cl(T_{id}^* \mathcal{A}_n)) \otimes \text{End}(M_N).$$

Since we can split $Cl(T_{id}^* \mathcal{A}_n)$ into irreducible representations of $Cl(T_{id}^* \mathcal{A}_n)$, and D_n acts on each of these, the spectral projections of D_n is in

$$A_n = \mathcal{K}(L^2(\mathcal{A}_n)) \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes End(M_N).$$

Recall from section 4.2 the splitting

$$T_{id}^* \mathcal{A}_{n+1} = T_{id}^* \mathcal{A}_n \oplus V_{n,n+1}.$$

More generally we can write

$$T_{id}^* \mathcal{A}_m = T_{id}^* \mathcal{A}_n \oplus V_{n,m},$$

and we thus have

$$Cl(T_{id}^* G^m) = Cl(V_{n,m}) \hat{\otimes} Cl(T_{id}^* \mathcal{A}_n). \quad (5)$$

We assume that the system of connections defining the Dirac operator satisfies the properties in 4.3.2. These properties ensure the following equation:

$$D_m(P_{n,m}^*(\xi) \otimes v) = P_{n,m}^*(D_n(\xi)) \otimes v, \quad \xi \in \mathcal{H}'_n, v \in Cl(V_{n,m}).$$

If ξ is an eigenvector with eigenvalue λ , then $P_{n,m}^*(\xi) \otimes v$ is an eigenvector with eigenvalue λ if $v \in Cl(V_{n,m})$. Therefore, if $P_{\lambda,n}$ is the spectral of eigenvalue λ of D_n the projection $P_{n,\infty}^*(P_{\lambda,n})$ is a subprojection of the spectral projection of eigenvalue λ of D . By construction $P_{n,\infty}^*(P_{\lambda,n}) \in \mathcal{N}$

◁

Proposition 4.4.1 *The spectral projections of the Dirac operator is contained in \mathcal{N} .*

Proof. For a spectral projection P_λ of eigenvalue λ of D we have

$$P_{n,\infty}^*(P_{\lambda,n}) \nearrow P_\lambda.$$

Hence P_λ is in the weak closure of A , i.e. $P_\lambda \in \mathcal{N}$.

◁

Corollary 4.4.2 *The operator D is affiliated with \mathcal{N} .*

4.5 The main theorem

We begin by enlarging our Hilbert space slightly. Let $\mathcal{H} = \mathcal{H}' \otimes Cl(1)$. We also enlarge \mathcal{N} by tensoring with $Cl(1)$. By abuse of notation we will also call the enlargement \mathcal{N} . The trace τ is also enlarged by tensoring with the normalized trace on $Cl(1)$ and denoted by τ .

The Dirac operator extends to an operator on \mathcal{H} . We write

$$D_n = \sum_{j < n, k, i} a^{j,k} \mathcal{E}_i^{j,k} \nabla_{\hat{\mathcal{E}}_i^{j,k}}^n + \sum_{k, i} a^{n,k} \mathcal{E}_i^{n,k} \nabla_{\hat{\mathcal{E}}_i^{n,k}}^n = D_n^+ + D_n^-.$$

After changing coordinates with Θ_n we have

$$D_n^- = \sum_{k, i} a^{n,k} \mathcal{E}_i^{n,k} \nabla_{\hat{\mathcal{E}}_i^{2k}}^n.$$

Therefore, when we use Θ_n to trivialize

$$\mathcal{A}_n = \mathcal{A}_{n-1} \times G^{2^{n-1}}$$

we see that D_n^- is an operator acting only in the fiber, i.e. in the $G^{2^{n-1}}$ part of \mathcal{A}_n . There is an embedding of

$$L^2(\mathcal{A}_{n-1}) \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes M_N \otimes Cl(1) \quad (6)$$

in $Ker(D_n^-)$ by identifying it as a subspace of \mathcal{H}_n . Let P_n be the projection onto the orthogonal complement of (6) in $Ker(D_n^-)$. Define

$$D_{n,p} = D_n + \sum_j b^j e P_j,$$

where $b_j \in i\mathbb{R}$, e is the generator of $Cl(1)$ and where we have used $P_{j,j+1}^*$ to push forward P_j . Note that $D_{n,p}$ is self adjoint. By construction

$$P_{n,n+1}^*(D_{n,p}(\xi)) = D_{n+1,p}(\xi),$$

and therefore $\{D_{n,p}\}$ defines a densely defined self adjoint operator D_p on \mathcal{H} . Since D is affiliated with \mathcal{N} by corollary 4.4.2, so is D_p by construction.

Theorem 4.5.1 *There exist sequences $\{a^{j,k}\}$ and $\{b_k\}$ such that $(\mathcal{B}_v, \mathcal{H}, D_p)$ is a semifinite spectral triple with respect to (\mathcal{N}, τ) .*

Proof. The selfadjointness and affiliation to \mathcal{N} of D_p are already taken care of. What remains is to prove that $[b, D_p] \in \mathcal{B}(\mathcal{H})$ and that $b(\lambda - D_p)^{-1} \in \mathcal{K}_\tau$ for all $b \in \mathcal{B}_v$.

The boundedness of the commutator: Let L be a loop in Γ_n . If we write

$$\mathcal{H}_n = L^2(G^{2^n}) \otimes Cl(T^* \mathcal{A}_n) \otimes M_N \otimes Cl(1)$$

the loop operator L acts by point wise matrix multiplication over G^{2^n} in the M_N factor, i.e. the matrix entries is of the form $f(g_1, \dots, g_{2^n})$. The action of L on \mathcal{H}_{n+1} is then matrix multiplication with entries of the form $f(g_1 g_2, \dots, g_{2^{n+1}-1} g_{2^{n+1}})$. Conjugating the operator with Θ_{n+1} we see, since

$$\Theta_{n+1}^{-1}(g_1, \dots, g_{2^{n+1}}) = (g_1 g_2^{-1}, g_2 g_3^{-1}, \dots, g_{2^n-1} g_{2^{n+1}}^{-1}, g_{2^{n+1}}),$$

that the result is independent of $g_2, g_4, \dots, g_{2^{n+1}}$, and thus $[L, D_{n+1}^-] = 0$. It therefore follows that $[L, P_{n+1}] = 0$. Hence $[L, D_{n+1,p}] = [L, D_{n,p}]$ and therefore $[L, D_p]$ is bounded.

To prove that there exist sequences $\{a^{j,k}\}$ and b_j such D_p has τ -compact resolvent we will prove that for any real sequence c_n converging to ∞ we can choose $a^{n,k}$ and b_n such that the new eigenvalues, modulo the extra multiplicity of the existing eigenvalue due to the growth of the Clifford bundle, introduced by going from $(D_{n-1,p})^2$ to $(D_{n,p})^2$ are bigger than c_n .

In the following we will omit the M_N part. This will play no role, since the Dirac operator does not act on the M_N part. First we rewrite the operator in the following way

$$D_{n,p} = \left(D_n^+ + \sum_{j \leq n-1} b_j e P_j \right) + (D_n^- + b_n e P_n) = D_{n,p}^+ + (D_n^- + b_n e P_n),$$

and

$$\begin{aligned} & (D_n^- + b_n e P_n) \\ &= \sum_{k,i} a^{n,k} \mathcal{E}_i^{n,k} \nabla_{\mathcal{E}_i^{n,k}}^n + b_n e P_n \\ &= a \left(\sum_{k,i} \frac{a^{n,k}}{a} \mathcal{E}_i^{n,k} \nabla_{\mathcal{E}_i^{n,k}}^n + \frac{b_n}{a} e P_n \right) =: a D_{n,p}^-. \end{aligned}$$

Using the coordinate change Θ_n we factorize

$$\begin{aligned} \mathcal{H}_n &= L^2(G^{2^{n-1}}) \otimes L^2(G^{2^{n-1}}) \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1) \\ &= \mathcal{H}^1 \otimes \mathcal{H}^2 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1), \end{aligned}$$

where $\mathcal{H}^1 = L^2(G^{2^{n-1}})$ corresponds to the coordinates $(g_1, g_3, \dots, g_{2^n-1})$ under Θ_n and \mathcal{H}^2 corresponds to the coordinates $(g_2, g_4, \dots, g_{2^n})$. In particular by lemma 4.3.1, $D_{n,p}^-$ acts trivially on \mathcal{H}^1 . Taking the square of $D_{n,p}$ we get

$$(D_{n,p})^2 = (D_{n,p}^+)^2 + a \{D_{n,p}^-, D_{n,p}^+\} + a^2 (D_{n,p}^-)^2. \quad (7)$$

Using lemma 4.3.1 it is easy to see that $\{D_{n,p}^-, D_{n,p}^+\}$ does not act on \mathcal{H}^1 .

Let ξ belong to the orthogonal complement of $\mathcal{H}^1 \otimes 1 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1)$. We know, due to the fact that D_p is self adjoint and commutes with the maps $P_{k,k+1}^*$ that this complement is an invariant subspace for D_p . Decompose ξ with respect to the above decomposition of \mathcal{H}_n into

$$\xi = \sum_k \xi_k^1 \otimes \xi_k^2, \quad (8)$$

where $\{\xi_k^1\}$ is an orthonormal basis for \mathcal{H}^1 and ξ_k^2 belongs to the orthogonal complement of

$$1 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1)$$

in

$$\mathcal{H}^2 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1).$$

Combining (7) and (8) we get

$$\begin{aligned} & \langle (D_{n,p})^2 \xi, \xi \rangle \\ &= \langle (D_{n,p}^+)^2 \xi, \xi \rangle + a \langle \{D_{n,p}^+, D_{n,p}^-\} \xi, \xi \rangle + a^2 \langle (D_{n,p}^-)^2 \xi, \xi \rangle \\ &\geq \langle \{D_{n,p}^+, D_{n,p}^-\} \xi, \xi \rangle + a^2 \langle (D_{n,p}^-)^2 \xi, \xi \rangle \\ &= \sum_k \langle \xi_k^1, \xi_k^1 \rangle \left(a \langle \{D_{n,p}^+, D_{n,p}^-\} \xi_k^2, \xi_k^2 \rangle + a^2 \langle (D_{n,p}^-)^2 \xi_k^2, \xi_k^2 \rangle \right) \end{aligned}$$

Since $(D_{n,p}^-)^2$ is an elliptic second order operator on $\mathcal{H}^2 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1)$ and the operator $\{D_{n,p}^+, D_{n,p}^-\}$ is a first order operator on $\mathcal{H}^2 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1)$, the operator

$$A = (D_{n,p}^-)^{-1} \{D_{n,p}^+, D_{n,p}^-\} (D_{n,p}^-)^{-1}$$

is bounded. We thus get

$$\begin{aligned} & \langle (D_{n,p})^2 \xi, \xi \rangle \\ &\geq \sum_k \langle \xi_k^1, \xi_k^1 \rangle \left(a \langle \{D_{n,p}^+, D_{n,p}^-\} \xi_k^2, \xi_k^2 \rangle + a^2 \langle (D_{n,p}^-)^2 \xi_k^2, \xi_k^2 \rangle \right) \\ &= \sum_k \langle \xi_k^1, \xi_k^1 \rangle (a \langle A D_{n,p}^- \xi_k^2, D_{n,p}^- \xi_k^2 \rangle + a^2 \langle (D_{n,p}^-)^2 \xi_k^2, \xi_k^2 \rangle) \\ &\geq \sum_k \langle \xi_k^1, \xi_k^1 \rangle \left(-a \|A\| \langle D_{n,p}^- \xi_k^2, D_{n,p}^- \xi_k^2 \rangle + a^2 \langle (D_{n,p}^-)^2 \xi_k^2, \xi_k^2 \rangle \right) \\ &\geq \sum_k \langle \xi_k^1, \xi_k^1 \rangle \lambda (a^2 - a \|A\|) \langle \xi_k^2, \xi_k^2 \rangle \\ &= \lambda (a^2 - a \|A\|) \|\xi\|^2 \end{aligned}$$

where λ is the lowest eigenvalue of $(D_{n,p}^-)^2$ on the complement of

$$\mathcal{H}^1 \otimes 1 \otimes Cl(T_{id}^* \mathcal{A}_n) \otimes Cl(1).$$

Hence by choosing a big enough we have $\langle (D_{n,p})^2 \xi, \xi \rangle \geq c_n \|\xi\|$. \triangleleft

The sequence $\{b_n\}$ is needed in the case when the operator

$$D_e = \sum_i E_i d_{\hat{E}_i},$$

where \hat{E}_i is an orthonormal frame of left translated vectorfields on one copy of G , has a non trivial kernel in the sense that the kernel is given by the span of $\{E_i\}_i$. This is clearly not the case for $U(1)$. In the appendix we will show that this is also not the case for $SU(2)$.

Proposition 4.5.2 *When D_b is constructed from a gauge admissible system of connections satisfying the demands in proposition 4.3.2, D_b is gauge invariant, i.e. $D_b = UD_bU^*$ for all $U \in \overline{\mathcal{G}}^\Delta$. (U acts trivially on the $Cl(1)$ -part.)*

Proof. From the proof of proposition 3.5.2 it follows that D_n^- is gauge invariant, i.e. invariant under U_n for all $U_n \in \mathcal{G}_n$. In particular the kernel of D_n^- is gauge invariant. Also the space (6) is invariant under \mathcal{G}_n and therefore the projection P_n onto the orthogonal complement of (6) in the kernel of D_n^- is gauge invariant under \mathcal{G}_n . From this the invariance follows. \triangleleft

Note that the system of trivial connections with respect to the trivializations given by $\{\mathcal{E}_i^{j,k}\}$ fulfills the demands of proposition 4.5.2.

5 Appendix

In this appendix we will first demonstrate the case of $U(1)$ to show what kind of growth conditions are needed on $a^{j,k}$. Secondly we show that for $SU(2)$ the perturbation with the P_n 's is not needed to obtain a semifinite spectral triple.

5.1 The $U(1)$ -case

We write $U(1) = \{e^{2\pi i\theta} | \theta \in [0, 1]\}$ and choose the metric such that

$$\langle \frac{d}{d\theta}, \frac{d}{d\theta} \rangle = 1.$$

The system of connections we will use is the system of trivial connections. The operator D_n has the form

$$\sum_{j \leq n, k, i} a^{j,k} \mathcal{E}_i^{j,k} d_{\mathcal{E}_i^{j,k}}$$

Since i can be only one in this formula, we will simply omit it. Also we will assume that $a^{j,k_1} = a^{j,k_2}$ for all k_1, k_2 , and simply denote $a_j := a^{j,k}$. All $\mathcal{E}^{j,k}$ commute. Hence

$$D_n^2 = - \sum_{j \leq n, k} a_j^2 (d_{\mathcal{E}^{j,k}})^2.$$

Since D_n^2 acts trivially in the Clifford bundle, and since the identity on the Clifford bundle is normalized to have trace 1, we will omit the Clifford bundle in the rest of this computation.

We will use the coordinate change Θ_n to rewrite D_n^2 . The rewritten operator will be denoted \tilde{D}_n^2 . According to lemma 4.3.1 the vector fields

$$\hat{\mathcal{E}}^{j,k} = c_1^{j,k} \partial_{\theta_1} + c_1^{j,k} \partial_{\theta_2} + c_2^{j,k} \partial_{\theta_3} + c_2^{j,k} \partial_{\theta_4} + \dots + c_{2^{n-1}}^{j,k} \partial_{\theta_{2^{n-1}}} + c_{2^{n-1}}^{j,k} \partial_{\theta_{2^n}}, \quad j < n,$$

gets mapped to

$$2 \sum_{l=1} c_l^{j,k} \partial_{\theta_1} + (c_1 + 2 \sum_{l=2} c_l^{j,k}) \partial_{\theta_2} + 2 \sum_{l=2} c_l^{j,k} \partial_{\theta_3} + \dots + 2 c_{2^{n-1}}^{j,k} \partial_{\theta_{2^{n-1}}} + c_{2^{n-1}}^{j,k} \partial_{\theta_{2^n}}$$

under Θ_n , and $\hat{\mathcal{E}}^{n,k}$ gets mapped to $-\frac{1}{2} \partial_{\theta_{2^k}}$. It follows from the construction that

$$\sum_{j < n, k} |c_{2^{n-1}}^{j,k}|^2 < 1.$$

In particular for a function

$$\xi(\theta_1, \dots, \theta_{2^n}) = e^{2\pi i \theta_{2^n} l}, \quad l \in \mathbb{Z}$$

we see that

$$\tilde{D}_n^2(\xi) = 4\pi^2 l^2 \left(\sum_{j < n, k} (c_{2^{n-1}}^{j,k} a_j)^2 + \frac{a_n^2}{4} \right) \xi,$$

and

$$\sum_{j < n, k} (c_{2^{n-1}}^{j,k} a_j)^2 + \frac{a_n^2}{4} \leq \max\{a_j^2\}_{j < n} + \frac{a_n^2}{4}.$$

From this follows, that D only has finite many eigenvalues with finite multiplicity module the semifinite trace in a bounded set of \mathbb{R} if

$$a_n \rightarrow \infty. \quad (9)$$

This is therefore a necessary condition for D to have τ -compact resolvent.

On the other hand the eigenfunctions for \tilde{D}_n^2 are on the form

$$\exp(2\pi i(\sum_{l \leq 2^n} n_l \theta_l)).$$

The eigenfunctions for \tilde{D}_{n-1}^2 are the functions on the form

$$\exp(2\pi i(\sum_{l \leq 2^{n-1}} n_{2l-1} \theta_l)).$$

Hence the smallest new eigenvalue appearing from going to from $n-1$ to n is bigger than a_n^2 . Hence D has τ -compact resolvent if and only if condition (9) is satisfied.

5.2 The $SU(2)$ -case

We will show that the Dirac operator on $SU(2)$ on the form

$$D_e = \sum_i E_i d_{\hat{E}_i},$$

where d is understood with respect to the trivialization given by $\{E_i\}$, has a trivial kernel in the sense that the kernel is given by the span of $\{E_i\}_i$. In other words the kernel consists of the constants with respect to the trivialization of the Clifford bundle given by $\{E_i\}_i$. According to the remark after the proof of 4.5.1 this ensures that the perturbation with the P_n 's is not needed.

For the analysis of we will use that states $|jm\rangle$, where $j \in \{0, \frac{1}{2}, 1, \dots\}$ and $m \in \{-j, -j+1, \dots, j-1, j\}$ form an orthonormal basis for $L^2(SU(2))$. Choose a basis $\hat{e}_1, \hat{e}_2, \hat{e}_3$ for $\mathfrak{su}(2)$ such that

$$[\hat{e}_i, \hat{e}_j] = \sum_k \varepsilon_{ijk} \hat{e}_k.$$

Define a metric on $\mathfrak{su}(2)$ by letting $\hat{e}_1, \hat{e}_2, \hat{e}_3$ be an orthonormal basis. Note this metric by left and right translation define a left and right invariant metric on $SU(2)$.

The action of the corresponding left translated vectorfields of $\hat{e}_1, \hat{e}_2, \hat{e}_3$ is best described by forming the raising and lowering operators

$$\hat{e}_\pm = \hat{e}_1 \pm i\hat{e}_2 .$$

The action of the translated vectorfields is given by

$$d_{\hat{E}_3}|jm\rangle = -im|jm\rangle, \quad d_{\hat{E}_\pm}|jm\rangle = -ic_\pm(m)|jm \pm 1\rangle,$$

where

$$c_\pm(m) = \sqrt{(j \mp m)(j \pm m + 1)}.$$

We will restrict the operator D_e to acting in one of the two irreducible representations of $Cl(T_{id}^*SU(2))$ instead of the full Clifford bundle. We chose the representation in M_2 given by

$$E_1 = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \quad E_3 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix},$$

The Dirac operator therefore has the form

$$D_e = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \cdot d_{\hat{E}_1} + \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix} \cdot d_{\hat{E}_2} + \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \cdot d_{\hat{E}_3}$$

which is rewritten to

$$D_e = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \cdot d_{\hat{E}_+} + \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} \cdot d_{\hat{E}_-} + \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \cdot d_{\hat{E}_3}$$

The square of D_e is calculated

$$\begin{aligned} D_e^2 &= - \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \cdot d_{\hat{E}_+} d_{\hat{E}_-} - \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \cdot d_{\hat{E}_-} d_{\hat{E}_+} - \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \cdot d_{\hat{E}_3} d_{\hat{E}_3} \\ &\quad + \begin{pmatrix} 0 & i \\ 0 & 0 \end{pmatrix} \cdot d_{[\hat{E}_3, \hat{E}_+]} + \begin{pmatrix} 0 & 0 \\ i & 0 \end{pmatrix} \cdot d_{[\hat{E}_3, \hat{E}_-]} \end{aligned}$$

Using how \hat{E}_\pm and \hat{E}_3 act we get

$$\begin{aligned}
D_e^2 \begin{pmatrix} a|jm\rangle \\ b|jm\rangle \end{pmatrix} &= \begin{pmatrix} a \\ 0 \end{pmatrix} \cdot (c_-)^2(m)|jm\rangle + \begin{pmatrix} 0 \\ b \end{pmatrix} \cdot (c_+)^2(m)|jm\rangle \\
&+ \begin{pmatrix} -ib \\ 0 \end{pmatrix} \cdot c_+(m)|jm+1\rangle + \begin{pmatrix} 0 \\ ia \end{pmatrix} \cdot c_-(m)|jm-1\rangle \\
&+ \begin{pmatrix} a \\ b \end{pmatrix} \cdot m^2|jm\rangle \\
&= \begin{pmatrix} a \\ b \end{pmatrix} \cdot j(j+1)|jm\rangle + \begin{pmatrix} a \\ -b \end{pmatrix} \cdot m|jm\rangle \\
&+ \begin{pmatrix} -ib \\ 0 \end{pmatrix} \cdot c_+(m)|jm+1\rangle + \begin{pmatrix} 0 \\ ia \end{pmatrix} \cdot c_-(m)|jm-1\rangle
\end{aligned}$$

For a given j the following subspaces

$$\begin{pmatrix} 0 \\ b \end{pmatrix} \cdot |jj\rangle \quad \begin{pmatrix} a \\ 0 \end{pmatrix} \cdot |j-j\rangle \quad \begin{pmatrix} a \\ 0 \end{pmatrix} \cdot |jm\rangle + \begin{pmatrix} 0 \\ b \end{pmatrix} \cdot |jm-1\rangle$$

are invariant under D_e^2 . Therefore analyzing D_e^2 is the same as analyzing matrices of the form

$$\begin{pmatrix} j(j+1) + m & -ic_-(m) \\ ic_-(m) & j(j+1) - m + 1 \end{pmatrix}.$$

A lower bound for the product of the diagonal term is j^4 . The off diagonal can be estimated by

$$c_-(m)^2 = (j+m)(j-m+1) = j^2 + j - m^2 + m \leq j^2 + 2j.$$

Hence for $j \geq \frac{3}{2}$ these matrices are clearly invertible. For $j = 1$ and $j = \frac{1}{2}$ it can easily be checked that these matrices are invertible. In particular we see that D_e^2 has trivial kernel.

Choosing the other irreducible representation of $Cl(T_{id}^*SU(2))$ corresponds to replacing E_3 with $-E_3$ and therefore D_e^2 in this representation also has trivial kernel.

References

- [1] J. Aastrup, J. Grimstrup, and R. Nest. On spectral triples in quantum gravity I. *In Preparation.*, 2008.

- [2] Johannes Aastrup and Jesper Møller Grimstrup. Spectral triples of holonomy loops. *Comm. Math. Phys.*, 264(3):657–681, 2006.
- [3] Johannes Aastrup and Jesper Møller Grimstrup. Intersecting Connes non-commutative geometry with quantum gravity. *Internat. J. Modern Phys. A*, 22(8-9):1589–1603, 2007.
- [4] Abhay Ashtekar and Jerzy Lewandowski. Representation theory of analytic holonomy C^* -algebras. In *Knots and quantum gravity (Riverside, CA, 1993)*, volume 1 of *Oxford Lecture Ser. Math. Appl.*, pages 21–61. Oxford Univ. Press, New York, 1994.
- [5] Abhay Ashtekar and Jerzy Lewandowski. Background independent quantum gravity: a status report. *Classical Quantum Gravity*, 21(15):R53–R152, 2004.
- [6] A. L. Carey, J. Phillips, and F. A. Sukochev. On unbounded p -summable Fredholm modules. *Adv. Math.*, 151(2):140–163, 2000.
- [7] Ali H. Chamseddine and Alain Connes. Universal formula for noncommutative geometry actions: unification of gravity and the standard model. *Phys. Rev. Lett.*, 77(24):4868–4871, 1996.
- [8] A.H. Chamsedinne, A. Connes, and M. Marcolli. Gravity and the standard model with neutrino mixing. *arXiv:hep-th/0610241*, 2006.
- [9] Alain Connes. *Noncommutative geometry*. Academic Press Inc., San Diego, CA, 1994.
- [10] Alain Connes. Gravity coupled with matter and the foundation of non-commutative geometry. *Comm. Math. Phys.*, 182(1):155–176, 1996.
- [11] Christian Fleischhack. Stratification of the generalized gauge orbit space. *Comm. Math. Phys.*, 214(3):607–649, 2000.
- [12] Christian Fleischhack. Hyphs and the Ashtekar-Lewandowski measure. *J. Geom. Phys.*, 45(3-4):231–251, 2003.
- [13] D. Marolf and J.M. Mourao. On the support of the Ashtekar-Lewandowski measure. *Comm. Math. Phys.*, 170:583–606, 1995.
- [14] T. Thiemann. Introduction to modern canonical quantum general relativity. *arXiv:gr-qc/0110034v1*, 2001.