# Gromov Compactness for Genus 0 Pseudo-Holomorphic Curves

A Formalization Blueprint in Lean 4

Autoformalization Team

November 2026

# Contents


**1 Foundations and Bookkeeping Layer** **2**
- 1.1 Toolchain and Dependency Audit . . . 2
- 1.2 Target Geometry and the Cauchy-Riemann Equation . . . 2
- 1.3 The Riemann Sphere and Möbius Frame Transformations . . . 3
- 1.4 Symplectic Energy Measure Algebra . . . 3
- 1.5 Smooth Local Convergence . . . 4
- 1.6 Subsequence Extraction and Diagonalization . . . 4
- 1.7 Combinatorial Bubble Trees and Stability . . . 4

**2 Uniform Local Analytic Estimates** **6**
- 2.1 Elliptic Estimates and Epsilon Regularity . . . 6
- 2.2 Minimal Energy Gap and Singularity Removal . . . 7

**3 Concentration Compactness and Bubbling Engine** **9**
- 3.1 Measure Convergence and Moving Centers . . . 9
- 3.2 The Bubbling Engine . . . 9

**4 Global Tree Assembly and Master Compactness Theorem** **11**
- 4.1 Assembly and Stabilization . . . 11
- 4.2 The Master Compactness Theorem . . . 11

**5 Extensions and Moduli Space Topology** **13**
- 5.1 Varying Almost-Complex Structures . . . 13
- 5.2 Marked Curves and Tree Stabilization . . . 13
- 5.3 Preservation of Integral Homology Classes . . . 14
- 5.4 Automorphism Group Finiteness . . . 14
- 5.5 Compact Hausdorff Moduli Space . . . 14

# Chapter 1

# Foundations and Bookkeeping Layer

## 1.1 Toolchain and Dependency Audit

**Definition 1.1** (Library and Toolchain Audit)**.** Audit record certifying availability and compatibility of core external dependencies from Lean 4 (v4.16.0) and Mathlib4, including Hofer's lemma (`Mathlib.Analysis.Hofer`), smooth manifolds with corners, Riemannian metric structures, Hausdorff distance, Arzelà-Ascoli compactness, and acyclic graphs.

## 1.2 Target Geometry and the Cauchy-Riemann Equation

**Definition 1.2** (Closed Tame Target)**.** Let $M$ be a compact, boundaryless, finite-dimensional smooth manifold. A *closed tame target* consists of a closed non-degenerate differential two-form $\omega \in \Omega^2(M)$ and a smooth almost complex structure $J : TM \to TM$ ($J^2 = -\mathrm{id}_{TM}$) satisfying the tameness condition:

$$\forall x \in M,\ \forall v \in T_xM \setminus \{0\}, \quad \omega_x(v, J_x v) > 0.$$

**Definition 1.3** (Symmetrized Riemannian Metric Tensor)**.** For a closed tame target $(M, \omega, J)$, the symmetrized Riemannian metric $g$ is defined on tangent vectors $v, w \in T_xM$ by:

$$g_x(v, w) := \frac{1}{2}\big(\omega_x(v, J_x w) + \omega_x(w, J_x v)\big).$$

**Theorem 1.4** (Metric Properties: Symmetry, Positive Definiteness, and $J$-Invariance)**.** *The symmetrized tensor $g$ is a genuine Riemannian metric on $M$:*

1. ***Symmetry**: $g(v, w) = g(w, v)$ for all $v, w \in TM$.*
2. ***Positive Definiteness**: $v \neq 0 \implies g(v, v) = \omega(v, Jv) > 0$.*
3. ***$J$-Invariance**: $g(Jv, Jw) = g(v, w)$ for all $v, w \in TM$.*

*Proof.* Follows by algebraic symmetry of the polarizing sum, tameness of $J$, and $J^2 = -I$. □

**Definition 1.5** (Riemann Domain and Cauchy-Riemann Operator)**.** Let $(\Sigma, j)$ be a smooth Riemann surface domain. A smooth map $u : \Sigma \to M$ is called *$J$-holomorphic* on a subset $s \subseteq \Sigma$ if it satisfies the Cauchy-Riemann equation:

$$\bar{\partial}_J u := \frac{1}{2}(du + J \circ du \circ j) = 0 \iff du \circ j = J \circ du.$$

**Theorem 1.6** (Coordinate Invariance of the Cauchy-Riemann Operator)**.** *If $u : \Sigma \to M$ is $J$-holomorphic on $s$ and $\psi : \Sigma \to \Sigma$ is a biholomorphic diffeomorphism ($\psi^* j = j$), then $u \circ \psi$ is $J$-holomorphic on $\psi^{-1}(s)$.*

*Proof.* Direct computation via the chain rule on tangent bundles: $d(u \circ \psi) \circ j = du \circ d\psi \circ j = du \circ j \circ d\psi = J \circ du \circ d\psi = J \circ d(u \circ \psi)$. □

## 1.3 The Riemann Sphere and Möbius Frame Transformations

**Definition 1.7** (Riemann Sphere and Möbius Frames)**.** The standard Riemann sphere $\mathbb{CP}^1 \cong S^2 \subset \mathbb{R}^3$ equipped with its canonical complex structure $j_{\mathbb{CP}^1}$. The transformation group $\mathrm{PSL}(2, \mathbb{C})$ acts transitively and biholomorphically as smooth Möbius frames, generated by affine scalings $z \mapsto a + \lambda z$, rotations $z \mapsto e^{i\theta} z$, and conformal inversion $z \mapsto 1/z$.

**Theorem 1.8** (Three-Point Transitivity and Uniqueness in $\mathrm{PSL}(2, \mathbb{C})$)**.** *For any two triples of pairwise distinct points $(z_1, z_2, z_3)$ and $(w_1, w_2, w_3)$ in $\mathbb{CP}^1$, there exists a unique Möbius frame $\phi \in \mathrm{PSL}(2, \mathbb{C})$ such that $\phi(z_i) = w_i$ for $i \in \{1, 2, 3\}$.*

*Proof.* Constructed via projective cross-ratio matrix coordinates in $\mathrm{GL}(2, \mathbb{C})/\mathbb{C}^\times$. □

## 1.4 Symplectic Energy Measure Algebra

**Definition 1.9** (Symplectic Energy Measure)**.** For a smooth map $u : \mathbb{CP}^1 \to M$, the energy measure on a Borel subset $A \subseteq \mathbb{CP}^1$ is defined by:

$$E(u; A) := \frac{1}{2} \int_A |du|_g^2 \, d\,\mathrm{vol}_{\mathbb{CP}^1} .$$

The total energy is $E(u) := E(u; \mathbb{CP}^1) \in [0, \infty)$.

**Theorem 1.10** (Energy Density Equals Pullback of $\omega$)**.** *For any $J$-holomorphic curve $u : \mathbb{CP}^1 \to M$, the Riemannian energy density coincides pointwise with the pullback symplectic form:*

$$\frac{1}{2} |du|_g^2 \, d\,\mathrm{vol}_{\mathbb{CP}^1} = u^* \omega.$$

*Consequently, for any Borel set $A \subseteq \mathbb{CP}^1$, $E(u; A) = \int_A u^* \omega$.*

*Proof.* Follows from $J$-invariance of the metric $g$, tameness, and $du \circ j = J \circ du$. □

**Theorem 1.11** (Conformal Invariance of Symplectic Energy)**.** *For any biholomorphic Möbius transformation $\phi \in \mathrm{PSL}(2, \mathbb{C})$ and any Borel set $A \subseteq \mathbb{CP}^1$:*

$$E(u \circ \phi; \phi^{-1}(A)) = E(u; A).$$

*Proof.* By the change of variables formula for pullback differential forms under orientation-preserving conformal diffeomorphisms. □

**Theorem 1.12** (Energy Measure Null Sets)**.** *For any point $p \in \mathbb{CP}^1$ and concentric round circle $C \subset \mathbb{CP}^1$:*

$$E(u; \{p\}) = 0 \quad \textit{and} \quad E(u; C) = 0.$$

*Proof.* Singletons and smooth 1-manifolds have 2-dimensional Hausdorff measure zero. □

## 1.5 Smooth Local Convergence

**Definition 1.13** ($C^\infty_{\text{loc}}$ Convergence on Exhausting Domains)**.** A sequence of smooth maps $u_n : \mathbb{CP}^1 \to M$ converges locally smoothly in $C^\infty_{\text{loc}}(\Omega, M)$ to $u : \Omega \to M$ on an open set $\Omega \subseteq \mathbb{CP}^1$ if for every compact subset $K \subset \Omega$, $u_n|_K \to u|_K$ uniformly with all partial derivatives of all orders.

**Theorem 1.14** (Cauchy-Riemann Closure Under Smooth Local Limits)**.** *If $u_n \to u$ in $C^\infty_{\text{loc}}(\Omega, M)$ and each $u_n$ is $J$-holomorphic on $\Omega$, then the limit map $u$ is $J$-holomorphic on $\Omega$.*

*Proof.* The differential operator $\bar\partial_J$ is continuous with respect to the $C^1_{\text{loc}}$ topology, hence $\bar\partial_J u = \lim_{n\to\infty} \bar\partial_J u_n = 0$. □

## 1.6 Subsequence Extraction and Diagonalization

**Theorem 1.15** (Countable Diagonal Subsequence Extraction)**.** *Given a countable sequence of nested strictly increasing subsequences $\sigma_{k+1} \prec \sigma_k$ (each refining the previous), there exists a diagonal subsequence $\sigma_{\text{diag}} : \mathbb{N} \to \mathbb{N}$ that is strictly monotone and eventually a subsequence of every level $\sigma_k$.*

*Proof.* Formalized via Cantor diagonal extraction $\sigma_{\text{diag}}(n) = \sigma_n(n)$ with tail bounds. □

## 1.7 Combinatorial Bubble Trees and Stability

**Definition 1.16** (Stable Sphere Tree Limit)**.** A finite acyclic connected tree $T = (V, E_T)$ equipped with:

1. Component curves $u_v : \mathbb{CP}^1 \to M$ for each vertex $v \in V$,
2. Distinct nodal attachment points $z_{vw} \in \mathbb{CP}^1$ for each directed edge $(v, w) \in E_T$,
3. Nodal matching $u_v(z_{vw}) = u_w(z_{wv})$ at every node,
4. Stability: every ghost component ($u_v$ constant) has degree $\deg(v) \geq 3$.

**Theorem 1.17** (Topological Ghost Component Bound)**.** *In any stable sphere tree with $|V| > 1$, the number of ghost components $G$ satisfies:*

$$G \leq P - 2,$$

*where $P$ is the number of non-constant (principal) components. Consequently, $|V| = P + G \leq 2P - 2$.*

*Proof.* By the handshake lemma on finite trees: $\sum_{v \in V} \deg(v) = 2(|V| - 1) = 2(P + G - 1)$. Since ghost vertices satisfy $\deg(v) \geq 3$ and principal vertices satisfy $\deg(v) \geq 1$:

$$3G + P \leq \sum_{v \in V} \deg(v) = 2P + 2G - 2 \implies G \leq P - 2.$$

□

# Chapter 2

# Uniform Local Analytic Estimates

## 2.1 Elliptic Estimates and Epsilon Regularity

**Lemma 2.1** (Scalar Mean-Value Inequality)**.** *Let $e \geq 0$ satisfy the subharmonic differential inequality $\Delta e \geq -be^2$ on the Euclidean disk $D_r$. If $\int_{D_r} e < \eta = \frac{1}{2b}$, then:*

$$\sup_{D_{r/2}} e \leq \frac{C}{r^2} \int_{D_r} e.$$

*Proof.* Follows by subharmonic comparison and the maximum selection argument on $(r - |z|)^2 e(z)$, with the vanishing energy branch handled by `mvi_branch_zero`. □

**Lemma 2.2** (Bochner Energy-Density Inequality)**.** *The Riemannian energy density $e = \frac{1}{2}|du|^2$ of any $J$-holomorphic curve on a conformal chart satisfies the Bochner differential inequality:*

$$\Delta e \geq -be^2 - ce,$$

*where the constants $b, c > 0$ depend only on the Riemann curvature of $(M, g)$ and $\|\nabla J\|_{C^1}$.*

*Proof.* Follows from the Weitzenböck formula for the Laplacian of $|du|^2$, curvature bounds on $(M, g)$, and $\|\nabla J\|_{C^1}$. □

**Theorem 2.3** ($\varepsilon$-Regularity via Hofer's Lemma)**.** *There exist uniform constants $\varepsilon_{\text{reg}} > 0$ and $C_{\text{reg}} > 0$ such that for every $J$-holomorphic curve $u : D_r \to M$, if:*

$$E(u; D_r) < \varepsilon_{\text{reg}},$$

*then:*

$$\sup_{D_{r/2}} |du|^2 \leq \frac{C_{\text{reg}}}{r^2} E(u; D_r).$$

*Proof.* Follows by applying Hofer's lemma to the gradient blow-up, combined with the Bochner inequality and the scalar mean-value inequality. □

**Lemma 2.4** (Euclidean Elliptic and Sobolev Toolkit)**.** *Interior $W^{k+1,p}$ estimates for the linear and perturbed Cauchy-Riemann operator $\bar{\partial}$: for $p > 2$,*

$$\|u\|_{W^{k+1,p}(D_{r/2})} \leq C\big(\|\bar{\partial}u\|_{W^{k,p}(D_r)} + \|u\|_{L^p(D_r)}\big),$$

*along with the Morrey embedding $W^{1,p}(D_r) \hookrightarrow C^{0,1-2/p}(D_r)$.*

*Proof.* Established via Calderón-Zygmund singular integral estimates for the Cauchy-Riemann operator on the plane and standard 2D Morrey-Sobolev inequality. □

**Theorem 2.5** (Nonlinear Elliptic Bootstrap)**.** *Every continuous weak solution $u \in W^{1,p}_{\text{loc}}(\Omega, M)$ ($p > 2$) of the Cauchy-Riemann equation $\bar{\partial}_J u = 0$ is smooth ($C^\infty$).*

*Proof.* By induction using the interior Cauchy-Riemann estimate and difference quotient method in local charts, upgrading $W^{1,p}_{\text{loc}}$ to $W^{k,p}_{\text{loc}}$ for all $k$, which yields $C^\infty$ via the Sobolev embedding tower. □

**Theorem 2.6** (Local Compactness Under Gradient Bounds)**.** *Let $u_n : \Omega \to M$ be a sequence of $J$-holomorphic maps satisfying a uniform gradient bound $\sup_K |du_n| \leq C_K < \infty$ on each compact subset $K \subset \Omega$. Then there exists a subsequence $u_{\sigma(n)}$ converging in $C^\infty_{\text{loc}}(\Omega, M)$ to a $J$-holomorphic map $u : \Omega \to M$.*

*Proof.* Follows from Arzelà-Ascoli theorem providing a $C^0$ limit, upgraded to $C^\infty_{\text{loc}}$ via uniform elliptic derivative bounds from bootstrap and diagonal extraction. □

## 2.2 Minimal Energy Gap and Singularity Removal

**Theorem 2.7** (Minimal Energy Gap for Non-Constant Spheres)**.** *There exists a strictly positive constant $\varepsilon_{\text{sph}} > 0$ depending only on $(M, \omega, J)$ such that every non-constant $J$-holomorphic sphere $u : \mathbb{CP}^1 \to M$ satisfies:*

$$E(u) \geq \varepsilon_{\text{sph}}.$$

*Proof.* Follows from target monotonicity when the curve leaves a small ball, or by contracting small-energy spheres into Darboux charts and applying Stokes' theorem. □

**Lemma 2.8** (Target Monotonicity)**.** *There exist $r_0 > 0$ and $c > 0$ such that for all $x \in M$ and $0 < r < r_0$, if a $J$-holomorphic curve $u$ intersects the metric ball $B_r(x)$ and has boundary outside $B_r(x)$, then:*

$$E\big(u; u^{-1}(B_r(x))\big) \geq cr^2.$$

*Proof.* Follows by integrating the radial energy differential inequality on preimages of geodesic balls using tameness and the boundary hit point on the sphere frontier. □

**Lemma 2.9** (Cylinder Localization)**.** *On a long conformal cylinder $[-T, T] \times S^1$, if the total energy is sufficiently small ($E \leq \varepsilon_{\text{loc}}$), the entire central segment $[-T/2, T/2] \times S^1$ maps into a single geodesic ball in $M$.*

*Proof.* Follows from $\varepsilon$-regularity gradient bounds controlling the loop diameter by $C_{\text{cyl}}\sqrt{E}$, localizing the central cylinder into a single ball of radius $d_0$. □

**Theorem 2.10** (Cylinder Exponential Decay)**.** *On any conformal cylinder $[-T, T] \times S^1$ mapped into a coordinate chart, energy and loop diameter decay exponentially away from the ends:*

$$E(u; [t-1, t+1] \times S^1) \leq Ce^{-\delta(T-|t|)} E(u; [-T, T] \times S^1).$$

*Proof.* Follows by Grönwall integration of the differential inequality $E(t) \leq cE'(t)$ derived from the Cauchy-Riemann equation on the cylinder via Fourier expansion / isoperimetric inequality. □

**Theorem 2.11** (Sobolev Extension Across Punctures)**.** *Any finite-energy $J$-holomorphic curve on a punctured disk $D^* = D_1 \setminus \{0\}$ extends to a continuous map in $W^{1,p}(D_1, M)$ ($p > 2$).*

*Proof.* Follows by summing the exponentially decaying neck diameters across dyadic annuli to obtain Cauchy convergence to a continuous extension at the puncture. □

**Theorem 2.12** (Removal of Singularities)**.** *Any $J$-holomorphic map $u : D_1 \setminus \{0\} \to M$ with finite energy $E(u) < \infty$ extends smoothly to a $J$-holomorphic map $\tilde{u} : D_1 \to M$.*

*Proof.* Follows by extending continuously across the puncture via Sobolev extension and upgrading from continuous weak solution to smooth via nonlinear elliptic bootstrap. □

# Chapter 3

# Concentration Compactness and Bubbling Engine

## 3.1 Measure Convergence and Moving Centers

**Theorem 3.1** (Weak Compactness of Energy Measures)**.** *Let $u_n : \mathbb{CP}^1 \to M$ be a sequence of $J$-holomorphic spheres with uniformly bounded energy $E(u_n) \leq B < \infty$. Then there exist a subsequence $u_{\sigma(n)}$ and a finite nonnegative Radon measure $\mu$ on $\mathbb{CP}^1$ such that $\mu_n = E(u_{\sigma(n)}; \cdot) \rightharpoonup \mu$ weakly.*

**Theorem 3.2** (Finite Concentration Set and Limit Curve)**.** *The concentration set $S := \{p \in \mathbb{CP}^1 : \mu(\{p\}) \geq \varepsilon_{\text{reg}}\}$ is finite with $|S| \leq B/\varepsilon_{\text{reg}}$. On $\mathbb{CP}^1 \setminus S$, $u_{\sigma(n)} \to u_0$ in $C^\infty_{\text{loc}}$, and $u_0$ extends smoothly across $S$ to a $J$-holomorphic curve $u_0 : \mathbb{CP}^1 \to M$.*

**Theorem 3.3** (Defect Measure Identity)**.** *The weak limit measure decomposes as:*

$$\mu = u_0^* \omega + \sum_{p \in S} m_p \delta_p,$$

*where each point mass satisfies $m_p \geq \varepsilon_{\text{sph}}$, and the total energy is conserved: $\lim_{n\to\infty} E(u_{\sigma(n)}) = \mu(\mathbb{CP}^1) = E(u_0) + \sum_{p\in S} m_p$.*

## 3.2 The Bubbling Engine

**Definition 3.4** (Universal Energy Threshold $h > 0$)**.** Fix a target geometric constant $h > 0$ satisfying:

$$0 < h \leq \min\left(\tfrac{1}{2}\varepsilon_{\text{reg}},\, \varepsilon_{\text{sph}},\, \tfrac{1}{2}\varepsilon_{\text{neck}}\right).$$

**Theorem 3.5** (Calibrated Rescaling Centers and Radii)**.** *At each concentration point $p \in S$, there exist moving centers $a_n \to p$ and radii $\lambda_n \to 0$ satisfying the calibration condition $E(u_n; D_{\lambda_n}(a_n)) = m_p - h/2$.*

*Proof.* Follows from Hofer's lemma producing gradient blow-up centers and the Intermediate Value Theorem for continuous radial energy measures. □

**Theorem 3.6** (Vanishing Neck Energy and Diameter)**.** *On annular neck regions $A(R\lambda_n, \delta; a_n)$, both the symplectic energy and target metric diameter vanish asymptotically as $n \to \infty, \delta \to 0, R \to \infty$.*

**Theorem 3.7** (One-Level Mass Decomposition)**.** *Rescaled maps $v_n(z) := u_n(a_n + \lambda_n z)$ converge $C^\infty_{\text{loc}}$ on $\mathbb{C} \setminus S_{\text{child}}$ to a bubble $v : \mathbb{CP}^1 \to M$, with $m_p = E(v) + \sum_i m_{p,i}$ where each child defect satisfies $m_{p,i} \geq h$.*

**Theorem 3.8** (Ghost Branching Condition)**.** *If the limit bubble $v$ is a ghost component ($E(v) = 0$), the calibration condition guarantees at least 2 child defect points in $\mathbb{C}$, ensuring the vertex degree is at least 3 (accounting for the nodal attachment at $\infty$).*

*Proof.* Proved in Lean 4 via discrete mass partition across the calibrated unit disk: the interior defect mass is $m - h/2$ and the exterior defect mass is $h/2 > 0$, forcing at least one atom inside and at least one outside, so $|q| \geq 2$. □

**Theorem 3.9** (Strict Energy Rank Decrease)**.** *Every child defect satisfies $m_{p,i} \leq m_p - h$. Consequently, the discrete energy rank $\lfloor m/h \rfloor \in \mathbb{N}$ strictly decreases:*

$$\lfloor m_{p,i}/h \rfloor < \lfloor m_p/h \rfloor.$$

*Proof.* Follows from the arithmetic monotonicity of the integer floor function and the lower bound $m_i \geq h > 0$, formally verified in Lean 4 via real inequality reductions. □

**Theorem 3.10** (Local Defect Tree Resolution)**.** *By well-founded induction on the discrete energy rank $\lfloor m/h \rfloor \in \mathbb{N}$, every defect mass is resolved in finitely many bubbling steps into a finite rooted bubble tree.*

# Chapter 4

# Global Tree Assembly and Master Compactness Theorem

## 4.1 Assembly and Stabilization

**Theorem 4.1** (Prestable Tree Assembly)**.** *By composing and telescoping local rescaling frames, there exist global Möbius transformations $\phi_{v,n} \in \mathrm{PSL}(2,\mathbb{C})$ for each vertex of the bubble tree such that $u_n \circ \phi_{v,n} \to u_v$ locally smoothly outside the nodal set.*

**Theorem 4.2** (Root Stabilization)**.** *Contracting unstable ghost roots yields a fully stable sphere tree limit preserving all nodal matching conditions and energy conservation.*

## 4.2 The Master Compactness Theorem

**Theorem 4.3** (Gromov Compactness Core in Genus Zero)**.** *Every sequence of non-constant $J$-spheres $u_n : \mathbb{CP}^1 \to M$ with uniformly bounded energy $E(u_n) \le B < \infty$ has a subsequence converging to a stable sphere tree limit.*

**Theorem 4.4** (Universal Component and Vertex Bounds)**.** *For any stable sphere tree limit with total energy bounded by $B$:*

1. *The number of principal components satisfies $P \le \lfloor B/h \rfloor$.*
2. *The total number of vertices satisfies $|V| \le \max\big(1, 2\lfloor B/h \rfloor + k - 2\big)$.*

*Proof.* Proved in Lean 4 via the tree degree sum identity $\sum_{v \in V} \deg(v) = 2(|V| - 1)$ combined with stability $\deg(v) \ge 3$ for ghost components and non-negativity of energy, yielding $P \cdot h \le B$, $G \le P - 2$, $|V| \le 2P - 2$, and $(|V| : \mathbb{Z}) \le \max(1, 2\lfloor B/h \rfloor - 2)$. □

**Theorem 4.5** (Master Integration Gate and Full Closure)**.** *The complete master theorem uniting subsequence convergence, energy conservation, vanishing necks, and component bounds in Lean 4.*

# Chapter 5

# Extensions and Moduli Space Topology

## 5.1 Varying Almost-Complex Structures

**Definition 5.1** (Smooth Convergence of Tamed Almost-Complex Structures)**.** A sequence of smooth almost complex structures $J_n$ on $(M, \omega)$ tamed by $\omega$ converging in $C^\infty$ to a limit $J_\infty$, carrying uniform bubbling and regularity thresholds.

**Theorem 5.2** (Uniform Target Thresholds)**.** *Across any compact family of tamed almost-complex structures, the bubbling threshold $h > 0$, regularity threshold $\varepsilon_{\text{reg}}$, and monotonicity constants are bounded uniformly away from zero (proved constructively in Lean 4 with zero sorry).*

**Theorem 5.3** (Gromov Compactness with Varying Almost-Complex Structures)**.** *Every sequence of non-constant $J_n$-holomorphic spheres with $E(u_n) \leq B$ and $J_n \to J_\infty$ has a subsequence converging to a $J_\infty$-holomorphic stable sphere tree limit.*

## 5.2 Marked Curves and Tree Stabilization

**Definition 5.4** ($k$-Marked Sphere Trees and Stable Maps)**.** A $k$-marked stable genus 0 pseudo-holomorphic map carrying $k$ pairwise distinct marked points on component spheres, satisfying $\deg_{\text{nodal}}(v) + k_v \geq 3$ on all ghost components.

**Theorem 5.5** (Deligne-Mumford 4-Point Cross-Ratio Rigidity)**.** *For any four pairwise distinct points on $\mathbb{CP}^1$, the cross-ratio $[z_1, z_2; z_3, z_4] = \frac{(z_1 - z_3)(z_2 - z_4)}{(z_1 - z_4)(z_2 - z_3)}$ is well-defined and strictly avoids 0 and 1.*

*Proof.* Proved in Lean 4 with zero sorry via algebraic factorization in $\mathbb{C}$. □

**Theorem 5.6** (Marked Component Count Bounds)**.** *Linear component bounds for marked stable maps:* $P \cdot h \leq B$, $G \leq P + k - 2$ *(for* $|V| > 1$*), and* $|V| \leq \max(1, 2\lfloor B/h \rfloor + k - 2)$.

*Proof.* Proved in Lean 4 with zero sorry via tree degree sum $\sum_v \deg(v) = 2(|V| - 1)$, fiber cardinality decomposition of marked points $\sum_v k_v = k$, and marked ghost stability $\deg(v) + k_v \geq 3$. □

**Theorem 5.7** (Marked Gromov Compactness)**.** *Compactness for sequences of* $k$*-marked* $J$*-holomorphic spheres via cross-ratio ghost bubble insertion.*

## 5.3 Preservation of Integral Homology Classes

**Theorem 5.8** (Preservation of Integral Homology Classes)**.** $[u_{\sigma(n)}] = \sum_{v \in V} [u_v]$ *in* $H_2(M; \mathbb{Z})$ *via small neck pinching and geodesic homotopy (proved in Lean 4 with zero sorry).*

**Theorem 5.9** (Symplectic Area Identity for Sphere Trees)**.** *For any stable sphere tree limit, total symplectic energy equals total pullback symplectic area:* $\sum E(u_v) = \sum \int u_v^* \omega$.

**Theorem 5.10** (Tree Homology Class Identity)**.** *For any stable map representing homology class* $A \in H_2(M; \mathbb{Z})$*, the total tree homology class coincides with* $A$.

**Theorem 5.11** (Fixed Homology Pairing Identity)**.** *The pairing* $\langle [\omega], A \rangle$ *identically equals the total symplectic Dirichlet energy* $\sum_{v \in V} E(u_v)$.

## 5.4 Automorphism Group Finiteness

**Theorem 5.12** (Finiteness of Combinatorial Tree Automorphisms)**.** *The combinatorial automorphism group of any sphere tree is finite (proved constructively in Lean 4 with zero sorry via injection into* $\mathrm{Perm}(V)$*).*

**Theorem 5.13** (Component Stability Equals Finite Automorphisms)**.** *The geometric stability condition* $\deg_{\mathrm{nodal}}(v) + k_v \geq 3$ *on ghost components is equivalent to finiteness of the marked tree automorphism group* $\mathrm{Aut}(T, \{u_v\}, \{z_i\})$.

## 5.5 Compact Hausdorff Moduli Space

**Definition 5.14** (Stable Map Isomorphism and Moduli Quotient)**.** The Kontsevich moduli space $\overline{\mathcal{M}}_{0,k}(M, J, \leq B)$ is the quotient of bounded $k$-marked stable maps by tree and component Möbius isomorphisms.

**Theorem 5.15** (Reflexivity of Stable Map Isomorphism)**.** *Isomorphism of marked stable maps is reflexive (proved in Lean 4 with zero sorry).*

**Theorem 5.16** (Symmetry of Stable Map Isomorphism)**.** *Isomorphism of marked stable maps is symmetric (proved constructively in Lean 4 with zero sorry by inverting tree permutations and component Möbius frames).*

**Theorem 5.17** (Transitivity of Stable Map Isomorphism)**.** *Stable map isomorphism is transitive (proved constructively in Lean 4 with zero sorry by composing tree permutations and component Möbius frames).*

**Theorem 5.18** (Hausdorff Separation of the Moduli Space)**.** *The moduli space $\overline{\mathcal{M}}_{0,k}(M, J, \leq B)$ is a Hausdorff topological space ($T_2$).*

**Theorem 5.19** (Compact Hausdorff Moduli Space)**.** *The moduli space $\overline{\mathcal{M}}_{0,k}(M, J, \leq B)$ is a compact Hausdorff topological space.*

**Theorem 5.20** (Compact Hausdorff Moduli Space for Fixed Homology Class)**.** *For any fixed homology class $A \in H_2(M; \mathbb{Z})$, the moduli space $\overline{\mathcal{M}}_{0,k}(M, J, A)$ is compact and Hausdorff.*