

# Multi-ancestry finemapping of gene clusters implicated in cardiovascular disease and mental health

Mark James Adams, INCR

2026-01-31

A genomic region containing a cluster of genes that regulate fatty acid desaturation has recently been found to have a strong association with major depressive disorder (Adams et al. 2025). However, it is unclear which variant in this region is the cause of the association. This project will use biobank whole genome-sequencing datasets to finemap common variants, catalogue rare variants, and understand shared genetic risk between cardiovascular disease and common psychiatric disorders.

## The FADS gene cluster

The fatty acid desaturase gene family (FADS1, FADS2, FADS3) is co-located in a gene-dense region on chromosome 11 (11q12-q13.1). FADS1 and FADS2 regulate the conversion of polyunsaturated fatty acids (omega-6 and -3) into long-chain forms, which have roles in a wide range of cellular processes and are implicated in cardiovascular, metabolic, and immunological health (Koletzko et al. 2019).

The role of FADS as the genes underlying the association with major depression is bolstered by a genetic causal analysis (Mendelian randomisation) of circulating metabolites (Davyson

et al. 2023), which showed that omega-3 levels and omega-6 to omega-3 ratios had a causal effect on depression. While this makes FADS1 and FADS2 the most plausible biological candidates, expression levels of other genes in this region (MYRF and TMEM258) also have possible links to depression. Further work is therefore required to pinpoint the causal variant driving the association.

## Genetic associations

Prior studies examining specific FADS genetic variants and depression (Sublette et al. 2016; Cribb et al. 2018) had small sample sizes ( $N \approx 200$ –1000). Large genome-wide association studies (GWASs) have identified a strong association signal in this region, but the most significant variant has been located in other nearby genes like DAGLA (Howard et al. 2019; Meng et al. 2024) and TMEM258 (Levey et al. 2021; Als et al. 2023). However, the most recent and largest multi-ancestry genetic study of depression found the top signal to be a single nucleotide intron variant in FADS2 (Figure 1).

## Finemapping

Because genetic variants in a region are in close physical linkage, association studies find a correlated statistical signal across variants that makes it difficult to determine which variant is the cause of the association. Finemapping is the process of comparing the linkage and association signals to determine the probability that each SNP is the causal SNP. Finemapping can be enhanced by considering prior causal probabilities based on functional annotations (such as a coding or regulatory function, or evidence of mutational constraint).

Previous investigations of this region used (1) association summary statistics from European-ancestry cohorts only and (2) summary statistics from all ancestry groups merged together. Pre-computed information from UK Biobank was used and it was assumed that the region only had a single causal variant. This latter assumption was stipulated because it does not require that the linkage disequilibrium estimates be population




Figure 1: Region plot of genome-wide association findings for the FADS gene cluster. Top: Tested variants with transformed association p-value. Bottom: Gene positions in the region.

accurate. The assumption of a single causal variant was also reasonable given that the conditional-and-joint analysis (which tests whether each other variant is still significant after controlling for the top associated variant in the region) did not indicate the presence of multiple causal variants.

However, this analysis did not identify a variant that met the finemapping identification criterion, which was a posterior inclusion probability (PIP)  $> 0.95$ . The top variants identified in the multi-ancestry analysis were rs28456 and rs174544.

### Rare variant burden

An analysis to look for the effect of rare variants on major depression and on metabolite levels has also been conducted (Nisbet et al. 2025). This study used short read whole genome sequencing (srWGS) datasets from All of Us and UK Biobank. It considered metabolic measures that had been associated with major depression (Omega 3, Omega 3 percentage, Docosahexaenoic acid, Omega 6/Omega 3 ratio and Degree of Unsaturation.). The authors found that rare variant burden was significantly associated with all tested metabolite measures but was not associated with major depression. This indicates that the GWAS association is likely to be caused by a common variant than being a proxy for a rare variant.

- [rs28456](#): an intron variant in the [FADS2](#) gene and the uncharacterised gene [LOC124902680](#); PIP = 0.122.
- [rs174544](#): a variant in 3' untranslated region of [FADS1](#); PIP = 0.181.

## Analyses

In this project I aimed to discover more about causal variants in the FADS region using multi-ancestry and whole-genome sequencing datasets. Initial analysis improved on previous methods in three ways:

- Rather than using summary statistics from merged ancestries, use summary statistics from the single largest ancestry group with a matched reference panel to test for multiple causal variants.
- Use disaggregated summary statistics from multiple ancestries to improve finemapping.

- Use a large, multi-ancestry-matched WGS dataset as an improve reference panel for finemapping.

All analyses were focused on the FADS gene region chr11:61400000-61700000 using summary statistics from Adams (2025) and Meng (2024). The ancestry clusters used were dervied from continental reference populations, namely African (AFR), Admixed American/Hispanic/Latino (AMR), East Asian (EAS), European (EUR), and South Asian (SAS). I conducted finemapping using PolyFun (Weissbrod et al. 2020).

## Test for multiple causal variants

## Multi-ancestry finemapping

## WGS reference dataset

## Acknowledgments

This work was supported by a BHF REA4 Pump Priming Award (RE/24/130012).

## References

- Adams MJ, Streit F, Meng X, et al (2025) Trans-ancestry genome-wide study of depression identifies 697 associations implicating cell types and pharmacotherapies. *Cell* 188:640–652. <https://doi.org/10.1016/j.cell.2024.12.002>
- Als TD, Kurki MI, Grove J, et al (2023) Depression pathophysiology, risk prediction of recurrence and comorbid psychiatric disorders using genome-wide analyses. *Nature Medicine* 29:1832–1844. <https://doi.org/10.1038/s41591-023-02352-1>

- Cribb L, Murphy J, Froud A, et al (2018) Erythrocyte polyunsaturated fatty acid composition is associated with depression and FADS genotype in caucasians. *Nutritional Neuroscience* 21:589–601. <https://doi.org/10.1080/1028415X.2017.1327685>
- Davyson E, Shen X, Gadd DA, et al (2023) Metabolomic investigation of major depressive disorder identifies a potentially causal association with polyunsaturated fatty acids. *Biological Psychiatry* 94:630–639. <https://doi.org/10.1016/j.biopsych.2023.01.027>
- Howard DM, Adams MJ, Clarke T-K, et al (2019) Genome-wide meta-analysis of depression identifies 102 independent variants and highlights the importance of the prefrontal brain regions. *Nature Neuroscience* 22:343–352. <https://doi.org/10.1038/s41593-018-0326-7>
- Koletzko B, Reischl E, Tanjung C, et al (2019) FADS1 and FADS2 polymorphisms modulate fatty acid metabolism and dietary impact on health. *Annual Review of Nutrition* 39:21–44. <https://doi.org/10.1146/annurev-nutr-082018-124250>
- Levey DF, Stein MB, Wendt FR, et al (2021) Bi-ancestral depression GWAS in the million veteran program and meta-analysis in >1.2 million individuals highlight new therapeutic directions. *Nature Neuroscience* 24:954–963. <https://doi.org/10.1038/s41593-021-00860-2>
- Meng X, Navoly G, Giannakopoulou O, et al (2024) Multi-ancestry genome-wide association study of major depression aids locus discovery, fine mapping, gene prioritization and causal inference. *Nature Genetics*. <https://doi.org/10.1038/s41588-023-01596-4>
- Nisbet L, Adams M, Davyson E, et al (2025) The effect of rare variants in the FADS gene cluster on major depression and associated metabolites. *World Congress of Psychiatric Genetics*, Cancún, Mexico

Sublette ME, Vaquero C, Baca-Garcia E, et al (2016) Lack of association of SNPs from the FADS1-FADS2 gene cluster with major depression or suicidal behavior. *Psychiatric Genetics* 26: <https://doi.org/10.1097/ypg.000000000000111>

Weissbrod O, Hormozdiari F, Benner C, et al (2020) Functionally informed fine-mapping and polygenic localization of complex trait heritability. *Nature Genetics* 52:1355–1363. <https://doi.org/10.1038/s41588-020-00735-5>