

REVIEW PAPER

Crosstalk between Rho of Plants GTPase signalling and plant hormones

Haoyu Tian, Ruohan Lyu, and Peishan Yi^{*} 

Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of Life Sciences, Sichuan University, Chengdu, 610064, P. R. China

^{*} Correspondence: yipeishan@scu.edu.cn

Received 16 January 2024; Editorial decision 8 April 2024; Accepted 12 April 2024

Editor: Karl-Josef Dietz, Bielefeld University, Germany

Abstract

Rho of Plants (ROPs) constitute a plant-specific subset of small guanine nucleotide-binding proteins within the Cdc42/Rho/Rac family. These versatile proteins regulate diverse cellular processes, including cell growth, cell division, cell morphogenesis, organ development, and stress responses. In recent years, the dynamic cellular and subcellular behaviours orchestrated by ROPs have unveiled a notable connection to hormone-mediated organ development and physiological responses, thereby expanding our knowledge of the functions and regulatory mechanisms of this signalling pathway. This review delineates advancements in understanding the interplay between plant hormones and the ROP signalling cascade, focusing primarily on the connections with auxin and abscisic acid pathways, alongside preliminary discoveries in cytokinin, brassinosteroid, and salicylic acid responses. It endeavours to shed light on the intricate, coordinated mechanisms bridging cell- and tissue-level signals that underlie plant cell behaviour, organ development, and physiological processes, and highlights future research prospects and challenges in this rapidly developing field.

Keywords: Abscisic acid, auxin, GTPase, plant hormone, Rho of Plants.

Introduction

Rho of Plants (ROPs) are plant homologues of the Cdc42/Rho/Rac family of small guanine nucleotide-binding proteins (GTPases) (Feiguelman *et al.*, 2018). They exist in two primary forms determined by their binding status with guanosine triphosphate (GTP) to guanosine diphosphate (GDP): an active GTP-binding form and an inactive GDP-binding form (Fu and Yang, 2001; Shichrur and Yalovsky, 2006; Kost, 2008). Guanine nucleotide exchange factor (GEF), GTPase-activating protein (GAP), and guanine nucleotide dissociation inhibitor (GDI) are three key regulatory factors that control the transition of ROP between its active and inactive states (Fig. 1A)

(Feiguelman *et al.*, 2018). RopGEFs promote the conversion of ROP-GDP to ROP-GTP. ROP-related GAPs enhance the GTP hydrolysis activity of ROPs, thus converting ROP-GTP to ROP-GDP. RopGDIs dissociate ROPs from the membrane, negatively regulating membrane localization and activation of ROPs. Plant ROPs comprise two subgroups (Winge *et al.*, 2000; Zheng and Yang, 2000; Fowler, 2010). Type-I ROPs are present in all land plants and are featured by a carboxyl-terminal prenylation motif that mediates membrane attachment. Their evolutionary origin dates back to early streptophyte lineages and coincides with the innovation of multicellularity




Fig. 1. Category and evolutionary origin of ROPs. (A) Domain organization of type-I and type-II ROPs and the cycling of ROPs between the GTP-bound and GDP-bound forms. Both groups comprise a GTPase domain (G domain), a polybasic region (PBR), and a carboxyl terminus. The carboxyl terminus of type-I ROPs is featured by a CAAX motif wherein C is an invariant cysteine, A represents an aliphatic amino acid, and X is a non-specific residue. The carboxyl terminus of type-II ROPs has a glycine (G) and cysteine (C)-containing GC-CG motif. Guanine nucleotide exchange factor (GEF) converts GDP-bound ROPs to the GTP-bound form. GTPase-activating protein (GAP) stimulates the GTPase activity of ROPs and facilitates the conversion of ROP-GTP to ROP-GDP. Guanine nucleotide dissociation inhibitor (GDI) extracts ROPs from the membrane to cytosol. (B) Evolutionary origin of type-I and type-II ROPs. The putative origins of major phytohormone signalling pathways and developmental patterns are shown to indicate their phylogenetic relationship with ROPs. The origin of phytohormone is based on the following literature: auxin (AUX) (Carrillo-Carrasco *et al.*, 2023), cytokinin (CK) (Powell and Heyl, 2023), abscisic acid (ABA) (Sun *et al.*, 2020), salicylic acid (SA) (Wang *et al.*, 2015; Jia *et al.*, 2023), jasmonic acid (JA) (Wang *et al.*, 2015; Blazquez *et al.*, 2020), gibberellin (GA) (Wang *et al.*, 2015; Blazquez *et al.*, 2020), and brassinosteroid (BR) (Wang *et al.*, 2015; Kim and Russinova, 2020). *Note that BR signalling might have been established early in *Klebsormidiophyceae* and charophytes because major components except BK11 are present in these lineages.

in plants (Fig. 1B) (Mulvey and Dolan, 2023b). Type-II ROPs are found only in seed plants and are believed to have evolved from type-I ROPs as a result of acquiring additional sequences that disrupt the prenylation motif (Winge *et al.*, 2000; Zheng and Yang, 2000; Fowler, 2010). Consequently, type-II ROPs are anchored to the membrane through S-acylation in the new carboxyl-terminal GC-CG box (Lavy *et al.*, 2002; Lavy and Yalovsky, 2006). Despite these differences, the catalytic GTPase domains of type-I and type-II ROPs are highly conserved. The functional divergence between them may largely result from differential transcriptional regulation and membrane targeting mechanisms (Yalovsky, 2015).

ROPs are best known for their crucial roles in regulating cell morphogenesis and organization and have been shown to trigger cytoskeleton-associated downstream signalling by activating various effectors (Feiguelman *et al.*, 2018; Ou and Yi, 2022; Li *et al.*, 2023). Present studies of ROP signalling mainly focus on simple cellular systems such as pollen tubes (Li *et al.*, 2023), root hairs (Mendrinna and Persson, 2015), leaf epidermal cells (Lin and Yang, 2020), and secondary cell walls in the xylem (H. Xu *et al.*, 2022). These have collectively revealed

ROP-dependent mechanisms in polarized cell growth, cell division, and cell morphogenesis (Oda and Fukuda, 2013; Ou and Yi, 2022; Yi and Goshima, 2022; Li *et al.*, 2023; Muller, 2023). The function of ROP signalling in regulating cell growth and morphology is conserved in basal land plants (Ito *et al.*, 2014; Burkart *et al.*, 2015; Bascom *et al.*, 2019; Hiwatashi *et al.*, 2019; Le Bail *et al.*, 2019; Cheng *et al.*, 2020; Orr *et al.*, 2020; Yi and Goshima, 2020; Bao *et al.*, 2022; Rong *et al.*, 2022; Mulvey and Dolan, 2023a; Ruan *et al.*, 2023). As these fields are beyond the scope of this review, we refer the readers to the following reviews for progress in these areas (Nibau *et al.*, 2006; Cheung and Wu, 2008; Kost, 2008; Craddock *et al.*, 2012; Yang and Lavagi, 2012; Oda and Fukuda, 2013; Lin *et al.*, 2015; Mendrinna and Persson, 2015; Qin and Dong, 2015; Yalovsky, 2015; Honkanen and Dolan, 2016; Feiguelman *et al.*, 2018; Lin and Yang, 2020; Liu *et al.*, 2021; Smokvariska *et al.*, 2021; Ou and Yi, 2022; Li *et al.*, 2023; Muller, 2023; Pan *et al.*, 2023). It is noteworthy that ROPs also control many developmental processes including seed dormancy and germination, embryo development, seedling growth, and the development of organs such as roots, inflorescence stems, leaves, and petals

(Li *et al.*, 2001, 2005, 2008, 2018; Tao *et al.*, 2002; Lavy *et al.*, 2007; Chen *et al.*, 2011, 2012; Lin *et al.*, 2012; Nagawa *et al.*, 2012; Nibau *et al.*, 2013; Poraty-Gavra *et al.*, 2013; Huang *et al.*, 2014; Zhao *et al.*, 2015; Ren *et al.*, 2016, 2017; Liu *et al.*, 2017, H. Liu *et al.*, 2023), although the underlying mechanisms in these processes are not fully understood.

Compared to the ROP pathway, plant hormones are recognized as central driving factors regulating organ development and morphogenesis. The emergence of terrestrial plants coincided with the complexity of multicellular tissues and an enhanced capacity to adapt to terrestrial environments. Therefore, it is conceivable that this process is closely linked to the origin and evolution of hormone signalling pathways (Rensing *et al.*, 2008; Bowman *et al.*, 2017; Blazquez *et al.*, 2020). Indeed, major hormone pathways involved in developmental regulation, such as auxin and cytokinin (CK) pathways, have emerged before the occurrence of land plants (Fig. 1B) (Carrillo-Carrasco *et al.*, 2023; Powell and Heyl, 2023). CK and abscisic acid (ABA) signalling pathways are among the earliest phytohormones whose occurrence correlates with the emergence of type-I ROPs and the transition from unicellularity to multicellularity (Sun *et al.*, 2020; Mulvey and Dolan, 2023b; Powell and Heyl, 2023). Other pathways mediated by jasmonic acid (JA) (Wang *et al.*, 2015; Blazquez *et al.*, 2020), salicylic acid (SA) (Wang *et al.*, 2015; Jia *et al.*, 2023), gibberellin (GA) (Wang *et al.*, 2015; Blazquez *et al.*, 2020), and brassinosteroid (BR) (Wang *et al.*, 2015; Kim and Russinova, 2020) appear to be established around the time of land colonization by plants or later, potentially in adaption to stress pressures and the need for complex organ construction in terrestrial environments.

Alterations in developmental patterns necessitate the coordination of changes in cell structure and tissue organization, and the establishment of intercellular communication modules. Classical hormone responses involve hormone synthesis, transport, receptor perception, signal transduction, and ultimately changes in gene expression levels (McSteen and Zhao, 2008; Blazquez *et al.*, 2020). Notably, hormone signalling, such as auxin, can also stimulate rapid responses independent of its transcriptional activity (Dubey *et al.*, 2021; Fiedler and Friml, 2023). ROP, as a membrane signalling switch regulating cellular behaviour, has been found to interact with hormone signals, such as auxin (Wu *et al.*, 2011; Lin *et al.*, 2015; Pan *et al.*, 2015), ABA (Hsu *et al.*, 2021), CK (Li *et al.*, 2013; H. Liu *et al.*, 2023), and BR (Zhang *et al.*, 2022). The interplay between ROPs and hormone signalling is gradually being unveiled, initiating a new era in understanding cell-autonomous regulation mechanisms and intercellular communication that collectively determine organogenesis. Nevertheless, this emerging field still faces challenges and lacks a unified model, partly because ROP and hormone pathways function in different ways: ROPs typically influence calcium signalling, cytoskeleton organization, and intracellular trafficking to regulate cell morphology and behaviours in a cell-autonomous manner (Feiguelman *et al.*, 2018; Ou and Yi, 2022; Li *et al.*, 2023; Muller, 2023), while

hormones control cell fate and cell differentiation through transcriptional regulation non-autonomously (Blazquez *et al.*, 2020). In this review, we present updates on our understanding of the crosstalk between ROP signalling and hormone pathways, and highlight directions and challenges for further studies in this exciting area.

Discovery of ROPs and their interaction with hormones

Genes encoding ROPs were cloned in the 1990s (Yang and Watson, 1993; Winge *et al.*, 1997, 2000) and were initially found to regulate pollen tube growth (Lin *et al.*, 1996; Lin and Yang, 1997; Kost *et al.*, 1999). Soon after the discovery, ROPs were found broadly expressed in complex organs and were required for organ development (Li *et al.*, 1998, 2001). Some phenotypes are reminiscent of effects induced by hormone treatment (Li *et al.*, 2001). Since then, an increasing number of studies have revealed reciprocal regulation between the ROP pathway and multiple hormone signals, such as auxin (Tao *et al.*, 2002, 2005), ABA (Lemichiez *et al.*, 2001; Zheng *et al.*, 2002), CK (Li *et al.*, 2013), ethylene (Zermiani *et al.*, 2015), SA (Rong *et al.*, 2016), and BR (Zhang *et al.*, 2022) (Table 1). The best studied examples are ROP-auxin interactions during the development of the interdigitated pavement cells in leaves and the roles of ROPs in ABA-regulated processes such as stomatal closure, seed dormancy, and root growth inhibition. In the following, we discuss the current understanding of ROP signalling and hormone crosstalk by focusing on auxin and ABA. We provide updates in these fields and also discuss recent findings related to other hormones and developmental processes. These discussions may offer insight into the mechanism that integrates cell-autonomous and cell-non-autonomous signalling pathways for establishing complex organs in plants.

Crosstalk between ROP signalling and auxin pathways

Auxin perception, transport, and function

As the first discovered plant hormone, auxin regulates nearly all processes of plant growth and development (Friml, 2022), including the establishment of the body axis and primordial formation in early embryos (Verma *et al.*, 2021), the formation and maintenance of root and shoot meristems (Pernisova and Vernoux, 2021; Roychoudhry and Kepinski, 2022), and gravitropic and phototropic growth (Han *et al.*, 2021; L. Li *et al.*, 2022). At the single-cell level, auxin promotes cell wall expansion and cell growth (Du *et al.*, 2020). Two classes of receptors mediate auxin perception: through binding with auxin, the intracellular receptors TIR1/AFB family F-box proteins target the Aux/IAA transcription repressors for degradation, leading to transcriptional expression of auxin-responsive genes

Table 1. ROP-related genes involved in hormone responses

Gene	Species	Process	Function	Reference
Auxin signalling				
<i>Ntrac1</i>	<i>Nicotiana tabacum</i>	Seedling growth; root development; root hair initiation and morphology	Activated by auxin; promotes Aux/IAA degradation and auxin-responsive gene expression	(Tao et al., 2002, 2005)
<i>rop2</i>	<i>Arabidopsis</i>	Morphology of pavement cells, trichomes, and other cell types	Activated by auxin-ABP1/ABL1/2-TMK1; induces lobe expansion by promoting RIC4-mediated F-actin assembly and PIN1 localization at the lobe membrane; inhibits RIC1-triggered MT organization; inhibits clathrin-mediated PIN1 internalization	(Fu et al., 2002, 2005; Xu et al., 2010, 2014; Nagawa et al., 2012; Yu et al., 2023)
<i>rop3</i>	<i>Arabidopsis</i>	Embryo development; root gravitropism; hypocotyl elongation; cotyledon development	Expression induced by auxin; promotes membrane localization of PIN1/3 but not PIN2 or AUX1; promotes auxin-responsive gene expression	(Huang et al., 2014)
<i>rop4</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis	Activated by auxin-ABP1/ABL1/2-TMK1; induces lobe expansion by promoting RIC4-mediated F-actin assembly and PIN1 localization at the lobe membrane; inhibits RIC1-triggered MT organization; inhibits clathrin-mediated PIN1 internalization	(Fu et al., 2005; Xu et al., 2010, 2014; Nagawa et al., 2012; Yu et al., 2023)
<i>rop6</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis; root gravitropism; vascular tissue patterning; lateral root initiation; inflorescence development	Expression induced by auxin; activated by auxin-ABP1/ABL1/2-TMK1; promotes pavement cell indentation by regulating RIC1-triggered MT organization; inhibits clathrin-mediated endocytosis of PIN1/2 in pavement cells; inhibits PIN2 internalization via RIC1 and actin in roots	(Fu et al., 2009; Xu et al., 2010, 2014; Chen et al., 2012; Lin et al., 2012; Poraty-Gavra et al., 2013; Yu et al., 2023)
<i>rop9</i>	<i>Arabidopsis</i>	Primary root growth; embryo development	Expression induced by auxin; inhibits or promotes auxin-responsive gene expression	(Nibau et al., 2013; Choi et al., 2014)
<i>ropgef1</i>	<i>Arabidopsis</i>	Root hair growth; embryo development; root gravitropic growth; (d) primary root growth	May be involved in auxin-induced root hair growth; promotes correct localization of PIN2/7 and AUX1; promotes auxin-responsive gene expression; promotes actin assembly	(Duan et al., 2010; Liu et al., 2017)
<i>ropgef4</i>	<i>Arabidopsis</i>	Root hair growth	Interacts with FER and may be involved in auxin-induced root hair growth by activating ROP2/6	(Duan et al., 2010; Huang et al., 2013)
<i>ropgef7</i>	<i>Arabidopsis</i>	Meristem maintenance in embryo and seedling roots	Expression induced by auxin; promotes PIN1 localization; promotes auxin-responsive gene expression; binds ROP3 and activate ROPs	(Chen et al., 2011)
<i>ropgef10</i>	<i>Arabidopsis</i>	Root hair initiation	Interacts with FER and may be involved in auxin-induced root hair initiation by activating ROP2/6	(Duan et al., 2010; Huang et al., 2013)
<i>spk1</i>	<i>Arabidopsis</i>	Primary root growth; lateral root initiation; root gravitropic growth	Mediates auxin-induced ROP6 activation; inhibits PIN2 internalization; promotes auxin-responsive gene expression	(Lin et al., 2012)
<i>ric1</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis; root growth; lateral root formation	Expression induced by auxin; acts as a ROP6 effector to promote MT ordering in pavement cells; inhibits clathrin-mediated endocytosis of PIN1/2 in pavement cells; inhibits PIN2 internalization via actin in roots; promotes auxin-responsive gene expression	(Fu et al., 2005, 2009; Xu et al., 2010, 2014; Chen et al., 2012; Lin et al., 2012; Choi et al., 2013)
<i>ric4</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis	Acts as a ROP2 effector to promote F-actin assembly; inhibits PIN1 internalization	(Fu et al., 2005; Nagawa et al., 2012; Xu et al., 2014)
<i>icr1</i>	<i>Arabidopsis</i>	Embryo development; root development; lateral root initiation	Expression induced by auxin treatment; promotes membrane localization of PIN1/2 likely via exocytosis and recycling; degraded by high auxin activity	(Lavy et al., 2007; Li et al., 2008; Hazak et al., 2010, 2014)
<i>fer</i>	<i>Arabidopsis</i>	Root hair initiation and growth	Promotes auxin-induced root hair growth by activating ROPs via interacting with Rop-GEF1/4/10	(Duan et al., 2010; Huang et al., 2013)

Table 1. Continued

Gene	Species	Process	Function	Reference
<i>makr2</i>	<i>Arabidopsis</i>	Primary root growth; root gravitropic growth	Promotes asymmetric PIN2 localization; acts genetically upstream of TMK1 and ROP6 to inhibit their function; inhibited by auxin-TMK1-mediated phosphorylation and dissociation from plasma membrane	(Marques-Bueno <i>et al.</i> , 2021)
<i>rbk1</i>	<i>Arabidopsis</i>	Root elongation; cotyledon expansion	Inhibits auxin signalling by phosphorylating ROP4/6 and likely inactivating them downstream of MKK3/MPK1	(Molendijk <i>et al.</i> , 2008; Enders <i>et al.</i> , 2017)
<i>cmi1</i>	<i>Arabidopsis</i>	Primary root growth; root hair growth; hypocotyl elongation; lateral root formation	Expression induced by auxin; recruited to MTs by ICR1 in Ca ²⁺ dependent manner; promotes auxin-induced Ca ²⁺ patterning; inhibits auxin-induced root growth retardation; inhibits auxin-responsive gene expression	(Hazak <i>et al.</i> , 2019)
<i>tor</i>	<i>Arabidopsis</i>	Shoot development; root development; leaf organogenesis	Activated by auxin and ROP2; phosphorylates E2Fa/b to promote translation re-initiation and cell proliferation	(Li <i>et al.</i> , 2017; Schepetilnikov <i>et al.</i> , 2017)
<i>hda6</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis	Suppresses the expression of auxin-responsive transgene reporters via transgene silencing; inhibits histone acylation of ROP6 promoter and ROP6 expression	(Murfett <i>et al.</i> , 2001; Du <i>et al.</i> , 2024)
<i>Ghrop6</i>	<i>Gossypium hirsutum</i>	Cotton fibre development	Inhibits GhPIN3 localization via GhRIC1	(Xi <i>et al.</i> , 2023)
<i>Osropgef10</i>	<i>Oryza sativa</i>	Crown root development	Expression induced by auxin; interacts with OsRAC3 to inhibit CK signalling	(H. Liu <i>et al.</i> , 2023)
<i>Osrac3</i>	<i>Oryza sativa</i>	Crown root development	Expression induced by auxin; activated by auxin; sequesters OsAHP1/2 to the plasma membrane to inhibit CK signalling	(H. Liu <i>et al.</i> , 2023)
<i>Mprop</i>	<i>Marchantia polymorpha</i>	Rhizoid growth; meristem notch formation; gemma dormancy	Promotes auxin synthesis gene expression; promotes cell division orientation	(Rong <i>et al.</i> , 2022)
<i>Ppric</i>	<i>Physcomitrium patens</i>	Caulonemal cell differentiation	Localizes at the plasma membrane and in the nucleus; nuclear PpRIC inhibits auxin-induced caulonemal cell differentiation	(Ntefidou <i>et al.</i> , 2023)
ABA signalling <i>rop2</i>	<i>Arabidopsis</i>	Stomatal opening and closure	Translocates to plasma membrane under light; inhibits light-induced stomatal opening likely via activating RIC7; translocates into the cytosol upon ABA treatment; inhibits ABA- and CO ₂ -induced stomatal closure likely via regulating endocytosis	(Jeon <i>et al.</i> , 2008; Hwang <i>et al.</i> , 2011)
<i>rop6</i>	<i>Arabidopsis</i>	Stomatal closure	Inactivated by ABA likely via ABI1; promotes actin assembly and organization	(Lemichez <i>et al.</i> , 2001)
<i>rop9</i>	<i>Arabidopsis</i>	Seed dormancy; root elongation	Expression inhibited by ABA; promotes or inhibits ABA-responsive gene expression	(Nibau <i>et al.</i> , 2013; Choi <i>et al.</i> , 2014)
<i>rop10</i>	<i>Arabidopsis</i>	Root elongation; stomatal closure; seed germination	Expression inhibited by ABA; inhibits ABA-responsive gene expression	(Zheng <i>et al.</i> , 2002; Xin <i>et al.</i> , 2005; Choi <i>et al.</i> , 2014)
<i>rop11</i>	<i>Arabidopsis</i>	Seed germination; seedling growth; stomatal closure; primary root growth	Translocates into the nucleus upon ABA treatment; binds to ABI1 and prevents it from being inactivated by ABA; inhibits ABA-responsive gene expression	(Li <i>et al.</i> , 2012a, 2012b; Yu <i>et al.</i> , 2012)
<i>ropgef1</i>	<i>Arabidopsis</i>	Stomatal closure; primary root growth; seed germination; seedling development	Inhibits ABA-induced stomatal closure by activating ROP11; translocates to cytosol for degradation upon ABA treatment; inhibits ABA response likely by interacting with ABI1; phosphorylated by CPK4 and likely CPK3/6/11 for degradation	(Li and Liu, 2012; Yu <i>et al.</i> , 2012; Li <i>et al.</i> , 2016, 2018)
<i>ropgef2</i>	<i>Arabidopsis</i>	Seed germination; seedling development	Associates with mitochondria and recruited to plasma membrane by ROP2/6/10; degraded upon ABA treatment	(Zhao <i>et al.</i> , 2015)
<i>ropgef4</i>	<i>Arabidopsis</i>	Stomatal closure; primary root growth	Inhibits ABA-induced stomatal closure by activating ROP11; may be degraded upon ABA treatment	(Li and Liu, 2012; Yu <i>et al.</i> , 2012; Li <i>et al.</i> , 2016)
<i>RopGEF10</i>	<i>Arabidopsis</i>	Stomatal closure; primary root growth	Inhibits ABA-induced stomatal closure by activating ROP11; may be degraded upon ABA treatment	(Li and Liu, 2012; Yu <i>et al.</i> , 2012; Li <i>et al.</i> , 2016)

Table 1. Continued

Gene	Species	Process	Function	Reference
<i>ric1</i>	<i>Arabidopsis</i>	Seed germination; root growth; lateral root formation	Expression induced by ABA; inhibits ABA-responsive gene expression	(Choi et al., 2013)
<i>ric7</i>	<i>Arabidopsis</i>	Stomatal opening	Translocates to plasma membrane by light; inhibits light-induced stomatal opening	(Jeon et al., 2008)
<i>fer</i>	<i>Arabidopsis</i>	Seedling development; primary root growth; stomatal closure	Expression inhibited by ABA; inhibits ABA-induced phenotypes likely via regulating RopGEF1/4/10-ROP11; increased phosphorylation level by RALF and ABA treatment and dephosphorylated by ABI2	(Duan et al., 2010; Yu et al., 2012; Chen et al., 2016)
<i>hda15</i>	<i>Arabidopsis</i>	Seed germination; drought tolerance	Expression induced by ABA; binds to MYB96 and inhibits the expression of ROP6/10/11 by deacetylating H3/H4 in the promoter	(Lee and Seo, 2019)
<i>myb96</i>	<i>Arabidopsis</i>	Seed germination; drought tolerance	Expression induced by ABA; binds to HDA15 and inhibits the expression of ROP6/10/11 by deacetylating H3/H4 in the promoter	(Seo et al., 2009; Lee and Seo, 2019)
<i>Pprop-gef1/2/3/4/5/6</i>	<i>Physcomitrium patens</i>	Tip cell growth; cell morphology	Expression inhibited by ABA treatment	(Beier et al., 2023)
<i>Pprop1/2/3/4</i>	<i>Physcomitrium patens</i>	Tip cell growth; cell morphology	Expression inhibited by ABA treatment	(Beier et al., 2023)
<i>Csrac1</i>	<i>Camellia sinensis</i>	Seed germination	Expression inhibited by ABA treatment; inhibits the inhibitory effect of ABA on seed germination	(X. Xu et al., 2022)
SA signalling				
<i>rop6</i>	<i>Arabidopsis</i>	Pathogen response	Inhibits SA-responsive gene expression; inhibits SA synthesis; inhibits pathogen entry	(Poraty-Gavra et al., 2013)
<i>ren1</i>	<i>Arabidopsis</i>	Pollen tube growth	Abundance at pollen tube tip inhibited by SA but increased by methylated SA, resulting in hyperactivation and inactivation of ROPs, respectively	(Rong et al., 2016)
<i>Tarop10</i>	<i>Triticum aestivum</i>	Pathogen response	Expression inhibited by SA; interacts with TaTrxh9 and negatively regulates resistance against the stripe rust	(Shi et al., 2021)
BR signalling				
<i>rop2</i>	<i>Arabidopsis</i>	Root gravitropic growth	Expression and localization enhanced by BR treatment; may promote BR-induced gravitropic growth via positively regulating PIN2 expression and localization	(Li et al., 2005)
<i>phgap1/2</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis	Stabilized by BIN2 phosphorylation at the indentation region and inactivates ROP2; degraded at the lobe region due to the inhibition of BIN2 by BR, leading to activation of ROP2	(Lauster et al., 2022; Zhang et al., 2022)
CK signalling				
<i>rop2/4</i>	<i>Arabidopsis</i>	Pavement cell morphogenesis	Activity decreased by ARR20, a positive regulator in CK signalling pathway	(Li et al., 2013)
<i>Osropgef10</i>	<i>Oryza sativa</i>	Crown root development	Expression induced by CK; binds to OsAHP1/2 and recruits them to the plasma membrane; promotes OsRR6 expression; inhibits OshK1 and OsRR22 expression	(H. Liu et al., 2023a)
<i>Osrac3</i>	<i>Oryza sativa</i>	Crown root development	Expression induced by CK; binds to OsAHP1/2 and recruits them to the plasma membrane	(H. Liu et al., 2023a)

(Weijers and Wagner, 2016; Leyser, 2018); extracellular receptor ABP1 functions together with the membrane receptor-like kinase TMK1 to mediate transcription-independent signal transduction (Xu *et al.*, 2014; Napier, 2021). Besides perception, the distribution of auxin is critical for its function. Auxin transport is regulated by the PIN-FORMED (PIN) family efflux transporters and AUX1/LAX influx carriers (Hammes *et al.*, 2022). Their polar localization on the plasma membrane generates an auxin gradient in tissues and differential downstream transcriptional responses, therefore leading to organ initiation and development (Hajny *et al.*, 2022).

Principal modes of ROP-auxin interactions

ROP-auxin interactions are found at both transcriptional and non-transcription levels (Fig. 2). In 2002, based on the similarity of phenotypes in transgenic plants overexpressing tobacco NtRAC1 (a homologue of ROPs) to those in auxin-related defects, Tao *et al.* (2002) found that ROPs could activate auxin responses and the activation process depends on the degradation of Aux/IAA repressors (Tao *et al.*, 2005). Later studies show that PIN transporters are the primary targets of ROPs (Fig. 2). For instance, the ROP effector ICR1 positively regulates exocytosis and polar localization of PINs (Lavy *et al.*, 2007; Li *et al.*, 2008; Hazak *et al.*, 2010). During pavement cell morphogenesis and root development, ROP2/4/6 inhibits the endocytosis of PIN proteins. These processes are mediated by the ROP effectors RIC4 and RIC1 which directly modulate and organize filamentous actin (F-actin) and microtubule (MT) networks (Fu *et al.*, 2002, 2005, 2009; Xu *et al.*, 2010; Chen *et al.*, 2012; Nagawa *et al.*, 2012) in a similar way as they do in the canonical ROP pathway found in tip-growing cells (Feiguelman *et al.*, 2018; Ou and Yi, 2022; Li *et al.*, 2023).

Perturbation of the function of ROP activators, including RopGEF7 and RopGEF1, alters PIN localization and auxin responses (Chen *et al.*, 2011; Liu *et al.*, 2017). RopGEF7 can bind ROP3 and likely activates it during embryo and seedling development as well as in root gravitropic growth (Huang *et al.*, 2014). These findings indicate that canonical ROP pathways commonly regulate membrane trafficking of auxin transporters to modulate auxin-dependent processes.

The regulation of the ROP pathway by auxin signalling can occur at multiple levels (Fig. 2). Firstly, auxin can promote the expression of various components of the ROP pathway genes, such as *ROP3/6/9* (Nibau *et al.*, 2013; Poraty-Gavra *et al.*, 2013; Huang *et al.*, 2014), *RopGEF7* (Chen *et al.*, 2011), and the ROP effector proteins *ICR1* and *RICs* (Hazak *et al.*, 2010; Choi *et al.*, 2013). Secondly, auxin can regulate the stability of ROP pathway proteins. For instance, *ICR1* is rapidly degraded by the TIR1/AFB pathway at the auxin maxima, thus forming a negative feedback regulation (Hazak *et al.*, 2010, 2014). Additionally, auxin can rapidly activate ROPs independent of transcriptional regulation, thereby regulating cell morphology and growth development (see below) (Xu *et al.*, 2010, 2014).

Auxin-ABP1/ABL1/2-TMK1 pathway

The activating effect of auxin on ROP was discovered as early as 20 years ago (Tao *et al.*, 2002). However, whether auxin activates ROP through transcriptional regulation or non-transcriptional regulation has only been recently resolved. Early evidence suggested that auxin might activate ROP through non-transcriptional mechanisms. Firstly, auxin can rapidly activate ROP2/6 within 30 s (Xu *et al.*, 2010). Transcriptional regulation mediated by auxin typically takes several minutes, a timeframe that seems insufficient to meet the demands of




Fig. 2. Crosstalk between ROP signalling and auxin pathways in flowering plants. (A) ROP signalling and auxin pathways in pavement cell morphogenesis. Note that BR has also been recently shown to participate in activating ROP2 through down-regulating PHGAP1/2 stability. (B) ROP signalling and auxin pathways in other developmental processes. Most of the presented network is derived from studies in root growth and root hair development. Stars indicate that redundant paralogues are likely involved and are not shown. Transcriptionally regulated factors are highlighted in the dashed oval. The lack of a direct interaction or regulatory mechanism is indicated with dashed lines.

such rapid responses (Badescu and Napier, 2006). Secondly, the activation of ROP depends on ABP1 (Xu *et al.*, 2010). ABP1 is an auxin receptor located in the endoplasmic reticulum and apoplast and has been found to function extracellularly for rapid auxin responses (Napier, 2021). Additionally, ABP1 can form an auxin-dependent complex with the receptor-like kinase TMK1 (Xu *et al.*, 2014). Despite Gao *et al.* (2015) finding that *abp1* single mutants have no impact on auxin response and organ development and previously reported defects are attributed to background mutations in other genes (Dai *et al.*, 2015; Enders *et al.*, 2015; Michalko *et al.*, 2015), raising questions about ABP1 as an extracellular receptor for auxin, the latest research has demonstrated that ABL1/2 are co-receptors of TMK1 for auxin perception and function partially redundantly with ABP1 in the apoplast, confirming that the ABP1/ABL1/2-TMK1 complex drives rapid auxin response on the plasma membrane (Yu *et al.*, 2023).

ABP1-mediated rapid responses can activate ROP2/6, which, in turn, regulates the localization of PIN proteins, influencing auxin distribution and the auxin response pathway (Fig. 2). In pavement cells, activated ROP2 and ROP6 are localized to the lobe and indentation regions, respectively (Fu *et al.*, 2005, 2009). ROP6, by activating the effector RIC1, promotes microtubule organization in the indentation region, mediating plasma membrane invagination (Fu *et al.*, 2009). ROP2 facilitates lobe expansion by activating the effector RIC4 for F-actin assembly and concurrently inhibiting RIC1 (Fu *et al.*, 2005). Interestingly, PIN proteins are polarly localized to the lobe region, not at the indentation region (Fig. 2A). This localization is achieved through ABP1-ROP2/4 and actin-mediated inhibition of PIN1 endocytosis (Xu *et al.*, 2010; Nagawa *et al.*, 2012). In root development, auxin and ABP1 similarly inhibit the endocytosis of PIN1/2, promoting auxin efflux to establish the correct auxin gradient (Paciorek *et al.*, 2005; Robert *et al.*, 2010). The inhibition of PIN1/2 endocytosis by auxin depends on ROP6 and the effector RIC1 but not on ROP2 (Chen *et al.*, 2012; Choi *et al.*, 2013). Therefore, despite the involvement of different ROPs and effector factors in regulation, the feedback mechanism involving auxin-dependent ROP activation and the inhibition of PIN endocytosis may be universal (Pan *et al.*, 2015). A similar mechanism has recently been discovered in cotton fibre cells (Xi *et al.*, 2023). In pavement cells, does auxin form a local concentration gradient through PIN, thereby influencing cell morphology? As a small molecule, auxin can rapidly diffuse in the apoplast, making it unlikely to distribute differentially in a small area. How does a global signal then selectively activate ROP2/4 and ROP6 in distinct regions? The involvement of a differential sensing ability of auxin in these regions seems unlikely, given that the distribution of TMK1 in the lobe and indentation regions does not exhibit significant differences (Xu *et al.*, 2014). Instead, the ROP2/4 and ROP6 pathways reciprocally inhibit each other and probably generate an interdigitated pattern through self-organization (Fu *et al.*, 2005; Xu *et al.*, 2011; Lin *et al.*, 2015).

The mechanism by which TMK1 activates ROP is currently unclear. One promising possibility is that TMK1 facilitates the clustering of ROPs into lipid nanodomains (Smokvarska *et al.*, 2021; Pan *et al.*, 2023). Membrane lipid nanodomains or lipid rafts are cholesterol and sphingolipid-enriched regions of the membrane (Li *et al.*, 2024). They provide a platform to concentrate membrane receptors and signalling components, thus nicely fulfilling the needs for polar localization and activation of ROPs. Indeed, auxin induces lipid ordering in the indentation region in pavement cells, leading to the formation of nanoclusters of TMK1 and ROP6 (Pan *et al.*, 2020). This process depends on cholesterol synthesis, S-acylation of ROP6, and TMK1/4. In the root, auxin similarly stimulates ROP6 nanoclustering and requires the interaction between the polybasic tail of ROP6 and phosphatidylserine (Platre *et al.*, 2019). Although the direct link between Auxin-ABP1-TMK1 and ROPs is still unknown, emerging evidence suggests the involvement of RopGEFs. In the root, the inhibition of PIN2 endocytosis by ROP6-RIC1 is directly regulated by the guanine exchange factor SPK1 (Lin *et al.*, 2012). As the receptor-like kinase FER directly interacts with RopGEF during auxin-induced root hair formation and activates ROPs (Duan *et al.*, 2010), and FER and TMK1 both belong to the family of receptor-like kinases (Dievart *et al.*, 2020), TMK1 might function similarly in pavement cells. The Arabidopsis genome encodes 14 RopGEFs and one SPK1-GEF (Fowler, 2010; Feiguelman *et al.*, 2018). The development of root hairs requires at least three RopGEF proteins, namely RopGEF1/4/10 (Duan *et al.*, 2010; Huang *et al.*, 2013). Further research is needed to clarify which RopGEFs are regulated by TMK1 and whether TMK1 directly interacts with and activates RopGEFs (Miyawaki and Yang, 2014; Feher and Lajko, 2015). Recently, the membrane-associated protein kinase MAK2 was found to mutually inhibit the auxin-TMK1 pathway during root gravitropic growth, regulating the ROP6-mediated asymmetric distribution of PIN2 (Marques-Bueno *et al.*, 2021). There might be other regulatory factors in the auxin-ABP1/ABL1/2-TMK1-ROP pathway. The formation of ROP nanoclusters has been reported in various cells such as root hairs (Fiona Fuchs *et al.*, 2021), pollen tubes (Fratini *et al.*, 2021), moss protonemal cells (Ruan *et al.*, 2023), and in response to osmotic stress and other physiological processes (Smokvarska *et al.*, 2020, 2023). It is possible that hormone and other membrane signal-triggered clustering of ROPs is a common mechanism in regulating ROP signalling and involves receptor-like kinase-mediated activation (Pan *et al.*, 2023).

Interaction of the auxin-ROP pathway with other signals

In addition to the pathways mentioned above, the crosstalk between the auxin-ROP pathway involves interactions with other signalling molecules (Fig. 2B). For instance, auxin can promote the generation of reactive oxygen species (ROS) to

regulate root hair growth. This process relies on the FER-ROP-GEF1/4/10-ROP pathway (Duan *et al.*, 2010; Huang *et al.*, 2013). Further research is warranted to investigate whether auxin functions through an extracellular pathway to regulate FER activity. Root hairs typically arise from local protrusions formed at the base by the outermost root epidermal cells (Vissenberg *et al.*, 2020). The initiation of root hairs is determined by ROP2/4/6 (Molendijk *et al.*, 2001; Jones *et al.*, 2002; Gendre *et al.*, 2019). Genetic analyses indicate that auxin synergistically regulates ROP localization in root hair cells with ethylene signalling (Fischer *et al.*, 2006). Ethylene treatment in auxin-related mutants promotes ectopic root hair formation, further supporting the potential crosstalk between auxin and ethylene signalling (Kiefer *et al.*, 2015). In the inhibition of root growth, ethylene functions upstream of auxin synthesis, transport, and the TIR1/AFB-Aux/IAA pathway, but does not require the ABP1-ROP6-RIC1 pathway (Wang *et al.*, 2018). Cell growth and organ development mediated by the auxin-ROP pathway are regulated by a kinase cascade composed of MKK3-MPK1-RBK1 (Enders *et al.*, 2017), of which the cysteine-rich receptor-like kinase RBK1 directly phosphorylates ROP4/6 and potentially inhibits its activity (Molendijk *et al.*, 2008). Nitric oxide has been reported to inhibit root growth. It functions to promote the S-acylation and membrane localization of ROP2 and inhibit PIN1 abundance in a ROP2-dependent manner (Kenesi *et al.*, 2023). The ROP effector ICR1 recruits the calcium-binding protein CMI1 to microtubules, mediating auxin-dependent calcium signalling response and root growth (Hazak *et al.*, 2019). In addition to regulating cytoskeleton and calcium signalling, ROP can also mediate auxin-dependent TOR activation. ROP2/4/6 directly binds to TOR and promotes auxin-stimulated TOR phosphorylation and activation, therefore driving translation re-initiation and cell proliferation (Li *et al.*, 2017; Schepetilnikov *et al.*, 2017). Notably, in animals, Rac1 similarly binds mTOR and regulates its membrane localization and activity (Saci *et al.*, 2011). This function appears to be specific to Rac but not played by other groups of the Rho superfamily in animals. Moreover, although Cdc42, Rho, and Rac all regulate actin organization, they are distinctly involved in generating filopodia, stress fibres, and lamellipodia (Hall, 1998). Because ROPs represent the only group of the Rho superfamily in plants (Zheng and Yang, 2000), whether the reported non-canonical functions of ROPs and their interactions with auxin signalling are common or specific to each member or subgroup remains to be addressed.

Cell type-specific auxin-ROP signalling

The interaction between auxin and the ROP pathway is a common phenomenon, but the cell-specific mechanisms are still under investigation. In maize subsidiary mother cells, ROP2/9 regulates cell polarity-dependent asymmetric division (Humphries *et al.*, 2011). The polarization of the subsidiary mother cell is regulated by auxin and correlates with

changes in the localization of PIN proteins (Livanos *et al.*, 2015). Therefore, auxin and the ROP pathway may coordinate in establishing polarity before the division of subsidiary mother cells. During root hair development in Arabidopsis, the nucleus undergoes two consecutive migrations in different directions. This process is regulated by auxin and the ROP pathway. However, genetic analysis suggests they may function independently (Nakamura *et al.*, 2018). The auxin-insensitive mutant *arx1* exhibits enhanced ROP2 levels and genetically interacts with ROP2 in regulating trichome branching, suggesting a crosstalk between the auxin pathway and ROP in regulating trichoblast cell morphology (L. Liu *et al.*, 2023). Some ROPs may regulate development by inhibiting, rather than activating, auxin responses. For instance, the knockdown of *rop9* enhances the effects of auxin in promoting lateral root formation and inhibiting primary root growth (Nibau *et al.*, 2013), although contradictory results were obtained in another study using *rop9* mutants (Choi *et al.*, 2014).

In the liverwort *Marchantia polymorpha*, auxin is synthesized in the meristematic tissue and regulates thallus development (Eklund *et al.*, 2015). The expression of the auxin synthesis gene *YUC2* and auxin-regulated thallus formation and dormancy are regulated by MpROP (Rong *et al.*, 2022). These observations suggest that the interaction between auxin and the ROP pathway has already evolved in the early development of terrestrial plant organs. In the moss *Physcomitrium patens*, auxin can promote the differentiation of caulonemal cells (Thelander *et al.*, 2018). During caulonemal cell growth, the ROP effector PpRIC localizes to the apical membrane and the nucleus (Ntefidou *et al.*, 2023). Nuclear-localized PpRIC has an inhibitory effect on auxin-induced caulonemal cell differentiation (Fig. 3), indicating a negative regulatory role of the ROP pathway in auxin response in moss (Ntefidou *et al.*, 2023). Similar to PpROP4 (Cheng *et al.*, 2020; Yi and Goshima, 2020), the moss PIN homologue PpPINA localizes to the apical tip of caulonemal cells (Viaene *et al.*, 2014), consistent with lower auxin response signals in apical cells (Thelander *et al.*, 2019). However, the mechanism by which PpRIC inhibits the auxin response remains unknown, as the loss of PpRIC does not affect the expression of auxin-responsive genes or the localization of PpPINA, and auxin does not affect the transcription and localization of PpRIC (Ntefidou *et al.*, 2023). Because cell length is not affected in *Ppric* mutants or overexpression lines, PpRIC appears to be an atypical ROP effector that does not participate in cell growth but plays an important role in cell differentiation. Interestingly, the polar localization of PpPINA is negatively and positively regulated by actin and MT, respectively (Tang *et al.*, 2023). There might be other ROP effectors that control auxin gradients through cytoskeletons and membrane trafficking (Fig. 3). In Arabidopsis root hairs, PIN2 exhibits a localization similar to moss PIN proteins (Tang *et al.*, 2023). However, unlike PIN, the auxin influx carrier AUX1 does not localize in root hairs although it is essential for root hair growth (Jones *et al.*, 2009). Mutations and overexpression of auxin transporters




Fig. 3. Interactions between ROP signalling and auxin in the moss *Physcomitrium patens*. The life cycle of *P. patens* is dominated by the development of filamentous protonemata and leafy shoots termed gametophores (not shown). Protonemata are generated through tip growth and comprise two types of tissues, namely caulonemata and chloronemata. Caulonemal cells have fewer and smaller chloroplasts than chloronemal cells and may differentiate into chloronemal cells over time during development. The transition from caulonemal cells to chloronemal cells is inhibited by auxin. The dashed box shows localization patterns of PpROP, PpRIC, and PpPIN in tip-growing protonema cells. Auxin exhibits a gradient down to the tip due to the polar localization of PpPIN. The ROP effector PpRIC is localized at the growing tip and displays a tip-to-base gradient in the nucleus. Nuclear PpRIC can inhibit auxin-induced caulonemal cell differentiation. Localization of PpPIN depends on microtubules (MTs) and actin. This process might be regulated by ROP signalling through uncharacterized effectors.

further indicate the necessity of polar transport of auxin for root hair growth (Velasquez *et al.*, 2016). Similar to root hairs (Pitts *et al.*, 1998), pollen tube growth is positively regulated by auxin (Chen and Zhao, 2008; Wu *et al.*, 2008; Gao *et al.*, 2019). In *Arabidopsis*, only PIN8 is expressed in pollen tubes and is found to localize at the endoplasmic reticulum (Ding *et al.*, 2012). Pollen tubes may have acquired the ability to store auxin in the endoplasmic reticulum and release it in demand since they cannot obtain auxin from other cells. Nevertheless, the interaction between ROP and auxin signals in tip-growing cells remains poorly understood (Pan *et al.*, 2015). Apart from the FER-RopGEF-ROP pathway (Duan *et al.*, 2010), it is likely that additional molecules play a role in auxin-ROP signalling.

Crosstalk between ROP signalling and ABA pathways

ABA core signalling pathway

ABA is another important hormone that regulates plant growth, development, and physiology (Chen *et al.*, 2020), and plays a

crucial role in abiotic stress responses (Waadt *et al.*, 2022). ABA treatment usually promotes seed maturation and dormancy (Ali *et al.*, 2022) and induces stomatal closure (Munemasa *et al.*, 2015; Hsu *et al.*, 2021). In development, ABA typically exerts inhibitory effects on plant growth, although positive regulatory roles have also been noted (Humplik *et al.*, 2017). The core signalling pathway of ABA involves PYLs-PP2C-SnRK2s (Cutler *et al.*, 2010). PYLs serve as intracellular receptors for ABA, inhibiting the activity of the phosphatase PP2C. PP2C, in turn, dephosphorylates and inhibits the activation of SnRK2s. In the presence of ABA, the inhibitory effect of PYLs on PP2C is relieved, leading to the activation of SnRK2s and expression of responsive genes as well as non-transcriptional signalling events (Humplik *et al.*, 2017; Chen *et al.*, 2020).

ROP-ABA interactions in regulating stomata closure

As ABA regulates stomatal closure through organizing the F-actin network (Kim *et al.*, 1995; Eun and Lee, 1997) and ROPs are key regulators of actin assembly (Fu *et al.*, 2002),

the interaction between ROP signalling and ABA was initially studied in stomata (Fig. 4A; Table 1). In 2001, Lemichez *et al.* (2001) demonstrated that ABA inactivates AtRAC1/AtROP6, thus promoting F-actin disassembly and stomatal closure. This process requires the PP2C member ABI1. The following studies provide more evidence in supporting ROP-ABA crosstalk and have revealed that ROPs mainly act as a negative regulator of ABA signalling. For instance, Arabidopsis ROP9 and ROP10 inhibit ABA-mediated stomatal closure, seed dormancy, and root growth inhibition (Zheng *et al.*, 2002; Choi *et al.*, 2014). ROP11 has a similar function to ROP10 but genetically acts in parallel pathways (Li *et al.*, 2012a). ABA-mediated stomatal closure is also negatively regulated by ROP2 (Hwang *et al.*, 2011). In the light response, light triggers the translocation of ROP2 and its effector RIC7 to the plasma membrane and promotes stomatal opening (Jeon *et al.*, 2008). In this process, ROP2 plays an inhibitory role in stomatal opening as opposed to its function in ABA signalling (Jeon *et al.*, 2008; Hwang *et al.*, 2011). Why does ROP2 negatively regulate both the closure and opening of stomata? Under light, actin filaments are radially oriented to facilitate stomatal opening; darkness or ABA

treatment disrupts actin organization, leading to stomatal closure (Kim *et al.*, 1995; Eun and Lee, 1997). One possible explanation is that the actin network must be balanced for sufficient strength and plasticity during stomatal opening. The assembly and arrangement of F-actin induced by ROPs promote the stomatal opening; however, excess assembly and high stability of F-actin can inhibit stomatal opening.

Mechanisms of ROP regulation on ABA signalling

In Arabidopsis, the active form of ROP11 directly binds to ABI1/2 *in vitro* and releases the inhibition of ABA and PYLs on ABI1/2 phosphatase activity (Li *et al.*, 2012b; Yu *et al.*, 2012). Therefore, PP2C may be an important direct target of ROPs. RopGEF1 has been shown to bind to the receptor-like kinase FER and activate ROPs (Duan *et al.*, 2010). FER and various RopGEFs, such as RopGEF1/2/4/10, all have negative regulatory functions in ABA-mediated responses, such as seedling de-greening, root growth inhibition, and seed dormancy




Fig. 4. Crosstalk between ROP signalling and ABA pathways. (A) ROP signalling and ABA pathways in stomatal closure and opening. Some of the regulatory nodes such as FER-RopGEFs are also functional in other developmental processes. (B) ROP signalling and ABA pathways in other developmental processes. Most of the presented network is derived from studies in seed dormancy, root growth inhibition, and seedling development. Stars indicate that redundant paralogues are likely involved and are not shown. Transcriptionally regulated factors are highlighted in the dashed ovals. The lack of a direct interaction or regulatory mechanism is indicated with dashed lines.

(Li and Liu, 2012; Yu *et al.*, 2012; Zhao *et al.*, 2015; Li *et al.*, 2016, 2018). Hence, the entire FER–RopGEF–ROP pathway may be involved in ABA responses. Consistent with this notion, RALF, a ligand for FER, like ABA, can inhibit root growth (Haruta *et al.*, 2014). At the molecular level, both RALF and ABA promote the phosphorylation of FER. The effect induced by ABA is caused by the inactivation of PP2C/ABI2 which dephosphorylates FER (Chen *et al.*, 2016). These findings indicate that FER integrates RALF and ABA signalling pathways through its phosphorylation levels to regulate ROP activity (Chen *et al.*, 2016). ROPs may also inhibit ABA signalling by acting on effectors. Loss of function of *ric1* enhances the inhibition of seed germination, later root formation, and primary root elongation by ABA (Choi *et al.*, 2013). The direct mechanism of action of ROP effectors remains to be studied. In *ric1* mutants, the expression of ABA-responsive genes is up-regulated (Choi *et al.*, 2013). Transcriptional regulation may be involved (Fig. 4B). In support of this notion, ABA-mediated transcriptional responses are also significantly altered in *rop10* mutants (Xin *et al.*, 2005).

Mechanisms of ABA regulation on ROP signalling

The primary effect of ABA on the ROP pathway is to promote the inactivation of ROPs (Fig. 4). Early biochemical and genetic analyses have shown that the active forms of ROP significantly decrease upon ABA treatment (Lemichiez *et al.*, 2001) and overexpressing constitutively-active ROPs could suppress ABA responses (Hwang *et al.*, 2011). The inactivation of ROP is associated with changes in its localization and the down-regulation of RopGEF protein stability. For example, ABA can induce the translocation of ROP11 from the plasma membrane to the nucleus, thereby relieving its inhibition of ABA signalling (Li *et al.*, 2012a). ABA treatment promotes the translocation of RopGEF1/2 to granular structures associated with multivesicular bodies, suggesting that ABA induces RopGEF degradation through the vacuolar pathway (Zhao *et al.*, 2015; Li *et al.*, 2016). Interestingly, RopGEF2 also localizes to mitochondria, and ROP2/6/10 can recruit it to the cell membrane, thus preventing its degradation (Zhao *et al.*, 2015). The recruitment of RopGEF2 to the plasma membrane by ROPs seems to require its N-terminal sequence. Unlike RopGEF2, the N-terminus of RopGEF1 is phosphorylated by CPK kinases, which promotes the degradation of RopGEF1 (Li *et al.*, 2018). Despite the regulation of protein localization and stability, ABA may inhibit the ROP pathway through transcriptional regulation (Fig. 4B; Table 1). In mosses, ABA has inhibitory effects on protonemal cell growth and the expression of ROPs and RopGEFs (Beier *et al.*, 2023). In Arabidopsis, ABA treatment leads to the down-regulation of ROP9 and ROP10 expression (Zheng *et al.*, 2002; Nibau *et al.*, 2013). The *rop9* RNAi

mutant is insensitive to ABA, suggesting that ROP9, unlike ROP10, functions as a positive regulator of ABA (Nibau *et al.*, 2013). Similarly, the expression of *Camellia sinensis* CsRAC1 is positively regulated by ABA. Overexpression of CsRAC1 can enhance the inhibitory effect of ABA on seed germination (X. Xu *et al.*, 2022). In Arabidopsis, the ABA response factor HDA15 and the transcription factor MYB96 form a complex that deacetylates histones in the promoters of ROP6/10/11, thereby inhibiting their expression (Seo *et al.*, 2009; Lee and Seo, 2019). Interestingly, the deacylation of the ROP6 promoter is also regulated by HDA6 during pavement cell development (Du *et al.*, 2024). HDA6 was originally identified as a negative factor in auxin-responsive transgene expression but does not affect the expression of endogenous auxin-inducible genes (Murfett *et al.*, 2001). Since the expression of at least some ROP-related genes, such as *RopGEF2* (Zhao *et al.*, 2015) and *ROP11* (Li *et al.*, 2012a), is not affected by ABA (Fig. 4B), whether transcriptional regulation of the ROP pathway by ABA is a universal mechanism remains to be further explored.

Interactions between the ROP pathway and other hormone signals

In early studies, developmental phenotypes resulting from dysregulation of ROP activity exhibited similarities to the effects of BR treatment, suggesting an interaction between BR signalling and the ROP pathway (Li *et al.*, 2001). Subsequent research revealed that BR promotes the polar localization of ROP2 and the redistribution of PIN2, thereby enhancing root gravitropic growth (Li *et al.*, 2005). This process requires the involvement of F-actin. Therefore, BR may regulate the distribution of auxin by regulating ROPs and PINs similar to auxin. In pavement cells, ROP2 and ROP6 are localized at the lobe and indentation regions, respectively, to facilitate puzzle-shape morphogenesis (Fu *et al.*, 2005, 2009). BIN2 kinase, one of the core components in BR signalling, phosphorylates ROP-related GAP proteins PHGAP1/2 in the indentation region and promotes their protein stability (Fig. 2A), leading to the inactivation of ROP2 (Lauster *et al.*, 2022; Zhang *et al.*, 2022). In the lobe region, BR inactivates BIN2, thus promoting the degradation of PHGAP1/2 and the activation of ROP2 to promote lobe expansion (Zhang *et al.*, 2022). Through mutant screening, Li *et al.* (2013) found that CK signalling mutants exhibited abnormal pavement cell morphogenesis. Genetic analysis indicates that ROP2/4 acts downstream of CK signalling and their activity is inhibited by CK (Li *et al.*, 2013). Recently, rice OsRopGEF10–OsRAC3 was found to inhibit CK signalling during crown root development. They directly bind to OsAHP1/2, sequester it to the cell membrane, and prevent it from activating downstream transcription factors in the nucleus, thereby negatively regulating CK responses (H. Liu *et al.*, 2023). OsRopGEF10 also promotes the expression

of OsRR6 to inhibit CK signalling (H. Liu *et al.*, 2023). Interestingly, the expression and activity of OsRAC3 are positively regulated by auxin, indicating a crosstalk between auxin and CK signalling. In pollen tubes, high concentrations of SA inhibit tip growth. This effect is attributed to the increase in ROP activity caused by the inhibition of ROP-related GAP REN1 by SA (Rong *et al.*, 2016). The inhibitory effect of SA on pollen tube growth does not rely on its receptors but rather involves endocytic pathways (Rong *et al.*, 2016). In wheat *Triticum aestivum*, TaROP10 negatively regulates stripe rust defence responses (Shi *et al.*, 2021). Its expression is suppressed by SA and ABA, suggesting a negative feedback regulation between the ROP pathway and SA responses. In Arabidopsis, the inactivation of ROP6 also negatively influences SA synthesis and the expression of SA-responsive genes (Poraty-Gavra *et al.*, 2013). These studies indicate that the interaction between ROP signalling and hormones such as BR, CK, and SA is a common phenomenon (Table 1). Further exploration of the roles of ROP in hormonal responses will be an important direction for understanding this versatile molecular switch.

Concluding remarks and future perspectives

Similar to yeast Cdc42, ROPs play a crucial role in polarity establishment in single cells such as pollen tubes and root hairs (Ou and Yi, 2022; Li *et al.*, 2023; Muller, 2023). However, as complex organs emerged in land plants, it is not surprising that ROP-regulated signalling has been increasingly utilized during tissue and organ development (Li *et al.*, 2001). ROP-related genes, in particular, have significantly expanded in seed plants, accompanied by a unique type-II ROP subfamily (Winge *et al.*, 2000; Brembu *et al.*, 2006; Fowler, 2010). How do ROPs adapt to multicellularity and regulate tissue and organ development? Increasing evidence indicates that ROP signalling interacts with a variety of hormonal pathways, thus linking cellular dynamics to a broader developmental context involving cell-cell communication and transcriptional regulation. Currently, it is well accepted that ROP signalling undergoes self-organization and determines cell polarity, growth direction, and division direction at the single-cell level (Yang and Lavagi, 2012; Lin *et al.*, 2015; Smokvarska *et al.*, 2021; Yi and Goshima, 2022; Pan *et al.*, 2023). In contrast, hormone signals, such as auxin, regulate the fate and behaviour of cell populations through intercellular communication at the tissue level (Leyser, 2011; Bhatia and Heisler, 2018; Hajny *et al.*, 2022). There is a clear gap in our understanding of organismal development between the cell and tissue levels. Exploring the mechanisms underlying ROP-hormone crosstalk would possibly help bridge this gap.

Although research on the interactions between plant hormones and the ROP signalling pathway is still incomplete, some studies, such as those related to pavement cells (Fig.

2A), have revealed certain patterns and provided a paradigm: auxin rapidly activates ROPs through a non-canonical ABP1/ABL1/2-TMK1 pathway; ROPs, in turn, act on cytoskeleton networks through effectors to inhibit the endocytosis of PIN proteins, therefore regulating the auxin gradient and auxin-dependent morphogenesis (Chen *et al.*, 2015; Lin *et al.*, 2015; Liu *et al.*, 2021). This mechanism is likely conserved in root growth regulation across species (Dubey *et al.*, 2021; Xi *et al.*, 2023). The auxin-ROP interaction network should be not limited to this simple model. For example, recent studies show that mechanical stress in the epidermis of cotyledons can be sensed by FER, which activates ROP6 along with RopGEF14 to regulate pavement cell interdigitation (Lin *et al.*, 2022; Tang *et al.*, 2022). FER and its interaction with RopGEFs have been found to participate in auxin-induced root hair formation (Duan *et al.*, 2010; Huang *et al.*, 2013). Auxin may regulate ROP activity through other membrane signalling components. Furthermore, the expression of many ROP pathway-related genes is regulated by auxin (Fig. 2B). Auxin-responsive genes typically use *cis*-regulatory elements for transcriptional regulation (Weijers and Wagner, 2016). Some of the ROP-related genes have been identified to contain potential auxin response elements in their promoters (Nibau *et al.*, 2013; Poraty-Gavra *et al.*, 2013; B. Li *et al.*, 2022), implying that transcriptional regulation of ROP signalling components by auxin might be another important mechanism in auxin-ROP crosstalk. An in-depth analysis of the function of *cis*-regulatory elements in ROP-related genes may help understand whether auxin and other hormones play a direct role in the transcriptional regulation of ROP signalling. In addition to controlling PIN localization, ROPs can also interact with translation regulatory factors, such as TOR proteins, to regulate gene expression (Li *et al.*, 2017; Schepetilnikov *et al.*, 2017). In other hormone response processes, ROPs directly interact with signal transduction factors, such as the ABA pathway components ABI1/2 (Li *et al.*, 2012b) and the rice CK pathway components OsAHP1/2 (H. Liu *et al.*, 2023). Further identification of ROP-interacting molecules involved in hormone response is key to deciphering the crosstalk between these two types of signalling pathways. In recent years, in particular, the functions of ROPs in cell division and stress responses have emerged (Kawano *et al.*, 2014; Engelhardt *et al.*, 2020; Waadt *et al.*, 2022; Yi and Goshima, 2022; Ganotra *et al.*, 2023). More components in CK and ABA pathways and ROP signalling may interact with each other and remain to be discovered.

Currently, studying ROP-hormone crosstalk in a multicellular context is still challenging. Firstly, hormone signals typically act on cell populations, with longer response times, while the molecular machinery of the ROP pathway often operates in single or a few cells, with relatively faster response kinetics. To unravel the fine interaction mechanisms between hormone signals and the ROP pathway, long-term time-lapse imaging of cell populations at the tissue level with high spatiotemporal

resolution is needed. This requires the use of brighter and more stable fluorescent markers, the optimization of sample preparation, and the precise recognition of cell identity. In higher plants, it is possible to overcome these problems by studying tissues that are easy to manipulate and image, such as leaf epidermal cells (Liu *et al.*, 2021), using recently developed bright and stable fluorescent proteins, such as mNeonGreen and StayGold (Shaner *et al.*, 2013; Hirano *et al.*, 2022), and applying low phototoxicity fluorescence imaging techniques such as light-sheet microscopy (Ovecka *et al.*, 2022). Secondly, exploring how the ROP pathway and hormone signals interact at the transcriptional regulation level may require transcriptomics analysis with higher spatiotemporal resolution. Single-cell sequencing technology developed in recent years holds promise in this regard (Shaw *et al.*, 2021). Additionally, using new plant models such as mosses and liverworts is another option. Mosses and liverworts have relatively simple tissue structures and typically consist of single layers or a few layers of cells, making them ideal for visualizing organ development at the single-cell level (Naramoto *et al.*, 2022). The main components of ROP and hormone signalling pathways are present in moss and liverwort genomes (Eklund *et al.*, 2010; Fowler, 2010; Wang *et al.*, 2015; Blazquez *et al.*, 2020; Guillory and Bonhomme, 2021), and their functional studies have received widespread attention in recent years (Ou and Yi, 2022; Flores-Sandoval *et al.*, 2023). Investigating ROP-hormone crosstalk in basal land plants may provide valuable insights into the principles underpinning multicellular morphogenesis.

Acknowledgements

We apologize to authors whose original publications could not be cited due to space limitations.

Author contributions

HT, RL, and PY prepared the figures and tables, and wrote the manuscript.

Conflict of interest

The authors declare no conflict of interest.

Funding

This work is supported by the Graduate Education and Teaching Reform Research Project (grant no. GSSCU2023058) and the Youth Science and Technology Leading Talent Cultivation Project from Sichuan University (grant no. 0082604151432), and the National Natural Science Foundation of China (grant no. 32270776) to PY.

Data availability

This paper does not include experimental data.

References

- Ali F, Qanmber G, Li F, Wang Z. 2022. Updated role of ABA in seed maturation, dormancy, and germination. *Journal of Advanced Research* **35**, 199–214.
- Badescu GO, Napier RM. 2006. Receptors for auxin: will it all end in TIRs? *Trends in Plant Science* **11**, 217–223.
- Bao L, Ren J, Nguyen M, Slusarczyk AS, Thole JM, Martinez SP, Huang J, Fujita T, Running MP. 2022. The cellular function of ROP GTPase prenylation is important for multicellularity in the moss *Physcomitrium patens*. *Development* **149**, dev200279.
- Bascom C Jr, Burkart GM, Mallett DR, O'Sullivan JE, Tomaszewski AJ, Walsh K, Bezanilla M. 2019. Systematic survey of the function of ROP regulators and effectors during tip growth in the moss *Physcomitrella patens*. *Journal of Experimental Botany* **70**, 447–457.
- Beier MP, Jinno C, Noda N, Nakamura K, Sugano S, Suzuki Y, Fujita T. 2023. ABA signaling converts stem cell fate by substantiating a tradeoff between cell polarity, growth and cell cycle progression and abiotic stress responses in the moss *Physcomitrium patens*. *Frontiers in Plant Science* **14**, 1303195.
- Bhatia N, Heisler MG. 2018. Self-organizing periodicity in development: organ positioning in plants. *Development* **145**, dev149336.
- Blazquez MA, Nelson DC, Weijers D. 2020. Evolution of plant hormone response pathways. *Annual Review of Plant Biology* **71**, 327–353.
- Bowman JL, Kohchi T, Yamato KT, *et al.* 2017. Insights into land plant evolution garnered from the *Marchantia polymorpha* genome. *Cell* **171**, 287–304.e15.
- Brembu T, Winge P, Bones AM, Yang Z. 2006. A RHOse by any other name: a comparative analysis of animal and plant Rho GTPases. *Cell Research* **16**, 435–445.
- Burkart GM, Baskin TI, Bezanilla M. 2015. A family of ROP proteins that suppresses actin dynamics, and is essential for polarized growth and cell adhesion. *Journal of Cell Science* **128**, 2553–2564.
- Carrillo-Carrasco VP, Hernandez-Garcia J, Mutte SK, Weijers D. 2023. The birth of a giant: evolutionary insights into the origin of auxin responses in plants. *The EMBO Journal* **42**, e113018.
- Chen D, Zhao J. 2008. Free IAA in stigmas and styles during pollen germination and pollen tube growth of *Nicotiana tabacum*. *Physiologia Plantarum* **134**, 202–215.
- Chen J, Wang F, Zheng S, Xu T, Yang Z. 2015. Pavement cells: a model system for non-transcriptional auxin signalling and crosstalks. *Journal of Experimental Botany* **66**, 4957–4970.
- Chen J, Yu F, Liu Y, *et al.* 2016. FERONIA interacts with ABI2-type phosphatases to facilitate signaling cross-talk between abscisic acid and RALF peptide in Arabidopsis. *Proceedings of the National Academy of Sciences, USA* **113**, E5519–E5527.
- Chen K, Li GJ, Bressan RA, Song CP, Zhu JK, Zhao Y. 2020. Absciscic acid dynamics, signaling, and functions in plants. *Journal of Integrative Plant Biology* **62**, 25–54.
- Chen M, Liu H, Kong J, *et al.* 2011. RopGEF7 regulates PLETHORA-dependent maintenance of the root stem cell niche in Arabidopsis. *The Plant Cell* **23**, 2880–2894.
- Chen X, Naramoto S, Robert S, Tejos R, Lofke C, Lin D, Yang Z, Friml J. 2012. ABP1 and ROP6 GTPase signaling regulate clathrin-mediated endocytosis in Arabidopsis roots. *Current Biology* **22**, 1326–1332.
- Cheng X, Mwaura BW, Chang Stauffer SR, Bezanilla M. 2020. A fully functional ROP fluorescent fusion protein reveals roles for this GTPase in subcellular and tissue-level patterning. *The Plant Cell* **32**, 3436–3451.
- Cheung AY, Wu HM. 2008. Structural and signaling networks for the polar cell growth machinery in pollen tubes. *Annual Review of Plant Biology* **59**, 547–572.
- Choi Y, Lee Y, Hwang J-U. 2014. Arabidopsis ROP9 and ROP10 GTPases differentially regulate auxin and ABA responses. *Journal of Plant Biology* **57**, 245–254.
- Choi Y, Lee Y, Kim SY, Lee Y, Hwang JU. 2013. Arabidopsis ROP-interactive CRIB motif-containing protein 1 (RIC1) positively regulates auxin

signalling and negatively regulates abscisic acid (ABA) signalling during root development. *Plant, Cell & Environment* **36**, 945–955.

Craddock C, Lavagi I, Yang Z. 2012. New insights into Rho signaling from plant ROP/Rac GTPases. *Trends in Cell Biology* **22**, 492–501.

Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. 2010. Absciscic acid: emergence of a core signaling network. *Annual Review of Plant Biology* **61**, 651–679.

Dai X, Zhang Y, Zhang D, Chen J, Gao X, Estelle M, Zhao Y. 2015. Embryonic lethality of *Arabidopsis abp1-1* is caused by deletion of the adjacent *BSM* gene. *Nature Plants* **1**, 15183.

Dievart A, Gottin C, Perin C, Ranwez V, Chantret N. 2020. Origin and diversity of plant receptor-like kinases. *Annual Review of Plant Biology* **71**, 131–156.

Ding Z, Wang B, Moreno I, et al. 2012. ER-localized auxin transporter PIN8 regulates auxin homeostasis and male gametophyte development in *Arabidopsis*. *Nature Communications* **3**, 941.

Du M, Spalding EP, Gray WM. 2020. Rapid auxin-mediated cell expansion. *Annual Review of Plant Biology* **71**, 379–402.

Du X, Gao Y, Zhang H, Xu X, Li Y, Zhao L, Luo M, Wang H. 2024. HDA6 modulates *Arabidopsis* pavement cell morphogenesis through epigenetic suppression of ROP6 GTPase expression and signaling. *New Phytologist* **241**, 2523–2539.

Duan Q, Kita D, Li C, Cheung AY, Wu HM. 2010. FERONIA receptor-like kinase regulates RHO GTPase signaling of root hair development. *Proceedings of the National Academy of Sciences, USA* **107**, 17821–17826.

Dubey SM, Serre NBC, Oulehlova D, Vittal P, Fendrych M. 2021. No time for transcription-rapid auxin responses in plants. *Cold Spring Harbor Perspectives in Biology* **13**, a039891.

Eklund DM, Svensson EM, Kost B. 2010. *Physcomitrella patens*: a model to investigate the role of RAC/ROP GTPase signalling in tip growth. *Journal of Experimental Botany* **61**, 1917–1937.

Eklund DM, Ishizaki K, Flores-Sandoval E, et al. 2015. Auxin produced by the indole-3-pyruvic acid pathway regulates development and gemmae dormancy in the liverwort *Marchantia polymorpha*. *The Plant Cell* **27**, 1650–1669.

Enders TA, Oh S, Yang Z, Montgomery BL, Strader LC. 2015. Genome sequencing of *Arabidopsis abp1-5* reveals second-site mutations that may affect phenotypes. *The Plant Cell* **27**, 1820–1826.

Enders TA, Frick EM, Strader LC. 2017. An *Arabidopsis* kinase cascade influences auxin-responsive cell expansion. *The Plant Journal* **92**, 68–81.

Engelhardt S, Trutzenberg A, Huckelhoven R. 2020. Regulation and functions of ROP GTPases in plant-microbe interactions. *Cells* **9**, 2016.

Eun SO, Lee Y. 1997. Actin filaments of guard cells are reorganized in response to light and abscisic acid. *Plant Physiology* **115**, 1491–1498.

Feher A, Lajko DB. 2015. Signals fly when kinases meet Rho-of-plants (ROP) small G-proteins. *Plant Science* **237**, 93–107.

Feiguelman G, Fu Y, Yalovsky S. 2018. ROP GTPases structure-function and signaling pathways. *Plant Physiology* **176**, 57–79.

Fiedler L, Friml J. 2023. Rapid auxin signaling: unknowns old and new. *Current Opinion in Plant Biology* **75**, 102443.

Fiona Fuchs VA, Denninger P, Župunski M, Jaillais Y, Engel U, Grossmann G. 2021. Nanodomain-mediated lateral sorting drives polarization of the small GTPase ROP2 in the plasma membrane of root hair cells. *BioRxiv* doi: [10.1101/2021.09.10.459822](https://doi.org/10.1101/2021.09.10.459822). [Preprint].

Fischer U, Ikeda Y, Ljung K, Serralbo O, Singh M, Heidstra R, Palme K, Scheres B, Grebe M. 2006. Vectorial information for *Arabidopsis* planar polarity is mediated by combined AUX1, EIN2, and GNOM activity. *Current Biology* **16**, 2143–2149.

Flores-Sandoval E, Nishihama R, Bowman JL. 2023. Hormonal and genetic control of pluripotency in bryophyte model systems. *Current Opinion in Plant Biology* **77**, 102486.

Fowler JE. 2010. Evolution of the ROP GTPase signaling module. In: Yalovsky S, Baluška F, Jones A, eds. *Integrated G proteins signaling in plants*. Berlin, Heidelberg: Springer, 305–327.

Fratini M, Krishnamoorthy P, Stenzel I, Riechmann M, Matzner M, Bacia K, Heilmann M, Heilmann I. 2021. Plasma membrane nano-organization specifies phosphoinositide effects on Rho-GTPases and actin dynamics in tobacco pollen tubes. *The Plant Cell* **33**, 642–670.

Friml J. 2022. Fourteen stations of auxin. *Cold Spring Harbor Perspectives in Biology* **14**, a039859.

Fu Y, Yang Z. 2001. Rop GTPase: a master switch of cell polarity development in plants. *Trends in Plant Science* **6**, 545–547.

Fu Y, Gu Y, Zheng Z, Wasteneys G, Yang Z. 2005. *Arabidopsis* interdigitating cell growth requires two antagonistic pathways with opposing action on cell morphogenesis. *Cell* **120**, 687–700.

Fu Y, Li H, Yang Z. 2002. The ROP2 GTPase controls the formation of cortical fine F-actin and the early phase of directional cell expansion during *Arabidopsis* organogenesis. *The Plant Cell* **14**, 777–794.

Fu Y, Xu T, Zhu L, Wen M, Yang Z. 2009. A ROP GTPase signaling pathway controls cortical microtubule ordering and cell expansion in *Arabidopsis*. *Current Biology* **19**, 1827–1832.

Ganotra J, Sharma B, Biswal B, Bhardwaj D, Tuteja N. 2023. Emerging role of small GTPases and their interactome in plants to combat abiotic and biotic stress. *Protoplasma* **260**, 1007–1029.

Gao C, Wang Y, Qu H. 2019. Study of auxin regulation of pollen tube growth through calcium channels in *Pyrus pyrifolia*. *Plant Growth Regulation* **89**, 99–108.

Gao Y, Zhang Y, Zhang D, Dai X, Estelle M, Zhao Y. 2015. Auxin binding protein 1 (ABP1) is not required for either auxin signaling or *Arabidopsis* development. *Proceedings of the National Academy of Sciences, USA* **112**, 2275–2280.

Gendre D, Baral A, Dang X, et al. 2019. Rho-of-plant activated root hair formation requires *Arabidopsis YIP4a/b* gene function. *Development* **146**, dev168559.

Guillory A, Bonhomme S. 2021. Phytohormone biosynthesis and signaling pathways of mosses. *Plant Molecular Biology* **107**, 245–277.

Hajny J, Tan S, Friml J. 2022. Auxin canalization: from speculative models toward molecular players. *Current Opinion in Plant Biology* **65**, 102174.

Hall A. 1998. Rho GTPases and the actin cytoskeleton. *Science* **279**, 509–514.

Hammes UZ, Murphy AS, Schwechheimer C. 2022. Auxin transporters - A biochemical view. *Cold Spring Harbor Perspectives in Biology* **14**, a039875.

Han H, Adamowski M, Qi L, Alotaibi SS, Friml J. 2021. PIN-mediated polar auxin transport regulations in plant tropic responses. *New Phytologist* **232**, 510–522.

Haruta M, Sabat G, Stecker K, Minkoff BB, Sussman MR. 2014. A peptide hormone and its receptor protein kinase regulate plant cell expansion. *Science* **343**, 408–411.

Hazak O, Bloch D, Poraty L, Sternberg H, Zhang J, Friml J, Yalovsky S. 2010. A Rho scaffold integrates the secretory system with feedback mechanisms in regulation of auxin distribution. *PLoS Biology* **8**, e1000282.

Hazak O, Obolski U, Prat T, Friml J, Hadany L, Yalovsky S. 2014. Bimodal regulation of ICR1 levels generates self-organizing auxin distribution. *Proceedings of the National Academy of Sciences, USA* **111**, E5471–E5479.

Hazak O, Mamon E, Lavy M, et al. 2019. A novel Ca²⁺-binding protein that can rapidly transduce auxin responses during root growth. *PLoS Biology* **17**, e3000085.

Hirano M, Ando R, Shimozone S, et al. 2022. A highly photostable and bright green fluorescent protein. *Nature Biotechnology* **40**, 1132–1142.

Hiwatashi T, Goh H, Yasui Y, et al. 2019. The RopGEF KARAPPO is essential for the initiation of vegetative reproduction in *Marchantia polymorpha*. *Current Biology* **29**, 3525–3531.e7.

Honkanen S, Dolan L. 2016. Growth regulation in tip-growing cells that develop on the epidermis. *Current Opinion in Plant Biology* **34**, 77–83.

Hsu PK, Dubeaux G, Takahashi Y, Schroeder JI. 2021. Signaling mechanisms in abscisic acid-mediated stomatal closure. *The Plant Journal* **105**, 307–321.

- Huang GQ, Li E, Ge FR, Li S, Wang Q, Zhang CQ, Zhang Y. 2013. Arabidopsis RopGEF4 and RopGEF10 are important for FERONIA-mediated developmental but not environmental regulation of root hair growth. *New Phytologist* **200**, 1089–1101.
- Huang JB, Liu H, Chen M, *et al.* 2014. ROP3 GTPase contributes to polar auxin transport and auxin responses and is important for embryogenesis and seedling growth in Arabidopsis. *The Plant Cell* **26**, 3501–3518.
- Humphries JA, Vejlupekova Z, Luo A, Meeley RB, Sylvester AW, Fowler JE, Smith LG. 2011. ROP GTPases act with the receptor-like protein PAN1 to polarize asymmetric cell division in maize. *The Plant Cell* **23**, 2273–2284.
- Humplik JF, Bergougnoux V, Van Volkenburgh E. 2017. To stimulate or inhibit? That is the question for the function of abscisic acid. *Trends in Plant Science* **22**, 830–841.
- Hwang JU, Jeon BW, Hong D, Lee Y. 2011. Active ROP2 GTPase inhibits ABA- and CO₂-induced stomatal closure. *Plant, Cell & Environment* **34**, 2172–2182.
- Ito K, Ren J, Fujita T. 2014. Conserved function of Rho-related Rop/RAC GTPase signaling in regulation of cell polarity in *Physcomitrella patens*. *Gene* **544**, 241–247.
- Jeon BW, Hwang JU, Hwang Y, *et al.* 2008. The Arabidopsis small G protein ROP2 is activated by light in guard cells and inhibits light-induced stomatal opening. *The Plant Cell* **20**, 75–87.
- Jia X, Wang L, Zhao H, Zhang Y, Chen Z, Xu L, Yi K. 2023. The origin and evolution of salicylic acid signaling and biosynthesis in plants. *Molecular Plant* **16**, 245–259.
- Jones AR, Kramer EM, Knox K, Swarup R, Bennett MJ, Lazarus CM, Leyser HM, Grierson CS. 2009. Auxin transport through non-hair cells sustains root-hair development. *Nature Cell Biology* **11**, 78–84.
- Jones MA, Shen JJ, Fu Y, Li H, Yang Z, Grierson CS. 2002. The Arabidopsis Rop2 GTPase is a positive regulator of both root hair initiation and tip growth. *The Plant Cell* **14**, 763–776.
- Kawano Y, Kaneko-Kawano T, Shimamoto K. 2014. Rho family GTPase-dependent immunity in plants and animals. *Frontiers in Plant Science* **5**, 522.
- Kenesi E, Kolbert Z, Kaszler N, *et al.* 2023. The ROP2 GTPase participates in nitric oxide (NO)-induced root shortening in Arabidopsis. *Plants* **12**, 750.
- Kiefer CS, Claes AR, Nzayisenga JC, Pietra S, Stanislas T, Huser A, Ikeda Y, Grebe M. 2015. Arabidopsis AIP1-2 restricted by WER-mediated patterning modulates planar polarity. *Development* **142**, 151–161.
- Kim EJ, Russinova E. 2020. Brassinosteroid signalling. *Current Biology* **30**, R294–R298.
- Kim M, Hepler PK, Eun SO, Ha KS, Lee Y. 1995. Actin filaments in mature guard cells are radially distributed and involved in stomatal movement. *Plant Physiology* **109**, 1077–1084.
- Kost B. 2008. Spatial control of Rho (Rac-Rop) signaling in tip-growing plant cells. *Trends in Cell Biology* **18**, 119–127.
- Kost B, Lemichez E, Spielhofer P, Hong Y, Tolia K, Carpenter C, Chua NH. 1999. Rac homologues and compartmentalized phosphatidylinositol 4, 5-bisphosphate act in a common pathway to regulate polar pollen tube growth. *The Journal of Cell Biology* **145**, 317–330.
- Lauster T, Stockle D, Gabor K, *et al.* 2022. Arabidopsis pavement cell shape formation involves spatially confined ROPGAP regulators. *Current Biology* **32**, 532–544.e7.
- Lavy M, Yalovsky S. 2006. Association of Arabidopsis type-II ROPs with the plasma membrane requires a conserved C-terminal sequence motif and a proximal polybasic domain. *The Plant Journal* **46**, 934–947.
- Lavy M, Bracha-Drori K, Sternberg H, Yalovsky S. 2002. A cell-specific, prenylation-independent mechanism regulates targeting of type II RACs. *The Plant Cell* **14**, 2431–2450.
- Lavy M, Bloch D, Hazak O, Gutman I, Poraty L, Sorek N, Sternberg H, Yalovsky S. 2007. A Novel ROP/RAC effector links cell polarity, root-meristem maintenance, and vesicle trafficking. *Current Biology* **17**, 947–952.
- Le Bail A, Schulmeister S, Perroud PF, Ntefidou M, Rensing SA, Kost B. 2019. Analysis of the localization of fluorescent PpROP1 and PpROP-GEF4 fusion proteins in moss protonemata based on genomic 'Knock-In' and estradiol-titratable expression. *Frontiers in Plant Science* **10**, 456.
- Lee HG, Seo PJ. 2019. MYB96 recruits the HDA15 protein to suppress negative regulators of ABA signaling in Arabidopsis. *Nature Communications* **10**, 1713.
- Lemichez E, Wu Y, Sanchez JP, Mettouchi A, Mathur J, Chua NH. 2001. Inactivation of AtRac1 by abscisic acid is essential for stomatal closure. *Genes & Development* **15**, 1808–1816.
- Leyser O. 2011. Auxin, self-organisation, and the colonial nature of plants. *Current Biology* **21**, R331–R337.
- Leyser O. 2018. Auxin signaling. *Plant Physiology* **176**, 465–479.
- Li B, Zhang L, Xi J, Hou L, Fu X, Pei Y, Zhang M. 2022. An unexpected regulatory sequence from rho-related GTPase6 confers fiber-specific expression in upland cotton. *International Journal of Molecular Sciences* **23**, 1087.
- Li C, Quintana Perez Y, Lamaze C, Blouin CM. 2024. Lipid nanodomains and receptor signaling: From actin-based organization to membrane mechanics. *Current Opinion in Cell Biology* **86**, 102308.
- Li E, Zhang YL, Qin Z, Xu M, Qiao Q, Li S, Li SW, Zhang Y. 2023. Signaling network controlling ROP-mediated tip growth in Arabidopsis and beyond. *Plant Communications* **4**, 100451.
- Li H, Shen JJ, Zheng ZL, Lin Y, Yang Z. 2001. The Rop GTPase switch controls multiple developmental processes in Arabidopsis. *Plant Physiology* **126**, 670–684.
- Li H, Wu G, Ware D, Davis KR, Yang Z. 1998. Arabidopsis Rho-related GTPases: differential gene expression in pollen and polar localization in fission yeast. *Plant Physiology* **118**, 407–417.
- Li H, Xu T, Lin D, *et al.* 2013. Cytokinin signaling regulates pavement cell morphogenesis in Arabidopsis. *Cell Research* **23**, 290–299.
- Li L, Gallei M, Friml J. 2022. Bending to auxin: fast acid growth for tropisms. *Trends in Plant Science* **27**, 440–449.
- Li L, Xu J, Xu ZH, Xue HW. 2005. Brassinosteroids stimulate plant tropisms through modulation of polar auxin transport in *Brassica* and *Arabidopsis*. *Plant Cell* **17**, 2738–2753.
- Li S, Gu Y, Yan A, Lord E, Yang ZB. 2008. RIP1 (ROP Interactive Partner 1)/ICR1 marks pollen germination sites and may act in the ROP1 pathway in the control of polarized pollen growth. *Molecular Plant* **1**, 1021–1035.
- Li X, Cai W, Liu Y, Li H, Fu L, Liu Z, Xu L, Liu H, Xu T, Xiong Y. 2017. Differential TOR activation and cell proliferation in Arabidopsis root and shoot apices. *Proceedings of the National Academy of Sciences, USA* **114**, 2765–2770.
- Li Z, Liu D. 2012. ROPGEF1 and ROPGEF4 are functional regulators of ROP11 GTPase in ABA-mediated stomatal closure in Arabidopsis. *FEBS Letters* **586**, 1253–1258.
- Li Z, Kang J, Sui N, Liu D. 2012a. ROP11 GTPase is a negative regulator of multiple ABA responses in Arabidopsis. *Journal of Integrative Plant Biology* **54**, 169–179.
- Li Z, Li Z, Gao X, Chinnusamy V, Bressan R, Wang ZX, Zhu JK, Wu JW, Liu D. 2012b. ROP11 GTPase negatively regulates ABA signaling by protecting ABI1 phosphatase activity from inhibition by the ABA receptor RCAR1/PYL9 in Arabidopsis. *Journal of Integrative Plant Biology* **54**, 180–188.
- Li Z, Takahashi Y, Scavo A, Brandt B, Nguyen D, Rieu P, Schroeder JI. 2018. Abscisic acid-induced degradation of Arabidopsis guanine nucleotide exchange factor requires calcium-dependent protein kinases. *Proceedings of the National Academy of Sciences, USA* **115**, E4522–E4531.
- Li Z, Waadt R, Schroeder JI. 2016. Release of GTP exchange factor mediated down-regulation of abscisic acid signal transduction through ABA-induced rapid degradation of RopGEFs. *PLoS Biology* **14**, e1002461.
- Lin W, Yang Z. 2020. Unlocking the mechanisms behind the formation of interlocking pavement cells. *Current Opinion in Plant Biology* **57**, 142–154.
- Lin Y, Yang Z. 1997. Inhibition of pollen tube elongation by microinjected anti-Rop1Ps antibodies suggests a crucial role for rho-type GTPases in the control of tip growth. *Plant Cell* **9**, 1647–1659.

- Lin D, Nagawa S, Chen J, *et al.* 2012. A ROP GTPase-dependent auxin signaling pathway regulates the subcellular distribution of PIN2 in *Arabidopsis* roots. *Current Biology* **22**, 1319–1325.
- Lin D, Ren H, Fu Y. 2015. ROP GTPase-mediated auxin signaling regulates pavement cell interdigitation in *Arabidopsis thaliana*. *Journal of Integrative Plant Biology* **57**, 31–39.
- Lin W, Tang W, Pan X, Huang A, Gao X, Anderson CT, Yang Z. 2022. *Arabidopsis* pavement cell morphogenesis requires FERONIA binding to pectin for activation of ROP GTPase signaling. *Current Biology* **32**, 497–507.e4.
- Lin Y, Wang Y, Zhu JK, Yang Z. 1996. Localization of a Rho GTPase implies a role in tip growth and movement of the generative cell in pollen tubes. *The Plant Cell* **8**, 293–303.
- Liu H, Huang J, Zhang X, *et al.* 2023. The RAC/ROP GTPase activator OsRopGEF10 functions in crown root development by regulating cytokinin signaling in rice. *The Plant Cell* **35**, 453–468.
- Liu L, Niu L, Ji K, Wang Y, Zhang C, Pan M, Wang W, Schiefelbein J, Yu F, An L. 2023. AXR1 modulates trichome morphogenesis through mediating ROP2 stability in *Arabidopsis*. *The Plant Journal* **116**, 756–772.
- Liu S, Jobert F, Rahnesan Z, Doyle SM, Robert S. 2021. Solving the puzzle of shape regulation in plant epidermal pavement cells. *Annual Review of Plant Biology* **72**, 525–550.
- Liu Y, Dong Q, Kita D, Huang JB, Liu G, Wu X, Zhu X, Cheung AY, Wu HM, Tao LZ. 2017. RopGEF1 plays a critical role in polar auxin transport in early development. *Plant Physiology* **175**, 157–171.
- Livanos P, Giannoutsou E, Apostolakis P, Galatis B. 2015. Auxin as an inducer of asymmetrical division generating the subsidiary cells in stomatal complexes of *Zea mays*. *Plant Signaling & Behavior* **10**, e984531.
- Marques-Bueno MM, Armengot L, Noack LC, *et al.* 2021. Auxin-regulated reversible inhibition of TMK1 signaling by MAK2 modulates the dynamics of root gravitropism. *Current Biology* **31**, 228–237.e10.
- McSteen P, Zhao Y. 2008. Plant hormones and signaling: common themes and new developments. *Developmental Cell* **14**, 467–473.
- Mendrinna A, Persson S. 2015. Root hair growth: it's a one way street. *F1000prime Reports* **7**, 23.
- Michalko J, Dravecka M, Bollenbach T, Friml J. 2015. Embryo-lethal phenotypes in early *abp1* mutants are due to disruption of the neighboring *BSM* gene. *F1000Research* **4**, 1104.
- Miyawaki KN, Yang Z. 2014. Extracellular signals and receptor-like kinases regulating ROP GTPases in plants. *Frontiers in Plant Science* **5**, 449.
- Molendijk AJ, Bischoff F, Rajendrakumar CS, Friml J, Braun M, Gilroy S, Palme K. 2001. *Arabidopsis thaliana* RoP GTPases are localized to tips of root hairs and control polar growth. *The EMBO Journal* **20**, 2779–2788.
- Molendijk AJ, Ruperti B, Singh MK, *et al.* 2008. A cysteine-rich receptor-like kinase NCRK and a pathogen-induced protein kinase RBK1 are Rop GTPase interactors. *The Plant Journal* **53**, 909–923.
- Muller S. 2023. Update: on selected ROP cell polarity mechanisms in plant cell morphogenesis. *Plant Physiology* **193**, 26–41.
- Mulvey H, Dolan L. 2023a. RHO GTPase of plants regulates polarized cell growth and cell division orientation during morphogenesis. *Current Biology* **33**, 2897–2911.e6.
- Mulvey H, Dolan L. 2023b. RHO of plant signaling was established early in streptophyte evolution. *Current Biology* **33**, 5515–5525.e4.
- Munemasa S, Hauser F, Park J, Waadt R, Brandt B, Schroeder JI. 2015. Mechanisms of abscisic acid-mediated control of stomatal aperture. *Current Opinion in Plant Biology* **28**, 154–162.
- Murfett J, Wang XJ, Hagen G, Guilfoyle TJ. 2001. Identification of *Arabidopsis* histone deacetylase HDA6 mutants that affect transgene expression. *The Plant Cell* **13**, 1047–1061.
- Nagawa S, Xu T, Lin D, Dhonukshe P, Zhang X, Friml J, Scheres B, Fu Y, Yang Z. 2012. ROP GTPase-dependent actin microfilaments promote PIN1 polarization by localized inhibition of clathrin-dependent endocytosis. *PLoS Biology* **10**, e1001299.
- Nakamura M, Claes AR, Grebe T, Hermkes R, Viotti C, Ikeda Y, Grebe M. 2018. Auxin and ROP GTPase signaling of polar nuclear migration in root epidermal hair cells. *Plant Physiology* **176**, 378–391.
- Napier R. 2021. The story of auxin-binding protein 1 (ABP1). *Cold Spring Harbor Perspectives in Biology* **13**, a039909.
- Naramoto S, Hata Y, Fujita T, Kozuka J. 2022. The bryophytes *Physcomitrium patens* and *Marchantia polymorpha* as model systems for studying evolutionary cell and developmental biology in plants. *The Plant Cell* **34**, 228–246.
- Nibau C, Wu HM, Cheung AY. 2006. RAC/ROP GTPases: 'hubs' for signal integration and diversification in plants. *Trends in Plant Science* **11**, 309–315.
- Nibau C, Tao L, Levasseur K, Wu HM, Cheung AY. 2013. The *Arabidopsis* small GTPase AtRAC7/ROP9 is a modulator of auxin and abscisic acid signalling. *Journal of Experimental Botany* **64**, 3425–3437.
- Ntefidou M, Eklund DM, Le Bail A, *et al.* 2023. *Physcomitrium patens* PpRIC, an ancestral CRIB-domain ROP effector, inhibits auxin-induced differentiation of apical initial cells. *Cell Reports* **42**, 112130.
- Oda Y, Fukuda H. 2013. Spatial organization of xylem cell walls by ROP GTPases and microtubule-associated proteins. *Current Opinion in Plant Biology* **16**, 743–748.
- Orr RG, Cheng X, Vidali L, Bezanilla M. 2020. Orchestrating cell morphology from the inside out - using polarized cell expansion in plants as a model. *Current Opinion in Cell Biology* **62**, 46–53.
- Ou H, Yi P. 2022. ROP GTPase-dependent polarity establishment during tip growth in plants. *New Phytologist* **236**, 49–57.
- Ovecka M, Sojka J, Ticha M, Komis G, Basheer J, Marchetti C, Samajova O, Kubenova L, Samaj J. 2022. Imaging plant cells and organs with light-sheet and super-resolution microscopy. *Plant Physiology* **188**, 683–702.
- Paciorek T, Zazimalova E, Ruthardt N, *et al.* 2005. Auxin inhibits endocytosis and promotes its own efflux from cells. *Nature* **435**, 1251–1256.
- Pan X, Chen J, Yang Z. 2015. Auxin regulation of cell polarity in plants. *Current Opinion in Plant Biology* **28**, 144–153.
- Pan X, Fang L, Liu J, *et al.* 2020. Auxin-induced signaling protein nanoclustering contributes to cell polarity formation. *Nature Communications* **11**, 3914.
- Pan X, Perez-Henriquez P, Van Norman JM, Yang Z. 2023. Membrane nanodomains: dynamic nanobuilding blocks of polarized cell growth. *Plant Physiology* **193**, 83–97.
- Pernisova M, Vernoux T. 2021. Auxin does the SAMba: auxin signaling in the shoot apical meristem. *Cold Spring Harbor Perspectives in Biology* **13**, a039925.
- Pitts RJ, Cernac A, Estelle M. 1998. Auxin and ethylene promote root hair elongation in *Arabidopsis*. *The Plant Journal* **16**, 553–560.
- Platre MP, Bayle V, Armengot L, *et al.* 2019. Developmental control of plant Rho GTPase nano-organization by the lipid phosphatidylserine. *Science* **364**, 57–62.
- Poraty-Gavra L, Zimmermann P, Haigis S, Bednarek P, Hazak O, Stelmakh OR, Sadot E, Schulze-Lefert P, Gruissem W, Yalovsky S. 2013. The *Arabidopsis* Rho of plants GTPase AtROP6 functions in developmental and pathogen response pathways. *Plant Physiology* **161**, 1172–1188.
- Powell AE, Heyl A. 2023. The origin and early evolution of cytokinin signaling. *Frontiers in Plant Science* **14**, 1142748.
- Qin Y, Dong J. 2015. Focusing on the focus: what else beyond the master switches for polar cell growth? *Molecular Plant* **8**, 582–594.
- Ren H, Dang X, Yang Y, Huang D, Liu M, Gao X, Lin D. 2016. SPIKE1 activates ROP GTPase to modulate petal growth and shape. *Plant Physiology* **172**, 358–371.
- Ren H, Dang X, Cai X, Yu P, Li Y, Zhang S, Liu M, Chen B, Lin D. 2017. Spatio-temporal orientation of microtubules controls conical cell shape in *Arabidopsis thaliana* petals. *PLoS Genetics* **13**, e1006851.
- Rensing SA, Lang D, Zimmer AD, *et al.* 2008. The *Physcomitrella* genome reveals evolutionary insights into the conquest of land by plants. *Science* **319**, 64–69.
- Robert S, Kleine-Vehn J, Barbez E, *et al.* 2010. ABP1 mediates auxin inhibition of clathrin-dependent endocytosis in *Arabidopsis*. *Cell* **143**, 111–121.

- Rong D, Luo N, Mollet JC, Liu X, Yang Z. 2016. Salicylic acid regulates pollen tip growth through an NPR3/NPR4-independent pathway. *Molecular Plant* **9**, 1478–1491.
- Rong D, Zhao S, Tang W, *et al.* 2022. ROP signaling regulates spatial pattern of cell division and specification of meristem notch. *Proceedings of the National Academy of Sciences, USA* **119**, e2117803119.
- Roychoudhry S, Kepinski S. 2022. Auxin in root development. *Cold Spring Harbor Perspectives in Biology* **14**, a039933.
- Ruan J, Lai L, Ou H, Yi P. 2023. Two subtypes of GTPase-activating proteins coordinate tip growth and cell size regulation in *Physcomitrium patens*. *Nature Communications* **14**, 7084.
- Saci A, Cantley LC, Carpenter CL. 2011. Rac1 regulates the activity of mTORC1 and mTORC2 and controls cellular size. *Molecular Cell* **42**, 50–61.
- Schepetilnikov M, Makarian J, Srour O, Geldreich A, Yang Z, Chicher J, Hammann P, Ryabova LA. 2017. GTPase ROP2 binds and promotes activation of target of rapamycin, TOR, in response to auxin. *The EMBO Journal* **36**, 886–903.
- Seo PJ, Xiang F, Qiao M, Park JY, Lee YN, Kim SG, Lee YH, Park WJ, Park CM. 2009. The MYB96 transcription factor mediates abscisic acid signaling during drought stress response in *Arabidopsis*. *Plant Physiology* **151**, 275–289.
- Shaner NC, Lambert GG, Chammas A, *et al.* 2013. A bright monomeric green fluorescent protein derived from *Branchiostoma lanceolatum*. *Nature Methods* **10**, 407–409.
- Shaw R, Tian X, Xu J. 2021. Single-cell transcriptome analysis in plants: advances and challenges. *Molecular Plant* **14**, 115–126.
- Shi B, Wang J, Gao H, Yang Q, Wang Y, Day B, Ma Q. 2021. The small GTP-binding protein TaRop10 interacts with TaTrxh9 and functions as a negative regulator of wheat resistance against the stripe rust. *Plant Science* **309**, 110937.
- Shichrur K, Yalovsky S. 2006. Turning ON the switch—RhoGEFs in plants. *Trends in Plant Science* **11**, 57–59.
- Smokvarska M, Francis C, Platre MP, *et al.* 2020. A plasma membrane nanodomain ensures signal specificity during osmotic signaling in plants. *Current Biology* **30**, 4654–4664.e4.
- Smokvarska M, Jaillais Y, Martiniere A. 2021. Function of membrane domains in rho-of-plant signaling. *Plant Physiology* **185**, 663–681.
- Smokvarska M, Bayle V, Maneta-Peyret L, *et al.* 2023. The receptor kinase FERONIA regulates phosphatidylserine localization at the cell surface to modulate ROP signaling. *Science Advances* **9**, eadd4791.
- Sun Y, Pri-Tal O, Michaeli D, Mosquna A. 2020. Evolution of abscisic acid signaling module and its perception. *Frontiers in Plant Science* **11**, 934.
- Tang H, Lu KJ, Zhang Y, Cheng YL, Tu SL, Friml J. 2023. Divergence of trafficking and polarization mechanisms for PIN auxin transporters during land plant evolution. *Plant Communications* **5**, 100669.
- Tang W, Lin W, Zhou X, Guo J, Dang X, Li B, Lin D, Yang Z. 2022. Mechano-transduction via the pectin-FERONIA complex activates ROP6 GTPase signaling in *Arabidopsis* pavement cell morphogenesis. *Current Biology* **32**, 508–517.e3.
- Tao LZ, Cheung AY, Wu HM. 2002. Plant Rac-like GTPases are activated by auxin and mediate auxin-responsive gene expression. *The Plant Cell* **14**, 2745–2760.
- Tao LZ, Cheung AY, Nibau C, Wu HM. 2005. RAC GTPases in tobacco and *Arabidopsis* mediate auxin-induced formation of proteolytically active nuclear protein bodies that contain AUX/IAA proteins. *The Plant Cell* **17**, 2369–2383.
- Thelander M, Landberg K, Sundberg E. 2018. Auxin-mediated developmental control in the moss *Physcomitrella patens*. *Journal of Experimental Botany* **69**, 277–290.
- Thelander M, Landberg K, Sundberg E. 2019. Minimal auxin sensing levels in vegetative moss stem cells revealed by a ratiometric reporter. *New Phytologist* **224**, 775–788.
- Velasquez SM, Barbez E, Kleine-Vehn J, Estevez JM. 2016. Auxin and cellular elongation. *Plant Physiology* **170**, 1206–1215.
- Verma S, Attuluri VPS, Robert HS. 2021. An essential function for auxin in embryo development. *Cold Spring Harbor Perspectives in Biology* **13**, a039966.
- Viaene T, Landberg K, Thelander M, *et al.* 2014. Directional auxin transport mechanisms in early diverging land plants. *Current Biology* **24**, 2786–2791.
- Vissenberg K, Claeijs N, Balcerowicz D, Schoenaers S. 2020. Hormonal regulation of root hair growth and responses to the environment in *Arabidopsis*. *Journal of Experimental Botany* **71**, 2412–2427.
- Waadt R, Seller CA, Hsu PK, Takahashi Y, Munemasa S, Schroeder JI. 2022. Plant hormone regulation of abiotic stress responses. *Nature Reviews Molecular Cell Biology* **23**, 680–694.
- Wang C, Liu Y, Li SS, Han GZ. 2015. Insights into the origin and evolution of the plant hormone signaling machinery. *Plant Physiology* **167**, 872–886.
- Wang Y, Ji Y, Fu Y, Guo H. 2018. Ethylene-induced microtubule reorientation is essential for fast inhibition of root elongation in *Arabidopsis*. *Journal of Integrative Plant Biology* **60**, 864–877.
- Weijers D, Wagner D. 2016. Transcriptional responses to the auxin hormone. *Annual Review of Plant Biology* **67**, 539–574.
- Winge P, Brembu T, Bones AM. 1997. Cloning and characterization of rac-like cDNAs from *Arabidopsis thaliana*. *Plant Molecular Biology* **35**, 483–495.
- Winge P, Brembu T, Kristensen R, Bones AM. 2000. Genetic structure and evolution of RAC-GTPases in *Arabidopsis thaliana*. *Genetics* **156**, 1959–1971.
- Wu HM, Hazak O, Cheung AY, Yalovsky S. 2011. RAC/ROP GTPases and auxin signaling. *Plant Cell* **23**, 1208–1218.
- Wu JZ, Lin Y, Zhang XL, Pang DW, Zhao J. 2008. IAA stimulates pollen tube growth and mediates the modification of its wall composition and structure in *Torenia fournieri*. *Journal of Experimental Botany* **59**, 2529–2543.
- Xi J, Zeng J, Fu X, *et al.* 2023. GhROP6 GTPase modulates auxin accumulation in cotton fibers by regulating cell-specific GhPIN3a localization. *Journal of Experimental Botany* **74**, 265–282.
- Xin Z, Zhao Y, Zheng ZL. 2005. Transcriptome analysis reveals specific modulation of abscisic acid signaling by ROP10 small GTPase in *Arabidopsis*. *Plant Physiology* **139**, 1350–1365.
- Xu H, Giannetti A, Sugiyama Y, Zheng W, Schneider R, Watanabe Y, Oda Y, Persson S. 2022. Secondary cell wall patterning-connecting the dots, pits and helices. *Open Biology* **12**, 210208.
- Xu T, Wen M, Nagawa S, Fu Y, Chen JG, Wu MJ, Perrot-Rechenmann C, Friml J, Jones AM, Yang Z. 2010. Cell surface- and Rho GTPase-based auxin signaling controls cellular interdigitation in *Arabidopsis*. *Cell* **143**, 99–110.
- Xu T, Dai N, Chen J, *et al.* 2014. Cell surface ABP1-TMK auxin-sensing complex activates ROP GTPase signaling. *Science* **343**, 1025–1028.
- Xu T, Nagawa S, Yang Z. 2011. Uniform auxin triggers the Rho GTPase-dependent formation of interdigitation patterns in pavement cells. *Small GTPases* **2**, 227–232.
- Xu X, Ye X, Xing A, Wu Z, Li X, Shu Z, Wang Y. 2022. *Camellia sinensis* small GTPase gene (*CsRAC1*) involves in response to salt stress, drought stress and ABA signaling pathway. *Gene* **821**, 146318.
- Yalovsky S. 2015. Protein lipid modifications and the regulation of ROP GTPase function. *Journal of Experimental Botany* **66**, 1617–1624.
- Yang Z, Lavagi I. 2012. Spatial control of plasma membrane domains: ROP GTPase-based symmetry breaking. *Current Opinion in Plant Biology* **15**, 601–607.
- Yang Z, Watson JC. 1993. Molecular cloning and characterization of rho, a ras-related small GTP-binding protein from the garden pea. *Proceedings of the National Academy of Sciences, USA* **90**, 8732–8736.
- Yi P, Goshima G. 2020. Rho of Plants GTPases and cytoskeletal elements control nuclear positioning and asymmetric cell division during *Physcomitrella patens* branching. *Current Biology* **30**, 2860–2868.e3.
- Yi P, Goshima G. 2022. Division site determination during asymmetric cell division in plants. *The Plant Cell* **34**, 2120–2139.
- Yu F, Qian L, Nibau C, *et al.* 2012. FERONIA receptor kinase pathway suppresses abscisic acid signaling in *Arabidopsis* by activating ABI2 phosphatase. *Proceedings of the National Academy of Sciences, USA* **109**, 14693–14698.

- Yu Y, Tang W, Lin W, et al.** 2023. ABLs and TMKs are co-receptors for extra-cellular auxin. *Cell* **186**, 5457–5471.e17.
- Zermiani M, Zonin E, Nonis A, et al.** 2015. Ethylene negatively regulates transcript abundance of ROP-GAP rheostat-encoding genes and affects apoplastic reactive oxygen species homeostasis in epicarps of cold stored apple fruits. *Journal of Experimental Botany* **66**, 7255–7270.
- Zhang C, Lauster T, Tang W, et al.** 2022. ROPGAP-dependent interaction between brassinosteroid and ROP2-GTPase signaling controls pavement cell shape in Arabidopsis. *Current Biology* **32**, 518–531.e6.
- Zhao S, Wu Y, He Y, Wang Y, Xiao J, Li L, Wang Y, Chen X, Xiong W, Wu Y.** 2015. RopGEF2 is involved in ABA-suppression of seed germination and post-germination growth of Arabidopsis. *The Plant Journal* **84**, 886–899.
- Zheng ZL, Yang Z.** 2000. The Rop GTPase: an emerging signaling switch in plants. *Plant Molecular Biology* **44**, 1–9.
- Zheng ZL, Nafisi M, Tam A, Li H, Crowell DN, Chary SN, Schroeder JI, Shen J, Yang Z.** 2002. Plasma membrane-associated ROP10 small GTPase is a specific negative regulator of abscisic acid responses in Arabidopsis. *The Plant Cell* **14**, 2787–2797.