

```

using Comrade
using Distributions
using Pathfinder
using AdvancedHMC
using Plots
# load eht-imaging we use this to load eht data
load_ehtim()
# To download the data visit https://doi.org/10.25739/g85n-f134
obs = ehtim.obsdata.load_uvfits("SR1_M87_2017_096_lo_hops_netcal_StokesI.uvfits")
obs.add_scans()
# kill 0-baselines since we don't care about
# large scale flux and make scan-average data
obs = obs.flag_uvdist(uv_min=0.1e9).avg_coherent(0.0, scan_avg=true)
# extract log closure amplitudes and closure phases
dlcamp = extract_lcamp(obs; count="min")
dcphase = extract_cphase(obs, count="min")
# form the likelihood
lklhd = RadioLikelihood(dlcamp, dcphase)
# build the model here we fit a ring with a azimuthal
# brightness variation and a Gaussian
function model( $\theta$ )
    (;radius, width,  $\alpha$ ,  $\beta$ , f,  $\sigma_G$ ,  $\tau_G$ ,  $\xi_G$ , xG, yG) =  $\theta$ 
    ring = f*smoothed(stretched(MRing(( $\alpha$ ,), ( $\beta$ ,)), radius, radius), width)
    g = (1-f)*shifted(rotated(stretched(Gaussian(),  $\sigma_G$ ,  $\sigma_G*(1+\tau_G)$ ),  $\xi_G$ ), xG, yG)
    return ring + g
end
# define the priors
prior = (
    radius = Uniform(pas2rad(10.0), pas2rad(30.0)),
    width = Uniform(pas2rad(1.0), pas2rad(10.0)),
     $\alpha$  = Uniform(-0.5, 0.5),  $\beta$  = Uniform(-0.5, 0.5),
    f = Uniform(0.0, 1.0),
     $\sigma_G$  = Uniform(pas2rad(1.0), pas2rad(40.0)),
     $\tau_G$  = Uniform(0.0, 0.75),
     $\xi_G$  = Uniform(0.0,  $1\pi$ ),
    xG = Uniform(-pas2rad(80.0), pas2rad(80.0)),
    yG = Uniform(-pas2rad(80.0), pas2rad(80.0))
)
# Now form the posterior
post = Posterior(lklhd, prior, model)
# We will use HMC to sample the posterior.
# First to reduce burn in we use pathfinder
q, phi, _ = multipathfinder(post, 100)
# now we sample using hmc
metric = DiagEuclideanMetric(dimension(post))
chain, stats = sample(post, HMC(;metric), 2000; nadapts=1000, init_params=phi[1])
# plot a draw from the posterior
plot(model(chain[end]))

```