

# Review of Statistics: Primer

Dr. Patrick Toche

Textbook:

**James H. Stock and Mark W. Watson**, *Introduction to Econometrics*, 4th Edition, Pearson.

Other references:

**Joshua D. Angrist and Jörn-Steffen Pischke**, *Mostly Harmless Econometrics: An Empiricist's Companion*, 1st Edition, Princeton University Press.

**Jeffrey M. Wooldridge**, *Introductory Econometrics: A Modern Approach*, 7th Edition, Cengage Learning.

The textbook comes with online resources and study guides. Other references will be given from time to time.

## Contents

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variances, covariances
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the  $z$ -score
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variances, covariances
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the  $z$ -score
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► **Single Random Variable**

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variances, covariances
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the  $z$ -score
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the  $z$ -score
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the standard normal
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the standard normal distribution
- finding probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)



## ► **Single Random Variable**

- discrete and continuous random variables
- expectations, mean, variance

## ► **Multiple Random Variables**

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► **The Normal Distribution**

- the properties of the normal distribution
- standardizing and the  $z$ -score
- the normal distribution

## ► **The Central Limit Theorem**

- the distribution of the sample mean
- the Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- Distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- Distribution of the sample mean
- Sample size:  $n \geq 30$
- Central Limit Theorem

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- The Central Limit Theorem
- The Central Limit Theorem
- The Central Limit Theorem
- The Central Limit Theorem

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## ▶ **Single Random Variable**

- discrete and continuous random variables
- expectations, mean, variance

## ▶ **Multiple Random Variables**

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ▶ **The Normal Distribution**

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ▶ **The Central Limit Theorem**

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)



## ► Single Random Variable

- discrete and continuous random variables
- expectations, mean, variance

## ► Multiple Random Variables

- conditional probabilities
- conditional means, variance, covariance
- independence, correlation

## ► The Normal Distribution

- properties of the normal distribution
- standardizing and the z-score
- computing probabilities

## ► The Central Limit Theorem

- distribution of the sample mean
- Law of Large Numbers (LLN)
- Central Limit Theorem (CLT)

## Single-Random Variables

## What is a random variable?

Random variable is a function

From the outcome to a number, a random variable has a distribution.

For any subset of the sample space, the distribution describes the probability that the random variable takes a value in that subset.

A coin has a 1 in 2 probability of landing head. The probability that the random variable  $X$  is in  $\{1\}$  is  $1/2$ . The probability that  $X$  is in  $\{1\}$  is  $1/2$ . The probability that  $X$  is in  $\{1\}$  is  $1/2$ .

## What is a random variable?

- ▶ **Example:** Flipping a coin.
- ▶ While the outcome is uncertain, a random variable has a **distribution**.
- ▶ For any subset of the sample space, the distribution describes the probability that the random variable takes a value in that subset.
- ▶ A fair coin has a 1 in 2 probability of landing head. The probability that the random variable  $X$  is in  $\{H\}$  is  $1/2$ . The probability that  $X$  is in  $\{T\}$  is  $1/2$ . The probability that  $X$  is in  $\{H, T\}$  is 1.

## What is a random variable?

- ▶ **Example:** Flipping a coin.
- ▶ While the outcome is uncertain, a random variable has a **distribution**.
- ▶ For any subset of the sample space, the distribution describes the probability that the random variable takes a value in that subset.
- ▶ A fair coin has a 1 in 2 probability of landing head. The probability that the random variable  $X$  is in  $\{H\}$  is  $1/2$ . The probability that  $X$  is in  $\{T\}$  is  $1/2$ . The probability that  $X$  is in  $\{H, T\}$  is 1.

## What is a random variable?

- ▶ **Example:** Flipping a coin.
- ▶ While the outcome is uncertain, a random variable has a **distribution**.
- ▶ For any subset of the sample space, the distribution describes the probability that the random variable takes a value in that subset.
- ▶ A fair coin has a 1 in 2 probability of landing head. The probability that the random variable  $X$  is in  $\{H\}$  is  $1/2$ . The probability that  $X$  is in  $\{T\}$  is  $1/2$ . The probability that  $X$  is in  $\{H, T\}$  is 1.

## What is a random variable?

- ▶ **Example:** Flipping a coin.
- ▶ While the outcome is uncertain, a random variable has a **distribution**.
- ▶ For any subset of the sample space, the distribution describes the probability that the random variable takes a value in that subset.
- ▶ A fair coin has a 1 in 2 probability of landing head. The probability that the random variable  $X$  is in  $\{H\}$  is  $1/2$ . The probability that  $X$  is in  $\{T\}$  is  $1/2$ . The probability that  $X$  is in  $\{H, T\}$  is 1.

# Random Variable

- ▶ Suppose we want to know about the education levels of people in California.
- ▶ Different people have different levels of education. The education level of a randomly selected person in the population is uncertain.
- ▶ The education level is a **random variable** with a distribution that describes the probability that a randomly selected person has an education level in certain range, e.g. 80% have a high school diploma, 30% have a college degree.
- ▶ In general, we do not know the exact distribution of the random variable. Statistics is a set of methods used to extract information from fixed samples and make inferences about the underlying distribution of the random variable.



# Random Variable

- ▶ Suppose we want to know about the education levels of people in California.
- ▶ Different people have different levels of education. The education level of a randomly selected person in the population is uncertain.
- ▶ The education level is a **random variable** with a distribution that describes the probability that a randomly selected person has an education level in certain range, e.g. 80% have a high school diploma, 30% have a college degree.
- ▶ In general, we do not know the exact distribution of the random variable. Statistics is a set of methods used to extract information from fixed samples and make inferences about the underlying distribution of the random variable.

# Random Variable

- ▶ Suppose we want to know about the education levels of people in California.
- ▶ Different people have different levels of education. The education level of a randomly selected person in the population is uncertain.
- ▶ The education level is a **random variable** with a distribution that describes the probability that a randomly selected person has an education level in certain range, e.g. 80% have a high school diploma, 30% have a college degree.
- ▶ In general, we do not know the exact distribution of the random variable. Statistics is a set of methods used to extract information from fixed samples and make inferences about the underlying distribution of the random variable.

# Random Variable

- ▶ Suppose we want to know about the education levels of people in California.
- ▶ Different people have different levels of education. The education level of a randomly selected person in the population is uncertain.
- ▶ The education level is a **random variable** with a distribution that describes the probability that a randomly selected person has an education level in certain range, e.g. 80% have a high school diploma, 30% have a college degree.
- ▶ In general, we do not know the exact distribution of the random variable. Statistics is a set of methods used to extract information from fixed samples and make inferences about the underlying distribution of the random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.

Flipping a coin and rolling a die are represented by discrete random variables.

Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.

Flipping a coin and rolling a die are represented by discrete random variables.

Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.

Flipping a coin and rolling a die are represented by discrete random variables.

Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.

▶ **Discrete random variable:** The sample space of  $X$  is countable.

▶ **Continuous random variable:** The sample space of  $X$  is uncountable.

Flipping a coin and rolling a die are represented by discrete random variables.

Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.



# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.
  - Flipping a coin and rolling a die are represented by discrete random variables.
  - Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.
  - Flipping a coin and rolling a die are represented by discrete random variables.
  - Racing the 100m dash is represented by a continuous random variable.

# Sample Space

- ▶ Let  $X$  denote a random variable. The **sample space**  $\Omega$  is the set of all possible values of  $X$ .
  - Coin flipping: The sample space is  $\{H, T\}$ .
  - Rolling a die: The sample space is  $\{1, 2, 3, 4, 5, 6\}$ .
  - Racing the 100m dash at a competition: The sample space may be the range  $[9.5, 11]$ , measured in seconds.
- ▶ **Discrete random variable:** The sample space of  $X$  is **countable**.
- ▶ **Continuous random variable:** The sample space of  $X$  is **uncountable**.
  - Flipping a coin and rolling a die are represented by discrete random variables.
  - Racing the 100m dash is represented by a continuous random variable.

# Probability

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  – an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_{i=1}^n \Pr(E_i)$ .

# Probability

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - ▶  $\Pr(\Omega) = 1$
  - ▶  $\Pr(\emptyset) = 0$
  - ▶  $0 \leq \Pr(X) \leq 1$
  - ▶ If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_{i=1}^n \Pr(E_i)$ .

# Probability

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - ▶  $\Pr(\Omega) = 1$
  - ▶  $\Pr(\emptyset) = 0$
  - ▶  $0 \leq \Pr(X) \leq 1$
  - ▶ If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_{i=1}^n \Pr(E_i)$

# Probability

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:

$$\Pr(\emptyset) = 0$$

$$\Pr(\Omega) = 1$$

$$0 \leq \Pr(X) \leq 1$$

$$\Pr(E_1 \cup E_2 \cup \dots \cup E_n) = \Pr(E_1) + \Pr(E_2) + \dots + \Pr(E_n) \text{ if } E_1, E_2, \dots, E_n \text{ are pairwise disjoint. Then } \Pr(\cup_{i=1}^n E_i) = \sum_{i=1}^n \Pr(E_i)$$

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_i \Pr(E_i)$ .



- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  – an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_i \Pr(E_i)$ .

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_i \Pr(E_i)$ .

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_i \Pr(E_i)$ .

- ▶ Let  $\Omega$  denote the sample space of random variable  $X$ .
- ▶ Let  $E_i$  denote any subset of the sample space  $\Omega$  — an event.
- ▶ We are interested in the probability  $\Pr(X)$  that  $X$  takes values in  $E_i \in \Omega$ .
- ▶ The distribution of the random variable  $X$  is a map from  $\cup_{i=1}^n E_i$  onto  $[0, 1]$ .
- ▶ **Axioms of probability:** The probability  $\Pr(X)$  satisfies:
  - $\Pr(\Omega) = 1$
  - $\Pr(\emptyset) = 0$
  - $0 \leq \Pr(X) \leq 1$
  - If  $E_1, E_2, \dots, E_n$  are pairwise disjoint, then  $\Pr(\cup_{i=1}^n E_i) = \sum_i \Pr(E_i)$ .

# Discrete Random Variables

- ▶ Let  $X$  be a **discrete random variable**. The distribution of  $X$  is described by the **probability mass function (pmf)**,  $p_X(X): E \subseteq \Omega \rightarrow [0, 1]$ .
- ▶ For each element  $x$  of the sample space,  $x \in \Omega$ , the probability mass function,  $p_X(\cdot)$ , describes the probability that  $X$  takes value  $x$ :  $p_X(x) = \Pr(X = x)$ .
- ▶ The pmf can be used to recover the probability that  $X$  takes values in any subset  $E$  of the sample space  $\Omega$ .

$$\Pr(E) = \sum_{x \in E} \Pr(X = x) = \sum_{x \in E} p_X(x)$$

# Discrete Random Variables

- ▶ Let  $X$  be a **discrete random variable**. The distribution of  $X$  is described by the **probability mass function (pmf)**,  $p_X(X): E \subseteq \Omega \rightarrow [0, 1]$ .
- ▶ For each element  $x$  of the sample space,  $x \in \Omega$ , the probability mass function,  $p_X(\cdot)$ , describes the probability that  $X$  takes value  $x$ :  $p_X(x) = \Pr(X = x)$ .
- ▶ The pmf can be used to recover the probability that  $X$  takes values in any subset  $E$  of the sample space  $\Omega$ .

$$\Pr(E) = \sum_{x \in E} \Pr(X = x) = \sum_{x \in E} p_X(x)$$

# Discrete Random Variables

- ▶ Let  $X$  be a **discrete random variable**. The distribution of  $X$  is described by the **probability mass function (pmf)**,  $p_X(X): E \subseteq \Omega \rightarrow [0, 1]$ .
- ▶ For each element  $x$  of the sample space,  $x \in \Omega$ , the probability mass function,  $p_X(\cdot)$ , describes the probability that  $X$  takes value  $x$ :  $p_X(x) = \Pr(X = x)$ .
- ▶ The pmf can be used to recover the probability that  $X$  takes values in any subset  $E$  of the sample space  $\Omega$ .

$$\Pr(E) = \sum_{x \in E} \Pr(X = x) = \sum_{x \in E} p_X(x)$$

# Discrete Random Variables

- ▶ Compute the probability of getting an value in one roll of a fair die.
- ▶ Let  $X$  denote the outcome of the die. The distribution of  $X$  has the following pmf:

$$p_X(x) = \begin{cases} \frac{1}{6} & \text{if } x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Use the pmf to compute  $\Pr(E)$  for  $E = \{2, 4, 6\}$ :

$$\begin{aligned} \Pr(E) &= \Pr(x \in \{2, 4, 6\}) = \sum_{x \in \{2, 4, 6\}} p_X(x) \\ &= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2} \end{aligned}$$



# Discrete Random Variables

- ▶ Compute the probability of getting an value in one roll of a fair die.
- ▶ Let  $X$  denote the outcome of the die. The distribution of  $X$  has the following pmf:

$$p_X(x) = \begin{cases} \frac{1}{6} & \text{if } x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Use the pmf to compute  $\Pr(E)$  for  $E = \{2, 4, 6\}$ :

$$\begin{aligned} \Pr(E) &= \Pr(x \in \{2, 4, 6\}) = \sum_{x \in \{2, 4, 6\}} p_X(x) \\ &= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2} \end{aligned}$$

# Discrete Random Variables

- ▶ Compute the probability of getting an value in one roll of a fair die.
- ▶ Let  $X$  denote the outcome of the die. The distribution of  $X$  has the following pmf:

$$p_X(x) = \begin{cases} \frac{1}{6} & \text{if } x \in \{1, 2, 3, 4, 5, 6\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Use the pmf to compute  $\Pr(E)$  for  $E = \{2, 4, 6\}$ :

$$\begin{aligned} \Pr(E) = \Pr(x \in \{2, 4, 6\}) &= \sum_{x \in \{2, 4, 6\}} p_X(x) \\ &= \frac{1}{6} + \frac{1}{6} + \frac{1}{6} = \frac{1}{2} \end{aligned}$$

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - ▶ If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - ▶ The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - ▶ What is the probability of drawing the number 2.1 on the interval of the real line  $(0, 4)$ ? Answer: 0.
  - ▶ What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number 2 from the interval  $(0, 1)$ ? Answer: 0.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
    - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - ▶ What is the probability of drawing a number  $x$  from the interval of the real line  $(a, b)$ ? Answer: 0.
  - ▶ What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number  $\pi$  from the subset of the real line  $[3, 4]$ ? Answer: zero.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number  $\pi$  from the subset of the real line  $[3, 4]$ ? Answer: zero.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.



# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number  $\pi$  from the subset of the real line  $[3, 4]$ ? Answer: zero.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number  $\pi$  from the subset of the real line  $[3, 4]$ ? Answer: zero.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables

- ▶ If  $X$  is a **continuous random variable**, the sample space  $\Omega$  is uncountable, so we cannot use a probability mass function to describe its distribution.
- ▶ If we attempted to assign a probability to each element in the sample space, we would be faced with one of two consequences:
  - If we assigned a finite probability to an uncountable subset of the sample space, the sum of the probabilities would tend to infinity.
  - The only way for the sum of probabilities to tend to 1 would be to assign a zero probability to each element of an uncountable subset of the sample space.
- ▶ Either way, it would not be useful and would lead to absurd calculations.
  - What is the probability of drawing the number  $\pi$  from the subset of the real line  $[3, 4]$ ? Answer: zero.
  - What is the probability of drawing any number from a subset of the real line? Answer: zero.
- ▶ We cannot use a pmf to describe the distribution of a continuous random variable.
- ▶ Instead, we reason in terms of non-overlapping dense subsets of the real line that are made arbitrarily small.

# Continuous Random Variables.

- ▶ If  $X$  is a **continuous random variable**, we use the probability density function (pdf),  $f_X(\cdot)$  to describe the distribution of  $X$ .
- ▶ The pdf is related to the probability measure  $\Pr$  via the following equation:

$$\Pr(a \leq X \leq b) = \int_a^b f_X(x) dx$$

- ▶ This identity can be used to calculate  $\Pr(E)$  for any set  $E \subseteq \Omega$ .

# Continuous Random Variables.

- ▶ If  $X$  is a **continuous random variable**, we use the probability density function (pdf),  $f_X(\cdot)$  to describe the distribution of  $X$ .
- ▶ The pdf is related to the probability measure  $\Pr$  via the following equation:

$$\Pr(a \leq X \leq b) = \int_a^b f_X(x) dx$$

- ▶ This identity can be used to calculate  $\Pr(E)$  for any set  $E \subseteq \Omega$ .

# Continuous Random Variables.

- ▶ If  $X$  is a **continuous random variable**, we use the probability density function (pdf),  $f_X(\cdot)$  to describe the distribution of  $X$ .
- ▶ The pdf is related to the probability measure  $\Pr$  via the following equation:

$$\Pr(a \leq X \leq b) = \int_a^b f_X(x) dx$$

- ▶ This identity can be used to calculate  $\Pr(E)$  for any set  $E \subseteq \Omega$ .

# Continuous Random Variables

- **Example:** Let  $X$  be a continuous random variable with pdf  $f_X$  given

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- This distribution is called the **uniform distribution** on  $[0, 1]$ .
- Calculate  $\Pr([0, 0.5])$ :

$$\begin{aligned} \Pr(X \in [0, 0.5]) &= \int_0^{0.5} f_X(x) \, dx \\ &= \int_0^{0.5} 1 \, dx \\ &= x \Big|_0^{0.5} \\ &= 0.5 - 0 = 0.5 \end{aligned}$$

# Continuous Random Variables

- **Example:** Let  $X$  be a continuous random variable with pdf  $f_X$  given

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- This distribution is called the **uniform distribution** on  $[0, 1]$ .

- Calculate  $\Pr([0, 0.5])$ :

$$\begin{aligned} \Pr(X \in [0, 0.5]) &= \int_0^{0.5} f_X(x) dx \\ &= \int_0^{0.5} 1 dx \\ &= x \Big|_0^{0.5} \\ &= 0.5 - 0 = 0.5 \end{aligned}$$



# Continuous Random Variables

- **Example:** Let  $X$  be a continuous random variable with pdf  $f_X$  given

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- This distribution is called the **uniform distribution** on  $[0, 1]$ .
- Calculate  $\Pr([0, 0.5])$ :

$$\begin{aligned} \Pr(X \in [0, 0.5]) &= \int_0^{0.5} f_X(x) dx \\ &= \int_0^{0.5} 1 dx \\ &= x \Big|_0^{0.5} \\ &= 0.5 - 0 = 0.5 \end{aligned}$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function ( $p(x)$ ).

$$Pr(X \in \{x\}) = p(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function ( $f(x)$ ).

$$Pr(X \in [a,b]) = \int_a^b f(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by a probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b p(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X = x) = p(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by a probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b p(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
  - The probability distribution is described by a probability density function (pdf).
- $$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$



## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$

## ► Random Variable:

- Describes an event whose outcome is unknown.
- To each outcome, associate a probability.

## ► Discrete Random Variable:

- Its sample space is countable.
- The probability distribution is described by a probability mass function (pmf).

$$\Pr(X \in \{x\}) = p_X(x)$$

## ► Continuous Random Variable:

- Its sample space is uncountable.
- The probability distribution is described by probability density function (pdf).

$$\Pr(X \in [a, b]) = \int_a^b f_X(x) dx$$

# Expectation

- ▶ The expectation can be interpreted as a generalization of the arithmetic mean.
- ▶ The weighted average of the random variable  $X$ , where the weights are the probability associated with each possible value of  $X$ .
- ▶ The expectation of  $X$  is denoted  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} x \cdot p_X(x)$$

$$\int_{\Omega} x \cdot f_X(x) dx$$

---

- ▶ Note that the difference between discrete and continuous is just summation vs. integral.
- ▶ In a population, the expectation may be denoted  $\mu_X$ . In a sample, it may be denoted  $\bar{X}$ .

# Expectation

- ▶ The expectation can be interpreted as a generalization of the arithmetic mean.
- ▶ The weighted average of the random variable  $X$ , where the weights are the probability associated with each possible value of  $X$ .
- ▶ The expectation of  $X$  is denoted  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} x \cdot p_X(x)$$

$$\int_{\Omega} x \cdot f_X(x) dx$$

---

- ▶ Note that the difference between discrete and continuous is just summation vs. integral.
- ▶ In a population, the expectation may be denoted  $\mu_X$ . In a sample, it may be denoted  $\bar{X}$ .

# Expectation

- ▶ The expectation can be interpreted as a generalization of the arithmetic mean.
- ▶ The weighted average of the random variable  $X$ , where the weights are the probability associated with each possible value of  $X$ .
- ▶ The expectation of  $X$  is denoted  $E[X]$ .

| Discrete R.V                         | Continuous R.V                    |
|--------------------------------------|-----------------------------------|
| $\sum_{x \in \Omega} x \cdot p_X(x)$ | $\int_{\Omega} x \cdot f_X(x) dx$ |

- ▶ Note that the difference between discrete and continuous is just summation vs. integral.
- ▶ In a population, the expectation may be denoted  $\mu_X$ . In a sample, it may be denoted  $\bar{X}$ .

# Expectation

- ▶ The expectation can be interpreted as a generalization of the arithmetic mean.
- ▶ The weighted average of the random variable  $X$ , where the weights are the probability associated with each possible value of  $X$ .
- ▶ The expectation of  $X$  is denoted  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} x \cdot p_X(x)$$

$$\int_{\Omega} x \cdot f_X(x) dx$$

---

- ▶ Note that the difference between discrete and continuous is just summation vs. integral.
- ▶ In a population, the expectation may be denoted  $\mu_X$ . In a sample, it may be denoted  $\bar{X}$ .

# Expectation

- ▶ The expectation can be interpreted as a generalization of the arithmetic mean.
- ▶ The weighted average of the random variable  $X$ , where the weights are the probability associated with each possible value of  $X$ .
- ▶ The expectation of  $X$  is denoted  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} x \cdot p_X(x)$$

$$\int_{\Omega} x \cdot f_X(x) dx$$

---

- ▶ Note that the difference between discrete and continuous is just summation vs. integral.
- ▶ In a population, the expectation may be denoted  $\mu_X$ . In a sample, it may be denoted  $\bar{X}$ .

# Expectation

► Consider a lottery that pays:

- \$100 with probability  $1/2$
- \$400 with probability  $1/4$
- \$0 with probability  $1/4$

► The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

► Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

► A gambler can expect to win \$150 by playing this lottery.



# Expectation

► Consider a lottery that pays:

- \$100 with probability  $1/2$
- \$400 with probability  $1/4$
- \$0 with probability  $1/4$

► The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

► Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

► A gambler can expect to win \$150 by playing this lottery.

# Expectation

► Consider a lottery that pays:

- \$100 with probability  $1/2$
- \$400 with probability  $1/4$
- \$0 with probability  $1/4$

► The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

► Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

► A gambler can expect to win \$150 by playing this lottery.

# Expectation

► Consider a lottery that pays:

- \$100 with probability  $1/2$
- \$400 with probability  $1/4$
- \$0 with probability  $1/4$

► The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

► Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

► A gambler can expect to win \$150 by playing this lottery.

# Expectation

- ▶ Consider a lottery that pays:
  - \$100 with probability  $1/2$
  - \$400 with probability  $1/4$
  - \$0 with probability  $1/4$
- ▶ The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

- ▶ A gambler can expect to win \$150 by playing this lottery.

# Expectation

- ▶ Consider a lottery that pays:
  - \$100 with probability  $1/2$
  - \$400 with probability  $1/4$
  - \$0 with probability  $1/4$
- ▶ The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

- ▶ A gambler can expect to win \$150 by playing this lottery.

# Expectation

- ▶ Consider a lottery that pays:
  - \$100 with probability  $1/2$
  - \$400 with probability  $1/4$
  - \$0 with probability  $1/4$
- ▶ The payout of this lottery can be represented by a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} 1/2 & \text{if } x = \{100\} \\ 1/4 & \text{if } x \in \{0, 400\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expected value of this lottery  $E[X]$  – The mean value of the lottery prize.

$$\begin{aligned} E[X] &= \sum_{x \in \{0, 100, 400\}} x \cdot p_X(x) \\ &= 0 \cdot \frac{1}{4} + 100 \cdot \frac{1}{2} + 400 \cdot \frac{1}{4} = 50 + 100 = 150 \end{aligned}$$

- ▶ A gambler can expect to win \$150 by playing this lottery.

# Expectation

- ▶ Let  $X$  be continuously distributed with sample space  $[0, 1]$  and pdf:

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expectation:

$$\begin{aligned} E[X] &= \int_{\Omega} x \cdot f_X(x) dx \\ &= \int_0^1 x \cdot 1 dx \\ &= \frac{x^2}{2} \Big|_0^1 = \frac{1^2}{2} - \frac{0^2}{2} = \frac{1}{2} \end{aligned}$$

- ▶ The expected value of  $X$  is 0.5. You can expect to win half a dollar from playing the lottery many, many times.

# Expectation

- ▶ Let  $X$  be continuously distributed with sample space  $[0, 1]$  and pdf:

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expectation:

$$\begin{aligned} \mathbb{E}[X] &= \int_{\Omega} x \cdot f_X(x) dx \\ &= \int_0^1 x \cdot 1 dx \\ &= \frac{x^2}{2} \Big|_0^1 = \frac{1^2}{2} - \frac{0^2}{2} = \frac{1}{2} \end{aligned}$$

- ▶ The expected value of  $X$  is 0.5. You can expect to win half a dollar from playing the lottery many, many times.



# Expectation

- ▶ Let  $X$  be continuously distributed with sample space  $[0, 1]$  and pdf:

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the expectation:

$$\begin{aligned} \mathbb{E}[X] &= \int_{\Omega} x \cdot f_X(x) dx \\ &= \int_0^1 x \cdot 1 dx \\ &= \frac{x^2}{2} \Big|_0^1 = \frac{1^2}{2} - \frac{0^2}{2} = \frac{1}{2} \end{aligned}$$

- ▶ The expected value of  $X$  is 0.5. You can expect to win half a dollar from playing the lottery many, many times.

# Expectation

- ▶ Consider the mean of any function  $g(X)$ , denoted  $E[g(X)]$ .
- ▶  $g(X)$  is a random variable with distribution derived from  $X$ .
- ▶ The formula for calculating  $E[g(X)]$  is basically the same as for calculating  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} g(x) \cdot p_X(x)$$

$$\int_{\Omega} g(x) \cdot f_X(x) dx$$

- 
- ▶ The pmf/pdf is multiplied by  $g(x)$  instead of just  $x$ .

# Expectation

- ▶ Consider the mean of any function  $g(X)$ , denoted  $E[g(X)]$ .
- ▶  $g(X)$  is a random variable with distribution derived from  $X$ .
- ▶ The formula for calculating  $E[g(X)]$  is basically the same as for calculating  $E[X]$ .

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} g(x) \cdot p_X(x)$$

$$\int_{\Omega} g(x) \cdot f_X(x) dx$$

- ▶ The pmf/pdf is multiplied by  $g(x)$  instead of just  $x$ .

# Expectation

- ▶ Consider the mean of any function  $g(X)$ , denoted  $E[g(X)]$ .
- ▶  $g(X)$  is a random variable with distribution derived from  $X$ .
- ▶ The formula for calculating  $E[g(X)]$  is basically the same as for calculating  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} g(x) \cdot p_X(x)$$

$$\int_{\Omega} g(x) \cdot f_X(x) dx$$

---

- ▶ The pmf/pdf is multiplied by  $g(x)$  instead of just  $x$ .

# Expectation

- ▶ Consider the mean of any function  $g(X)$ , denoted  $E[g(X)]$ .
- ▶  $g(X)$  is a random variable with distribution derived from  $X$ .
- ▶ The formula for calculating  $E[g(X)]$  is basically the same as for calculating  $E[X]$ .

---

Discrete R.V

Continuous R.V

$$\sum_{x \in \Omega} g(x) \cdot p_X(x)$$

$$\int_{\Omega} g(x) \cdot f_X(x) dx$$

---

- ▶ The pmf/pdf is multiplied by  $g(x)$  instead of just  $x$ .

► **Linearity of Expectations:**

$$\mathbb{E}[ag(X) + bh(X)] = a \mathbb{E}[g(X)] + b \mathbb{E}[h(X)]$$

for any  $a, b \in \mathbb{R}$ .

- This property applies to two different random variables  $X$  and  $Y$ :

$$\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$$

- ▶ **Linearity of Expectations:**

$$\mathbb{E}[ag(X) + bh(X)] = a \mathbb{E}[g(X)] + b \mathbb{E}[h(X)]$$

for any  $a, b \in \mathbb{R}$ .

- ▶ This property applies to two different random variables  $X$  and  $Y$ :

$$\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$$

# Expectation

- Suppose that  $X$  has pdf

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- Calculate the **second moment** of  $X$ ,  $E[X^2]$ :

$$\begin{aligned} E[X^2] &= \int_{\Omega} x^2 \cdot f_X(x) dx \\ &= \int_0^1 x^2 \cdot 1 dx \\ &= \left. \frac{x^3}{3} \right|_0^1 = \frac{1^3}{3} - \frac{0^3}{3} = \frac{1}{3} \end{aligned}$$



# Expectation

- ▶ Suppose that  $X$  has pdf

$$f_X(x) = \begin{cases} 1 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Calculate the **second moment** of  $X$ ,  $E[X^2]$ :

$$\begin{aligned} E[X^2] &= \int_{\Omega} x^2 \cdot f_X(x) dx \\ &= \int_0^1 x^2 \cdot 1 dx \\ &= \left. \frac{x^3}{3} \right|_0^1 = \frac{1^3}{3} - \frac{0^3}{3} = \frac{1}{3} \end{aligned}$$

# Variance

- ▶ The variance of a random variable  $X$  is a measure of the spread of the random variable  $X$  — a measure of how far on average  $X$  is from its mean.

$$\text{var}(X) = \mathbb{E}[(X - \mathbb{E}[X])^2]$$

- ▶ Using linearity of the expectation the expression above can be simplified:

$$\begin{aligned}\text{var}(X) &= \mathbb{E}[X^2 - 2X\mathbb{E}[X] + (\mathbb{E}[X])^2] \\ &= \mathbb{E}[X^2] - 2\mathbb{E}[X]\mathbb{E}[X] + \mathbb{E}[X]^2 \\ &= \mathbb{E}[X^2] - (\mathbb{E}[X])^2\end{aligned}$$

- ▶ From the definition, the variance is always positive  $\text{var}(X) \geq 0$ .
- ▶ The last expression is often convenient.

# Variance

- ▶ The variance of a random variable  $X$  is a measure of the spread of the random variable  $X$  — a measure of how far on average  $X$  is from its mean.

$$\text{var}(X) = \text{E}[(X - \text{E}[X])^2]$$

- ▶ Using linearity of the expectation the expression above can be simplified:

$$\begin{aligned}\text{var}(X) &= \text{E}[X^2 - 2X \text{E}[X] + (\text{E}[X])^2] \\ &= \text{E}[X^2] - 2 \text{E}[X] \text{E}[X] + \text{E}[X]^2 \\ &= \text{E}[X^2] - (\text{E}[X])^2\end{aligned}$$

- ▶ From the definition, the variance is always positive  $\text{var}(X) \geq 0$ .
- ▶ The last expression is often convenient.

# Variance

- ▶ The variance of a random variable  $X$  is a measure of the spread of the random variable  $X$  – a measure of how far on average  $X$  is from its mean.

$$\text{var}(X) = \text{E}[(X - \text{E}[X])^2]$$

- ▶ Using linearity of the expectation the expression above can be simplified:

$$\begin{aligned}\text{var}(X) &= \text{E}[X^2 - 2X \text{E}[X] + (\text{E}[X])^2] \\ &= \text{E}[X^2] - 2 \text{E}[X] \text{E}[X] + \text{E}[X]^2 \\ &= \text{E}[X^2] - (\text{E}[X])^2\end{aligned}$$

- ▶ From the definition, the variance is always positive  $\text{var}(X) \geq 0$ .

- ▶ The last expression is often convenient.

# Variance

- ▶ The variance of a random variable  $X$  is a measure of the spread of the random variable  $X$  — a measure of how far on average  $X$  is from its mean.

$$\text{var}(X) = \text{E}[(X - \text{E}[X])^2]$$

- ▶ Using linearity of the expectation the expression above can be simplified:

$$\begin{aligned}\text{var}(X) &= \text{E}[X^2 - 2X \text{E}[X] + (\text{E}[X])^2] \\ &= \text{E}[X^2] - 2 \text{E}[X] \text{E}[X] + \text{E}[X]^2 \\ &= \text{E}[X^2] - (\text{E}[X])^2\end{aligned}$$

- ▶ From the definition, the variance is always positive  $\text{var}(X) \geq 0$ .
- ▶ The last expression is often convenient.

# Variance

- ▶ The linearity of the expectation and the formula  $\text{var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$  imply several convenient properties of the variance.

- ▶ For any constants  $a, b \in \mathbb{R}$  and any random variable  $X$ , we have:

$$\text{var}(X + a) = \text{var}(X)$$

$$\text{var}(aX) = a^2 \text{var}(X)$$

- ▶ Combining these gives:

$$\text{var}(aX + b) = a^2 \text{var}(X)$$

# Variance

- ▶ The linearity of the expectation and the formula  $\text{var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$  imply several convenient properties of the variance.
- ▶ For any constants  $a, b \in \mathbb{R}$  and any random variable  $X$ , we have:

$$\text{var}(X + a) = \text{var}(X)$$

$$\text{var}(aX) = a^2 \text{var}(X)$$

- ▶ Combining these gives:

$$\text{var}(aX + b) = a^2 \text{var}(X)$$

# Variance

- ▶ The linearity of the expectation and the formula  $\text{var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$  imply several convenient properties of the variance.
- ▶ For any constants  $a, b \in \mathbb{R}$  and any random variable  $X$ , we have:

$$\text{var}(X + a) = \text{var}(X)$$

$$\text{var}(aX) = a^2 \text{var}(X)$$


- ▶ Combining these gives:

$$\text{var}(aX + b) = a^2 \text{var}(X)$$



# Variance

Changes in the mean  $\mu$  do not affect the variance  $\sigma$ :



# Variance

- ▶ Suppose we have a lottery that pays \$200 with probability  $1/2$  and nothing otherwise. The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The expected payout of this lottery is  $E[X] = \$100$ .
- ▶ How much we can expect our winnings to deviate from the expected value?
- ▶ This can be computed directly:

$$\text{var}(X) = E[(X - E[X])^2]$$

or indirectly:

$$\text{var}(X) = E[X^2] - (E[X])^2$$

# Variance

- ▶ Suppose we have a lottery that pays \$200 with probability  $1/2$  and nothing otherwise. The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The expected payout of this lottery is  $E[X] = \$100$ .
- ▶ How much we can expect our winnings to deviate from the expected value?
- ▶ This can be computed directly:

$$\text{var}(X) = E[(X - E[X])^2]$$

or indirectly:

$$\text{var}(X) = E[X^2] - (E[X])^2$$

# Variance

- ▶ Suppose we have a lottery that pays \$200 with probability  $1/2$  and nothing otherwise. The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The expected payout of this lottery is  $E[X] = \$100$ .
- ▶ How much we can expect our winnings to deviate from the expected value?
- ▶ This can be computed directly:

$$\text{var}(X) = E[(X - E[X])^2]$$

or indirectly:

$$\text{var}(X) = E[X^2] - (E[X])^2$$

# Variance

- ▶ Suppose we have a lottery that pays \$200 with probability  $1/2$  and nothing otherwise. The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ The expected payout of this lottery is  $E[X] = \$100$ .
- ▶ How much we can expect our winnings to deviate from the expected value?
- ▶ This can be computed directly:

$$\text{var}(X) = E[(X - E[X])^2]$$

or indirectly:

$$\text{var}(X) = E[X^2] - (E[X])^2$$

- ▶ The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Direct Calculation:

$$\begin{aligned} \text{var}(X) &= E[(X - E[X])^2] \\ &= E[(X - 100)^2] \\ &= \sum_{x \in \{0, 200\}} (x - 100)^2 \cdot p_X(x) \\ &= (0 - 100)^2 \cdot \frac{1}{2} + (200 - 100)^2 \cdot \frac{1}{2} \\ &= 10,000 \end{aligned}$$

- ▶ The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Direct Calculation:

$$\begin{aligned} \text{var}(X) &= E[(X - E[X])^2] \\ &= E[(X - 100)^2] \\ &= \sum_{x \in \{0, 200\}} (x - 100)^2 \cdot p_X(x) \\ &= (0 - 100)^2 \cdot \frac{1}{2} + (200 - 100)^2 \cdot \frac{1}{2} \\ &= 10,000 \end{aligned}$$

- ▶ The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Indirect Calculation:

$$\begin{aligned} \text{var}(X) &= E[X^2] - (E[X])^2 \\ &= \sum_{x \in \{0, 200\}} x^2 \cdot p_X(x) - (100)^2 \\ &= 0^2 \cdot \frac{1}{2} + 200^2 \cdot \frac{1}{2} - 100^2 \\ &= 10,000 \end{aligned}$$



- ▶ The payout of this lottery is a random variable  $X$  with pmf:

$$p_X(x) = \begin{cases} \frac{1}{2} & \text{if } x \in \{0, 200\} \\ 0 & \text{otherwise} \end{cases}$$

- ▶ Indirect Calculation:

$$\begin{aligned} \text{var}(X) &= E[X^2] - (E[X])^2 \\ &= \sum_{x \in \{0, 200\}} x^2 \cdot p_X(x) - (100)^2 \\ &= 0^2 \cdot \frac{1}{2} + 200^2 \cdot \frac{1}{2} - 100^2 \\ &= 10,000 \end{aligned}$$

# Standard Deviation

- ▶ **Standard deviation of  $X$ :**

$$\sigma_X = \sqrt{\sigma_X^2} = \sqrt{\text{var}(X)}$$

- ▶ The standard deviation is the square root of the variance.

# Standard Deviation

- ▶ **Standard deviation of  $X$ :**

$$\sigma_X = \sqrt{\sigma_X^2} = \sqrt{\text{var}(X)}$$

- ▶ The standard deviation is the square root of the variance.

## Multiple Random Variables

# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome
- ▶ Note that:
  - not everyone has the same education/income
  - not everyone who takes a medicine will have the same health outcome

# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome
- ▶ Note that:
  - not everyone has the same education/income
  - not everyone who takes a new drug will have the same health outcome

# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome

▶ Note that:

▶ We don't always want to know the joint distribution of two variables

▶ We don't always want to know the joint distribution of two variables

# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome
- ▶ Note that:
  - not everyone has the same education/income
  - not everyone who takes a medicine will have the same health outcome



# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome
- ▶ Note that:
  - not everyone has the same education/income
  - not everyone who takes a medicine will have the same health outcome

# Bivariate Random Variables

- ▶ Consider the relationship between two random variables.
  - We care about the relationship between education and income
  - We care about the relationship between consumption of a medicine and a health outcome
- ▶ Note that:
  - not everyone has the same education/income
  - not everyone who takes a medicine will have the same health outcome

# Bivariate Random Variables

- ▶ Let  $(X, Y)$  be a pair of joint random variables. Let  $\Omega$  denote the sample space and  $\Pr(\cdot)$  a probability measure defined on subsets of the sample space  $E \subseteq \Omega$ , that is  $\Pr(\cdot): \Omega \rightarrow [0, 1]$ .
- ▶ Example: Let  $X$  denote income and  $Y$  denote age. The probability that a randomly selected person from the population has an income between \$0 and \$100,000 and is between 40 and 42 years old is denoted:

$$\Pr(\{0 \leq X \leq 100,000, 40 \leq Y \leq 42\})$$

- ▶ For two random variables,  $X, Y$ , we can represent the probability mass function (pmf) as a table.

## Bivariate Random Variables

- ▶ Let  $(X, Y)$  be a pair of joint random variables. Let  $\Omega$  denote the sample space and  $\Pr(\cdot)$  a probability measure defined on subsets of the sample space  $E \subseteq \Omega$ , that is  $\Pr(\cdot): \Omega \rightarrow [0, 1]$ .
- ▶ Example: Let  $X$  denote income and  $Y$  denote age. The probability that a randomly selected person from the population has an income between \$0 and \$100,000 and is between 40 and 42 years old is denoted:

$$\Pr(\{0 \leq X \leq 100,000, 40 \leq Y \leq 42\})$$

- ▶ For two random variables,  $X, Y$ , we can represent the probability mass function (pmf) as a table.

# Bivariate Random Variables

- ▶ Let  $(X, Y)$  be a pair of joint random variables. Let  $\Omega$  denote the sample space and  $\Pr(\cdot)$  a probability measure defined on subsets of the sample space  $E \subseteq \Omega$ , that is  $\Pr(\cdot): \Omega \rightarrow [0, 1]$ .
- ▶ Example: Let  $X$  denote income and  $Y$  denote age. The probability that a randomly selected person from the population has an income between \$0 and \$100,000 and is between 40 and 42 years old is denoted:

$$\Pr(\{0 \leq X \leq 100,000, 40 \leq Y \leq 42\})$$

- ▶ For two random variables,  $X, Y$ , we can represent the probability mass function (pmf) as a table.

## Discrete Random Variables

- ▶ Let  $X$  be a random variable that describes whether a person gets 4 hours of sleep a night; 8 hours a sleep a night; or 12 hours of sleep a night. Let  $Y$  be a random variable that describes whether a person drinks 1 or 2 cups of coffee a day.
- ▶ The joint pmf of  $X$  and  $Y$  can be described with the table below

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- ▶ The probability that a randomly selected person gets 8 hours of sleep and drinks 1 cup of coffee a day is 1/3.
- ▶ Exercise: What is the probability that a randomly selected person gets 8 hours of sleep?

## Discrete Random Variables

- ▶ Let  $X$  be a random variable that describes whether a person gets 4 hours of sleep a night; 8 hours a sleep a night; or 12 hours of sleep a night. Let  $Y$  be a random variable that describes whether a person drinks 1 or 2 cups of coffee a day.
- ▶ The joint pmf of  $X$  and  $Y$  can be described with the table below

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- ▶ The probability that a randomly selected person gets 8 hours of sleep and drinks 1 cup of coffee a day is 1/3.
- ▶ Exercise: What is the probability that a randomly selected person gets 8 hours of sleep?

## Discrete Random Variables

- ▶ Let  $X$  be a random variable that describes whether a person gets 4 hours of sleep a night; 8 hours a sleep a night; or 12 hours of sleep a night. Let  $Y$  be a random variable that describes whether a person drinks 1 or 2 cups of coffee a day.
- ▶ The joint pmf of  $X$  and  $Y$  can be described with the table below

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- ▶ The probability that a randomly selected person gets 8 hours of sleep and drinks 1 cup of coffee a day is  $1/3$ .
- ▶ Exercise: What is the probability that a randomly selected person gets 8 hours of sleep?



## Discrete Random Variables

- ▶ Let  $X$  be a random variable that describes whether a person gets 4 hours of sleep a night; 8 hours a sleep a night; or 12 hours of sleep a night. Let  $Y$  be a random variable that describes whether a person drinks 1 or 2 cups of coffee a day.
- ▶ The joint pmf of  $X$  and  $Y$  can be described with the table below

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- ▶ The probability that a randomly selected person gets 8 hours of sleep and drinks 1 cup of coffee a day is 1/3.
- ▶ Exercise: What is the probability that a randomly selected person gets 8 hours of sleep?

# Continuous Random Variables

- ▶ As with single continuous random variables, the distribution of a continuous random variable is defined by a probability density function,  $f_{XY}(x, y)$ .
- ▶ As before, the joint pdf will be related to the joint probability measure  $\Pr(\cdot)$

$$\Pr(\{a \leq X \leq b, c \leq Y \leq d\}) = \int_a^b \int_c^d f_{XY}(x, y) dy dx$$

# Continuous Random Variables

- ▶ As with single continuous random variables, the distribution of a continuous random variable is defined by a probability density function,  $f_{XY}(x, y)$ .
- ▶ As before, the joint pdf will be related to the joint probability measure  $\Pr(\cdot)$

$$\Pr(\{a \leq X \leq b, c \leq Y \leq d\}) = \int_a^b \int_c^d f_{XY}(x, y) dy dx$$

## Continuous Random Variables

- Consider two sprinters in the 100m dash. Let  $X$  denote the finish time of the favorite competitor and  $Y$  denote the finish time of the underdog competitor. Suppose their times follow the following joint pdf:

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- Calculate the probability that the favorite competitor runs faster than 10 seconds and that the underdog competitor runs faster than 10.5 seconds,  $\Pr(\{X \leq 10, Y \leq 10.5\})$ .

$$\begin{aligned} \Pr(\{X \leq 10, Y \leq 10.5\}) &= \int_{9.5}^{10} \int_{10}^{10.5} f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10} \int_{10}^{10.5} 1 dy dx \\ &= \int_{9.5}^{10} 0.5 dx \\ &= 0.5 \times 0.5 \\ &= 0.25 \end{aligned}$$

## Continuous Random Variables

- Consider two sprinters in the 100m dash. Let  $X$  denote the finish time of the favorite competitor and  $Y$  denote the finish time of the underdog competitor. Suppose their times follow the following joint pdf:

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- Calculate the probability that the favorite competitor runs faster than 10 seconds and that the underdog competitor runs faster than 10.5 seconds,  $\Pr(\{X \leq 10, Y \leq 10.5\})$ .

$$\begin{aligned} \Pr(\{X \leq 10, Y \leq 10.5\}) &= \int_{9.5}^{10} \int_{10}^{10.5} f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10} \int_{10}^{10.5} 1 dy dx \\ &= \int_{9.5}^{10} 0.5 dx \\ &= 0.5 \times 0.5 \\ &= 0.25 \end{aligned}$$

# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .

- ▶ Examples of useful functions  $g(x, y)$ :

- Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average distance between  $X$  and  $Y$ :

$$g(x, y) = |x - y| \implies E[g(X, Y)] = E[|X - Y|]$$

- Indicator of a event:

$$g(x, y) = \mathbb{I}(x \leq y) \implies E[g(X, Y)] = P(X \leq Y)$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$

# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .
- ▶ Examples of useful functions  $g(x, y)$ :

- Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average difference between  $X$  and  $Y$ :

$$g(x, y) = x - y \implies E[g(X, Y)] = E[X - Y]$$

- Range of interest:

$$g(x, y) = \mathbb{1}\{x \leq a, y \leq b\} \implies E[g(X, Y)] = \Pr(\{X \leq a, Y \leq b\})$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$

# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .
- ▶ Examples of useful functions  $g(x, y)$ :
  - Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average difference between  $X$  and  $Y$ :

$$g(x, y) = x - y \implies E[g(X, Y)] = E[X - Y]$$

- Range of interest:

$$g(x, y) = \mathbb{1}\{x \leq a, y \leq b\} \implies E[g(X, Y)] = \Pr(\{X \leq a, Y \leq b\})$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$



# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .
- ▶ Examples of useful functions  $g(x, y)$ :
  - Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average difference between  $X$  and  $Y$ :

$$g(x, y) = x - y \implies E[g(X, Y)] = E[X - Y]$$

- Range of interest:

$$g(x, y) = \mathbb{1}\{x \leq a, y \leq b\} \implies E[g(X, Y)] = \Pr(\{X \leq a, Y \leq b\})$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$

# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .
- ▶ Examples of useful functions  $g(x, y)$ :

- Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average difference between  $X$  and  $Y$ :

$$g(x, y) = x - y \implies E[g(X, Y)] = E[X - Y]$$

- Range of interest:

$$g(x, y) = \mathbb{1}\{x \leq a, y \leq b\} \implies E[g(X, Y)] = \Pr(\{X \leq a, Y \leq b\})$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$

# Expectations

- ▶ Consider the average or expected value that some function,  $g(X, Y)$ , of the joint random variables. Calculate  $E[g(X, Y)]$ .
- ▶ Examples of useful functions  $g(x, y)$ :

- Expected value of  $X$ :

$$g(x, y) = x \implies E[g(X, Y)] = E[X]$$

- Average difference between  $X$  and  $Y$ :

$$g(x, y) = x - y \implies E[g(X, Y)] = E[X - Y]$$

- Range of interest:

$$g(x, y) = \mathbb{1}\{x \leq a, y \leq b\} \implies E[g(X, Y)] = \Pr(\{X \leq a, Y \leq b\})$$

- Covariance between  $X$  and  $Y$ :

$$g(x, y) = (x - \mu_X)(y - \mu_Y) \implies E[g(X, Y)] = \text{cov}(X, Y)$$

# Expectations

- ▶ The formula for calculating expected value is the same as before:

| Discrete R.V                               | Continuous R.V                           |
|--|--|
| $\sum_{a,b \in \Omega} g(a,b) p_{XY}(a,b)$ | $\int_X \int_Y g(a,b) f_{XY}(a,b) db da$ |

- ▶ The function is evaluated at each point in the outcome space and weighted by the probability associated with that outcome.
- ▶ By linearity of the expectation, for any two functions  $g(x,y)$  and  $h(x,y)$  and any  $a, b \in \mathbb{R}$ :

$$E[a \cdot g(X, Y) + b \cdot h(X, Y)] = a E[g(X, Y)] + b E[h(X, Y)]$$

- ▶ For instance,

$$E[aX + bY] = a E[X] + b E[Y]$$

# Expectations

- ▶ The formula for calculating expected value is the same as before:

|  |  |
|--|--|
| Discrete R.V                               | Continuous R.V                           |
| $\sum_{a,b \in \Omega} g(a,b) p_{XY}(a,b)$ | $\int_X \int_Y g(a,b) f_{XY}(a,b) db da$ |

- ▶ The function is evaluated at each point in the outcome space and weighted by the probability associated with that outcome.

- ▶ By linearity of the expectation, for any two functions  $g(x,y)$  and  $h(x,y)$  and any  $a, b \in \mathbb{R}$ :

$$E[a \cdot g(X, Y) + b \cdot h(X, Y)] = a E[g(X, Y)] + b E[h(X, Y)]$$

- ▶ For instance,

$$E[aX + bY] = a E[X] + b E[Y]$$

# Expectations

- ▶ The formula for calculating expected value is the same as before:

| Discrete R.V                               | Continuous R.V                           |
|--|--|
| $\sum_{a,b \in \Omega} g(a,b) p_{XY}(a,b)$ | $\int_X \int_Y g(a,b) f_{XY}(a,b) db da$ |

- ▶ The function is evaluated at each point in the outcome space and weighted by the probability associated with that outcome.
- ▶ By linearity of the expectation, for any two functions  $g(x, y)$  and  $h(x, y)$  and any  $a, b \in \mathbb{R}$ :

$$\mathbb{E}[a \cdot g(X, Y) + b \cdot h(X, Y)] = a \mathbb{E}[g(X, Y)] + b \mathbb{E}[h(X, Y)]$$

- ▶ For instance,

$$\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$$

# Expectations

- ▶ The formula for calculating expected value is the same as before:

| Discrete R.V                               | Continuous R.V                           |
|--|--|
| $\sum_{a,b \in \Omega} g(a,b) p_{XY}(a,b)$ | $\int_X \int_Y g(a,b) f_{XY}(a,b) db da$ |

- ▶ The function is evaluated at each point in the outcome space and weighted by the probability associated with that outcome.
- ▶ By linearity of the expectation, for any two functions  $g(x, y)$  and  $h(x, y)$  and any  $a, b \in \mathbb{R}$ :

$$\mathbb{E}[a \cdot g(X, Y) + b \cdot h(X, Y)] = a \mathbb{E}[g(X, Y)] + b \mathbb{E}[h(X, Y)]$$

- ▶ For instance,

$$\mathbb{E}[aX + bY] = a \mathbb{E}[X] + b \mathbb{E}[Y]$$

# Expectations

- Consider the 100m dash example again.

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5, 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- Calculate  $E[X - Y]$ , the expected difference in finishing times between the favorite competitor and the underdog:

$$\begin{aligned} E[X - Y] &= \int_{9.5}^{10.5} \int_{10}^{11} (x - y) f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} \int_{10}^{11} x dy dx - \int_{9.5}^{10.5} \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \left( y \Big|_{10}^{11} \right) dx - \int_{9.5}^{10.5} \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= 1 \cdot \frac{x^2}{2} \Big|_{9.5}^{10.5} - \frac{21}{2} \cdot x \Big|_{9.5}^{10.5} \\ &= -0.5 \end{aligned}$$



# Expectations

- Consider the 100m dash example again.

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5, 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- Calculate  $E[X - Y]$ , the expected difference in finishing times between the favorite competitor and the underdog:

$$\begin{aligned} E[X - Y] &= \int_{9.5}^{10.5} \int_{10}^{11} (x - y) f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} \int_{10}^{11} x dy dx - \int_{9.5}^{10.5} \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \left( y \Big|_{10}^{11} \right) dx - \int_{9.5}^{10.5} \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= 1 \cdot \frac{x^2}{2} \Big|_{9.5}^{10.5} - \frac{21}{2} \cdot x \Big|_{9.5}^{10.5} \\ &= -0.5 \end{aligned}$$

# Covariance

- ▶ The covariance measures how much the variables  $X$  and  $Y$  co-vary, i.e. how they vibrate together.

- ▶ The **covariance** between  $X$  and  $Y$  is:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ Simplify the expression:

$$\begin{aligned}\text{cov}(X, Y) &= E[XY - X E[Y] - Y E[X] + E[X] E[Y]] \\ &= E[XY] - E[X] E[Y] - E[Y] E[X] + E[X] E[Y] \\ &= E[XY] - E[X] E[Y]\end{aligned}$$

- ▶ The population covariance is often denoted  $\sigma_{XY}$ .
- ▶ The sample covariance is often denoted  $s_{XY}$ .

# Covariance

- ▶ The covariance measures how much the variables  $X$  and  $Y$  co-vary, i.e. how they vibrate together.
- ▶ The **covariance** between  $X$  and  $Y$  is:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ Simplify the expression:

$$\begin{aligned}\text{cov}(X, Y) &= E[XY - X E[Y] - Y E[X] + E[X] E[Y]] \\ &= E[XY] - E[X] E[Y] - E[Y] E[X] + E[X] E[Y] \\ &= E[XY] - E[X] E[Y]\end{aligned}$$

- ▶ The population covariance is often denoted  $\sigma_{XY}$ .
- ▶ The sample covariance is often denoted  $s_{XY}$ .

# Covariance

- ▶ The covariance measures how much the variables  $X$  and  $Y$  co-vary, i.e. how they vibrate together.
- ▶ The **covariance** between  $X$  and  $Y$  is:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ Simplify the expression:

$$\begin{aligned}\text{cov}(X, Y) &= E[XY - X E[Y] - Y E[X] + E[X] E[Y]] \\ &= E[XY] - E[X] E[Y] - E[Y] E[X] + E[X] E[Y] \\ &= E[XY] - E[X] E[Y]\end{aligned}$$

- ▶ The population covariance is often denoted  $\sigma_{XY}$ .
- ▶ The sample covariance is often denoted  $s_{XY}$ .

# Covariance

- ▶ The covariance measures how much the variables  $X$  and  $Y$  co-vary, i.e. how they vibrate together.
- ▶ The **covariance** between  $X$  and  $Y$  is:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ Simplify the expression:

$$\begin{aligned}\text{cov}(X, Y) &= E[XY - X E[Y] - Y E[X] + E[X] E[Y]] \\ &= E[XY] - E[X] E[Y] - E[Y] E[X] + E[X] E[Y] \\ &= E[XY] - E[X] E[Y]\end{aligned}$$

- ▶ The population covariance is often denoted  $\sigma_{XY}$ .
- ▶ The sample covariance is often denoted  $s_{XY}$ .

# Covariance

- ▶ The covariance measures how much the variables  $X$  and  $Y$  co-vary, i.e. how they vibrate together.
- ▶ The **covariance** between  $X$  and  $Y$  is:

$$\text{cov}(X, Y) = E[(X - E[X])(Y - E[Y])]$$

- ▶ Simplify the expression:

$$\begin{aligned}\text{cov}(X, Y) &= E[XY - X E[Y] - Y E[X] + E[X] E[Y]] \\ &= E[XY] - E[X] E[Y] - E[Y] E[X] + E[X] E[Y] \\ &= E[XY] - E[X] E[Y]\end{aligned}$$

- ▶ The population covariance is often denoted  $\sigma_{XY}$ .
- ▶ The sample covariance is often denoted  $s_{XY}$ .

- Covariance between  $X$  and  $Y$ :

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- We know  $E[X] = 10$  and  $E[Y] = 10.5$ , so calculate  $E[XY]$ :

$$\begin{aligned} E[XY] &= \int_{9.5}^{10.5} \int_{10}^{11} xy f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} x \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \cdot \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= \frac{21}{2} \cdot \left( \frac{x^2}{2} \Big|_{9.5}^{10.5} \right) = 105 \end{aligned}$$

- Thus,  $\text{cov}(X, Y) = E[XY] - E[X] E[Y] = 105 - 10 \cdot 10.5 = 105 - 105 = 0$ .
- There is no measurable association between competitors  $A$  and  $B$ .

# Covariance

- Covariance between  $X$  and  $Y$ :

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- We know  $E[X] = 10$  and  $E[Y] = 10.5$ , so calculate  $E[XY]$ :

$$\begin{aligned} E[XY] &= \int_{9.5}^{10.5} \int_{10}^{11} xy f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} x \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \cdot \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= \frac{21}{2} \cdot \left( \frac{x^2}{2} \Big|_{9.5}^{10.5} \right) = 105 \end{aligned}$$

- Thus,  $\text{cov}(X, Y) = E[XY] - E[X] E[Y] = 105 - 10 \cdot 10.5 = 105 - 105 = 0$ .
- There is no measurable association between competitors  $A$  and  $B$ .



- Covariance between  $X$  and  $Y$ :

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- We know  $E[X] = 10$  and  $E[Y] = 10.5$ , so calculate  $E[XY]$ :

$$\begin{aligned} E[XY] &= \int_{9.5}^{10.5} \int_{10}^{11} xy f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} x \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \cdot \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= \frac{21}{2} \cdot \left( \frac{x^2}{2} \Big|_{9.5}^{10.5} \right) = 105 \end{aligned}$$

- Thus,  $\text{cov}(X, Y) = E[XY] - E[X] E[Y] = 105 - 10 \cdot 10.5 = 105 - 105 = 0$ .

► There is no measurable association between competitors  $A$  and  $B$ .

# Covariance

- Covariance between  $X$  and  $Y$ :

$$f_{XY}(x, y) = \begin{cases} 1 & \text{if } 9.5 \leq x \leq 10.5 \text{ or } 10 \leq y \leq 11 \\ 0 & \text{otherwise} \end{cases}$$

- We know  $E[X] = 10$  and  $E[Y] = 10.5$ , so calculate  $E[XY]$ :

$$\begin{aligned} E[XY] &= \int_{9.5}^{10.5} \int_{10}^{11} xy f_{XY}(x, y) dy dx \\ &= \int_{9.5}^{10.5} x \int_{10}^{11} y dy dx \\ &= \int_{9.5}^{10.5} x \cdot \left( \frac{y^2}{2} \Big|_{10}^{11} \right) dx \\ &= \frac{21}{2} \cdot \left( \frac{x^2}{2} \Big|_{9.5}^{10.5} \right) = 105 \end{aligned}$$

- Thus,  $\text{cov}(X, Y) = E[XY] - E[X] E[Y] = 105 - 10 \cdot 10.5 = 105 - 105 = 0$ .
- There is no measurable association between competitors  $A$  and  $B$ .

- Important properties of the covariance follow from  $\text{cov}(X, Y) = E[XY] - E[X]E[Y]$ :

1. **Linearity:**  $\text{cov}(aX, bY) = ab \text{cov}(X, Y)$

$$\text{cov}(aX, bY) = E[(aX)(bY)] - E[aX]E[bY] = ab(E[XY] - E[X]E[Y])$$

2. **Symmetry:**  $\text{cov}(X, Y) = \text{cov}(Y, X)$  and  $\text{cov}(X, X) = \text{var}(X)$

$$\text{cov}(X, X) = E[XX] - E[X]E[X] = E[X^2] - (E[X])^2 = \text{var}(X)$$

3. **Addition Rule:**  $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)$

$$\begin{aligned}\text{var}(X + Y) &= E[(X + Y)^2] - E[(X + Y)]^2 \\&= E[X^2 + 2XY + Y^2] - (E[X] + E[Y])^2 \\&= \underbrace{E[X^2]}_{\text{var}(X)} + 2 \underbrace{E[XY]}_{\text{cov}(X, Y)} + \underbrace{E[Y^2]}_{\text{var}(Y)} - \underbrace{(E[X])^2}_{\text{var}(X)} - 2 \underbrace{E[X]E[Y]}_{\text{cov}(X, Y)} - \underbrace{(E[Y])^2}_{\text{var}(Y)} \\&= \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)\end{aligned}$$

- Important properties of the covariance follow from  $\text{cov}(X, Y) = E[XY] - E[X] E[Y]$ :

1. **Linearity:**  $\text{cov}(aX, bY) = ab \text{cov}(X, Y)$

$$\text{cov}(aX, bY) = E[(aX)(bY)] - E[aX] E[bY] = ab (E[XY] - E[X] E[Y])$$

2. **Symmetry:**  $\text{cov}(X, Y) = \text{cov}(Y, X)$  and  $\text{cov}(X, X) = \text{var}(X)$

$$\text{cov}(X, X) = E[XX] - E[X] E[X] = E[X^2] - (E[X])^2 = \text{var}(X)$$

3. **Addition Rule:**  $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)$

$$\begin{aligned}\text{var}(X + Y) &= E[(X + Y)^2] - E[(X + Y)]^2 \\&= E[X^2 + 2XY + Y^2] - (E[X] + E[Y])^2 \\&= \underbrace{E[X^2]}_{\text{var}(X)} + 2 \underbrace{E[XY]}_{\text{cov}(X, Y)} + \underbrace{E[Y^2]}_{\text{var}(Y)} - \underbrace{(E[X])^2}_{\text{var}(X)} - 2 \underbrace{E[X] E[Y]}_{\text{cov}(X, Y)} - \underbrace{(E[Y])^2}_{\text{var}(Y)} \\&= \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)\end{aligned}$$

# Covariance

- Important properties of the covariance follow from  $\text{cov}(X, Y) = E[XY] - E[X] E[Y]$ :

1. **Linearity:**  $\text{cov}(aX, bY) = ab \text{cov}(X, Y)$

$$\text{cov}(aX, bY) = E[(aX)(bY)] - E[aX] E[bY] = ab (E[XY] - E[X] E[Y])$$

2. **Symmetry:**  $\text{cov}(X, Y) = \text{cov}(Y, X)$  and  $\text{cov}(X, X) = \text{var}(X)$

$$\text{cov}(X, X) = E[XX] - E[X] E[X] = E[X^2] - (E[X])^2 = \text{var}(X)$$

3. **Addition Rule:**  $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)$

$$\begin{aligned}\text{var}(X + Y) &= E[(X + Y)^2] - E[(X + Y)]^2 \\&= E[X^2 + 2XY + Y^2] - (E[X] + E[Y])^2 \\&= \underbrace{E[X^2]}_{\text{var}(X)} + 2 \underbrace{E[XY]}_{\text{cov}(X, Y)} + \underbrace{E[Y^2]}_{\text{var}(Y)} - \underbrace{(E[X])^2}_{\text{var}(X)} - 2 \underbrace{E[X] E[Y]}_{\text{cov}(X, Y)} - \underbrace{(E[Y])^2}_{\text{var}(Y)} \\&= \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)\end{aligned}$$

- Important properties of the covariance follow from  $\text{cov}(X, Y) = E[XY] - E[X] E[Y]$ :

1. **Linearity:**  $\text{cov}(aX, bY) = ab \text{cov}(X, Y)$

$$\text{cov}(aX, bY) = E[(aX)(bY)] - E[aX] E[bY] = ab (E[XY] - E[X] E[Y])$$

2. **Symmetry:**  $\text{cov}(X, Y) = \text{cov}(Y, X)$  and  $\text{cov}(X, X) = \text{var}(X)$

$$\text{cov}(X, X) = E[XX] - E[X] E[X] = E[X^2] - (E[X])^2 = \text{var}(X)$$

3. **Addition Rule:**  $\text{var}(X + Y) = \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)$

$$\begin{aligned}\text{var}(X + Y) &= E[(X + Y)^2] - E[(X + Y)]^2 \\&= E[X^2 + 2XY + Y^2] - (E[X] + E[Y])^2 \\&= \underbrace{E[X^2]}_{\text{var}(X)} + 2 \underbrace{E[XY]}_{\text{cov}(X, Y)} + \underbrace{E[Y^2]}_{\text{var}(Y)} - \underbrace{(E[X])^2}_{\text{var}(X)} - 2 \underbrace{E[X] E[Y]}_{\text{cov}(X, Y)} - \underbrace{(E[Y])^2}_{\text{var}(Y)} \\&= \text{var}(X) + \text{var}(Y) + 2 \text{cov}(X, Y)\end{aligned}$$

- Useful rule:

$$\begin{aligned}\text{var}(aX + bY) &= \text{var}(aX) + \text{var}(bY) + 2 \text{cov}(aX, bY) \\ &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y)\end{aligned}$$

# Correlation

- ▶ The population **correlation coefficient** is denoted  $\rho$ :

$$\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$$

where

$$\sigma_{XY} = \text{cov}(X, Y), \quad \sigma_X = \sqrt{\text{var}(X)}, \quad \sigma_Y = \sqrt{\text{var}(Y)}$$

- ▶ The sample **correlation coefficient** is denoted  $r$ :

$$r_{XY} = \frac{s_{XY}}{s_X s_Y}$$



# Correlation

- ▶ The population **correlation coefficient** is denoted  $\rho$ :

$$\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y}$$

where

$$\sigma_{XY} = \text{cov}(X, Y), \quad \sigma_X = \sqrt{\text{var}(X)}, \quad \sigma_Y = \sqrt{\text{var}(Y)}$$

- ▶ The sample **correlation coefficient** is denoted  $r$ :

$$r_{XY} = \frac{s_{XY}}{s_X s_Y}$$

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:
  - ▶ The average income of college graduates:  
 $E[\text{Income} | \text{Education} = \text{College Graduate}]$
  - ▶ The average sales price of a house with 3 bedrooms:  $E[\text{Sales Price} | \text{Bedrooms} = 3]$
  - ▶ The average response for smokers:  
 $E[\text{Response} | \text{Smoker} = 1]$
- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
  - ▶ Examples:
    - ▶ The average income of college graduates.
    - ▶ The average income of college graduates with a certain GPA.
    - ▶ The average income of graduates with a certain GPA who are employed.
    - ▶ The average income of graduates with a certain GPA who are employed in a certain region.
    - ▶ The average income of graduates with a certain GPA who are employed in a certain region and have a certain level of education.
- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:

- The average income of college graduates:

$$E[\text{Income} \mid \text{Education} = \text{College Graduate}]$$

- The average sales price of a home with floor size 1200 sq. ft:

$$E[\text{Sales Price} \mid \text{Sqft} = 1200]$$

- The average lifespan for smokers:

$$E[\text{Lifespan} \mid \text{Smoker} = 1]$$

- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:
  - The average income of college graduates:

$$E[\text{Income} \mid \text{Education} = \text{College Graduate}]$$

- The average sales price of a home with floor size 1200 sq. ft:

$$E[\text{Sales Price} \mid \text{Sqft} = 1200]$$

- The average lifespan for smokers:

$$E[\text{Lifespan} \mid \text{Smoker} = 1]$$

- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:

- The average income of college graduates:

$$E[\text{Income} \mid \text{Education} = \text{College Graduate}]$$

- The average sales price of a home with floor size 1200 sq. ft:

$$E[\text{Sales Price} \mid \text{Sqft} = 1200]$$

- The average lifespan for smokers:

$$E[\text{Lifespan} \mid \text{Smoker} = 1]$$

- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:

- The average income of college graduates:

$$E[\text{Income} \mid \text{Education} = \text{College Graduate}]$$

- The average sales price of a home with floor size 1200 sq. ft:

$$E[\text{Sales Price} \mid \text{Sqft} = 1200]$$

- The average lifespan for smokers:

$$E[\text{Lifespan} \mid \text{Smoker} = 1]$$

- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.

# Conditioning and Independence

- ▶ Given two joint random variables,  $X$  and  $Y$ , we may be interested in characteristics of the distribution of  $Y$  conditional on  $X$  taking a certain value.
- ▶ Conditional expectation of  $Y$  given  $X = x$   $E[Y|X = x]$ :
- ▶ Examples:

- The average income of college graduates:

$$E[\text{Income} \mid \text{Education} = \text{College Graduate}]$$

- The average sales price of a home with floor size 1200 sq. ft:

$$E[\text{Sales Price} \mid \text{Sqft} = 1200]$$

- The average lifespan for smokers:

$$E[\text{Lifespan} \mid \text{Smoker} = 1]$$

- ▶ Knowing the conditional expectation is particularly useful for predictions when we observe the  $X$  variable before we observe the  $Y$  variable.



# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

$$\Pr(Y = y | X = x) = \frac{\Pr(X = x, Y = y)}{\Pr(X = x)} = \frac{\sum_z \Pr(X = x, Y = y, Z = z)}{\sum_z \Pr(X = x, Y = y, Z = z)}$$

where the marginal distribution of  $X$  at value  $x$  is

$$\Pr(X = x) = \sum_y \Pr(X = x, Y = y)$$

$$f_X(x) = \int f_{X,Y}(x, y) dy$$

- Divide by the marginal distribution  $X$  to find it - when  $Y$  is varied

# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

$$\begin{aligned} \Pr(Y = y) &= \sum_x \Pr(X = x, Y = y) && \text{Discrete RV} \\ \Pr(Y = y) &= \sum_x \Pr(Y = y | X = x) \Pr(X = x) && \text{Discrete RV} \\ \Pr(Y = y) &= \sum_x \Pr(Y = y | X = x) \Pr(X = x) && \text{Discrete RV} \\ \Pr(Y = y) &= \sum_x \Pr(Y = y | X = x) \Pr(X = x) && \text{Discrete RV} \end{aligned}$$

where the marginal distribution of  $X$  is given by

$$\Pr(X = x) = \sum_y \Pr(X = x, Y = y) \text{ for a discrete RV}$$

$$f_X(x) = \int_{-\infty}^{\infty} f_{X,Y}(x,y) dy \text{ for a continuous RV}$$

- Example: If the marginal distribution of  $X$  is  $\Gamma$  and if  $Y$  is uniform

# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

$$\begin{aligned} \Pr(X = x) &= \sum_{y \in \mathcal{Y}} \Pr(X = x, Y = y) \\ \Pr(Y = y | X = x) &= \frac{\Pr(X = x, Y = y)}{\Pr(X = x)} \end{aligned}$$

Example: conditional distribution of  $X$  given  $Y$

$$\Pr(X = x | Y = y) = \frac{\Pr(X = x, Y = y)}{\Pr(Y = y)}$$

$$\Pr(X = x | Y = y) = \frac{\Pr(X = x, Y = y)}{\sum_{x' \in \mathcal{X}} \Pr(X = x', Y = y)}$$

► Example: the marginal distribution of  $X$  is  $\Pr(X = x) = \sum_{y \in \mathcal{Y}} \Pr(X = x, Y = y)$

# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

$$E[Y|X = x] = \frac{\begin{array}{c} \text{Discrete R.V} \\ \sum_y y \cdot p_{XY}(y, x) \end{array}}{\sum_y p_{XY}(x, y)} \quad \frac{\begin{array}{c} \text{Continuous R.V} \\ \int_Y y \cdot f_{XY}(x, y) dy \end{array}}{\int_Y f_{XY}(x, y) dy}$$

where the marginal distribution of  $X$  at value  $x$  is:

$$\Pr(X = x) = \sum_y p_{XY}(x, y) \text{ for a discrete rv.}$$

$$f_X(x) = \int_Y f_{XY}(x, y) dy \text{ for a continuous rv.}$$

► The value of the marginal distribution of  $X$  is fixed at a value  $x$ , while  $Y$  is varied.

# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

$$E[Y|X = x] = \frac{\begin{array}{c} \text{Discrete R.V} \\ \sum_y y \cdot p_{XY}(y, x) \end{array}}{\sum_y p_{XY}(x, y)} \quad \frac{\begin{array}{c} \text{Continuous R.V} \\ \int_Y y \cdot f_{XY}(x, y) dy \end{array}}{\int_Y f_{XY}(x, y) dy}$$

where the marginal distribution of  $X$  at value  $x$  is:

$$\Pr(X = x) = \sum_y p_{XY}(x, y) \text{ for a discrete rv.}$$

$$f_X(x) = \int_Y f_{XY}(x, y) dy \text{ for a continuous rv.}$$

► The value of the marginal distribution of  $X$  is fixed at a value  $x$ , while  $Y$  is varied.

# Conditioning and Independence

## ► Conditional expectation:

1. Calculate the marginal probability  $\Pr(X = x)$ .
2. Fix the  $X$  variable at value  $X = x$ .
3. Divide by the probability that  $X = x$ .

---

|                | Discrete R.V  | Continuous R.V  |
|----------------|---|---|
| $E[Y X = x] =$ | $\frac{\sum_y y \cdot p_{XY}(y, x)}{\sum_y p_{XY}(x, y)}$ | $\frac{\int_Y y \cdot f_{XY}(x, y) dy}{\int_Y f_{XY}(x, y) dy}$ |

---

where the marginal distribution of  $X$  at value  $x$  is:

$$\Pr(X = x) = \sum_y p_{XY}(x, y) \text{ for a discrete r.v.}$$

$$f_X(x) = \int_Y f_{XY}(x, y) dy \text{ for a continuous r.v.}$$

- To compute the marginal distribution,  $X$  is fixed at value  $x$ , while  $Y$  is varied.

# Conditioning and Independence

- Let  $Y$  be hours of sleep and  $X$  cups of coffee drunk per day. The joint pmf  $p(x, y)$  is:

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- Compute the expected number of hours of sleep for someone who drinks 2 cups of coffee.
1. Calculate the probability that one random person drinks 2 cups of coffee:

$$\Pr(X = x) = p_{XY}(x, y) = \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

2. Fix  $X = 2$  and calculate  $\sum_y y \cdot p_{XY}(2, y)$ :

$$\sum_y y \cdot p_{XY}(2, y) = 4 \cdot \frac{1}{6} + 8 \cdot \frac{1}{3} + 12 \cdot 0 = \frac{10}{3}$$

3. Calculate  $E[Y|X = 2]$

$$E[Y|X = 2] = \frac{10}{3} \cdot \frac{2}{1} = \frac{20}{3} \approx 6.67$$

# Conditioning and Independence

- Let  $Y$  be hours of sleep and  $X$  cups of coffee drunk per day. The joint pmf  $p(x, y)$  is:

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- Compute the expected number of hours of sleep for someone who drinks 2 cups of coffee.
1. Calculate the probability that one random person drinks 2 cups of coffee:

$$\Pr(X = 2) = p_{XY}(2, y) = \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

2. Fix  $X = 2$  and calculate  $\sum_y y \cdot p_{XY}(2, y)$ :

$$\sum_y y \cdot p_{XY}(2, y) = 4 \cdot \frac{1}{6} + 8 \cdot \frac{1}{3} + 12 \cdot 0 = \frac{10}{3}$$

3. Calculate  $E[Y|X = 2]$

$$E[Y|X = 2] = \frac{10}{3} \cdot \frac{2}{1} = \frac{20}{3} \approx 6.67$$



# Conditioning and Independence

- Let  $Y$  be hours of sleep and  $X$  cups of coffee drunk per day. The joint pmf  $p(x, y)$  is:

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- Compute the expected number of hours of sleep for someone who drinks 2 cups of coffee.
1. Calculate the probability that one random person drinks 2 cups of coffee:

$$\Pr(X = x) = p_{XY}(x, y) = \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

2. Fix  $X = 2$  and calculate  $\sum_y y \cdot p_{XY}(2, y)$ :

$$\sum_y y \cdot p_{XY}(2, y) = 4 \cdot \frac{1}{6} + 8 \cdot \frac{1}{3} + 12 \cdot 0 = \frac{10}{3}$$

3. Calculate  $E[Y|X = 2]$

$$E[Y|X = 2] = \frac{10}{3} \cdot \frac{2}{1} = \frac{20}{3} \approx 6.67$$

## Conditioning and Independence

- Let  $Y$  be hours of sleep and  $X$  cups of coffee drunk per day. The joint pmf  $p(x, y)$  is:

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | 1/6    |
| 8 hours   | 1/3   | 1/3    |
| 12 hours  | 1/6   | 0      |

- Compute the expected number of hours of sleep for someone who drinks 2 cups of coffee.
1. Calculate the probability that one random person drinks 2 cups of coffee:

$$\Pr(X = x) = p_{XY}(x, y) = \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

2. Fix  $X = 2$  and calculate  $\sum_y y \cdot p_{XY}(2, y)$ :

$$\sum_y y \cdot p_{XY}(2, y) = 4 \cdot \frac{1}{6} + 8 \cdot \frac{1}{3} + 12 \cdot 0 = \frac{10}{3}$$

3. Calculate  $E[Y|X = 2]$

$$E[Y|X = 2] = \frac{10}{3} \cdot \frac{2}{1} = \frac{20}{3} \approx 6.67$$

## Conditioning and Independence

- Let  $Y$  be hours of sleep and  $X$  cups of coffee drunk per day. The joint pmf  $p(x, y)$  is:

| $p(x, y)$ | 1 cup | 2 cups |
|-----------|-------|--------|
| 4 hours   | 0     | $1/6$  |
| 8 hours   | $1/3$ | $1/3$  |
| 12 hours  | $1/6$ | 0      |

- Compute the expected number of hours of sleep for someone who drinks 2 cups of coffee.
1. Calculate the probability that one random person drinks 2 cups of coffee:

$$\Pr(X = x) = p_{XY}(x, y) = \frac{1}{6} + \frac{1}{3} = \frac{1}{2}$$

2. Fix  $X = 2$  and calculate  $\sum_y y \cdot p_{XY}(2, y)$ :

$$\sum_y y \cdot p_{XY}(2, y) = 4 \cdot \frac{1}{6} + 8 \cdot \frac{1}{3} + 12 \cdot 0 = \frac{10}{3}$$

3. Calculate  $E[Y|X = 2]$

$$E[Y|X = 2] = \frac{10}{3} \cdot \frac{2}{1} = \frac{20}{3} \approx 6.67$$

# Conditioning and Independence

► If  $X$  does not help predict  $Y$ , we say that  $X$  is **independent** of  $Y$  and denote  $X \perp\!\!\!\perp Y$ .

► Examples of independent random variables:

• Knowing that one coin flip came up heads doesn't help predict the next coin flip or whether coin flipping is independent.

• Knowing the number that comes up in a game of roulette doesn't help to choose a winning game strategy.

# Conditioning and Independence

- ▶ If  $X$  does not help predict  $Y$ , we say that  $X$  is **independent** of  $Y$  and denote  $X \perp\!\!\!\perp Y$ .
- ▶ Examples of independent random variables:
  - Knowing that one coin flip came up heads doesn't help predict the next coin flip: successive coin flips are independent.
  - Knowing the numbers that came up in a game of roulette cannot help to devise a better game strategy.

# Conditioning and Independence

- ▶ If  $X$  does not help predict  $Y$ , we say that  $X$  is **independent** of  $Y$  and denote  $X \perp\!\!\!\perp Y$ .
- ▶ Examples of independent random variables:
  - Knowing that one coin flip came up heads doesn't help predict the next coin flip: successive coin flips are independent.
  - Knowing the numbers that came up in a game of roulette cannot help to devise a better game strategy.

# Conditioning and Independence

- ▶ If  $X$  does not help predict  $Y$ , we say that  $X$  is **independent** of  $Y$  and denote  $X \perp\!\!\!\perp Y$ .
- ▶ Examples of independent random variables:
  - Knowing that one coin flip came up heads doesn't help predict the next coin flip: successive coin flips are independent.
  - Knowing the numbers that came up in a game of roulette cannot help to devise a better game strategy.

# Conditioning and Independence

- ▶ If  $X$  and  $Y$  are independent,  $X \perp\!\!\!\perp Y$ , then:

$$\Pr(a \leq X \leq b, c \leq Y \leq d) = \Pr(a \leq X \leq b) \cdot \Pr(c \leq Y \leq d) \quad \forall (a, b, c, d)$$

$$\mathbb{E}[g(Y)|X = x] = \mathbb{E}[g(Y)] \quad \forall x \in \Omega, g(\cdot) : \Omega \rightarrow \mathbb{R}$$

- ▶ Examples of variables that seem independent but may not be:

• a teacher and average height of class

• gender of students and average height of class



# Conditioning and Independence

- ▶ If  $X$  and  $Y$  are independent,  $X \perp\!\!\!\perp Y$ , then:

$$\Pr(a \leq X \leq b, c \leq Y \leq d) = \Pr(a \leq X \leq b) \cdot \Pr(c \leq Y \leq d) \quad \forall (a, b, c, d)$$

$$\mathbb{E}[g(Y)|X = x] = \mathbb{E}[g(Y)] \quad \forall x \in \Omega, g(\cdot) : \Omega \rightarrow \mathbb{R}$$

- ▶ Examples of variables that seem independent but may not be:

- weather and average house prices.
- academic achievements and average house prices.

# Conditioning and Independence

- ▶ If  $X$  and  $Y$  are independent,  $X \perp\!\!\!\perp Y$ , then:

$$\Pr(a \leq X \leq b, c \leq Y \leq d) = \Pr(a \leq X \leq b) \cdot \Pr(c \leq Y \leq d) \quad \forall (a, b, c, d)$$

$$\mathbb{E}[g(Y)|X = x] = \mathbb{E}[g(Y)] \quad \forall x \in \Omega, g(\cdot) : \Omega \rightarrow \mathbb{R}$$

- ▶ Examples of variables that seem independent but may not be:
  - weather and average house prices.
  - academic achievements and average house prices.

# Conditioning and Independence

- ▶ If  $X$  and  $Y$  are independent,  $X \perp\!\!\!\perp Y$ , then:

$$\Pr(a \leq X \leq b, c \leq Y \leq d) = \Pr(a \leq X \leq b) \cdot \Pr(c \leq Y \leq d) \quad \forall (a, b, c, d)$$

$$\mathbb{E}[g(Y)|X = x] = \mathbb{E}[g(Y)] \quad \forall x \in \Omega, g(\cdot) : \Omega \rightarrow \mathbb{R}$$

- ▶ Examples of variables that seem independent but may not be:
  - weather and average house prices.
  - academic achievements and average house prices.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- We often describe the "average" value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables "move with" each other.

## ► Conditioning and Independence:

- The conditional expectation is the average value of  $Y$  for individuals who have  $X$  as its value.
- If knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- We often describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

- The conditional expectation is the average value of  $Y$  for individuals who have  $X$  as its value.
- Knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- The *expectation* describes the “average” value of a function of the joint random variables.
- The *covariance function* is a particular expectation that is used to describe the joint variables’ “linear with” properties.

## ► Conditioning and Independence:

- The *conditional expectation* is the average value of  $Y$  for  $X$  that take values from  $\mathcal{X}$  as given.
- Two random variables  $X$  and  $Y$  are *independent* if any information on the distribution of  $Y$  we get from  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

The conditional expectation is the average value of  $Y$  for  $X$  individuals who have  $X=x$ .  
Knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

The conditional expectation is the average value of  $Y$  for  $X$  that take value  $x$ . It is denoted  $E(Y|X=x)$ . Knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.



## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

The conditional distribution of  $X$  given  $Y=y$  is the distribution of  $X$  given that  $Y=y$ . The conditional distribution of  $Y$  given  $X=x$  is the distribution of  $Y$  given that  $X=x$ .

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

- The conditional expectation is the average value of  $Y$  for individuals who have  $X = x$ .
- If knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

- The conditional expectation is the average value of  $Y$  for individuals who have  $X = x$ .
- If knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## ► Multiple Random Variables:

- Describe multiple events whose outcomes are unknown.
- Have probabilities that the outcomes jointly take values in arbitrary subsets of the joint sample space.

## ► Expectations:

- As before describe the “average” value of a function of the joint random variables.
- The covariance function is a particular expectation we are interested in as it describes how two variables “move with” each other.

## ► Conditioning and Independence:

- The conditional expectation is the average value of  $Y$  for individuals who have  $X = x$ .
- If knowing  $X$  does not give us any information on the distribution of  $Y$  we say that  $X$  and  $Y$  are independent.

## The Normal Distribution

# The Normal Distribution

## Normal Distribution


A random variable  $X$  follows a normal distribution with mean  $\mu$  and variance  $\sigma^2$  if it is continuously distributed with probability density function (pdf) given by:

$$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

It is denoted  $X \sim N(\mu, \sigma^2)$ .

- **Standard normal distribution:**  $Z \sim N(0, 1)$ . That is,  $\mu = 0$  and  $\sigma^2 = 1$ .

# The Normal Distribution



# Properties of Normal Distribution

## Property 1:

- ▶ If  $X \sim N(\mu, \sigma^2)$  then  $(X - \mu)/\sigma \sim N(0, 1)$ .
- ▶ Thus, we can express probabilities for any normal random variable in terms of  $Z \sim N(0, 1)$ .
- ▶ Exercise: Show that if  $X \sim N(2, 100)$  then  $\Pr(X \geq 22) = \Pr(Z \geq 2)$ .

$$\begin{aligned}\Pr(X \geq 22) &= \Pr\left(\frac{X - 2}{10} \geq \frac{22 - 2}{10}\right) \\ &= \Pr(Z \geq 2)\end{aligned}$$

Here we use the fact that  $\mu = 2$  and  $\sigma = \sqrt{100} = 10$ .

- ▶ We calculate  $\Pr(Z \geq 2)$  by looking up a table or using dedicated software.



# Properties of Normal Distribution

## Property 1:

- ▶ If  $X \sim N(\mu, \sigma^2)$  then  $(X - \mu)/\sigma \sim N(0, 1)$ .
- ▶ Thus, we can express probabilities for any normal random variable in terms of  $Z \sim N(0, 1)$ .
- ▶ Exercise: Show that if  $X \sim N(2, 100)$  then  $\Pr(X \geq 22) = \Pr(Z \geq 2)$ .

$$\begin{aligned}\Pr(X \geq 22) &= \Pr\left(\frac{X - 2}{10} \geq \frac{22 - 2}{10}\right) \\ &= \Pr(Z \geq 2)\end{aligned}$$

Here we use the fact that  $\mu = 2$  and  $\sigma = \sqrt{100} = 10$ .

- ▶ We calculate  $\Pr(Z \geq 2)$  by looking up a table or using dedicated software.

# Properties of Normal Distribution

## Property 1:

- ▶ If  $X \sim N(\mu, \sigma^2)$  then  $(X - \mu)/\sigma \sim N(0, 1)$ .
- ▶ Thus, we can express probabilities for any normal random variable in terms of  $Z \sim N(0, 1)$ .
- ▶ Exercise: Show that if  $X \sim N(2, 100)$  then  $\Pr(X \geq 22) = \Pr(Z \geq 2)$ .

$$\begin{aligned}\Pr(X \geq 22) &= \Pr\left(\frac{X - 2}{10} \geq \frac{22 - 2}{10}\right) \\ &= \Pr(Z \geq 2)\end{aligned}$$

Here we use the fact that  $\mu = 2$  and  $\sigma = \sqrt{100} = 10$ .

- ▶ We calculate  $\Pr(Z \geq 2)$  by looking up a table or using dedicated software.

# Properties of Normal Distribution

## Property 1:

- ▶ If  $X \sim N(\mu, \sigma^2)$  then  $(X - \mu)/\sigma \sim N(0, 1)$ .
- ▶ Thus, we can express probabilities for any normal random variable in terms of  $Z \sim N(0, 1)$ .
- ▶ Exercise: Show that if  $X \sim N(2, 100)$  then  $\Pr(X \geq 22) = \Pr(Z \geq 2)$ .

$$\begin{aligned}\Pr(X \geq 22) &= \Pr\left(\frac{X - 2}{10} \geq \frac{22 - 2}{10}\right) \\ &= \Pr(Z \geq 2)\end{aligned}$$

Here we use the fact that  $\mu = 2$  and  $\sigma = \sqrt{100} = 10$ .

- ▶ We calculate  $\Pr(Z \geq 2)$  by looking up a table or using dedicated software.

# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .

- ▶ Calculate  $E[W]$ :

$$\begin{aligned} &\Rightarrow \text{By linearity of expectation: } E[aX + bY] = aE[X] + bE[Y] \\ &\Rightarrow E[W] = E[aX + bY] = a\mu_X + b\mu_Y \end{aligned}$$

- ▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned} \text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY} \end{aligned}$$

- ▶ Putting it together:

$$W \sim N(a\mu_X + b\mu_Y, a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY})$$

# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .

- ▶ Calculate  $E[W]$ :

- By linearity of expectation we have that  $E[aX + bY] = a E[X] + b E[Y]$
- So,  $\mu_W = E[W] = a\mu_X + b\mu_Y$ .

- ▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned}\text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY}\end{aligned}$$

- ▶ Putting it together:

$$W \sim N(a\mu_X + b\mu_Y, a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY})$$

# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .
- ▶ Calculate  $E[W]$ :
  - By linearity of expectation we have that  $E[aX + bY] = a E[X] + b E[Y]$
  - So,  $\mu_W = E[W] = a\mu_X + b\mu_Y$ .

- ▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned}\text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY}\end{aligned}$$

- ▶ Putting it together:

$$W \sim N(a\mu_X + b\mu_Y, a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY})$$

# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .
- ▶ Calculate  $E[W]$ :
  - By linearity of expectation we have that  $E[aX + bY] = a E[X] + b E[Y]$
  - So,  $\mu_W = E[W] = a\mu_X + b\mu_Y$ .

▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned}\text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY}\end{aligned}$$

▶ Putting it together:

$$W \sim N(a\mu_X + b\mu_Y, a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY})$$

# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .
- ▶ Calculate  $E[W]$ :
  - By linearity of expectation we have that  $E[aX + bY] = a E[X] + b E[Y]$
  - So,  $\mu_W = E[W] = a\mu_X + b\mu_Y$ .
- ▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned}\text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY}\end{aligned}$$

- ▶ Putting it together:

$$W \sim N(a\mu_X + b\mu_Y, a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY})$$



# Properties of Normal Distribution

## Property 2:

- ▶ If  $X \sim N(\mu_X, \sigma_X^2)$  and  $Y \sim N(\mu_Y, \sigma_Y^2)$  are jointly normal, then  $W = aX + bY$  is also normally distributed for any  $a, b \in \mathbb{R}$ .
- ▶ Calculate  $E[W]$ :
  - By linearity of expectation we have that  $E[aX + bY] = a E[X] + b E[Y]$
  - So,  $\mu_W = E[W] = a\mu_X + b\mu_Y$ .

- ▶ Calculate  $\text{var}(W)$ :

$$\begin{aligned}\text{var}(aX + bY) &= a^2 \text{var}(X) + b^2 \text{var}(Y) + 2ab \text{cov}(X, Y) \\ \implies \text{var}(W) &= \text{var}(aX + bY) = a^2 \sigma_X^2 + b^2 \sigma_Y^2 + 2ab \sigma_{XY}\end{aligned}$$

- ▶ Putting it together:

$$W \sim N\left(a\mu_X + b\mu_Y, a^2\sigma_X^2 + b^2\sigma_Y^2 + 2ab\sigma_{XY}\right)$$

# Properties of Normal Distribution

## Property 3:

- ▶ The distribution of  $X \sim N(0, \sigma^2)$  is symmetric around zero:

$$\Pr(X \geq x) = \Pr(X \leq -x)$$

- ▶ Thus, we can compute:

$$\Pr(|X| \geq c) = \Pr(X \geq c) + \Pr(X \leq -c) = 2 \Pr(X \geq c)$$

# Properties of Normal Distribution


## Property 3:

- ▶ The distribution of  $X \sim N(0, \sigma^2)$  is symmetric around zero:

$$\Pr(X \geq x) = \Pr(X \leq -x)$$

- ▶ Thus, we can compute:

$$\Pr(|X| \geq c) = \Pr(X \geq c) + \Pr(X \leq -c) = 2 \Pr(X \geq c)$$



## The Sample Mean

# The Sample Mean

- **Example:** In an election the Green candidate is running against the Red candidate. We randomly select  $n = 100$  voters from the population and ask them who they plan on voting for. Answers are recorded as

$$X_i = \begin{cases} 1 & \text{if they plan to vote Green} \\ 0 & \text{if they plan to vote Red} \end{cases}$$

- The sample mean  $\bar{X}_n$  and sample variance  $s_n^2$  are:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i = 0.55$$

$$s_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = 0.25$$

## Questions:

1. What is the Green candidate's win probability?
2. How confident will we be in this estimate?
3. How confident will we be in the uncertainty that we have?

# The Sample Mean

- **Example:** In an election the Green candidate is running against the Red candidate. We randomly select  $n = 100$  voters from the population and ask them who they plan on voting for. Answers are recorded as

$$X_i = \begin{cases} 1 & \text{if they plan to vote Green} \\ 0 & \text{if they plan to vote Red} \end{cases}$$

- The sample mean  $\bar{X}_n$  and sample variance  $s_n^2$  are:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i = 0.55$$

$$s_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = 0.25$$

## Questions:

- Will the Green candidate win the election?
- Would another poll confirm these results?
- How can we measure the uncertainty the sample mean?

# The Sample Mean

- **Example:** In an election the Green candidate is running against the Red candidate. We randomly select  $n = 100$  voters from the population and ask them who they plan on voting for. Answers are recorded as

$$X_i = \begin{cases} 1 & \text{if they plan to vote Green} \\ 0 & \text{if they plan to vote Red} \end{cases}$$

- The sample mean  $\bar{X}_n$  and sample variance  $s_n^2$  are:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i = 0.55$$

$$s_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = 0.25$$

## Questions:

- Will the Green candidate win the election?
- Would another poll confirm these results?
- How can we measure the uncertainty the sample mean?

# The Sample Mean

- **Example:** In an election the Green candidate is running against the Red candidate. We randomly select  $n = 100$  voters from the population and ask them who they plan on voting for. Answers are recorded as

$$X_i = \begin{cases} 1 & \text{if they plan to vote Green} \\ 0 & \text{if they plan to vote Red} \end{cases}$$

- The sample mean  $\bar{X}_n$  and sample variance  $s_n^2$  are:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i = 0.55$$

$$s_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = 0.25$$

## Questions:

- Will the Green candidate win the election?
- Would another poll confirm these results?
- How can we measure the uncertainty the sample mean?



# The Sample Mean

- **Example:** In an election the Green candidate is running against the Red candidate. We randomly select  $n = 100$  voters from the population and ask them who they plan on voting for. Answers are recorded as

$$X_i = \begin{cases} 1 & \text{if they plan to vote Green} \\ 0 & \text{if they plan to vote Red} \end{cases}$$

- The sample mean  $\bar{X}_n$  and sample variance  $s_n^2$  are:

$$\bar{X}_n = \frac{1}{n} \sum_{i=1}^n X_i = 0.55$$

$$s_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X}_n)^2 = 0.25$$

## Questions:

- Will the Green candidate win the election?
- Would another poll confirm these results?
- How can we measure the uncertainty the sample mean?

# The Sample Mean: As a Random Variable

- ▶ **Random Sampling:**  $\bar{X}_n$  and  $s_n^2$  are random variables.
- ▶ Each time a random sample is drawn, different members of the population are drawn.
- ▶ We want to use this random sample to learn about the population.
  - The sample  $\{X_i\}_{i=1}^n$  is made up of  $n$  different random variables.
  - Each  $X_i$  is sampled from the same population distributed as  $X$ , so that

$$\begin{aligned}E[X_i] &= \mu_X \\ \text{var}(X_i) &= \sigma_X^2\end{aligned}$$

# The Sample Mean: As a Random Variable

- ▶ **Random Sampling:**  $\bar{X}_n$  and  $s_n^2$  are random variables.
- ▶ Each time a random sample is drawn, different members of the population are drawn.
- ▶ We want to use this random sample to learn about the population.

→ The sample  $\{X_i\}_{i=1}^n$  is made up of  $n$  iid random variables.

→ Each  $X_i$  is sampled from the same population distribution as the

$$E[X_i] = \mu$$

$$\text{var}(X_i) = \sigma^2$$

# The Sample Mean: As a Random Variable

- ▶ **Random Sampling:**  $\bar{X}_n$  and  $s_n^2$  are random variables.
- ▶ Each time a random sample is drawn, different members of the population are drawn.
- ▶ We want to use this random sample to learn about the population.
  - The sample  $\{X_i\}_{i=1}^n$  is made up of  $n$  different random variables.
  - Each  $X_i$  is sampled from the same population distribution, so that

$$\begin{aligned}E[X_i] &= \mu_X \\ \text{var}(X_i) &= \sigma_X^2\end{aligned}$$

# The Sample Mean: As a Random Variable

- ▶ **Random Sampling:**  $\bar{X}_n$  and  $s_n^2$  are random variables.
- ▶ Each time a random sample is drawn, different members of the population are drawn.
- ▶ We want to use this random sample to learn about the population.
  - The sample  $\{X_i\}_{i=1}^n$  is made up of  $n$  different random variables.
  - Each  $X_i$  is sampled from the same population distribution, so that

$$\begin{aligned} \mathbb{E}[X_i] &= \mu_X \\ \text{var}(X_i) &= \sigma_X^2 \end{aligned}$$

# The Sample Mean: As a Random Variable

- ▶ **Random Sampling:**  $\bar{X}_n$  and  $s_n^2$  are random variables.
- ▶ Each time a random sample is drawn, different members of the population are drawn.
- ▶ We want to use this random sample to learn about the population.
  - The sample  $\{X_i\}_{i=1}^n$  is made up of  $n$  different random variables.
  - Each  $X_i$  is sampled from the same population distribution, so that

$$\begin{aligned}E[X_i] &= \mu_X \\ \text{var}(X_i) &= \sigma_X^2\end{aligned}$$

# The Sample Mean: As a Random Variable

- **Random sampling:** Learn about the population from one sample.

|                       | Population      | Sample      |
|-----------------------|-----------------|-------------|
| Measure of location   | $E[X]$          | $\bar{X}_n$ |
| Measure of dispersion | $\text{var}(X)$ | $s_n^2$     |

1. **Expectation:** of  $\bar{X}_n$  is given  $E[\bar{X}_n] = E[X]$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = E[X]$$

2. **Variance:**  $\bar{X}_n$  is given  $\text{var}(\bar{X}_n) = \sigma_X^2/n$ .

$$\text{var}(\bar{X}_n) = \text{var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(X) = \sigma_X^2/n$$

- Note that  $\text{var}(\bar{X}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . This is the basis of the **Law of Large Numbers** which states that  $\bar{X}_n \rightarrow \mu_X$  as  $n \rightarrow \infty$ .

# The Sample Mean: As a Random Variable

- **Random sampling:** Learn about the population from one sample.

|                       | Population      | Sample      |
|-----------------------|-----------------|-------------|
| Measure of location   | $E[X]$          | $\bar{X}_n$ |
| Measure of dispersion | $\text{var}(X)$ | $s_n^2$     |

1. **Expectation:** of  $\bar{X}_n$  is given  $E[\bar{X}_n] = E[X]$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = E[X]$$

2. **Variance:**  $\bar{X}_n$  is given  $\text{var}(\bar{X}_n) = \sigma_X^2/n$ .

$$\text{var}(\bar{X}_n) = \text{var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(X) = \sigma_X^2/n$$

- Note that  $\text{var}(\bar{X}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . This is the basis of the **Law of Large Numbers** which states that  $\bar{X}_n \rightarrow \mu_X$  as  $n \rightarrow \infty$ .



# The Sample Mean: As a Random Variable

- **Random sampling:** Learn about the population from one sample.

|                       | Population      | Sample      |
|-----------------------|-----------------|-------------|
| Measure of location   | $E[X]$          | $\bar{X}_n$ |
| Measure of dispersion | $\text{var}(X)$ | $s_n^2$     |

1. **Expectation:** of  $\bar{X}_n$  is given  $E[\bar{X}_n] = E[X]$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = E[X]$$

2. **Variance:**  $\bar{X}_n$  is given  $\text{var}(\bar{X}_n) = \sigma_X^2/n$ .

$$\text{var}(\bar{X}_n) = \text{var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(X) = \sigma_X^2/n$$

- Note that  $\text{var}(\bar{X}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . This is the basis of the **Law of Large Numbers** which states that  $\bar{X}_n \rightarrow \mu_X$  as  $n \rightarrow \infty$ .

# The Sample Mean: As a Random Variable

- **Random sampling:** Learn about the population from one sample.

|                       | Population      | Sample      |
|-----------------------|-----------------|-------------|
| Measure of location   | $E[X]$          | $\bar{X}_n$ |
| Measure of dispersion | $\text{var}(X)$ | $s_n^2$     |

1. **Expectation:** of  $\bar{X}_n$  is given  $E[\bar{X}_n] = E[X]$ .

$$E[\bar{X}_n] = E\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n E[X_i] = E[X]$$

2. **Variance:**  $\bar{X}_n$  is given  $\text{var}(\bar{X}_n) = \sigma_X^2/n$ .

$$\text{var}(\bar{X}_n) = \text{var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \sum_{i=1}^n \text{var}(X) = \sigma_X^2/n$$

- Note that  $\text{var}(\bar{X}_n) \rightarrow 0$  as  $n \rightarrow \infty$ . This is the basis of the **Law of Large Numbers** which states that  $\bar{X}_n \rightarrow \mu_X$  as  $n \rightarrow \infty$ .

# The Sample Mean: Central Limit Theorem

## ► Distribution of $\bar{X}_n$ :

- Needed to make inferences about  $E[X]$  and compute probabilities like

$$\Pr \left( |\bar{X}_n - E[X]| > 0.05 \right)$$

- **Central Limit Theorem:** For  $n$  sufficiently large,

$$\bar{X}_n \sim N(\mu_x, \sigma_X^2/n)$$

- How large is “sufficiently large”?
- In practice, the central limit theorem provides a good approximation for the distribution of  $\bar{X}_n$  when  $n > 30$ .

# The Sample Mean: Central Limit Theorem

- ▶ **Distribution of  $\bar{X}_n$ :**

- ▶ Needed to make inferences about  $E[X]$  and compute probabilities like

$$\Pr\left(|\bar{X}_n - E[X]| > 0.05\right)$$

- ▶ **Central Limit Theorem:** For  $n$  sufficiently large,

$$\bar{X}_n \sim N(\mu_x, \sigma_X^2/n)$$

- ▶ How large is “sufficiently large”?
- ▶ In practice, the central limit theorem provides a good approximation for the distribution of  $\bar{X}_n$  when  $n > 30$ .

# The Sample Mean: Central Limit Theorem

- ▶ **Distribution of  $\bar{X}_n$ :**

- ▶ Needed to make inferences about  $E[X]$  and compute probabilities like

$$\Pr\left(|\bar{X}_n - E[X]| > 0.05\right)$$

- ▶ **Central Limit Theorem:** For  $n$  sufficiently large,

$$\bar{X}_n \sim N(\mu_x, \sigma_X^2/n)$$

- ▶ How large is “sufficiently large”?
- ▶ In practice, the central limit theorem provides a good approximation for the distribution of  $\bar{X}_n$  when  $n > 30$ .

# The Sample Mean: Central Limit Theorem

- ▶ **Distribution of  $\bar{X}_n$ :**

- ▶ Needed to make inferences about  $E[X]$  and compute probabilities like

$$\Pr\left(|\bar{X}_n - E[X]| > 0.05\right)$$

- ▶ **Central Limit Theorem:** For  $n$  sufficiently large,

$$\bar{X}_n \sim N(\mu_x, \sigma_X^2/n)$$

- ▶ How large is “sufficiently large”?

- ▶ In practice, the central limit theorem provides a good approximation for the distribution of  $\bar{X}_n$  when  $n > 30$ .

# The Sample Mean: Central Limit Theorem

- ▶ **Distribution of  $\bar{X}_n$ :**

- ▶ Needed to make inferences about  $E[X]$  and compute probabilities like

$$\Pr\left(|\bar{X}_n - E[X]| > 0.05\right)$$

- ▶ **Central Limit Theorem:** For  $n$  sufficiently large,

$$\bar{X}_n \sim N(\mu_x, \sigma_X^2/n)$$

- ▶ How large is “sufficiently large”?
- ▶ In practice, the central limit theorem provides a good approximation for the distribution of  $\bar{X}_n$  when  $n > 30$ .

# The Sample Mean: Central Limit Theorem

- For  $n$  sufficiently large,

$$\frac{\bar{X}_n - E[\bar{X}_n]}{\sqrt{\text{var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \sim N(0, 1)$$

- For  $n$  sufficiently large, we can approximate the population standard deviation  $\sigma_X$  with its sample estimate  $s_n$ :

$$\frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \approx \frac{\bar{X}_n - \mu_X}{s_n / \sqrt{n}} \sim N(0, 1)$$

- The distribution of the sample mean  $\bar{X}_n$  can be used for inference about  $\mu_X$ . We can build confidence intervals and conduct hypothesis tests.



# The Sample Mean: Central Limit Theorem

- ▶ For  $n$  sufficiently large,

$$\frac{\bar{X}_n - E[\bar{X}_n]}{\sqrt{\text{var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \sim N(0, 1)$$

- ▶ For  $n$  sufficiently large, we can approximate the population standard deviation  $\sigma_X$  with its sample estimate  $s_n$ :

$$\frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \approx \frac{\bar{X}_n - \mu_X}{s_n / \sqrt{n}} \sim N(0, 1)$$

- ▶ The distribution of the sample mean  $\bar{X}_n$  can be used for inference about  $\mu_X$ . We can build confidence intervals and conduct hypothesis tests.

# The Sample Mean: Central Limit Theorem

- ▶ For  $n$  sufficiently large,

$$\frac{\bar{X}_n - E[\bar{X}_n]}{\sqrt{\text{var}(\bar{X}_n)}} = \frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \sim N(0, 1)$$

- ▶ For  $n$  sufficiently large, we can approximate the population standard deviation  $\sigma_X$  with its sample estimate  $s_n$ :

$$\frac{\bar{X}_n - \mu_X}{\sigma_X / \sqrt{n}} \approx \frac{\bar{X}_n - \mu_X}{s_n / \sqrt{n}} \sim N(0, 1)$$

- ▶ The distribution of the sample mean  $\bar{X}_n$  can be used for inference about  $\mu_X$ . We can build confidence intervals and conduct hypothesis tests.

# The Sample Mean: Central Limit Theorem

- ▶ Consider the polling example with  $n = 100$
- ▶ What is the probability that the sample has mean  $\bar{X}_n = 0.55$  and variance  $s_n^2 = 0.25$ , if the true proportion of Green voters in the population is  $E[X] = 0.5$ ?
- ▶ By the central limit theorem:

$$\begin{aligned}\Pr(\bar{X}_n \geq 0.55) &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - \mu_X}{s_n/\sqrt{n}}\right) \\ &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - 0.5}{0.5/10}\right) \\ &= \Pr(Z \geq 1) \\ &\approx 0.159\end{aligned}$$

# The Sample Mean: Central Limit Theorem

- ▶ Consider the polling example with  $n = 100$
- ▶ What is the probability that the sample has mean  $\bar{X}_n = 0.55$  and variance  $s_n^2 = 0.25$ , if the true proportion of Green voters in the population is  $E[X] = 0.5$ ?
- ▶ By the central limit theorem:

$$\begin{aligned}\Pr(\bar{X}_n \geq 0.55) &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - \mu_X}{s_n/\sqrt{n}}\right) \\ &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - 0.5}{0.5/10}\right) \\ &= \Pr(Z \geq 1) \\ &\approx 0.159\end{aligned}$$

# The Sample Mean: Central Limit Theorem

- ▶ Consider the polling example with  $n = 100$
- ▶ What is the probability that the sample has mean  $\bar{X}_n = 0.55$  and variance  $s_n^2 = 0.25$ , if the true proportion of Green voters in the population is  $E[X] = 0.5$ ?
- ▶ By the central limit theorem:

$$\begin{aligned}\Pr(\bar{X}_n \geq 0.55) &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - \mu_X}{s_n/\sqrt{n}}\right) \\ &= \Pr\left(\frac{\bar{X}_n - \mu_X}{s_n/\sqrt{n}} \geq \frac{0.55 - 0.5}{0.5/10}\right) \\ &= \Pr(Z \geq 1) \\ &\approx 0.159\end{aligned}$$