Art Of Problem Solving - AMC 10 July 2nd, 2021

Patrick & James Toche

Revised: July 2, 2021

Abstract

Notes on the AMC-10 Course by Art Of Problem Solving (AOPS). Copyright restrictions may apply. Written for personal use. Please report typos and errors over at https://github.com/ptoche/Math/tree/master/aops.

1.

For how many integers n between 1 and 100 does $x^2 + x - n$ factor into the product of two linear factors with integer coefficients?

(A) 0 (B) 1 (C) 2 (D) 9 (E) 10

2.

Suppose that a and b are nonzero real numbers, and that the equation $x^2 + ax + b = 0$ has solutions a and b. Then the pair (a, b) is

(A) (-2,1) (B) (-1,2) (C) (1,-2) (D) (2,-1) (E) (4,4)

3.

Let f be the function defined by $f(x) = ax^2 - \sqrt{2}$ for some positive a. If $f(f(\sqrt{2})) = -\sqrt{2}$, then a =

(A) $\frac{2-\sqrt{2}}{2}$ (B) $\frac{1}{2}$ (C) $2-\sqrt{2}$ (D) $\frac{\sqrt{2}}{2}$ (E) $\frac{2+\sqrt{2}}{2}$

4.

Both roots of the quadratic equation $x^2 - 63x + k = 0$ are prime numbers. The number of possible values of k is

(A) 0 (B) 1 (C) 2 (D) 4 (E) more than four

5.

Let @ denote the "averaged with" operation: $a@b = \frac{a+b}{2}$. Which of the following distributive laws holds for all numbers x, y, and z?

I. x@(y+z) = (x@y) + (x@z)

III. x + (y@z) = (x + y)@(x + z)

III. x@(y@z) = (x@y)@(x@z)

(A) I only (B) II only (C) III only (D) I and III only (E) II and III only

6.

If $f(x) = ax^4 - bx^2 + x + 5$ and f(-3) = 2, then f(3) =

(A) -5 (B) -2 (C) 1 (D) 3 (E) 8

7.

What is the sum of the reciprocals of the roots of the equation

 $\frac{2003}{2004}x + 1 + \frac{1}{x} = 0?$

 $(A) - \frac{2004}{2003}$ (B) -1 (C) $\frac{2003}{2004}$ (D) 1 (E) $\frac{2004}{2003}$

8.

Let f be a polynomial function such that, for all real x,

$$f(x^2 + 1) = x^4 + 5x^2 + 3$$

For all real x, $f(x^2 - 1)$ is

(A) $x^4 + 5x^2 + 1$ (B) $x^4 + x^2 - 3$ (C) $x^4 - 5x^2 + 1$ (D) $x^4 + x^2 + 3$ (E) none of these

9.

The polynomial $x^3 - ax^2 + bx - 2010$ has three positive integer roots. What is the smallest possible value of a?

(A) 78 (B) 88 (C) 98 (D) 108 (E) 118

10.

Let f be a function for which $f(x/3) = x^2 + x + 1$. Find the sum of all values of z for which f(3z) = 7.

(A) -1/3 (B) -1/9 (C) 0 (D) 5/9 (E) 5/3