

University of Edinburgh

An Analysis of the Turbulence Spectrum at Varying Magnetic Prandtl Number

Clara Despard

s1244462

Supervisor: Arjun Berera

05/04/17



Aims of the Project



- ▶ Magnetohydrodynamics (MHD) describes the movement of an electrically conducting fluid, including plasmas and liquid metals.
- ▶ Numerical simulations of MHD turbulence can improve our understanding of the behaviour of these fluids in the centre of the earth or the solar wind.

Aims of the Project



- ▶ Magnetohydrodynamics (MHD) describes the movement of an electrically conducting fluid, including plasmas and liquid metals.
- ▶ Numerical simulations of MHD turbulence can improve our understanding of the behaviour of these fluids in the centre of the earth or the solar wind.
- ▶ This project focused on varying the magnetic Prandtl number of the fluid and studying the effects that had on the system.

Aims

Fluid Dynamics

Navier-Stokes Equation

Magnetohydrodynamics

Definition

MHD Equations

Turbulence

Length Scales and the Energy Cascade

Direct Numerical Simulations

Forcing Regimes

Results

Summary



- ▶ Fluid motion is modeled as a continuum.
- ▶ The motion is governed by the Navier-Stokes equation, and depends on various parameters:



- ▶ Fluid motion is modeled as a continuum.
- ▶ The motion is governed by the Navier-Stokes equation, and depends on various parameters:

\mathbf{u} velocity

- ▶ Fluid motion is modeled as a continuum.
- ▶ The motion is governed by the Navier-Stokes equation, and depends on various parameters:

\mathbf{u} velocity

P pressure

- ▶ Fluid motion is modeled as a continuum.
- ▶ The motion is governed by the Navier-Stokes equation, and depends on various parameters:

\mathbf{u} velocity ρ density

P pressure

- ▶ Fluid motion is modeled as a continuum.
- ▶ The motion is governed by the Navier-Stokes equation, and depends on various parameters:

\mathbf{u} velocity ρ density

P pressure ν viscosity

- ▶ The fluid is also assumed to be incompressible:

$$\nabla \cdot \mathbf{u} = 0$$

Navier-Stokes Equation:

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{F}_u$$

Navier-Stokes Equation:

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla P + \nu \nabla^2 \mathbf{u} + \mathbf{F}_u$$

Reynold's Number:

$$Re = \frac{UL}{\nu}$$



- ▶ Magnetohydrodynamics describes the flow of a fluid which conducts an electric charge.
- ▶ The movement of the fluid, and thus the charge, creates a magnetic field.
- ▶ Dynamics are thus governed by Navier-Stokes equations combined with Maxwell's Equations.



$$\eta = \frac{1}{\mu_0 \sigma_0}$$

- ▶ Magnetic diffusivity is proportional to the resistivity of the fluid.
- ▶ It is a measure of how strongly the material opposes the flow of current.

MHD equations



$$\nabla \cdot \mathbf{u} = 0$$

MHD equations



$$\nabla \cdot \mathbf{u} = 0$$

$$\nabla \cdot \mathbf{b} = 0$$

$$\nabla \cdot \mathbf{u} = 0$$

$$\nabla \cdot \mathbf{b} = 0$$

MHD equations

$$\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} = -\frac{1}{\rho} \nabla P + \frac{1}{\rho} (\nabla \times \mathbf{b}) \times \mathbf{b} + \nu \nabla^2 \mathbf{u} + \mathbf{F}_u$$

$$\frac{\partial \mathbf{b}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{b} = (\mathbf{b} \cdot \nabla) \mathbf{u} + \eta \nabla^2 \mathbf{b} + \mathbf{F}_b$$

Magnetic Prandtl Number



Magnetic Reynolds number:

$$Re_m = \frac{UL}{\eta}$$

Magnetic Reynolds number:

$$Re_m = \frac{UL}{\eta}$$

- ▶ The magnetic Prandtl number is a ratio of the viscous diffusion to the magnetic diffusion

Magnetic Reynolds number:

$$Re_m = \frac{UL}{\eta}$$

- ▶ The magnetic Prandtl number is a ratio of the viscous diffusion to the magnetic diffusion

Magnetic Prandtl number:

$$Pr_m = \frac{\nu}{\eta} = \frac{Re_m}{Re}$$

Turbulence




- ▶ Turbulence is chaotic behaviour exhibited by a fluid.
- ▶ Occurs when the Reynolds number is above a critical value.

Turbulence



9

- ▶ Turbulence is chaotic behaviour exhibited by a fluid.
- ▶ Occurs when the Reynolds number is above a critical value.



[D. F. Harlacher, H. Klimach, and S. Roller. Turbulence simulation at large scale. inSiDe, 10, 2012]



- ▶ Energy enters the system at large scales.
- ▶ This length scale is called the **integral length scale**, and features in our definition of the Reynold's number.



- ▶ Energy enters the system at large scales.
- ▶ This length scale is called the **integral length scale**, and features in our definition of the Reynold's number.
- ▶ Energy is transferred to smaller scales in turbulent motion.

- ▶ Energy enters the system at large scales.
- ▶ This length scale is called the **integral length scale**, and features in our definition of the Reynold's number.
- ▶ Energy is transferred to smaller scales in turbulent motion.
- ▶ Kinetic energy is dissipated at small scales due to the viscosity, and the magnetic energy is dissipated due to the resistivity.

Kolmogorov microscale



11

- ▶ The length scale where energy is dissipated is called the **Kolmogorov microscale**.

Kolmogorov microscale



- ▶ The length scale where energy is dissipated is called the **Kolmogorov microscale**.
- ▶ This scale has a corresponding velocity, and therefore a corresponding (Kolmogorov) Reynold's number.



- ▶ The length scale where energy is dissipated is called the **Kolmogorov microscale**.
- ▶ This scale has a corresponding velocity, and therefore a corresponding (Kolmogorov) Reynold's number.
- ▶ By definition, since no turbulent motions can occur at scales smaller than this, this Reynold's number **MUST** equal 1.
- ▶ From this, the microscale can be defined as:

- ▶ The length scale where energy is dissipated is called the **Kolmogorov microscale**.
- ▶ This scale has a corresponding velocity, and therefore a corresponding (Kolmogorov) Reynold's number.
- ▶ By definition, since no turbulent motions can occur at scales smaller than this, this Reynold's number **MUST** equal 1.
- ▶ From this, the microscale can be defined as:

$$l_D^u = \left(\frac{\nu^3}{\epsilon_u} \right)^{\frac{1}{4}}$$

- ▶ The length scale where energy is dissipated is called the **Kolmogorov microscale**.
- ▶ This scale has a corresponding velocity, and therefore a corresponding (Kolmogorov) Reynold's number.
- ▶ By definition, since no turbulent motions can occur at scales smaller than this, this Reynold's number **MUST** equal 1.
- ▶ From this, the microscale can be defined as:

$$l_D^u = \left(\frac{\nu^3}{\epsilon_u} \right)^{\frac{1}{4}}$$

$$l_D^b = \left(\frac{\eta^3}{\epsilon_b} \right)^{\frac{1}{4}}$$



- ▶ In Fourier space, this translates to energy entering the system at low wavenumbers and dissipating at high wavenumbers.

- In Fourier space, this translates to energy entering the system at low wavenumbers and dissipating at high wavenumbers.

$$k_D^u = \frac{1}{l_D} = \left(\frac{\epsilon_u}{\nu^3} \right)^{\frac{1}{4}}$$

- In Fourier space, this translates to energy entering the system at low wavenumbers and dissipating at high wavenumbers.

$$k_D^u = \frac{1}{l_D} = \left(\frac{\epsilon_u}{\nu^3} \right)^{\frac{1}{4}}$$

$$k_D^b = \frac{1}{l_D} = \left(\frac{\epsilon_b}{\eta^3} \right)^{\frac{1}{4}}$$

- ▶ The transfer of energy from low to high k is known as the Energy Cascade
- ▶ The rate of transfer is given by Kolmogorov's $\frac{5}{3}$ law for the energy spectrum:


$$E(k) = C\epsilon^{\frac{2}{3}}k^{-\frac{5}{3}}$$

Energy Cascade



- ▶ The transfer of energy from low to high k is known as the Energy Cascade
- ▶ The rate of transfer is given by Kolmogorov's $\frac{5}{3}$ law for the energy spectrum:

$$E(k) = C\epsilon^{\frac{2}{3}} k^{-\frac{5}{3}}$$






- ▶ A small scale dynamo occurs when energy is transferred to the magnetic field at small scales (high k).
- ▶ More energy is transferred at higher Prandtl numbers.

Energy Transfer



- ▶ A small scale dynamo occurs when energy is transferred to the magnetic field at small scales (high k).
- ▶ More energy is transferred at higher Prandtl numbers.



Homogeneous and Isotropic turbulence



- ▶ **Homogeneous**: invariant under translations in space
- ▶ **Isotropic**: invariant under rotations and reflections

Direct Numerical Simulations



- ▶ Direct numerical simulations simulate all scales of the motion down to the kolmogorov microscale.
- ▶ The code used in this Project was written by S. Yoffe in 2012 and extended to MHD by M. Linkmann in 2015.

Forcing Regimes




- ▶ Turbulence is dissipative and, without adding energy in to the system, it will decay
- ▶ Stationary turbulence can be obtained by adding a forcing term


- ▶ Turbulence is dissipative and, without adding energy in to the system, it will decay
- ▶ Stationary turbulence can be obtained by adding a forcing term

$$\mathbf{f}_u(\mathbf{k}, t) = \begin{cases} \left(\frac{\epsilon_W}{2E_f} \right) \hat{\mathbf{u}}(\mathbf{k}, t) & 0 < |\mathbf{k}| \leq k_f \\ 0 & \text{otherwise} \end{cases}$$

Forcing Regimes




Energy Time Graphs




(a) Decaying Simulation

Energy Time Graphs




(a) Decaying Simulation



(b) Stationary Simulation


Dissipation Ratio



Dissipation Ratio





20



Dissipation Ratio



21



[A. Brandenburg. Magnetic prandtl number dependence of the kinetic-to-magnetic dissipation ratio. The Astrophysical Journal, 791(1):12, 2014.]

Dissipation Ratio



A relationship of $\frac{\epsilon_u}{\epsilon_b} = 0.7Pr_m^{0.64}$ was found which is in good agreement with the relationship of $\frac{\epsilon_u}{\epsilon_b} = 0.6Pr_m^{0.55}$ found in Brandenburg et al. for decaying simulations.

Dissipation Ratio



A relationship of $\frac{\epsilon_u}{\epsilon_b} = 0.7Pr_m^{0.64}$ was found which is in good agreement with the relationship of $\frac{\epsilon_u}{\epsilon_b} = 0.6Pr_m^{0.55}$ found in Brandenburg et al. for decaying simulations.

No relationship between $\frac{\epsilon_u}{\epsilon_b}$ and Pr_m could be found for the stationary simulations

Explanations and further research



- ▶ Unlikely due to the forcing regimes used, as we used the same method as Sahoo et. al.

Explanations and further research



- ▶ Unlikely due to the forcing regimes used, as we used the same method as Sahoo et. al.
- ▶ Stationary simulations were run at a higher viscosity than the decaying simulations. It is possible that there was not enough turbulent motion for the relationship to arise.

Explanations and further research



- ▶ Unlikely due to the forcing regimes used, as we used the same method as Sahoo et. al.
- ▶ Stationary simulations were run at a higher viscosity than the decaying simulations. It is possible that there was not enough turbulent motion for the relationship to arise.
- ▶ Further studies at different viscosity and magnetic diffusivity would need to be done to determine if this is the case.

Summary

