

RESEARCH ONE-PAGER

GLP-2X

Dual-receptor agonist · GLP-1 / GIP · metabolic research

10 mg, 15 mg, and 30 mg lyophilized vials

OVERVIEW

GLP-2X is HarmonyX's dual incretin research peptide, studied on GLP-1 and GIP receptors. It is the two-lever version of the metabolic story GLP-3X takes to three.

IN PLAIN LANGUAGE

Think of GLP-2X as fullness plus a second metabolic lever. People look at it for appetite, blood-sugar handling, and body-composition research without the third (glucagon) receptor that GLP-3X adds.

WHAT RESEARCHERS LOOK AT

- Satiety and appetite-pathway research
- Blood-sugar and incretin signaling
- Body-composition and energy-intake models
- A dual-agonist comparison point next to GLP-3X
- Metabolic flexibility questions in research literature

LABORATORY RECONSTITUTION

Reconstitute aseptically with bacteriostatic water or the diluent specified in the laboratory SOP. Invert gently until dissolved; do not shake vigorously. Larger milligram vials need more diluent than a 10 mg peptide — confirm the volume for the vial on the bench before starting. Aliquot if the design requires repeated sampling. Offered as 10 mg, 15 mg, and 30 mg vials.

STORAGE

Lyophilized: refrigerate, protected from light and moisture. After reconstitution: refrigerate and follow laboratory SOP (commonly 14–30 days with bacteriostatic water). Avoid repeated freeze–thaw cycles.

RESEARCH USE ONLY

The language above is a plain-English reading of published research interests. It is not medical advice and not a claim that this product treats, prevents, or diagnoses any condition. This material is not a drug or cosmetic, and it is not intended for human or veterinary use. HarmonyX supplies it to qualified research purchasers for laboratory work. Certificates of analysis are available on request.

Questions on this vial? Paul and Thabby review every request.

research@harmony-x.com · **360-910-6274**