

RESEARCH ONE-PAGER

CJC-1295

GHRH analog · extended GH-axis signaling

10 mg lyophilized vials

OVERVIEW

CJC-1295 is a GHRH analog studied for stretching out growth-hormone signaling. Where a short GHRH fragment is a brief knock on the door, CJC-1295 is studied as a longer conversation with the GH axis.

IN PLAIN LANGUAGE

In plain language, CJC-1295 is looked at for a longer-lasting GH signal. Researchers reach for it when the question is “what happens if we keep the GH axis engaged,” often next to a pulse peptide such as Ipamorelin.

WHAT RESEARCHERS LOOK AT

- Longer GH-axis signaling than a short GHRH fragment
- Amplitude and duration of growth-hormone release
- Recovery, sleep, and body-composition research models
- A common partner to Ipamorelin in combination vials
- IGF-1-related downstream signaling in research literature

LABORATORY RECONSTITUTION

Reconstitute aseptically with bacteriostatic water. Invert gently until dissolved; do not shake vigorously. A common laboratory starting point is 2 mL bacteriostatic water per 10 mg of peptide (5 mg/mL). Match the volume to the vial on the bench and confirm against the laboratory SOP. Aliquot if the design requires repeated sampling. Offered as 10 mg lyophilized vials.

STORAGE

Lyophilized: refrigerate, protected from light and moisture. After reconstitution: refrigerate and follow laboratory SOP (commonly 14–30 days with bacteriostatic water). Avoid repeated freeze–thaw cycles.

RESEARCH USE ONLY

The language above is a plain-English reading of published research interests. It is not medical advice and not a claim that this product treats, prevents, or diagnoses any condition. This material is not a drug or cosmetic, and it is not intended for human or veterinary use. HarmonyX supplies it to qualified research purchasers for laboratory work. Certificates of analysis are available on request.

Questions on this vial? Paul and Thabby review every request.

research@harmony-x.com · 360-910-6274