

RESEARCH ONE-PAGER

NAD⁺ Buffered

Buffered NAD⁺ · mitochondrial and DNA-repair research

1000 mg vials

OVERVIEW

Buffered NAD⁺ is the same coenzyme as standard NAD⁺, prepared for more comfortable handling and stability on the bench. The research questions are the same: mitochondria, DNA repair, cellular energy.

IN PLAIN LANGUAGE

Same NAD⁺ story — cellular fuel and repair — in a buffered format that labs often find easier to work with. Choose it when the protocol wants NAD⁺ biology with a gentler reconstitution profile.

WHAT RESEARCHERS LOOK AT

- The same mitochondrial and DNA-repair research as NAD⁺
- Buffered for handling and stability
- Cellular energy and aging-biology models
- A practical alternative when standard NAD⁺ is harsh to work with
- Still a coenzyme, not a peptide

LABORATORY RECONSTITUTION

Reconstitute aseptically with bacteriostatic water or the diluent specified in the laboratory SOP. Invert gently until dissolved; do not shake vigorously. Larger milligram vials need more diluent than a 10 mg peptide — confirm the volume for the vial on the bench before starting. Aliquot if the design requires repeated sampling. Offered as 1000 mg buffered. Confirm diluent with the laboratory SOP.

STORAGE

Lyophilized: refrigerate, protected from light and moisture. After reconstitution: refrigerate and follow laboratory SOP (commonly 14–30 days with bacteriostatic water). Avoid repeated freeze–thaw cycles.

RESEARCH USE ONLY

The language above is a plain-English reading of published research interests. It is not medical advice and not a claim that this product treats, prevents, or diagnoses any condition. This material is not a drug or cosmetic, and it is not intended for human or veterinary use. HarmonyX supplies it to qualified research purchasers for laboratory work. Certificates of analysis are available on request.

Questions on this vial? Paul and Thabby review every request.

research@harmony-x.com · 360-910-6274